

Partial Differential Equations

SECOND EDITION

This page is intentionally left blank.

Partial Differential Equations

SECOND EDITION

Lawrence C. Evans

*Department of Mathematics
University of California, Berkeley*

Graduate Studies
in Mathematics

Volume 19



American Mathematical Society
Providence, Rhode Island

Editorial Board

James E. Humphreys (Chair)

David J. Saltman

David Sattinger

Julius L. Shaneson

2010 *Mathematics Subject Classification*. Primary 35-XX; Secondary 49-XX, 47Hxx.

For additional information and updates on this book, visit
www.ams.org/bookpages/gsm-19

Library of Congress Cataloging-in-Publication Data

Evans, Lawrence C., 1949–

Partial differential equations / Lawrence C. Evans. — 2nd ed.

p. cm. — (Graduate studies in mathematics ; v. 19)

Includes bibliographical references and index.

ISBN 978-0-8218-4974-3 (alk. paper)

1. Differential equations, Partial. I. Title.

QA377.E95 2010

515'.353–dc22

2009044716

Copying and reprinting. Individual readers of this publication, and nonprofit libraries acting for them, are permitted to make fair use of the material, such as to copy a chapter for use in teaching or research. Permission is granted to quote brief passages from this publication in reviews, provided the customary acknowledgment of the source is given.

Republication, systematic copying, or multiple reproduction of any material in this publication is permitted only under license from the American Mathematical Society. Requests for such permission should be addressed to the Acquisitions Department, American Mathematical Society, 201 Charles Street, Providence, Rhode Island 02904-2294 USA. Requests can also be made by e-mail to reprint-permission@ams.org.

First Edition © 1998 by the American Mathematical Society. All rights reserved.

Reprinted with corrections 1999, 2002

Second Edition © 2010 by the American Mathematical Society. All rights reserved.

The American Mathematical Society retains all rights
except those granted to the United States Government.

Printed in the United States of America.

∞ The paper used in this book is acid-free and falls within the guidelines
established to ensure permanence and durability.

Visit the AMS home page at <http://www.ams.org/>

10 9 8 7 6 5 4 3 2 1 15 14 13 12 11 10

I dedicate this book to the memory of my parents,
LAWRENCE S. EVANS and LOUISE J. EVANS.

This page is intentionally left blank.

CONTENTS

Preface to second edition	xvii
Preface to first edition	xix
1. Introduction	1
1.1. Partial differential equations	1
1.2. Examples	3
1.2.1. Single partial differential equations	3
1.2.2. Systems of partial differential equations	6
1.3. Strategies for studying PDE	6
1.3.1. Well-posed problems, classical solutions	7
1.3.2. Weak solutions and regularity	7
1.3.3. Typical difficulties	9
1.4. Overview	9
1.5. Problems	12
1.6. References	13

PART I: REPRESENTATION FORMULAS FOR SOLUTIONS

2. Four Important Linear PDE	17
2.1. Transport equation	18
2.1.1. Initial-value problem	18
2.1.2. Nonhomogeneous problem	19
2.2. Laplace's equation	20

2.2.1. Fundamental solution	21
2.2.2. Mean-value formulas	25
2.2.3. Properties of harmonic functions	26
2.2.4. Green's function	33
2.2.5. Energy methods	41
2.3. Heat equation	44
2.3.1. Fundamental solution	45
2.3.2. Mean-value formula	51
2.3.3. Properties of solutions	55
2.3.4. Energy methods	62
2.4. Wave equation	65
2.4.1. Solution by spherical means	67
2.4.2. Nonhomogeneous problem	80
2.4.3. Energy methods	82
2.5. Problems	84
2.6. References	90
3. Nonlinear First-Order PDE	91
3.1. Complete integrals, envelopes	92
3.1.1. Complete integrals	92
3.1.2. New solutions from envelopes	94
3.2. Characteristics	96
3.2.1. Derivation of characteristic ODE	96
3.2.2. Examples	99
3.2.3. Boundary conditions	102
3.2.4. Local solution	105
3.2.5. Applications	109
3.3. Introduction to Hamilton–Jacobi equations	114
3.3.1. Calculus of variations, Hamilton's ODE	115
3.3.2. Legendre transform, Hopf–Lax formula	120
3.3.3. Weak solutions, uniqueness	128
3.4. Introduction to conservation laws	135
3.4.1. Shocks, entropy condition	136
3.4.2. Lax–Oleinik formula	143
3.4.3. Weak solutions, uniqueness	148

3.4.4. Riemann's problem	153
3.4.5. Long time behavior	156
3.5. Problems	161
3.6. References	165
4. Other Ways to Represent Solutions	167
4.1. Separation of variables	167
4.1.1. Examples	168
4.1.2. Application: Turing instability	172
4.2. Similarity solutions	176
4.2.1. Plane and traveling waves, solitons	176
4.2.2. Similarity under scaling	185
4.3. Transform methods	187
4.3.1. Fourier transform	187
4.3.2. Radon transform	196
4.3.3. Laplace transform	203
4.4. Converting nonlinear into linear PDE	206
4.4.1. Cole–Hopf transformation	206
4.4.2. Potential functions	208
4.4.3. Hodograph and Legendre transforms	209
4.5. Asymptotics	211
4.5.1. Singular perturbations	211
4.5.2. Laplace's method	216
4.5.3. Geometric optics, stationary phase	218
4.5.4. Homogenization	229
4.6. Power series	232
4.6.1. Noncharacteristic surfaces	232
4.6.2. Real analytic functions	237
4.6.3. Cauchy–Kovalevskaya Theorem	239
4.7. Problems	244
4.8. References	249

PART II: THEORY FOR LINEAR PARTIAL DIFFERENTIAL EQUATIONS

5. Sobolev Spaces	253
5.1. Hölder spaces	254

5.2. Sobolev spaces	255
5.2.1. Weak derivatives	255
5.2.2. Definition of Sobolev spaces	258
5.2.3. Elementary properties	261
5.3. Approximation	264
5.3.1. Interior approximation by smooth functions	264
5.3.2. Approximation by smooth functions	265
5.3.3. Global approximation by smooth functions	266
5.4. Extensions	268
5.5. Traces	271
5.6. Sobolev inequalities	275
5.6.1. Gagliardo–Nirenberg–Sobolev inequality	276
5.6.2. Morrey’s inequality	280
5.6.3. General Sobolev inequalities	284
5.7. Compactness	286
5.8. Additional topics	289
5.8.1. Poincaré’s inequalities	289
5.8.2. Difference quotients	291
5.8.3. Differentiability a.e.	295
5.8.4. Hardy’s inequality	296
5.8.5. Fourier transform methods	297
5.9. Other spaces of functions	299
5.9.1. The space H^{-1}	299
5.9.2. Spaces involving time	301
5.10. Problems	305
5.11. References	309
6. Second-Order Elliptic Equations	311
6.1. Definitions	311
6.1.1. Elliptic equations	311
6.1.2. Weak solutions	313
6.2. Existence of weak solutions	315
6.2.1. Lax–Milgram Theorem	315
6.2.2. Energy estimates	317
6.2.3. Fredholm alternative	320

6.3. Regularity	326
6.3.1. Interior regularity	327
6.3.2. Boundary regularity	334
6.4. Maximum principles	344
6.4.1. Weak maximum principle	344
6.4.2. Strong maximum principle	347
6.4.3. Harnack's inequality	351
6.5. Eigenvalues and eigenfunctions	354
6.5.1. Eigenvalues of symmetric elliptic operators	354
6.5.2. Eigenvalues of nonsymmetric elliptic operators	360
6.6. Problems	365
6.7. References	370
7. Linear Evolution Equations	371
7.1. Second-order parabolic equations	371
7.1.1. Definitions	372
7.1.2. Existence of weak solutions	375
7.1.3. Regularity	380
7.1.4. Maximum principles	389
7.2. Second-order hyperbolic equations	398
7.2.1. Definitions	398
7.2.2. Existence of weak solutions	401
7.2.3. Regularity	408
7.2.4. Propagation of disturbances	414
7.2.5. Equations in two variables	418
7.3. Hyperbolic systems of first-order equations	421
7.3.1. Definitions	421
7.3.2. Symmetric hyperbolic systems	423
7.3.3. Systems with constant coefficients	429
7.4. Semigroup theory	433
7.4.1. Definitions, elementary properties	434
7.4.2. Generating contraction semigroups	439
7.4.3. Applications	441
7.5. Problems	446
7.6. References	449

**PART III: THEORY FOR NONLINEAR PARTIAL
DIFFERENTIAL EQUATIONS**

8. The Calculus of Variations	453
8.1. Introduction	453
8.1.1. Basic ideas	453
8.1.2. First variation, Euler–Lagrange equation	454
8.1.3. Second variation	458
8.1.4. Systems	459
8.2. Existence of minimizers	465
8.2.1. Coercivity, lower semicontinuity	465
8.2.2. Convexity	467
8.2.3. Weak solutions of Euler–Lagrange equation ...	472
8.2.4. Systems	475
8.2.5. Local minimizers	480
8.3. Regularity	482
8.3.1. Second derivative estimates	483
8.3.2. Remarks on higher regularity	486
8.4. Constraints	488
8.4.1. Nonlinear eigenvalue problems	488
8.4.2. Unilateral constraints, variational inequalities .	492
8.4.3. Harmonic maps	495
8.4.4. Incompressibility	497
8.5. Critical points	501
8.5.1. Mountain Pass Theorem	501
8.5.2. Application to semilinear elliptic PDE	507
8.6. Invariance, Noether’s Theorem	511
8.6.1. Invariant variational problems	512
8.6.2. Noether’s Theorem	513
8.7. Problems	520
8.8. References	525
9. Nonvariational Techniques	527
9.1. Monotonicity methods	527
9.2. Fixed point methods	533
9.2.1. Banach’s Fixed Point Theorem	534

9.2.2. Schauder's, Schaefer's Fixed Point Theorems . . .	538
9.3. Method of subsolutions and supersolutions	543
9.4. Nonexistence of solutions	547
9.4.1. Blow-up	547
9.4.2. Derrick–Pohozaev identity	551
9.5. Geometric properties of solutions	554
9.5.1. Star-shaped level sets	554
9.5.2. Radial symmetry	555
9.6. Gradient flows	560
9.6.1. Convex functions on Hilbert spaces	560
9.6.2. Subdifferentials and nonlinear semigroups	565
9.6.3. Applications	571
9.7. Problems	573
9.8. References	577
10. Hamilton–Jacobi Equations	579
10.1. Introduction, viscosity solutions	579
10.1.1. Definitions	581
10.1.2. Consistency	583
10.2. Uniqueness	586
10.3. Control theory, dynamic programming	590
10.3.1. Introduction to optimal control theory	591
10.3.2. Dynamic programming	592
10.3.3. Hamilton–Jacobi–Bellman equation	594
10.3.4. Hopf–Lax formula revisited	600
10.4. Problems	603
10.5. References	606
11. Systems of Conservation Laws	609
11.1. Introduction	609
11.1.1. Integral solutions	612
11.1.2. Traveling waves, hyperbolic systems	615
11.2. Riemann's problem	621
11.2.1. Simple waves	621
11.2.2. Rarefaction waves	624
11.2.3. Shock waves, contact discontinuities	625

11.2.4. Local solution of Riemann's problem	632
11.3. Systems of two conservation laws	635
11.3.1. Riemann invariants	635
11.3.2. Nonexistence of smooth solutions	639
11.4. Entropy criteria	641
11.4.1. Vanishing viscosity, traveling waves	642
11.4.2. Entropy/entropy-flux pairs	646
11.4.3. Uniqueness for scalar conservation laws	649
11.5. Problems	654
11.6. References	657
12. Nonlinear Wave Equations	659
12.1. Introduction	659
12.1.1. Conservation of energy	660
12.1.2. Finite propagation speed	660
12.2. Existence of solutions	663
12.2.1. Lipschitz nonlinearities	663
12.2.2. Short time existence	666
12.3. Semilinear wave equations	670
12.3.1. Sign conditions	670
12.3.2. Three space dimensions	674
12.3.3. Subcritical power nonlinearities	676
12.4. Critical power nonlinearity	679
12.5. Nonexistence of solutions	686
12.5.1. Nonexistence for negative energy	687
12.5.2. Nonexistence for small initial data	689
12.6. Problems	691
12.7. References	696

APPENDICES

Appendix A: Notation	697
A.1. Notation for matrices	697
A.2. Geometric notation	698
A.3. Notation for functions	699
A.4. Vector-valued functions	703
A.5. Notation for estimates	703

A.6. Some comments about notation	704
Appendix B: Inequalities	705
B.1. Convex functions	705
B.2. Useful inequalities	706
Appendix C: Calculus	710
C.1. Boundaries	710
C.2. Gauss–Green Theorem	711
C.3. Polar coordinates, coarea formula	712
C.4. Moving regions	713
C.5. Convolution and smoothing	713
C.6. Inverse Function Theorem	716
C.7. Implicit Function Theorem	717
C.8. Uniform convergence	718
Appendix D: Functional Analysis	719
D.1. Banach spaces	719
D.2. Hilbert spaces	720
D.3. Bounded linear operators	721
D.4. Weak convergence	723
D.5. Compact operators, Fredholm theory	724
D.6. Symmetric operators	728
Appendix E: Measure Theory	729
E.1. Lebesgue measure	729
E.2. Measurable functions and integration	730
E.3. Convergence theorems for integrals	731
E.4. Differentiation	732
E.5. Banach space-valued functions	733
Bibliography	735
Index	741

This page is intentionally left blank.

Preface to the second edition

Let me thank everyone who over the past decade has provided me with suggestions and corrections for improving the first edition of this book. I am extraordinarily grateful. Although I have not always followed these many pieces of advice and criticism, I have thought carefully about them all. So many people have helped me out that it is unfortunately no longer feasible to list all their names. I have also received extraordinary help from everyone at the AMS, especially Sergei Gelfand, Stephen Moye and Arlene O'Sean. The NSF has generously supported my research during the writing of both the original edition of the book and this revision. I will continue to maintain lists of errors on my homepage, accessible through the math.berkeley.edu website.

When you write a big book on a big subject, the temptation is to include everything. A critic famously once imagined Tolstoy during the writing of *War and Peace*: “The book is long, but even if it were twice as long, if it were three times as long, there would always be scenes that have been omitted, and these Tolstoy, waking up in the middle of the night, must have regretted. There must have been a night when it occurred to him that he had not included a yacht race...” (G. Moore, *Avowals*).

This image notwithstanding, I have tried to pack into this second edition as many fascinating new topics in partial differential equations (PDE) as I could manage, most notably in the new Chapter 12 on nonlinear wave equations. There are new sections on Noether's Theorem and on local minimizers in the calculus of variations, on the Radon transform, on Turing instabilities for reaction-diffusion systems, etc. I have rewritten and expanded the

previous discussions on blow-up of solutions, on group and phase velocities, and on several further subjects. I have also updated and greatly increased citations to books in the bibliography and have moved references to research articles to within the text. There are countless further minor modifications in notation and wording. Most importantly, I have added about 80 new exercises, most quite interesting and some rather elaborate. There are now over 200 in total.

And there is a yacht race among the problems for Chapter 10.

LCE

January, 2010

Berkeley

Preface to the first edition

I present in this book a wide-ranging survey of many important topics in the theory of partial differential equations (PDE), with particular emphasis on various modern approaches. I have made a huge number of editorial decisions about what to keep and what to toss out, and can only claim that this selection seems to me about right. I of course include the usual formulas for solutions of the usual linear PDE, but also devote large amounts of exposition to energy methods within Sobolev spaces, to the calculus of variations, to conservation laws, etc.

My general working principles in the writing have been these:

a. PDE theory is (mostly) not restricted to two independent variables. Many texts describe PDE as if functions of the two variables (x, y) or (x, t) were all that matter. This emphasis seems to me misleading, as modern discoveries concerning many types of equations, both linear and nonlinear, have allowed for the rigorous treatment of these in any number of dimensions. I also find it unsatisfactory to “classify” partial differential equations: this is possible in two variables, but creates the false impression that there is some kind of general and useful classification scheme available in general.

b. Many interesting equations are nonlinear. My view is that overall we know too much about linear PDE and too little about nonlinear PDE. I have accordingly introduced nonlinear concepts early in the text and have tried hard to emphasize everywhere nonlinear analogues of the linear theory.

c. Understanding generalized solutions is fundamental. Many of the partial differential equations we study, especially nonlinear first-order equations, do not in general possess smooth solutions. It is therefore essential to devise some kind of proper notion of generalized or weak solution. This is an important but subtle undertaking, and much of the hardest material in this book concerns the uniqueness of appropriately defined weak solutions.

d. PDE theory is not a branch of functional analysis. Whereas certain classes of equations can profitably be viewed as generating abstract operators between Banach spaces, the insistence on an overly abstract viewpoint, and consequent ignoring of deep calculus and measure theoretic estimates, is ultimately limiting.

e. Notation is a nightmare. I have really tried to introduce consistent notation, which works for all the important classes of equations studied. This attempt is sometimes at variance with notational conventions within a given subarea.

f. Good theory is (almost) as useful as exact formulas. I incorporate this principle into the overall organization of the text, which is subdivided into three parts, roughly mimicking the historical development of PDE theory itself. Part I concerns the search for explicit formulas for solutions, and Part II the abandoning of this quest in favor of general theory asserting the existence and other properties of solutions for linear equations. Part III is the mostly modern endeavor of fashioning general theory for important classes of nonlinear PDE.

Let me also explicitly comment here that I intend the development within each section to be rigorous and complete (exceptions being the frankly heuristic treatment of asymptotics in §4.5 and an occasional reference to a research paper). This means that even locally within each chapter the topics do not necessarily progress logically from “easy” to “hard” concepts. There are many difficult proofs and computations early on, but as compensation many easier ideas later. The student should certainly omit on first reading some of the more arcane proofs.

I wish next to emphasize that this is a *textbook*, and not a reference book. I have tried everywhere to present the essential ideas in the clearest possible settings, and therefore have almost never established sharp versions of any of the theorems. Research articles and advanced monographs, many of them listed in the Bibliography, provide such precision and generality. My goal has rather been to explain, as best I can, the many fundamental ideas of the subject within fairly simple contexts.

I have greatly profited from the comments and thoughtful suggestions of many of my colleagues, friends and students, in particular: S. Antman, J. Bang, X. Chen, A. Chorin, M. Christ, J. Cima, P. Colella, J. Cooper, M. Crandall, B. Driver, M. Feldman, M. Fitzpatrick, R. Gariepy, J. Goldstein, D. Gomes, O. Hald, W. Han, W. Hrusa, T. Ilmanen, I. Ishii, I. Israel, R. Jerrard, C. Jones, B. Kawohl, S. Koike, J. Lewis, T.-P. Liu, H. Lopes, J. McLaughlin, K. Miller, J. Morford, J. Neu, M. Portilheiro, J. Ralston, F. Rezakhanlou, W. Schlag, D. Serre, P. Souganidis, J. Strain, W. Strauss, M. Struwe, R. Temam, B. Tvedt, J.-L. Vazquez, M. Weinstein, P. Wolfe, and Y. Zheng.

I especially thank Tai-Ping Liu for many years ago writing out for me the first draft of what is now Chapter 11.

I am extremely grateful for the suggestions and lists of mistakes from earlier drafts of this book sent to me by many readers, and I encourage others to send me their comments, at evans@math.berkeley.edu. I have come to realize that I must be more than slightly mad to try to write a book of this length and complexity, but I am not yet crazy enough to think that I have made no mistakes. **I will therefore maintain a listing of errors which come to light, and will make this accessible through the math.berkeley.edu homepage.**

Faye Yeager at UC Berkeley has done a really magnificent job typing and updating these notes, and Jaya Nagendra heroically typed an earlier version at the University of Maryland. My deepest thanks to both.

I have been supported by the NSF during much of the writing, most recently under grant DMS-9424342.

LCE

August, 1997

Berkeley

This page is intentionally left blank.

INTRODUCTION

- 1.1 Partial differential equations
- 1.2 Examples
- 1.3 Strategies for studying PDE
- 1.4 Overview
- 1.5 Problems
- 1.6 References

This chapter surveys the principal theoretical issues concerning the solving of partial differential equations.

To follow the subsequent discussion, the reader should first of all turn to Appendix A and look over the notation presented there, particularly the multiindex notation for partial derivatives.

1.1. PARTIAL DIFFERENTIAL EQUATIONS

A *partial differential equation* (PDE) is an equation involving an unknown function of two or more variables and certain of its partial derivatives.

Using the notation explained in Appendix A, we can write out symbolically a typical PDE, as follows. Fix an integer $k \geq 1$ and let U denote an open subset of \mathbb{R}^n .

DEFINITION. *An expression of the form*

$$(1) \quad F(D^k u(x), D^{k-1} u(x), \dots, Du(x), u(x), x) = 0 \quad (x \in U)$$

is called a k^{th} -order partial differential equation, where

$$F : \mathbb{R}^{n^k} \times \mathbb{R}^{n^{k-1}} \times \dots \times \mathbb{R}^n \times \mathbb{R} \times U \rightarrow \mathbb{R}$$

is given and

$$u : U \rightarrow \mathbb{R}$$

is the unknown.

We solve the PDE if we find all u verifying (1), possibly only among those functions satisfying certain auxiliary boundary conditions on some part Γ of ∂U . By finding the solutions we mean, ideally, obtaining simple, explicit solutions, or, failing that, deducing the existence and other properties of solutions.

DEFINITIONS.

(i) The partial differential equation (1) is called linear if it has the form

$$\sum_{|\alpha| \leq k} a_\alpha(x) D^\alpha u = f(x)$$

for given functions a_α ($|\alpha| \leq k$), f . This linear PDE is homogeneous if $f \equiv 0$.

(ii) The PDE (1) is semilinear if it has the form

$$\sum_{|\alpha|=k} a_\alpha(x) D^\alpha u + a_0(D^{k-1}u, \dots, Du, u, x) = 0.$$

(iii) The PDE (1) is quasilinear if it has the form

$$\sum_{|\alpha|=k} a_\alpha(D^{k-1}u, \dots, Du, u, x) D^\alpha u + a_0(D^{k-1}u, \dots, Du, u, x) = 0.$$

(iv) The PDE (1) is fully nonlinear if it depends nonlinearly upon the highest order derivatives.

A system of partial differential equations is, informally speaking, a collection of several PDE for several unknown functions.

DEFINITION. An expression of the form

$$(2) \quad \mathbf{F}(D^k \mathbf{u}(x), D^{k-1} \mathbf{u}(x), \dots, D\mathbf{u}(x), \mathbf{u}(x), x) = \mathbf{0} \quad (x \in U)$$

is called a k^{th} -order system of partial differential equations, where

$$\mathbf{F} : \mathbb{R}^{mn^k} \times \mathbb{R}^{mn^{k-1}} \times \dots \times \mathbb{R}^{mn} \times \mathbb{R}^m \times U \rightarrow \mathbb{R}^m$$

is given and

$$\mathbf{u} : U \rightarrow \mathbb{R}^m, \quad \mathbf{u} = (u^1, \dots, u^m)$$

is the unknown.

Here we are supposing that the system comprises the same number m of scalar equations as unknowns (u^1, \dots, u^m) . This is the most common circumstance, although other systems may have fewer or more equations than unknowns. Systems are classified in the obvious way as being linear, semilinear, etc.

NOTATION. We write “PDE” as an abbreviation for both the singular “partial differential equation” and the plural “partial differential equations”.

1.2. EXAMPLES

There is no general theory known concerning the solvability of all partial differential equations. Such a theory is extremely unlikely to exist, given the rich variety of physical, geometric, and probabilistic phenomena which can be modeled by PDE. Instead, research focuses on various particular partial differential equations that are important for applications within and outside of mathematics, with the hope that insight from the origins of these PDE can give clues as to their solutions.

Following is a list of many specific partial differential equations of interest in current research. This listing is intended merely to familiarize the reader with the names and forms of various famous PDE. To display most clearly the mathematical structure of these equations, we have mostly set relevant physical constants to unity. We will later discuss the origin and interpretation of many of these PDE.

Throughout $x \in U$, where U is an open subset of \mathbb{R}^n , and $t \geq 0$. Also $Du = D_x u = (u_{x_1}, \dots, u_{x_n})$ denotes the gradient of u with respect to the spatial variable $x = (x_1, \dots, x_n)$. The variable t always denotes time.

1.2.1. Single partial differential equations.

a. Linear equations.

1. *Laplace’s equation*

$$\Delta u = \sum_{i=1}^n u_{x_i x_i} = 0.$$

2. *Helmholtz’s (or eigenvalue) equation*

$$-\Delta u = \lambda u.$$

3. *Linear transport equation*

$$u_t + \sum_{i=1}^n b^i u_{x_i} = 0.$$

4. *Liouville’s equation*

$$u_t - \sum_{i=1}^n (b^i u)_{x_i} = 0.$$

5. *Heat (or diffusion) equation*

$$u_t - \Delta u = 0.$$

6. *Schrödinger's equation*

$$iu_t + \Delta u = 0.$$

7. *Kolmogorov's equation*

$$u_t - \sum_{i,j=1}^n a^{ij} u_{x_i x_j} + \sum_{i=1}^n b^i u_{x_i} = 0.$$

8. *Fokker-Planck equation*

$$u_t - \sum_{i,j=1}^n (a^{ij} u)_{x_i x_j} - \sum_{i=1}^n (b^i u)_{x_i} = 0.$$

9. *Wave equation*

$$u_{tt} - \Delta u = 0.$$

10. *Klein-Gordon equation*

$$u_{tt} - \Delta u + m^2 u = 0.$$

11. *Telegraph equation*

$$u_{tt} + 2du_t - u_{xx} = 0.$$

12. *General wave equation*

$$u_{tt} - \sum_{i,j=1}^n a^{ij} u_{x_i x_j} + \sum_{i=1}^n b^i u_{x_i} = 0.$$

13. *Airy's equation*

$$u_t + u_{xxx} = 0.$$

14. *Beam equation*

$$u_{tt} + u_{xxxx} = 0.$$

b. Nonlinear equations.1. *Eikonal equation*

$$|Du| = 1.$$

2. *Nonlinear Poisson equation*

$$-\Delta u = f(u).$$

3. *p-Laplacian equation*

$$\operatorname{div}(|Du|^{p-2} Du) = 0.$$

4. *Minimal surface equation*

$$\operatorname{div} \left(\frac{Du}{(1 + |Du|^2)^{1/2}} \right) = 0.$$

5. *Monge–Ampère equation*

$$\det(D^2u) = f.$$

6. *Hamilton–Jacobi equation*

$$u_t + H(Du, x) = 0.$$

7. *Scalar conservation law*

$$u_t + \operatorname{div} \mathbf{F}(u) = 0.$$

8. *Inviscid Burgers' equation*

$$u_t + uu_x = 0.$$

9. *Scalar reaction-diffusion equation*

$$u_t - \Delta u = f(u).$$

10. *Porous medium equation*

$$u_t - \Delta(u^\gamma) = 0.$$

11. *Nonlinear wave equation*

$$u_{tt} - \Delta u + f(u) = 0.$$

12. *Korteweg–de Vries (KdV) equation*

$$u_t + uu_x + u_{xxx} = 0.$$

13. *Nonlinear Schrödinger equation*

$$iu_t + \Delta u = f(|u|^2)u.$$

1.2.2. Systems of partial differential equations.

a. Linear systems.

1. *Equilibrium equations of linear elasticity*

$$\mu\Delta\mathbf{u} + (\lambda + \mu)D(\operatorname{div}\mathbf{u}) = \mathbf{0}.$$

2. *Evolution equations of linear elasticity*

$$\mathbf{u}_{tt} - \mu\Delta\mathbf{u} - (\lambda + \mu)D(\operatorname{div}\mathbf{u}) = \mathbf{0}.$$

3. *Maxwell's equations*

$$\begin{cases} \mathbf{E}_t = \operatorname{curl}\mathbf{B} \\ \mathbf{B}_t = -\operatorname{curl}\mathbf{E} \\ \operatorname{div}\mathbf{B} = \operatorname{div}\mathbf{E} = 0. \end{cases}$$

b. Nonlinear systems.

1. *System of conservation laws*

$$\mathbf{u}_t + \operatorname{div}\mathbf{F}(\mathbf{u}) = \mathbf{0}.$$

2. *Reaction-diffusion system*

$$\mathbf{u}_t - \Delta\mathbf{u} = \mathbf{f}(\mathbf{u}).$$

3. *Euler's equations for incompressible, inviscid flow*

$$\begin{cases} \mathbf{u}_t + \mathbf{u} \cdot D\mathbf{u} = -Dp \\ \operatorname{div}\mathbf{u} = 0. \end{cases}$$

4. *Navier–Stokes equations for incompressible, viscous flow*

$$\begin{cases} \mathbf{u}_t + \mathbf{u} \cdot D\mathbf{u} - \Delta\mathbf{u} = -Dp \\ \operatorname{div}\mathbf{u} = 0. \end{cases}$$

See Zwillinger [Zw] for a much more extensive listing of interesting PDE.

1.3. STRATEGIES FOR STUDYING PDE

As explained in §1.1 our goal is the discovery of ways to solve partial differential equations of various sorts, but—as should now be clear in view of the

many diverse examples set forth in §1.2—this is no easy task. And indeed the very question of what it means to “solve” a given PDE can be subtle, depending in large part on the particular structure of the problem at hand.

1.3.1. Well-posed problems, classical solutions.

The informal notion of a well-posed problem captures many of the desirable features of what it means to solve a PDE. We say that a given problem for a partial differential equation is *well-posed* if

- (i) the problem in fact has a solution;
- (ii) this solution is unique;

and

- (iii) the solution depends continuously on the data given in the problem.

The last condition is particularly important for problems arising from physical applications: we would prefer that our (unique) solution changes only a little when the conditions specifying the problem change a little. (For many problems, on the other hand, uniqueness is not to be expected. In these cases the primary mathematical tasks are to classify and characterize the solutions.)

Now clearly it would be desirable to “solve” PDE in such a way that (i)–(iii) hold. But notice that we still have not carefully defined what we mean by a “solution”. Should we ask, for example, that a “solution” u must be real analytic or at least infinitely differentiable? This might be desirable, but perhaps we are asking too much. Maybe it would be wiser to require a solution of a PDE of order k to be at least k times continuously differentiable. Then at least all the derivatives which appear in the statement of the PDE will exist and be continuous, although maybe certain higher derivatives will not exist. Let us informally call a solution with this much smoothness a *classical* solution of the PDE: this is certainly the most obvious notion of solution.

So by solving a partial differential equation in the classical sense we mean if possible to write down a formula for a classical solution satisfying (i)–(iii) above, or at least to show such a solution exists, and to deduce various of its properties.

1.3.2. Weak solutions and regularity.

But can we achieve this? The answer is that certain specific partial differential equations (e.g. Laplace’s equation) can be solved in the classical sense, but many others, if not most others, cannot. Consider for instance

the scalar conservation law

$$u_t + F(u)_x = 0.$$

We will see in §3.4 that this PDE governs various one-dimensional phenomena involving fluid dynamics, and in particular models the formation and propagation of shock waves. Now a shock wave is a curve of discontinuity of the solution u ; and so if we wish to study conservation laws, and recover the underlying physics, we must surely allow for solutions u which are not continuously differentiable or even continuous. In general, as we shall see, the conservation law has no classical solutions but is well-posed if we allow for properly defined *generalized* or *weak solutions*.

This is all to say that we may be forced by the structure of the particular equation to abandon the search for smooth, classical solutions. We must instead, while still hoping to achieve the well-posedness conditions (i)–(iii), investigate a wider class of candidates for solutions. And in fact, even for those PDE which turn out to be classically solvable, it is often most expedient initially to search for some appropriate kind of weak solution.

The point is this: if from the outset we demand that our solutions be very regular, say k -times continuously differentiable, then we are usually going to have a really hard time finding them, as our proofs must then necessarily include possibly intricate demonstrations that the functions we are building are in fact smooth enough. A far more reasonable strategy is to consider as separate the *existence* and the *smoothness* (or *regularity*) problems. The idea is to define for a given PDE a reasonably wide notion of a *weak solution*, with the expectation that since we are not asking too much by way of smoothness of this weak solution, it may be easier to establish its existence, uniqueness, and continuous dependence on the given data. Thus, to repeat, it is often wise to aim at proving well-posedness in some appropriate class of weak or generalized solutions.

Now, as noted above, for various partial differential equations this is the best that can be done. For other equations we can hope that our weak solution may turn out after all to be smooth enough to qualify as a classical solution. This leads to the question of *regularity* of weak solutions. As we will see, it is often the case that the existence of weak solutions depends upon rather simple estimates plus ideas of functional analysis, whereas the regularity of the weak solutions, when true, usually rests upon many intricate calculus estimates.

Let me explicitly note here that once we are past Part I (Chapters 2–4), our efforts will be largely devoted to proving mathematically the existence of solutions to various sorts of partial differential equations, and not so much

to deriving formulas for these solutions. This may seem wasted or misguided effort, but in fact mathematicians are like theologians: we regard existence as the prime attribute of what we study. But unlike theologians, we need not always rely upon faith alone.

1.3.3. Typical difficulties.

Following are some vague but general principles, which may be useful to keep in mind:

- (i) Nonlinear equations are more difficult than linear equations; and, indeed, the more the nonlinearity affects the higher derivatives, the more difficult the PDE is.
- (ii) Higher-order PDE are more difficult than lower-order PDE.
- (iii) Systems are harder than single equations.
- (iv) Partial differential equations entailing many independent variables are harder than PDE entailing few independent variables.
- (v) For most partial differential equations it is not possible to write out explicit formulas for solutions.

None of these assertions is without important exceptions.

1.4. OVERVIEW

This textbook is divided into three major Parts.

PART I: Representation Formulas for Solutions

Here we identify those important partial differential equations for which in certain circumstances explicit or more-or-less explicit formulas can be had for solutions. The general progression of the exposition is from direct formulas for certain linear equations to far less concrete representation formulas, of a sort, for various nonlinear PDE.

Chapter 2 is a detailed study of four exactly solvable partial differential equations: the linear transport equation, Laplace's equation, the heat equation, and the wave equation. These PDE, which serve as archetypes for the more complicated equations introduced later, admit directly computable solutions, at least in the case that there is no domain whose boundary geometry complicates matters. The explicit formulas are augmented by various indirect, but easy and attractive, "energy"-type arguments, which serve as motivation for the developments in Chapters 6, 7 and thereafter.

Chapter 3 continues the theme of searching for explicit formulas, now for general first-order nonlinear PDE. The key insight is that such PDE

can, locally at least, be transformed into systems of ordinary differential equations (ODE), the characteristic equations. We stipulate that once the problem becomes “only” the question of integrating a system of ODE, it is in principle solved, sometimes quite explicitly. The derivation of the characteristic equations given in the text is very simple and does not require any geometric insights. It is in truth so easy to derive the characteristic equations that no real purpose is had by dealing with the quasilinear case first.

We introduce also the Hopf–Lax formula for Hamilton–Jacobi equations (§3.3) and the Lax–Oleinik formula for scalar conservation laws (§3.4). (Some knowledge of measure theory is useful here but is not essential.) These sections provide an early acquaintance with the global theory of these important nonlinear PDE and so motivate the later Chapters 10 and 11.

Chapter 4 is a grab bag of techniques for explicitly (or kind of explicitly) solving various linear and nonlinear partial differential equations, and the reader should study only whatever seems interesting. The section on the Fourier transform is, however, essential. The Cauchy–Kovalevskaya Theorem appears at the very end. Although this is basically the only general existence theorem in the subject, and thus logically should perhaps be regarded as central, in practice these power series methods are not so prevalent.

PART II: Theory for Linear Partial Differential Equations

Next we abandon the search for explicit formulas and instead rely on functional analysis and relatively easy “energy” estimates to prove the existence of weak solutions to various linear PDE. We investigate also the uniqueness and regularity of such solutions and deduce various other properties.

Chapter 5 is an introduction to Sobolev spaces, the proper setting for the study of many linear and nonlinear partial differential equations via energy methods. This is a hard chapter, the real worth of which is only later revealed, and requires some basic knowledge of Lebesgue measure theory. However, the requirements are not really so great, and the review in Appendix E should suffice. In my opinion there is no particular advantage in considering only the Sobolev spaces with exponent $p = 2$, and indeed insisting upon this obscures the two central inequalities, those of Gagliardo–Nirenberg–Sobolev (§5.6.1) and of Morrey (§5.6.2).

In Chapter 6 we vastly generalize our knowledge of Laplace’s equation to other second-order elliptic equations. Here we work through a rather complete treatment of existence, uniqueness and regularity theory for solutions, including the maximum principle, and also a reasonable introduction to the

study of eigenvalues, including a discussion of the principal eigenvalue for nonselfadjoint operators.

Chapter 7 expands the energy methods to a variety of linear partial differential equations characterizing evolutions in time. We broaden our earlier investigation of the heat equation to general second-order parabolic PDE and of the wave equation to general second-order hyperbolic PDE. We study as well linear first-order hyperbolic systems, with the aim of motivating the developments concerning nonlinear systems of conservation laws in Chapter 11. The concluding section 7.4 presents the alternative functional analytic method of semigroups for building solutions.

(Missing from this long Part II on linear partial differential equations is any discussion of distribution theory or potential theory. These are important topics, but for our purposes seem dispensable, even in a book of such length. These omissions do not slow us up much and make room for more nonlinear theory.)

PART III: Theory for Nonlinear Partial Differential Equations

This section parallels for nonlinear PDE the development in Part II but is far less unified in its approach, as the various types of nonlinearity must be treated in quite different ways.

Chapter 8 commences the general study of nonlinear partial differential equations with an extensive discussion of the calculus of variations. Here we set forth a careful derivation of the direct method for deducing the existence of minimizers and discuss also a variety of variational systems and constrained problems, as well as minimax methods. Variational theory is the most useful and accessible of the methods for nonlinear PDE, and so this chapter is fundamental.

Chapter 9 is, rather like Chapter 4 earlier, a gathering of assorted other techniques of use for nonlinear elliptic and parabolic partial differential equations. We encounter here monotonicity and fixed point methods and a variety of other devices, mostly involving the maximum principle. We study as well certain nice aspects of nonlinear semigroup theory, to complement the linear semigroup theory from Chapter 7.

Chapter 10 is an introduction to the modern theory of Hamilton–Jacobi PDE and in particular to the notion of “viscosity solutions”. We encounter also the connections with the optimal control of ODE, through dynamic programming.

Chapter 11 picks up from Chapter 3 the discussion of conservation laws, now systems of conservation laws. Unlike the general theoretical developments in Chapters 5–9, for which Sobolev spaces provide the proper abstract framework, we are forced to employ here direct linear algebra and calculus computations. We pay particular attention to the solution of Riemann's problem and to entropy criteria.

Chapter 12, an introduction to nonlinear wave equations, is new with this edition. We provide long time and short time existence theorems for certain quasilinear wave equations and an in-depth examination of semilinear wave equations, especially for subcritical and critical power nonlinearities in three space dimensions. To complement these existence theorems, the final section identifies various criteria ensuring nonexistence of solutions.

Appendices A–E provide for the reader's convenience some background material, with selected proofs, on inequalities, linear functional analysis, measure theory, etc.

The Bibliography is an updated and extensive listing of interesting PDE books to consult for further information. Since this is a textbook and not a reference monograph, I have mostly not attempted to track down and document the original sources for the myriads of ideas and methods we will encounter. The mathematical literature for partial differential equations is truly vast, but the books cited in the Bibliography should at least provide a starting point for locating the primary sources. (Citations to selected research papers appear throughout the text.)

1.5. PROBLEMS

1. Classify each of the partial differential equations in §1.2 as follows:
 - (a) Is the PDE linear, semilinear, quasilinear or fully nonlinear?
 - (b) What is the order of the PDE?
2. Let k be a positive integer. Show that a smooth function defined on \mathbb{R}^n has in general

$$\binom{n+k-1}{k} = \binom{n+k-1}{n-1}$$

distinct partial derivatives of order k .

(Hint: This is the number of ways of inserting $n-1$ dividers $|$ within a row of k symbols \circ : for example, $\circ \circ || \circ \circ \circ | \circ | \circ \circ \circ || \circ \circ \circ \circ$.) Explain why each such pattern corresponds to precisely one of the partial derivatives of order k .)

The next exercises provide some practice with the multiindex notation introduced in Appendix A.

3. Prove the *Multinomial Theorem*:

$$(x_1 + \dots + x_n)^k = \sum_{|\alpha|=k} \binom{|\alpha|}{\alpha} x^\alpha,$$

where $\binom{|\alpha|}{\alpha} := \frac{|\alpha|!}{\alpha!}$, $\alpha! = \alpha_1! \alpha_2! \dots \alpha_n!$, and $x^\alpha = x_1^{\alpha_1} \dots x_n^{\alpha_n}$. The sum is taken over all multiindices $\alpha = (\alpha_1, \dots, \alpha_n)$ with $|\alpha| = k$.

4. Prove *Leibniz's formula*:

$$D^\alpha(uv) = \sum_{\beta \leq \alpha} \binom{\alpha}{\beta} D^\beta u D^{\alpha-\beta} v,$$

where $u, v : \mathbb{R}^n \rightarrow \mathbb{R}$ are smooth, $\binom{\alpha}{\beta} := \frac{\alpha!}{\beta!(\alpha-\beta)!}$, and $\beta \leq \alpha$ means $\beta_i \leq \alpha_i$ ($i = 1, \dots, n$).

5. Assume that $f : \mathbb{R}^n \rightarrow \mathbb{R}$ is smooth. Prove

$$f(x) = \sum_{|\alpha| \leq k} \frac{1}{\alpha!} D^\alpha f(0) x^\alpha + O(|x|^{k+1}) \quad \text{as } x \rightarrow 0$$

for each $k = 1, 2, \dots$. This is *Taylor's formula* in multiindex notation.

(Hint: Fix $x \in \mathbb{R}^n$ and consider the function of one variable $g(t) := f(tx)$.)

1.6. REFERENCES

Klainerman's article [Kl] is a nice modern overview of the field of partial differential equations.

Good general texts and monographs on PDE include Arnold [Ar2], Courant–Hilbert [C-H], DiBenedetto [DB1], Folland [F1], Friedman [Fr2], Garabedian [G], John [J2], Jost [Jo], McOwen [MO], Mikhailov [M], Petrovsky [Py], Rauch [R], Renardy–Rogers [R-R], Smirnov [Sm], Smoller [S], Strauss [St2], Taylor [Ta], Thoe–Zachmanoglou [T-Z], Zauderer [Za], and many others. The prefaces to Arnold [Ar2] and to Bernstein [Bt] are interesting reading. Zwillinger's handbook [Zw] on differential equations is a useful compendium of methods for PDE.

This page is intentionally left blank.

Part I

**REPRESENTATION
FORMULAS FOR
SOLUTIONS**

This page is intentionally left blank.

FOUR IMPORTANT LINEAR PARTIAL DIFFERENTIAL EQUATIONS

- 2.1 Transport equation
- 2.2 Laplace's equation
- 2.3 Heat equation
- 2.4 Wave equation
- 2.5 Problems
- 2.6 References

In this chapter we introduce four fundamental linear partial differential equations for which various explicit formulas for solutions are available. These are

the transport equation	$u_t + b \cdot Du = 0$	(§2.1),
Laplace's equation	$\Delta u = 0$	(§2.2),
the heat equation	$u_t - \Delta u = 0$	(§2.3),
the wave equation	$u_{tt} - \Delta u = 0$	(§2.4).

Before going further, the reader should review the discussions of inequalities, integration by parts, Green's formulas, convolutions, etc., in Appendices B and C and later refer back to these as necessary.

2.1. TRANSPORT EQUATION

One of the simplest partial differential equations is the *transport equation* with constant coefficients. This is the PDE

$$(1) \quad u_t + b \cdot Du = 0 \quad \text{in } \mathbb{R}^n \times (0, \infty),$$

where b is a fixed vector in \mathbb{R}^n , $b = (b_1, \dots, b_n)$, and $u : \mathbb{R}^n \times [0, \infty) \rightarrow \mathbb{R}$ is the unknown, $u = u(x, t)$. Here $x = (x_1, \dots, x_n) \in \mathbb{R}^n$ denotes a typical point in space, and $t \geq 0$ denotes a typical time. We write $Du = D_x u = (u_{x_1}, \dots, u_{x_n})$ for the gradient of u with respect to the spatial variables x .

Which functions u solve (1)? To answer, let us suppose for the moment we are given some smooth solution u and try to compute it. To do so, we first must recognize that the partial differential equation (1) asserts that a particular directional derivative of u vanishes. We exploit this insight by fixing any point $(x, t) \in \mathbb{R}^n \times (0, \infty)$ and defining

$$z(s) := u(x + sb, t + s) \quad (s \in \mathbb{R}).$$

We then calculate

$$\dot{z}(s) = Du(x + sb, t + s) \cdot b + u_t(x + sb, t + s) = 0 \quad \left(\dot{\cdot} = \frac{d}{ds} \right),$$

the second equality holding owing to (1). Thus $z(\cdot)$ is a constant function of s , and consequently for each point (x, t) , u is constant on the line through (x, t) with the direction $(b, 1) \in \mathbb{R}^{n+1}$. Hence if we know the value of u at any point on each such line, we know its value everywhere in $\mathbb{R}^n \times (0, \infty)$.

2.1.1. Initial-value problem.

For definiteness therefore, let us consider the initial-value problem

$$(2) \quad \begin{cases} u_t + b \cdot Du = 0 & \text{in } \mathbb{R}^n \times (0, \infty) \\ u = g & \text{on } \mathbb{R}^n \times \{t = 0\}. \end{cases}$$

Here $b \in \mathbb{R}^n$ and $g : \mathbb{R}^n \rightarrow \mathbb{R}$ are known, and the problem is to compute u . Given (x, t) as above, the line through (x, t) with direction $(b, 1)$ is represented parametrically by $(x + sb, t + s)$ ($s \in \mathbb{R}$). This line hits the plane $\Gamma := \mathbb{R}^n \times \{t = 0\}$ when $s = -t$, at the point $(x - tb, 0)$. Since u is constant on the line and $u(x - tb, 0) = g(x - tb)$, we deduce

$$(3) \quad u(x, t) = g(x - tb) \quad (x \in \mathbb{R}^n, t \geq 0).$$

So, if (2) has a sufficiently regular solution u , it must certainly be given by (3). And conversely, it is easy to check directly that if g is C^1 , then u defined by (3) is indeed a solution of (2).

Weak solutions. If g is not C^1 , then there is obviously no C^1 solution of (2). But even in this case formula (3) certainly provides a strong, and in fact the only reasonable, candidate for a solution. We may thus informally declare $u(x, t) = g(x - tb)$ ($x \in \mathbb{R}^n$, $t \geq 0$) to be a *weak solution* of (2), even should g not be C^1 . This all makes sense even if g and thus u are discontinuous. Such a notion, that a nonsmooth or even discontinuous function may sometimes solve a PDE, will come up again later when we study nonlinear transport phenomena in §3.4.

2.1.2. Nonhomogeneous problem.

Next let us look at the associated nonhomogeneous problem

$$(4) \quad \begin{cases} u_t + b \cdot Du = f & \text{in } \mathbb{R}^n \times (0, \infty) \\ u = g & \text{on } \mathbb{R}^n \times \{t = 0\}. \end{cases}$$

As before fix $(x, t) \in \mathbb{R}^{n+1}$ and, inspired by the calculation above, set $z(s) := u(x + sb, t + s)$ for $s \in \mathbb{R}$. Then

$$\dot{z}(s) = Du(x + sb, t + s) \cdot b + u_t(x + sb, t + s) = f(x + sb, t + s).$$

Consequently

$$\begin{aligned} u(x, t) - g(x - tb) &= z(0) - z(-t) = \int_{-t}^0 \dot{z}(s) ds \\ &= \int_{-t}^0 f(x + sb, t + s) ds \\ &= \int_0^t f(x + (s - t)b, s) ds, \end{aligned}$$

and so

$$(5) \quad u(x, t) = g(x - tb) + \int_0^t f(x + (s - t)b, s) ds \quad (x \in \mathbb{R}^n, t \geq 0)$$

solves the initial-value problem (4).

We will later employ this formula to solve the one-dimensional wave equation, in §2.4.1.

Remark. Observe that we have derived our solutions (3), (5) by in effect converting the partial differential equations into ordinary differential equations. This procedure is a special case of the *method of characteristics*, developed later in §3.2.

2.2. LAPLACE'S EQUATION

Among the most important of all partial differential equations are undoubtedly *Laplace's equation*

$$(1) \quad \Delta u = 0$$

and *Poisson's equation*

$$(2) \quad -\Delta u = f. *$$

In both (1) and (2), $x \in U$ and the unknown is $u : \bar{U} \rightarrow \mathbb{R}$, $u = u(x)$, where $U \subset \mathbb{R}^n$ is a given open set. In (2) the function $f : U \rightarrow \mathbb{R}$ is also given. Remember from §A.3 that the *Laplacian* of u is $\Delta u = \sum_{i=1}^n u_{x_i x_i}$.

DEFINITION. A C^2 function u satisfying (1) is called a *harmonic function*.

Physical interpretation. Laplace's equation comes up in a wide variety of physical contexts. In a typical interpretation u denotes the density of some quantity (e.g. a chemical concentration) in equilibrium. Then if V is any smooth subregion within U , the net flux of u through ∂V is zero:

$$\int_{\partial V} \mathbf{F} \cdot \boldsymbol{\nu} \, dS = 0,$$

\mathbf{F} denoting the flux density and $\boldsymbol{\nu}$ the unit outer normal field. In view of the Gauss–Green Theorem (§C.2), we have

$$\int_V \operatorname{div} \mathbf{F} \, dx = \int_{\partial V} \mathbf{F} \cdot \boldsymbol{\nu} \, dS = 0,$$

and so

$$(3) \quad \operatorname{div} \mathbf{F} = 0 \quad \text{in } U,$$

since V was arbitrary. In many instances it is physically reasonable to assume the flux \mathbf{F} is proportional to the gradient Du but points in the opposite direction (since the flow is from regions of higher to lower concentration). Thus

$$(4) \quad \mathbf{F} = -aDu \quad (a > 0).$$

*I prefer to write (2) with the minus sign, to be consistent with the notation for general second-order elliptic operators in Chapter 6.

Substituting into (3), we obtain Laplace's equation

$$\operatorname{div}(Du) = \Delta u = 0.$$

If u denotes the

$$\left\{ \begin{array}{l} \text{chemical concentration} \\ \text{temperature} \\ \text{electrostatic potential,} \end{array} \right.$$

equation (4) is

$$\left\{ \begin{array}{l} \text{Fick's law of diffusion} \\ \text{Fourier's law of heat conduction} \\ \text{Ohm's law of electrical conduction.} \end{array} \right.$$

See Feynman–Leighton–Sands [F-L-S, Chapter 12] for a discussion of the ubiquity of Laplace's equation in mathematical physics. Laplace's equation arises as well in the study of analytic functions and the probabilistic investigation of Brownian motion.

2.2.1. Fundamental solution.

a. Derivation of fundamental solution. One good strategy for investigating any partial differential equation is first to identify some explicit solutions and then, provided the PDE is linear, to assemble more complicated solutions out of the specific ones previously noted. Furthermore, in looking for explicit solutions, it is often wise to restrict attention to classes of functions with certain symmetry properties. Since Laplace's equation is invariant under rotations (Problem 2), it consequently seems advisable to search first for *radial* solutions, that is, functions of $r = |x|$.

Let us therefore attempt to find a solution u of Laplace's equation (1) in $U = \mathbb{R}^n$, having the form

$$u(x) = v(r),$$

where $r = |x| = (x_1^2 + \cdots + x_n^2)^{1/2}$ and v is to be selected (if possible) so that $\Delta u = 0$ holds. First note for $i = 1, \dots, n$ that

$$\frac{\partial r}{\partial x_i} = \frac{1}{2} (x_1^2 + \cdots + x_n^2)^{-1/2} 2x_i = \frac{x_i}{r} \quad (x \neq 0).$$

We thus have

$$u_{x_i} = v'(r) \frac{x_i}{r}, \quad u_{x_i x_i} = v''(r) \frac{x_i^2}{r^2} + v'(r) \left(\frac{1}{r} - \frac{x_i^2}{r^3} \right)$$

for $i = 1, \dots, n$, and so

$$\Delta u = v''(r) + \frac{n-1}{r}v'(r).$$

Hence $\Delta u = 0$ if and only if

$$(5) \quad v'' + \frac{n-1}{r}v' = 0.$$

If $v' \neq 0$, we deduce

$$\log(|v'|)' = \frac{v''}{v'} = \frac{1-n}{r},$$

and hence $v'(r) = \frac{a}{r^{n-1}}$ for some constant a . Consequently if $r > 0$, we have

$$v(r) = \begin{cases} b \log r + c & (n = 2) \\ \frac{b}{r^{n-2}} + c & (n \geq 3), \end{cases}$$

where b and c are constants.

These considerations motivate the following

DEFINITION. *The function*

$$(6) \quad \Phi(x) := \begin{cases} -\frac{1}{2\pi} \log |x| & (n = 2) \\ \frac{1}{n(n-2)\alpha(n)} \frac{1}{|x|^{n-2}} & (n \geq 3), \end{cases}$$

defined for $x \in \mathbb{R}^n$, $x \neq 0$, is the fundamental solution of Laplace's equation.

The reason for the particular choices of the constants in (6) will be apparent in a moment. (Recall from §A.2 that $\alpha(n)$ denotes the volume of the unit ball in \mathbb{R}^n .)

We will sometimes slightly abuse notation and write $\Phi(x) = \Phi(|x|)$ to emphasize that the fundamental solution is radial. Observe also that we have the estimates

$$(7) \quad |D\Phi(x)| \leq \frac{C}{|x|^{n-1}}, \quad |D^2\Phi(x)| \leq \frac{C}{|x|^n} \quad (x \neq 0)$$

for some constant $C > 0$.

b. Poisson's equation. By construction the function $x \mapsto \Phi(x)$ is harmonic for $x \neq 0$. If we shift the origin to a new point y , the PDE (1) is unchanged; and so $x \mapsto \Phi(x - y)$ is also harmonic as a function of x , $x \neq y$. Let us now take $f : \mathbb{R}^n \rightarrow \mathbb{R}$ and note that the mapping $x \mapsto \Phi(x - y)f(y)$ ($x \neq y$) is harmonic for each point $y \in \mathbb{R}^n$, and thus so is the sum of finitely many such expressions built for different points y .

This reasoning might suggest that the convolution

$$(8) \quad \begin{aligned} u(x) &= \int_{\mathbb{R}^n} \Phi(x-y) f(y) dy \\ &= \begin{cases} -\frac{1}{2\pi} \int_{\mathbb{R}^2} \log(|x-y|) f(y) dy & (n=2) \\ \frac{1}{n(n-2)\alpha(n)} \int_{\mathbb{R}^n} \frac{f(y)}{|x-y|^{n-2}} dy & (n \geq 3) \end{cases} \end{aligned}$$

will solve Laplace's equation (1). *However, this is wrong.* Indeed, as intimated by estimate (7), $D^2\Phi(x-y)$ is *not* summable near the singularity at $y=x$, and so naive differentiation through the integral sign is unjustified (and incorrect). We must proceed more carefully in calculating Δu .

Let us for simplicity now assume $f \in C_c^2(\mathbb{R}^n)$; that is, f is twice continuously differentiable, with compact support.

THEOREM 1 (Solving Poisson's equation). *Define u by (8). Then*

$$(i) \quad u \in C^2(\mathbb{R}^n)$$

and

$$(ii) \quad -\Delta u = f \quad \text{in } \mathbb{R}^n.$$

We consequently see that (8) provides us with a formula for a solution of Poisson's equation (2) in \mathbb{R}^n .

Proof. 1. We have

$$(9) \quad u(x) = \int_{\mathbb{R}^n} \Phi(x-y) f(y) dy = \int_{\mathbb{R}^n} \Phi(y) f(x-y) dy;$$

hence

$$\frac{u(x+he_i) - u(x)}{h} = \int_{\mathbb{R}^n} \Phi(y) \left[\frac{f(x+he_i-y) - f(x-y)}{h} \right] dy,$$

where $h \neq 0$ and $e_i = (0, \dots, 1, \dots, 0)$, the 1 in the i^{th} -slot. But

$$\frac{f(x+he_i-y) - f(x-y)}{h} \rightarrow f_{x_i}(x-y)$$

uniformly on \mathbb{R}^n as $h \rightarrow 0$, and thus

$$u_{x_i}(x) = \int_{\mathbb{R}^n} \Phi(y) f_{x_i}(x-y) dy \quad (i=1, \dots, n).$$

Similarly

$$(10) \quad u_{x_i x_j}(x) = \int_{\mathbb{R}^n} \Phi(y) f_{x_i x_j}(x-y) dy \quad (i, j=1, \dots, n).$$

As the expression on the right-hand side of (10) is continuous in the variable x , we see $u \in C^2(\mathbb{R}^n)$.

2. Since Φ blows up at 0, we will need for subsequent calculations to isolate this singularity inside a small ball. So fix $\varepsilon > 0$. Then

$$(11) \quad \begin{aligned} \Delta u(x) &= \int_{B(0,\varepsilon)} \Phi(y) \Delta_x f(x-y) dy + \int_{\mathbb{R}^n - B(0,\varepsilon)} \Phi(y) \Delta_x f(x-y) dy \\ &=: I_\varepsilon + J_\varepsilon. \end{aligned}$$

Now

$$(12) \quad |I_\varepsilon| \leq C \|D^2 f\|_{L^\infty(\mathbb{R}^n)} \int_{B(0,\varepsilon)} |\Phi(y)| dy \leq \begin{cases} C\varepsilon^2 |\log \varepsilon| & (n=2) \\ C\varepsilon^2 & (n \geq 3). \end{cases}$$

An integration by parts (see §C.2) yields

$$(13) \quad \begin{aligned} J_\varepsilon &= \int_{\mathbb{R}^n - B(0,\varepsilon)} \Phi(y) \Delta_y f(x-y) dy \\ &= - \int_{\mathbb{R}^n - B(0,\varepsilon)} D\Phi(y) \cdot D_y f(x-y) dy \\ &\quad + \int_{\partial B(0,\varepsilon)} \Phi(y) \frac{\partial f}{\partial \nu}(x-y) dS(y) \\ &=: K_\varepsilon + L_\varepsilon, \end{aligned}$$

ν denoting the *inward* pointing unit normal along $\partial B(0,\varepsilon)$. We readily check

$$(14) \quad |L_\varepsilon| \leq \|Df\|_{L^\infty(\mathbb{R}^n)} \int_{\partial B(0,\varepsilon)} |\Phi(y)| dS(y) \leq \begin{cases} C\varepsilon |\log \varepsilon| & (n=2) \\ C\varepsilon & (n \geq 3). \end{cases}$$

3. We continue by integrating by parts once again in the term K_ε , to discover

$$\begin{aligned} K_\varepsilon &= \int_{\mathbb{R}^n - B(0,\varepsilon)} \Delta \Phi(y) f(x-y) dy - \int_{\partial B(0,\varepsilon)} \frac{\partial \Phi}{\partial \nu}(y) f(x-y) dS(y) \\ &= - \int_{\partial B(0,\varepsilon)} \frac{\partial \Phi}{\partial \nu}(y) f(x-y) dS(y), \end{aligned}$$

since Φ is harmonic away from the origin. Now $D\Phi(y) = \frac{-1}{n\alpha(n)} \frac{y}{|y|^n}$ ($y \neq 0$) and $\nu = \frac{-y}{|y|} = -\frac{y}{\varepsilon}$ on $\partial B(0,\varepsilon)$. Consequently $\frac{\partial \Phi}{\partial \nu}(y) = \nu \cdot D\Phi(y) = \frac{1}{n\alpha(n)\varepsilon^{n-1}}$ on $\partial B(0,\varepsilon)$. Since $n\alpha(n)\varepsilon^{n-1}$ is the surface area of the sphere $\partial B(0,\varepsilon)$, we have

$$(15) \quad \begin{aligned} K_\varepsilon &= -\frac{1}{n\alpha(n)\varepsilon^{n-1}} \int_{\partial B(0,\varepsilon)} f(x-y) dS(y) \\ &= -\int_{\partial B(x,\varepsilon)} f(y) dS(y) \rightarrow -f(x) \quad \text{as } \varepsilon \rightarrow 0. \end{aligned}$$

(Remember from §A.3 that a slash through an integral denotes an average.)

4. Combining now (11)–(15) and letting $\varepsilon \rightarrow 0$, we find $-\Delta u(x) = f(x)$, as asserted. \square

Theorem 1 is in fact valid under far less stringent smoothness requirements for f : see Gilbarg–Trudinger [G-T].

Interpretation of fundamental solution. We sometimes write

$$-\Delta\Phi = \delta_0 \quad \text{in } \mathbb{R}^n,$$

δ_0 denoting the Dirac measure on \mathbb{R}^n giving unit mass to the point 0. Adopting this notation, we may formally compute

$$\begin{aligned} -\Delta u(x) &= \int_{\mathbb{R}^n} -\Delta_x \Phi(x-y) f(y) dy \\ &= \int_{\mathbb{R}^n} \delta_x f(y) dy = f(x) \quad (x \in \mathbb{R}^n), \end{aligned}$$

in accordance with Theorem 1. This corrects the faulty calculation (9).

2.2.2. Mean-value formulas.

Consider now an open set $U \subset \mathbb{R}^n$ and suppose u is a harmonic function within U . We next derive the important *mean-value formulas*, which declare that $u(x)$ equals both the average of u over the sphere $\partial B(x, r)$ and the average of u over the entire ball $B(x, r)$, provided $B(x, r) \subset U$. These implicit formulas involving u generate a remarkable number of consequences, as we will momentarily see.

THEOREM 2 (Mean-value formulas for Laplace's equation). *If $u \in C^2(U)$ is harmonic, then*

$$(16) \quad u(x) = \int_{\partial B(x,r)} u dS = \int_{B(x,r)} u dy$$

for each ball $B(x, r) \subset U$.

Proof. 1. Set

$$\phi(r) := \int_{\partial B(x,r)} u(y) dS(y) = \int_{\partial B(0,1)} u(x + rz) dS(z).$$

Then

$$\phi'(r) = \int_{\partial B(0,1)} Du(x + rz) \cdot z dS(z),$$

and consequently, using Green's formulas from §C.2, we compute

$$\begin{aligned}\phi'(r) &= \int_{\partial B(x,r)} Du(y) \cdot \frac{y-x}{r} dS(y) \\ &= \int_{\partial B(x,r)} \frac{\partial u}{\partial \nu} dS(y) \\ &= \frac{r}{n} \int_{B(x,r)} \Delta u(y) dy = 0.\end{aligned}$$

Hence ϕ is constant, and so

$$\phi(r) = \lim_{t \rightarrow 0} \phi(t) = \lim_{t \rightarrow 0} \int_{\partial B(x,t)} u(y) dS(y) = u(x).$$

2. Observe next that our employing polar coordinates, as in §C.3, gives

$$\begin{aligned}\int_{B(x,r)} u dy &= \int_0^r \left(\int_{\partial B(x,s)} u dS \right) ds \\ &= u(x) \int_0^r n\alpha(n)s^{n-1} ds = \alpha(n)r^n u(x).\end{aligned} \quad \square$$

THEOREM 3 (Converse to mean-value property). *If $u \in C^2(U)$ satisfies*

$$u(x) = \int_{\partial B(x,r)} u dS$$

for each ball $B(x,r) \subset U$, then u is harmonic.

Proof. If $\Delta u \not\equiv 0$, there exists some ball $B(x,r) \subset U$ such that, say, $\Delta u > 0$ within $B(x,r)$. But then for ϕ as above,

$$0 = \phi'(r) = \frac{r}{n} \int_{B(x,r)} \Delta u(y) dy > 0,$$

a contradiction. □

2.2.3. Properties of harmonic functions.

We now present a sequence of interesting deductions about harmonic functions, all based upon the mean-value formulas. Assume for the following that $U \subset \mathbb{R}^n$ is open and bounded.

a. Strong maximum principle, uniqueness. We begin with the assertion that a harmonic function must attain its maximum on the boundary and cannot attain its maximum in the interior of a connected region unless it is constant.

THEOREM 4 (Strong maximum principle). *Suppose $u \in C^2(U) \cap C(\bar{U})$ is harmonic within U .*

(i) *Then*

$$\max_{\bar{U}} u = \max_{\partial U} u.$$

(ii) *Furthermore, if U is connected and there exists a point $x_0 \in U$ such that*

$$u(x_0) = \max_{\bar{U}} u,$$

then

u is constant within U .

Assertion (i) is the *maximum principle* for Laplace's equation and (ii) is the *strong maximum principle*. Replacing u by $-u$, we recover also similar assertions with "min" replacing "max".

Proof. Suppose there exists a point $x_0 \in U$ with $u(x_0) = M := \max_{\bar{U}} u$. Then for $0 < r < \text{dist}(x_0, \partial U)$, the mean-value property asserts

$$M = u(x_0) = \int_{B(x_0, r)} u \, dy \leq M.$$

As equality holds only if $u \equiv M$ within $B(x_0, r)$, we see $u(y) = M$ for all $y \in B(x_0, r)$. Hence the set $\{x \in U \mid u(x) = M\}$ is both open and relatively closed in U and thus equals U if U is connected. This proves assertion (ii), from which (i) follows. \square

Positivity. The strong maximum principle asserts in particular that if U is connected and $u \in C^2(U) \cap C(\bar{U})$ satisfies

$$\begin{cases} \Delta u = 0 & \text{in } U \\ u = g & \text{on } \partial U, \end{cases}$$

where $g \geq 0$, then u is positive everywhere in U if g is positive somewhere on ∂U .

An important application of the maximum principle is establishing the uniqueness of solutions to certain boundary-value problems for Poisson's equation.

THEOREM 5 (Uniqueness). *Let $g \in C(\partial U)$, $f \in C(U)$. Then there exists at most one solution $u \in C^2(U) \cap C(\bar{U})$ of the boundary-value problem*

$$(17) \quad \begin{cases} -\Delta u = f & \text{in } U \\ u = g & \text{on } \partial U. \end{cases}$$

Proof. If u and \tilde{u} both satisfy (17), apply Theorem 4 to the harmonic functions $w := \pm(u - \tilde{u})$. □

b. Regularity. Next we prove that if $u \in C^2$ is harmonic, then necessarily $u \in C^\infty$. Thus *harmonic functions are automatically infinitely differentiable*. This sort of assertion is called a *regularity* theorem. The interesting point is that the algebraic structure of Laplace's equation $\Delta u = \sum_{i=1}^n u_{x_i x_i} = 0$ leads to the analytic deduction that all the partial derivatives of u exist, even those which do not appear in the PDE.

THEOREM 6 (Smoothness). *If $u \in C(U)$ satisfies the mean-value property (16) for each ball $B(x, r) \subset U$, then*

$$u \in C^\infty(U).$$

Note carefully that u may not be smooth, or even continuous, up to ∂U .

Proof. Let η be a standard mollifier, as described in §C.4, and recall that η is a radial function. Set $u^\varepsilon := \eta_\varepsilon * u$ in $U_\varepsilon = \{x \in U \mid \text{dist}(x, \partial U) > \varepsilon\}$. As shown in §C.4, $u^\varepsilon \in C^\infty(U_\varepsilon)$.

We will prove u is smooth by demonstrating that in fact $u \equiv u^\varepsilon$ on U_ε . Indeed if $x \in U_\varepsilon$, then

$$\begin{aligned} u^\varepsilon(x) &= \int_U \eta_\varepsilon(x-y)u(y) dy \\ &= \frac{1}{\varepsilon^n} \int_{B(x,\varepsilon)} \eta\left(\frac{|x-y|}{\varepsilon}\right) u(y) dy \\ &= \frac{1}{\varepsilon^n} \int_0^\varepsilon \eta\left(\frac{r}{\varepsilon}\right) \left(\int_{\partial B(x,r)} u dS \right) dr \\ &= \frac{1}{\varepsilon^n} u(x) \int_0^\varepsilon \eta\left(\frac{r}{\varepsilon}\right) n\alpha(n)r^{n-1} dr \quad \text{by (16)} \\ &= u(x) \int_{B(0,\varepsilon)} \eta_\varepsilon dy = u(x). \end{aligned}$$

Thus $u^\varepsilon \equiv u$ in U_ε , and so $u \in C^\infty(U_\varepsilon)$ for each $\varepsilon > 0$. □

c. Local estimates for harmonic functions. Now we employ the mean-value formulas to derive careful estimates on the various partial derivatives of a harmonic function. The precise structure of these estimates will be needed below, when we prove analyticity.

THEOREM 7 (Estimates on derivatives). *Assume u is harmonic in U . Then*

$$(18) \quad |D^\alpha u(x_0)| \leq \frac{C_k}{r^{n+k}} \|u\|_{L^1(B(x_0, r))}$$

for each ball $B(x_0, r) \subset U$ and each multiindex α of order $|\alpha| = k$.

Here

$$(19) \quad C_0 = \frac{1}{\alpha(n)}, \quad C_k = \frac{(2^{n+1}nk)^k}{\alpha(n)} \quad (k = 1, \dots).$$

Proof. 1. We establish (18), (19) by induction on k , the case $k = 0$ being immediate from the mean-value formula (16). For $k = 1$, we note upon differentiating Laplace's equation that u_{x_i} ($i = 1, \dots, n$) is harmonic. Consequently

$$(20) \quad \begin{aligned} |u_{x_i}(x_0)| &= \left| \int_{B(x_0, r/2)} u_{x_i} dx \right| \\ &= \left| \frac{2^n}{\alpha(n)r^n} \int_{\partial B(x_0, r/2)} u \nu_i dS \right| \\ &\leq \frac{2n}{r} \|u\|_{L^\infty(\partial B(x_0, \frac{r}{2}))}. \end{aligned}$$

Now if $x \in \partial B(x_0, r/2)$, then $B(x, r/2) \subset B(x_0, r) \subset U$, and so

$$|u(x)| \leq \frac{1}{\alpha(n)} \left(\frac{2}{r}\right)^n \|u\|_{L^1(B(x_0, r))}$$

by (18), (19) for $k = 0$. Combining the inequalities above, we deduce

$$|D^\alpha u(x_0)| \leq \frac{2^{n+1}n}{\alpha(n)} \frac{1}{r^{n+1}} \|u\|_{L^1(B(x_0, r))}$$

if $|\alpha| = 1$. This verifies (18), (19) for $k = 1$.

2. Assume now $k \geq 2$ and (18), (19) are valid for all balls in U and each multiindex of order less than or equal to $k - 1$. Fix $B(x_0, r) \subset U$ and let α

be a multiindex with $|\alpha| = k$. Then $D^\alpha u = (D^\beta u)_{x_i}$ for some $i \in \{1, \dots, n\}$, $|\beta| = k - 1$. By calculations similar to those in (20), we establish that

$$|D^\alpha u(x_0)| \leq \frac{nk}{r} \|D^\beta u\|_{L^\infty(\partial B(x_0, \frac{r}{k}))}.$$

If $x \in \partial B(x_0, \frac{r}{k})$, then $B(x, \frac{k-1}{k}r) \subset B(x_0, r) \subset U$. Thus (18), (19) for $k - 1$ imply

$$|D^\beta u(x)| \leq \frac{(2^{n+1}n(k-1))^{k-1}}{\alpha(n) \left(\frac{k-1}{k}r\right)^{n+k-1}} \|u\|_{L^1(B(x_0, r))}.$$

Combining the two previous estimates yields the bound

$$(21) \quad |D^\alpha u(x_0)| \leq \frac{(2^{n+1}nk)^k}{\alpha(n)r^{n+k}} \|u\|_{L^1(B(x_0, r))}.$$

This confirms (18), (19) for $|\alpha| = k$. □

d. Liouville's Theorem. We assert now that there are no nontrivial bounded harmonic functions on all of \mathbb{R}^n .

THEOREM 8 (Liouville's Theorem). *Suppose $u : \mathbb{R}^n \rightarrow \mathbb{R}$ is harmonic and bounded. Then u is constant.*

Proof. Fix $x_0 \in \mathbb{R}^n$, $r > 0$, and apply Theorem 7 on $B(x_0, r)$:

$$\begin{aligned} |Du(x_0)| &\leq \frac{\sqrt{n}C_1}{r^{n+1}} \|u\|_{L^1(B(x_0, r))} \\ &\leq \frac{\sqrt{n}C_1\alpha(n)}{r} \|u\|_{L^\infty(\mathbb{R}^n)} \rightarrow 0, \end{aligned}$$

as $r \rightarrow \infty$. Thus $Du \equiv 0$, and so u is constant. □

THEOREM 9 (Representation formula). *Let $f \in C_c^2(\mathbb{R}^n)$, $n \geq 3$. Then any bounded solution of*

$$-\Delta u = f \quad \text{in } \mathbb{R}^n$$

has the form

$$u(x) = \int_{\mathbb{R}^n} \Phi(x-y)f(y) dy + C \quad (x \in \mathbb{R}^n)$$

for some constant C .

Proof. Since $\Phi(x) \rightarrow 0$ as $|x| \rightarrow \infty$ for $n \geq 3$, $\tilde{u}(x) := \int_{\mathbb{R}^n} \Phi(x-y)f(y) dy$ is a bounded solution of $-\Delta u = f$ in \mathbb{R}^n . If u is another solution, $w := u - \tilde{u}$ is constant, according to Liouville's Theorem. \square

Remark. If $n = 2$, $\Phi(x) = -\frac{1}{2\pi} \log|x|$ is unbounded as $|x| \rightarrow \infty$ and so may be $\int_{\mathbb{R}^2} \Phi(x-y)f(y) dy$.

e. Analyticity. We refine Theorem 6:

THEOREM 10 (Analyticity). *Assume u is harmonic in U . Then u is analytic in U .*

Proof. 1. Fix any point $x_0 \in U$. We must show u can be represented by a convergent power series in some neighborhood of x_0 .

Let $r := \frac{1}{4} \text{dist}(x_0, \partial U)$. Then $M := \frac{1}{\alpha(n)r^n} \|u\|_{L^1(B(x_0, 2r))} < \infty$.

2. Since $B(x, r) \subset B(x_0, 2r) \subset U$ for each $x \in B(x_0, r)$, Theorem 7 provides the bound

$$\|D^\alpha u\|_{L^\infty(B(x_0, r))} \leq M \left(\frac{2^{n+1}n}{r} \right)^{|\alpha|} |\alpha|^{|\alpha|}.$$

Now $\frac{k^k}{k!} < e^k$ for all positive integers k , and hence

$$|\alpha|^{|\alpha|} \leq e^{|\alpha|} |\alpha|!$$

for all multiindices α . Furthermore, the Multinomial Theorem (§1.5) implies

$$n^k = (1 + \cdots + 1)^k = \sum_{|\alpha|=k} \frac{|\alpha|!}{\alpha!},$$

whence

$$|\alpha|! \leq n^{|\alpha|} \alpha!.$$

Combining the previous inequalities yields the estimate

$$(22) \quad \|D^\alpha u\|_{L^\infty(B(x_0, r))} \leq CM \left(\frac{2^{n+1}n^2e}{r} \right)^{|\alpha|} \alpha!.$$

3. The Taylor series for u at x_0 is

$$\sum_{\alpha} \frac{D^\alpha u(x_0)}{\alpha!} (x - x_0)^\alpha,$$

the sum taken over all multiindices. We assert this power series converges, provided

$$(23) \quad |x - x_0| < \frac{r}{2^{n+2}n^3e}.$$

To verify this, let us compute for each N the remainder term:

$$\begin{aligned} R_N(x) &:= u(x) - \sum_{k=0}^{N-1} \sum_{|\alpha|=k} \frac{D^\alpha u(x_0)(x - x_0)^\alpha}{\alpha!} \\ &= \sum_{|\alpha|=N} \frac{D^\alpha u(x_0 + t(x - x_0))(x - x_0)^\alpha}{\alpha!} \end{aligned}$$

for some $0 \leq t \leq 1$, t depending on x . We establish this formula by writing out the first N terms and the error in the Taylor expansion about 0 for the function of one variable $g(t) := u(x_0 + t(x - x_0))$, at $t = 1$. Employing (22), (23), we can estimate

$$\begin{aligned} |R_N(x)| &\leq CM \sum_{|\alpha|=N} \left(\frac{2^{n+1}n^2e}{r} \right)^N \left(\frac{r}{2^{n+2}n^3e} \right)^N \\ &\leq CMn^N \frac{1}{(2n)^N} = \frac{CM}{2^N} \rightarrow 0 \quad \text{as } N \rightarrow \infty. \quad \square \end{aligned}$$

See §4.6.2 for more on analytic functions and partial differential equations.

f. Harnack's inequality. Recall from §A.2 that we write $V \subset\subset U$ to mean $V \subset \bar{V} \subset U$ and \bar{V} is compact.

THEOREM 11 (Harnack's inequality). *For each connected open set $V \subset\subset U$, there exists a positive constant C , depending only on V , such that*

$$\sup_V u \leq C \inf_V u$$

for all nonnegative harmonic functions u in U .

Thus in particular

$$\frac{1}{C}u(y) \leq u(x) \leq Cu(y)$$

for all points $x, y \in V$. These inequalities assert that *the values of a non-negative harmonic function within V are all comparable*: u cannot be very small (or very large) at any point of V unless u is very small (or very large) everywhere in V . The intuitive idea is that since V is a positive distance away from ∂U , there is “room for the averaging effects of Laplace's equation to occur”.

Proof. Let $r := \frac{1}{4} \text{dist}(V, \partial U)$. Choose $x, y \in V$, $|x - y| \leq r$. Then

$$\begin{aligned} u(x) &= \int_{B(x, 2r)} u \, dz \geq \frac{1}{\alpha(n)2^n r^n} \int_{B(y, r)} u \, dz \\ &= \frac{1}{2^n} \int_{B(y, r)} u \, dz = \frac{1}{2^n} u(y). \end{aligned}$$

Thus $2^n u(y) \geq u(x) \geq \frac{1}{2^n} u(y)$ if $x, y \in V$, $|x - y| \leq r$.

Since V is connected and \bar{V} is compact, we can cover \bar{V} by a chain of finitely many balls $\{B_i\}_{i=1}^N$, each of which has radius $\frac{r}{2}$ and $B_i \cap B_{i-1} \neq \emptyset$ for $i = 2, \dots, N$. Then

$$u(x) \geq \frac{1}{2^{n(N+1)}} u(y)$$

for all $x, y \in V$. □

2.2.4. Green's function.

Assume now $U \subset \mathbb{R}^n$ is open, bounded, and ∂U is C^1 . We propose next to obtain a general representation formula for the solution of Poisson's equation

$$-\Delta u = f \quad \text{in } U,$$

subject to the prescribed boundary condition

$$u = g \quad \text{on } \partial U.$$

a. Derivation of Green's function. Suppose $u \in C^2(\bar{U})$ is an arbitrary function. Fix $x \in U$, choose $\varepsilon > 0$ so small that $B(x, \varepsilon) \subset U$, and apply Green's formula from §C.2 on the region $V_\varepsilon := U - B(x, \varepsilon)$ to $u(y)$ and $\Phi(y - x)$. We thereby compute

$$\begin{aligned} (24) \quad & \int_{V_\varepsilon} u(y) \Delta \Phi(y - x) - \Phi(y - x) \Delta u(y) \, dy \\ &= \int_{\partial V_\varepsilon} u(y) \frac{\partial \Phi}{\partial \nu}(y - x) - \Phi(y - x) \frac{\partial u}{\partial \nu}(y) \, dS(y), \end{aligned}$$

ν denoting the outer unit normal vector on ∂V_ε . Recall next $\Delta \Phi(x - y) = 0$ for $x \neq y$. We observe also

$$\left| \int_{\partial B(x, \varepsilon)} \Phi(y - x) \frac{\partial u}{\partial \nu}(y) \, dS(y) \right| \leq C \varepsilon^{n-1} \max_{\partial B(0, \varepsilon)} |\Phi| = o(1)$$

as $\varepsilon \rightarrow 0$. Furthermore the calculations in the proof of Theorem 1 show

$$\int_{\partial B(x, \varepsilon)} u(y) \frac{\partial \Phi}{\partial \nu}(y - x) \, dS(y) = \int_{\partial B(x, \varepsilon)} u(y) \, dS(y) \rightarrow u(x)$$

as $\varepsilon \rightarrow 0$. Hence our sending $\varepsilon \rightarrow 0$ in (24) yields the formula

$$(25) \quad u(x) = \int_{\partial U} \Phi(y-x) \frac{\partial u}{\partial \nu}(y) - u(y) \frac{\partial \Phi}{\partial \nu}(y-x) dS(y) \\ - \int_U \Phi(y-x) \Delta u(y) dy.$$

This identity is valid for any point $x \in U$ and any function $u \in C^2(\bar{U})$.

Now formula (25) would permit us to solve for $u(x)$ if we knew the values of Δu within U and the values of $u, \partial u/\partial \nu$ along ∂U . However, for our application to Poisson's equation with prescribed boundary values for u , the normal derivative $\partial u/\partial \nu$ along ∂U is unknown to us. We must therefore somehow modify (25) to remove this term.

The idea is now to introduce for fixed x a *corrector* function $\phi^x = \phi^x(y)$, solving the boundary-value problem

$$(26) \quad \begin{cases} \Delta \phi^x = 0 & \text{in } U \\ \phi^x = \Phi(y-x) & \text{on } \partial U. \end{cases}$$

Let us apply Green's formula once more, to compute

$$(27) \quad - \int_U \phi^x(y) \Delta u(y) dy = \int_{\partial U} u(y) \frac{\partial \phi^x}{\partial \nu}(y) - \phi^x(y) \frac{\partial u}{\partial \nu}(y) dS(y) \\ = \int_{\partial U} u(y) \frac{\partial \phi^x}{\partial \nu}(y) - \Phi(y-x) \frac{\partial u}{\partial \nu}(y) dS(y).$$

We introduce next this

DEFINITION. Green's function for the region U is

$$G(x, y) := \Phi(y-x) - \phi^x(y) \quad (x, y \in U, x \neq y).$$

Adopting this terminology and adding (27) to (25), we find

$$(28) \quad u(x) = - \int_{\partial U} u(y) \frac{\partial G}{\partial \nu}(x, y) dS(y) - \int_U G(x, y) \Delta u(y) dy \quad (x \in U),$$

where

$$\frac{\partial G}{\partial \nu}(x, y) = D_y G(x, y) \cdot \nu(y)$$

is the outer normal derivative of G with respect to the variable y . Observe that the term $\partial u/\partial \nu$ does not appear in equation (28): we introduced the corrector ϕ^x precisely to achieve this.

Suppose now $u \in C^2(\bar{U})$ solves the boundary-value problem

$$(29) \quad \begin{cases} -\Delta u = f & \text{in } U \\ u = g & \text{on } \partial U, \end{cases}$$

for given continuous functions f, g . Plugging into (28), we obtain

THEOREM 12 (Representation formula using Green's function). *If $u \in C^2(\bar{U})$ solves problem (29), then*

$$(30) \quad u(x) = - \int_{\partial U} g(y) \frac{\partial G}{\partial \nu}(x, y) dS(y) + \int_U f(y) G(x, y) dy \quad (x \in U).$$

Here we have a formula for the solution of the boundary-value problem (29), provided we can construct Green's function G for the given domain U . This is in general a difficult matter and can be done only when U has simple geometry. Subsequent subsections identify some special cases for which an explicit calculation of G is possible.

Interpreting Green's function. Fix $x \in U$. Then regarding G as a function of y , we may symbolically write

$$\begin{cases} -\Delta G = \delta_x & \text{in } U \\ G = 0 & \text{on } \partial U, \end{cases}$$

δ_x denoting the Dirac measure giving unit mass to the point x .

Before moving on to specific examples, let us record the general assertion that G is symmetric in the variables x and y :

THEOREM 13 (Symmetry of Green's function). *For all $x, y \in U$, $x \neq y$, we have*

$$G(y, x) = G(x, y).$$

Proof. Fix $x, y \in U$, $x \neq y$. Write

$$v(z) := G(x, z), \quad w(z) := G(y, z) \quad (z \in U).$$

Then $\Delta v(z) = 0$ ($z \neq x$), $\Delta w(z) = 0$ ($z \neq y$) and $w = v = 0$ on ∂U . Thus our applying Green's identity on $V := U - [B(x, \varepsilon) \cup B(y, \varepsilon)]$ for sufficiently small $\varepsilon > 0$ yields

$$(31) \quad \int_{\partial B(x, \varepsilon)} \frac{\partial v}{\partial \nu} w - \frac{\partial w}{\partial \nu} v dS(z) = \int_{\partial B(y, \varepsilon)} \frac{\partial w}{\partial \nu} v - \frac{\partial v}{\partial \nu} w dS(z),$$

ν denoting the inward pointing unit vector field on $\partial B(x, \varepsilon) \cup \partial B(y, \varepsilon)$. Now w is smooth near x , whence

$$\left| \int_{\partial B(x, \varepsilon)} \frac{\partial w}{\partial \nu} v dS \right| \leq C \varepsilon^{n-1} \sup_{\partial B(x, \varepsilon)} |v| = o(1) \quad \text{as } \varepsilon \rightarrow 0.$$

On the other hand, $v(z) = \Phi(z - x) - \phi^x(z)$, where ϕ^x is smooth in U . Thus

$$\lim_{\varepsilon \rightarrow 0} \int_{\partial B(x, \varepsilon)} \frac{\partial v}{\partial \nu} w \, dS = \lim_{\varepsilon \rightarrow 0} \int_{\partial B(x, \varepsilon)} \frac{\partial \Phi}{\partial \nu} (x - z) w(z) \, dS = w(x),$$

by calculations as in the proof of Theorem 1. Thus the left-hand side of (31) converges to $w(x)$ as $\varepsilon \rightarrow 0$. Likewise the right-hand side converges to $v(y)$. Consequently

$$G(y, x) = w(x) = v(y) = G(x, y). \quad \square$$

b. Green's function for a half-space. In this and the next subsection we will build Green's functions for two regions with simple geometry, namely the half-space \mathbb{R}_+^n and the unit ball $B(0, 1)$. Everything depends upon our explicitly solving the corrector problem (26) in these regions, and this in turn depends upon some clever geometric reflection tricks.

First let us consider the half-space

$$\mathbb{R}_+^n = \{x = (x_1, \dots, x_n) \in \mathbb{R}^n \mid x_n > 0\}.$$

Although this region is unbounded, and so the calculations in the previous section do not directly apply, we will attempt nevertheless to build Green's function using the ideas developed earlier. Later of course we must check directly that the corresponding representation formula is valid.

DEFINITION. If $x = (x_1, \dots, x_{n-1}, x_n) \in \mathbb{R}_+^n$, its reflection in the plane $\partial\mathbb{R}_+^n$ is the point

$$\tilde{x} = (x_1, \dots, x_{n-1}, -x_n).$$

We will solve problem (26) for the half-space by setting

$$\phi^x(y) := \Phi(y - \tilde{x}) = \Phi(y_1 - x_1, \dots, y_{n-1} - x_{n-1}, y_n + x_n) \quad (x, y \in \mathbb{R}_+^n).$$

The idea is that the corrector ϕ^x is built from Φ by “reflecting the singularity” from $x \in \mathbb{R}_+^n$ to $\tilde{x} \notin \mathbb{R}_+^n$. We note

$$\phi^x(y) = \Phi(y - x) \quad \text{if } y \in \partial\mathbb{R}_+^n,$$

and thus

$$\begin{cases} \Delta\phi^x = 0 & \text{in } \mathbb{R}_+^n \\ \phi^x = \Phi(y - x) & \text{on } \partial\mathbb{R}_+^n, \end{cases}$$

as required.

DEFINITION. Green's function for the half-space \mathbb{R}_+^n is

$$G(x, y) := \Phi(y - x) - \Phi(y - \tilde{x}) \quad (x, y \in \mathbb{R}_+^n, x \neq y).$$

Then

$$\begin{aligned} G_{y_n}(x, y) &= \Phi_{y_n}(y - x) - \Phi_{y_n}(y - \tilde{x}) \\ &= \frac{-1}{n\alpha(n)} \left[\frac{y_n - x_n}{|y - x|^n} - \frac{y_n + x_n}{|y - \tilde{x}|^n} \right]. \end{aligned}$$

Consequently if $y \in \partial\mathbb{R}_+^n$,

$$\frac{\partial G}{\partial \nu}(x, y) = -G_{y_n}(x, y) = -\frac{2x_n}{n\alpha(n)} \frac{1}{|x - y|^n}.$$

Suppose now u solves the boundary-value problem

$$(32) \quad \begin{cases} \Delta u = 0 & \text{in } \mathbb{R}_+^n \\ u = g & \text{on } \partial\mathbb{R}_+^n. \end{cases}$$

Then from (30) we expect

$$(33) \quad u(x) = \frac{2x_n}{n\alpha(n)} \int_{\partial\mathbb{R}_+^n} \frac{g(y)}{|x - y|^n} dy \quad (x \in \mathbb{R}_+^n)$$

to be a representation formula for our solution. The function

$$K(x, y) := \frac{2x_n}{n\alpha(n)} \frac{1}{|x - y|^n} \quad (x \in \mathbb{R}_+^n, y \in \partial\mathbb{R}_+^n)$$

is *Poisson's kernel* for \mathbb{R}_+^n , and (33) is *Poisson's formula*.

We must now check directly that formula (33) does indeed provide us with a solution of the boundary-value problem (32).

THEOREM 14 (Poisson's formula for half-space). *Assume $g \in C(\mathbb{R}^{n-1}) \cap L^\infty(\mathbb{R}^{n-1})$, and define u by (33). Then*

$$(i) \quad u \in C^\infty(\mathbb{R}_+^n) \cap L^\infty(\mathbb{R}_+^n),$$

$$(ii) \quad \Delta u = 0 \quad \text{in } \mathbb{R}_+^n,$$

and

$$(iii) \quad \lim_{\substack{x \rightarrow x^0 \\ x \in \mathbb{R}_+^n}} u(x) = g(x^0) \quad \text{for each point } x^0 \in \partial\mathbb{R}_+^n.$$

Proof. 1. For each fixed x , the mapping $y \mapsto G(x, y)$ is harmonic, except for $y = x$. As $G(x, y) = G(y, x)$, $x \mapsto G(x, y)$ is harmonic, except for $x = y$. Thus $x \mapsto -\frac{\partial G}{\partial y_n}(x, y) = K(x, y)$ is harmonic for $x \in \mathbb{R}_+^n$, $y \in \partial\mathbb{R}_+^n$.

2. A direct calculation, the details of which we omit, verifies

$$(34) \quad 1 = \int_{\partial\mathbb{R}_+^n} K(x, y) dy$$

for each $x \in \mathbb{R}_+^n$. As g is bounded, u defined by (33) is likewise bounded. Since $x \mapsto K(x, y)$ is smooth for $x \neq y$, we easily verify as well $u \in C^\infty(\mathbb{R}_+^n)$, with

$$\Delta u(x) = \int_{\partial\mathbb{R}_+^n} \Delta_x K(x, y) g(y) dy = 0 \quad (x \in \mathbb{R}_+^n).$$

3. Now fix $x^0 \in \partial\mathbb{R}_+^n$, $\varepsilon > 0$. Choose $\delta > 0$ so small that

$$(35) \quad |g(y) - g(x^0)| < \varepsilon \quad \text{if } |y - x^0| < \delta, y \in \partial\mathbb{R}_+^n.$$

Then if $|x - x^0| < \frac{\delta}{2}$, $x \in \mathbb{R}_+^n$,

$$(36) \quad \begin{aligned} |u(x) - g(x^0)| &= \left| \int_{\partial\mathbb{R}_+^n} K(x, y) [g(y) - g(x^0)] dy \right| \\ &\leq \int_{\partial\mathbb{R}_+^n \cap B(x^0, \delta)} K(x, y) |g(y) - g(x^0)| dy \\ &\quad + \int_{\partial\mathbb{R}_+^n - B(x^0, \delta)} K(x, y) |g(y) - g(x^0)| dy \\ &=: I + J. \end{aligned}$$

Now (34), (35) imply

$$I \leq \varepsilon \int_{\partial\mathbb{R}_+^n} K(x, y) dy = \varepsilon.$$

Furthermore if $|x - x^0| \leq \frac{\delta}{2}$ and $|y - x^0| \geq \delta$, we have

$$|y - x^0| \leq |y - x| + \frac{\delta}{2} \leq |y - x| + \frac{1}{2}|y - x^0|;$$

and so $|y - x| \geq \frac{1}{2}|y - x^0|$. Thus

$$\begin{aligned} J &\leq 2\|g\|_{L^\infty} \int_{\partial\mathbb{R}_+^n - B(x^0, \delta)} K(x, y) dy \\ &\leq \frac{2^{n+2}\|g\|_{L^\infty} x_n}{n\alpha(n)} \int_{\partial\mathbb{R}_+^n - B(x^0, \delta)} |y - x^0|^{-n} dy \\ &\rightarrow 0 \quad \text{as } x_n \rightarrow 0^+. \end{aligned}$$

Combining this calculation with estimate (36), we deduce $|u(x) - g(x^0)| \leq 2\varepsilon$, provided $|x - x^0|$ is sufficiently small. \square

c. Green's function for a ball. To construct Green's function for the unit ball $B(0, 1)$, we will again employ a kind of reflection, this time through the sphere $\partial B(0, 1)$.

DEFINITION. If $x \in \mathbb{R}^n - \{0\}$, the point

$$\tilde{x} = \frac{x}{|x|^2}$$

is called the point dual to x with respect to $\partial B(0, 1)$. The mapping $x \mapsto \tilde{x}$ is inversion through the unit sphere $\partial B(0, 1)$.

We now employ inversion through the sphere to compute Green's function for the unit ball $U = B^0(0, 1)$. Fix $x \in B^0(0, 1)$. Remember that we must find a corrector function $\phi^x = \phi^x(y)$ solving

$$(37) \quad \begin{cases} \Delta \phi^x = 0 & \text{in } B^0(0, 1) \\ \phi^x = \Phi(y - x) & \text{on } \partial B(0, 1); \end{cases}$$

then Green's function will be

$$(38) \quad G(x, y) = \Phi(y - x) - \phi^x(y).$$

The idea now is to "invert the singularity" from $x \in B^0(0, 1)$ to $\tilde{x} \notin B(0, 1)$. Assume for the moment $n \geq 3$. Now the mapping $y \mapsto \Phi(y - \tilde{x})$ is harmonic for $y \neq \tilde{x}$. Thus $y \mapsto |x|^{2-n} \Phi(y - \tilde{x})$ is harmonic for $y \neq \tilde{x}$, and so

$$(39) \quad \phi^x(y) := \Phi(|x|(y - \tilde{x}))$$

is harmonic in U . Furthermore, if $y \in \partial B(0, 1)$ and $x \neq 0$,

$$\begin{aligned} |x|^2 |y - \tilde{x}|^2 &= |x|^2 \left(|y|^2 - \frac{2y \cdot x}{|x|^2} + \frac{1}{|x|^2} \right) \\ &= |x|^2 - 2y \cdot x + 1 = |x - y|^2. \end{aligned}$$

Thus $(|x||y - \tilde{x}|)^{-(n-2)} = |x - y|^{-(n-2)}$. Consequently

$$(40) \quad \phi^x(y) = \Phi(y - x) \quad (y \in \partial B(0, 1)),$$

as required.

DEFINITION. Green's function for the unit ball is

$$(41) \quad G(x, y) := \Phi(y - x) - \Phi(|x|(y - \tilde{x})) \quad (x, y \in B(0, 1), x \neq y).$$

The same formula is valid for $n = 2$ as well.

Assume now u solves the boundary-value problem

$$(42) \quad \begin{cases} \Delta u = 0 & \text{in } B^0(0, 1) \\ u = g & \text{in } \partial B(0, 1). \end{cases}$$

Then using (30), we see

$$(43) \quad u(x) = - \int_{\partial B(0,1)} g(y) \frac{\partial G}{\partial \nu}(x, y) dS(y).$$

According to formula (41),

$$G_{y_i}(x, y) = \Phi_{y_i}(y - x) - \Phi(|x|(y - \tilde{x}))_{y_i}.$$

But

$$\Phi_{y_i}(y - x) = \frac{1}{n\alpha(n)} \frac{x_i - y_i}{|x - y|^n},$$

and furthermore

$$\Phi(|x|(y - \tilde{x}))_{y_i} = \frac{-1}{n\alpha(n)} \frac{y_i|x|^2 - x_i}{(|x||y - \tilde{x}|)^n} = -\frac{1}{n\alpha(n)} \frac{y_i|x|^2 - x_i}{|x - y|^n}$$

if $y \in \partial B(0, 1)$. Accordingly

$$\begin{aligned} \frac{\partial G}{\partial \nu}(x, y) &= \sum_{i=1}^n y_i G_{y_i}(x, y) \\ &= \frac{-1}{n\alpha(n)} \frac{1}{|x - y|^n} \sum_{i=1}^n y_i ((y_i - x_i) - y_i|x|^2 + x_i) \\ &= \frac{-1}{n\alpha(n)} \frac{1 - |x|^2}{|x - y|^n}. \end{aligned}$$

Hence formula (43) yields the representation formula

$$u(x) = \frac{1 - |x|^2}{n\alpha(n)} \int_{\partial B(0,1)} \frac{g(y)}{|x - y|^n} dS(y).$$

Suppose now instead of (42) u solves the boundary-value problem

$$(44) \quad \begin{cases} \Delta u = 0 & \text{in } B^0(0, r) \\ u = g & \text{on } \partial B(0, r) \end{cases}$$

for $r > 0$. Then $\tilde{u}(x) = u(rx)$ solves (42), with $\tilde{g}(x) = g(rx)$ replacing g . We change variables to obtain *Poisson's formula*

$$(45) \quad u(x) = \frac{r^2 - |x|^2}{n\alpha(n)r} \int_{\partial B(0,r)} \frac{g(y)}{|x-y|^n} dS(y) \quad (x \in B^0(0,r)).$$

The function

$$K(x, y) := \frac{r^2 - |x|^2}{n\alpha(n)r} \frac{1}{|x-y|^n} \quad (x \in B^0(0,r), y \in \partial B(0,r))$$

is *Poisson's kernel* for the ball $B(0,r)$.

We have established (45) under the assumption that a smooth solution of (44) exists. We next assert that this formula in fact gives a solution:

THEOREM 15 (Poisson's formula for ball). *Assume $g \in C(\partial B(0,r))$ and define u by (45). Then*

$$(i) \quad u \in C^\infty(B^0(0,r)),$$

$$(ii) \quad \Delta u = 0 \quad \text{in } B^0(0,r),$$

and

$$(iii) \quad \lim_{\substack{x \rightarrow x^0 \\ x \in B^0(0,r)}} u(x) = g(x^0) \quad \text{for each point } x^0 \in \partial B(0,r).$$

The proof is similar to that for Theorem 14 and is left as an exercise.

2.2.5. Energy methods.

Most of our analysis of harmonic functions thus far has depended upon fairly explicit representation formulas entailing the fundamental solution, Green's functions, etc. In this concluding subsection we illustrate some "energy" methods, which is to say techniques involving the L^2 -norms of various expressions. These ideas foreshadow later theoretical developments in Parts II and III.

a. Uniqueness. Consider first the boundary-value problem

$$(46) \quad \begin{cases} -\Delta u = f & \text{in } U \\ u = g & \text{on } \partial U. \end{cases}$$

We have already employed the maximum principle in §2.2.3 to show uniqueness, but now we set forth a simple alternative proof. Assume U is open, bounded, and ∂U is C^1 .

THEOREM 16 (Uniqueness). *There exists at most one solution $u \in C^2(\bar{U})$ of (46).*

Proof. Assume \tilde{u} is another solution and set $w := u - \tilde{u}$. Then $\Delta w = 0$ in U , and so an integration by parts shows

$$0 = - \int_U w \Delta w \, dx = \int_U |Dw|^2 \, dx.$$

Thus $Dw \equiv 0$ in U , and, since $w = 0$ on ∂U , we deduce $w = u - \tilde{u} \equiv 0$ in U . \square

b. Dirichlet's principle. Next let us demonstrate that a solution of the boundary-value problem (46) for Poisson's equation can be characterized as the minimizer of an appropriate functional. For this, we define the *energy functional*

$$I[w] := \int_U \frac{1}{2} |Dw|^2 - wf \, dx,$$

w belonging to the *admissible set*

$$\mathcal{A} := \{w \in C^2(\bar{U}) \mid w = g \text{ on } \partial U\}.$$

THEOREM 17 (Dirichlet's principle). *Assume $u \in C^2(\bar{U})$ solves (46). Then*

$$(47) \quad I[u] = \min_{w \in \mathcal{A}} I[w].$$

Conversely, if $u \in \mathcal{A}$ satisfies (47), then u solves the boundary-value problem (46).

In other words if $u \in \mathcal{A}$, the PDE $-\Delta u = f$ is equivalent to the statement that u minimizes the energy $I[\cdot]$.

Proof. 1. Choose $w \in \mathcal{A}$. Then (46) implies

$$0 = \int_U (-\Delta u - f)(u - w) \, dx.$$

An integration by parts yields

$$0 = \int_U Du \cdot D(u - w) - f(u - w) \, dx,$$

and there is no boundary term since $u - w = g - g \equiv 0$ on ∂U . Hence

$$\begin{aligned} \int_U |Du|^2 - uf \, dx &= \int_U Du \cdot Dw - wf \, dx \\ &\leq \int_U \frac{1}{2} |Du|^2 \, dx + \int_U \frac{1}{2} |Dw|^2 - wf \, dx, \end{aligned}$$

where we employed the estimates

$$|Du \cdot Dw| \leq |Du| |Dw| \leq \frac{1}{2} |Du|^2 + \frac{1}{2} |Dw|^2,$$

following from the Cauchy–Schwarz and Cauchy inequalities (§B.2). Rearranging, we conclude

$$(48) \quad I[u] \leq I[w] \quad (w \in \mathcal{A}).$$

Since $u \in \mathcal{A}$, (47) follows from (48).

2. Now, conversely, suppose (47) holds. Fix any $v \in C_c^\infty(U)$ and write

$$i(\tau) := I[u + \tau v] \quad (\tau \in \mathbb{R}).$$

Since $u + \tau v \in \mathcal{A}$ for each τ , the scalar function $i(\cdot)$ has a minimum at zero, and thus

$$i'(0) = 0 \quad \left(' = \frac{d}{d\tau} \right),$$

provided this derivative exists. But

$$\begin{aligned} i(\tau) &= \int_U \frac{1}{2} |Du + \tau Dv|^2 - (u + \tau v) f \, dx \\ &= \int_U \frac{1}{2} |Du|^2 + \tau Du \cdot Dv + \frac{\tau^2}{2} |Dv|^2 - (u + \tau v) f \, dx. \end{aligned}$$

Consequently

$$0 = i'(0) = \int_U Du \cdot Dv - vf \, dx = \int_U (-\Delta u - f)v \, dx.$$

This identity is valid for each function $v \in C_c^\infty(U)$ and so $-\Delta u = f$ in U . \square

Dirichlet's principle is an instance of the *calculus of variations* applied to Laplace's equation. See Chapter 8 for more.

2.3. HEAT EQUATION

Next we study the *heat equation*

$$(1) \quad u_t - \Delta u = 0$$

and the *nonhomogeneous heat equation*

$$(2) \quad u_t - \Delta u = f,$$

subject to appropriate initial and boundary conditions. Here $t > 0$ and $x \in U$, where $U \subset \mathbb{R}^n$ is open. The unknown is $u : \bar{U} \times [0, \infty) \rightarrow \mathbb{R}$, $u = u(x, t)$, and the Laplacian Δ is taken with respect to the spatial variables $x = (x_1, \dots, x_n)$: $\Delta u = \Delta_x u = \sum_{i=1}^n u_{x_i x_i}$. In (2) the function $f : U \times [0, \infty) \rightarrow \mathbb{R}$ is given.

A guiding principle is that any assertion about harmonic functions yields an analogous (but more complicated) statement about solutions of the heat equation. Accordingly our development will largely parallel the corresponding theory for Laplace's equation.

Physical interpretation. The heat equation, also known as the *diffusion equation*, describes in typical applications the evolution in time of the density u of some quantity such as heat, chemical concentration, etc. If $V \subset U$ is any smooth subregion, the rate of change of the total quantity within V equals the negative of the net flux through ∂V :

$$\frac{d}{dt} \int_V u \, dx = - \int_{\partial V} \mathbf{F} \cdot \boldsymbol{\nu} \, dS,$$

\mathbf{F} being the flux density. Thus

$$(3) \quad u_t = - \operatorname{div} \mathbf{F},$$

as V was arbitrary. In many situations \mathbf{F} is proportional to the gradient of u but points in the opposite direction (since the flow is from regions of higher to lower concentration):

$$\mathbf{F} = -aDu \quad (a > 0).$$

Substituting into (3), we obtain the PDE

$$u_t = a \operatorname{div}(Du) = a\Delta u,$$

which for $a = 1$ is the heat equation.

The heat equation appears as well in the study of Brownian motion.

2.3.1. Fundamental solution.

a. Derivation of the fundamental solution. As noted in §2.2.1 an important first step in studying any PDE is often to come up with some specific solutions.

We observe that the heat equation involves one derivative with respect to the time variable t , but two derivatives with respect to the space variables x_i ($i = 1, \dots, n$). Consequently we see that if u solves (1), then so does $u(\lambda x, \lambda^2 t)$ for $\lambda \in \mathbb{R}$. This scaling indicates the ratio $\frac{r^2}{t}$ ($r = |x|$) is important for the heat equation and suggests that we search for a solution of (1) having the form $u(x, t) = v\left(\frac{x}{t}\right) = v\left(\frac{|x|^2}{t}\right)$ ($t > 0$, $x \in \mathbb{R}^n$), for some function v as yet undetermined.

Although this approach eventually leads to what we want (see Problem 13), it is quicker to seek a solution u having the special structure

$$(4) \quad u(x, t) = \frac{1}{t^\alpha} v\left(\frac{x}{t^\beta}\right) \quad (x \in \mathbb{R}^n, t > 0),$$

where the constants α, β and the function $v : \mathbb{R}^n \rightarrow \mathbb{R}$ must be found. We come to (4) if we look for a solution u of the heat equation invariant under the *dilation scaling*

$$u(x, t) \mapsto \lambda^\alpha u(\lambda^\beta x, \lambda t).$$

That is, we ask that

$$u(x, t) = \lambda^\alpha u(\lambda^\beta x, \lambda t)$$

for all $\lambda > 0$, $x \in \mathbb{R}^n$, $t > 0$. Setting $\lambda = t^{-1}$, we derive (4) for $v(y) := u(y, 1)$.

Let us insert (4) into (1) and thereafter compute

$$(5) \quad \alpha t^{-(\alpha+1)} v(y) + \beta t^{-(\alpha+1)} y \cdot Dv(y) + t^{-(\alpha+2\beta)} \Delta v(y) = 0$$

for $y := t^{-\beta} x$. In order to transform (5) into an expression involving the variable y alone, we take $\beta = \frac{1}{2}$. Then the terms with t are identical, and so (5) reduces to

$$(6) \quad \alpha v + \frac{1}{2} y \cdot Dv + \Delta v = 0.$$

We simplify further by guessing v to be radial; that is, $v(y) = w(|y|)$ for some $w : \mathbb{R} \rightarrow \mathbb{R}$. Thereupon (6) becomes

$$\alpha w + \frac{1}{2} r w' + w'' + \frac{n-1}{r} w' = 0,$$

for $r = |y|$, $' = \frac{d}{dr}$. Now if we set $\alpha = \frac{n}{2}$, this simplifies to read

$$(r^{n-1}w')' + \frac{1}{2}(r^n w)' = 0.$$

Thus

$$r^{n-1}w' + \frac{1}{2}r^n w = a$$

for some constant a . Assuming $\lim_{r \rightarrow \infty} w, w' = 0$, we conclude $a = 0$, whence

$$w' = -\frac{1}{2}rw.$$

But then for some constant b

$$(7) \quad w = be^{-\frac{r^2}{4}}.$$

Combining (4), (7) and our choices for α, β , we conclude that $\frac{b}{t^{n/2}}e^{-\frac{|x|^2}{4t}}$ solves the heat equation (1).

This computation motivates the following

DEFINITION. *The function*

$$\Phi(x, t) := \begin{cases} \frac{1}{(4\pi t)^{n/2}} e^{-\frac{|x|^2}{4t}} & (x \in \mathbb{R}^n, t > 0) \\ 0 & (x \in \mathbb{R}^n, t < 0) \end{cases}$$

is called the fundamental solution of the heat equation.

Notice that Φ is singular at the point $(0, 0)$. We will sometimes write $\Phi(x, t) = \Phi(|x|, t)$ to emphasize that the fundamental solution is radial in the variable x . The choice of the normalizing constant $(4\pi)^{-n/2}$ is dictated by the following

LEMMA (Integral of fundamental solution). *For each time $t > 0$,*

$$\int_{\mathbb{R}^n} \Phi(x, t) dx = 1.$$

Proof. We calculate

$$\begin{aligned} \int_{\mathbb{R}^n} \Phi(x, t) dx &= \frac{1}{(4\pi t)^{n/2}} \int_{\mathbb{R}^n} e^{-\frac{|x|^2}{4t}} dx \\ &= \frac{1}{\pi^{n/2}} \int_{\mathbb{R}^n} e^{-|z|^2} dz \\ &= \frac{1}{\pi^{n/2}} \prod_{i=1}^n \int_{-\infty}^{\infty} e^{-z_i^2} dz_i = 1. \end{aligned}$$

□

A different derivation of the fundamental solution of the heat equation appears in §4.3.1.

b. Initial-value problem. We now employ Φ to fashion a solution to the *initial-value* (or *Cauchy*) *problem*

$$(8) \quad \begin{cases} u_t - \Delta u = 0 & \text{in } \mathbb{R}^n \times (0, \infty) \\ u = g & \text{on } \mathbb{R}^n \times \{t = 0\}. \end{cases}$$

Let us note that the function $(x, t) \mapsto \Phi(x, t)$ solves the heat equation away from the singularity at $(0, 0)$, and thus so does $(x, t) \mapsto \Phi(x - y, t)$ for each fixed $y \in \mathbb{R}^n$. Consequently the convolution

$$(9) \quad \begin{aligned} u(x, t) &= \int_{\mathbb{R}^n} \Phi(x - y, t) g(y) dy \\ &= \frac{1}{(4\pi t)^{n/2}} \int_{\mathbb{R}^n} e^{-\frac{|x-y|^2}{4t}} g(y) dy \quad (x \in \mathbb{R}^n, t > 0) \end{aligned}$$

should also be a solution.

THEOREM 1 (Solution of initial-value problem). *Assume $g \in C(\mathbb{R}^n) \cap L^\infty(\mathbb{R}^n)$, and define u by (9). Then*

- (i) $u \in C^\infty(\mathbb{R}^n \times (0, \infty))$,
- (ii) $u_t(x, t) - \Delta u(x, t) = 0 \quad (x \in \mathbb{R}^n, t > 0)$,

and

- (iii) $\lim_{\substack{(x,t) \rightarrow (x^0,0) \\ x \in \mathbb{R}^n, t > 0}} u(x, t) = g(x^0)$ for each point $x^0 \in \mathbb{R}^n$.

Proof. 1. Since the function $\frac{1}{t^{n/2}} e^{-\frac{|x|^2}{4t}}$ is infinitely differentiable, with uniformly bounded derivatives of all orders, on $\mathbb{R}^n \times [\delta, \infty)$ for each $\delta > 0$, we see that $u \in C^\infty(\mathbb{R}^n \times (0, \infty))$. Furthermore

$$(10) \quad \begin{aligned} u_t(x, t) - \Delta u(x, t) &= \int_{\mathbb{R}^n} [(\Phi_t - \Delta_x \Phi)(x - y, t)] g(y) dy \\ &= 0 \quad (x \in \mathbb{R}^n, t > 0), \end{aligned}$$

since Φ itself solves the heat equation.

2. Fix $x^0 \in \mathbb{R}^n$, $\varepsilon > 0$. Choose $\delta > 0$ such that

$$(11) \quad |g(y) - g(x^0)| < \varepsilon \quad \text{if } |y - x^0| < \delta, \quad y \in \mathbb{R}^n.$$

Then if $|x - x^0| < \frac{\delta}{2}$, we have, according to the lemma,

$$\begin{aligned} |u(x, t) - g(x^0)| &= \left| \int_{\mathbb{R}^n} \Phi(x - y, t) [g(y) - g(x^0)] dy \right| \\ &\leq \int_{B(x^0, \delta)} \Phi(x - y, t) |g(y) - g(x^0)| dy \\ &\quad + \int_{\mathbb{R}^n - B(x^0, \delta)} \Phi(x - y, t) |g(y) - g(x^0)| dy \\ &=: I + J. \end{aligned}$$

Now

$$I \leq \varepsilon \int_{\mathbb{R}^n} \Phi(x - y, t) dy = \varepsilon,$$

owing to (11) and the lemma. Furthermore, if $|x - x^0| \leq \frac{\delta}{2}$ and $|y - x^0| \geq \delta$, then

$$|y - x^0| \leq |y - x| + \frac{\delta}{2} \leq |y - x| + \frac{1}{2}|y - x^0|.$$

Thus $|y - x| \geq \frac{1}{2}|y - x^0|$. Consequently

$$\begin{aligned} J &\leq 2\|g\|_{L^\infty} \int_{\mathbb{R}^n - B(x^0, \delta)} \Phi(x - y, t) dy \\ &\leq \frac{C}{t^{n/2}} \int_{\mathbb{R}^n - B(x^0, \delta)} e^{-\frac{|x-y|^2}{4t}} dy \\ &\leq \frac{C}{t^{n/2}} \int_{\mathbb{R}^n - B(x^0, \delta)} e^{-\frac{|y-x^0|^2}{16t}} dy \\ &= C \int_{\mathbb{R}^n - B(x^0, \delta/\sqrt{t})} e^{-\frac{|z|^2}{16}} dz \rightarrow 0 \quad \text{as } t \rightarrow 0^+. \end{aligned}$$

Hence if $|x - x^0| < \frac{\delta}{2}$ and $t > 0$ is small enough, $|u(x, t) - g(x^0)| < 2\varepsilon$. \square

Interpretation of fundamental solution. In view of Theorem 1 we sometimes write

$$\begin{cases} \Phi_t - \Delta\Phi = 0 & \text{in } \mathbb{R}^n \times (0, \infty) \\ \Phi = \delta_0 & \text{on } \mathbb{R}^n \times \{t = 0\}, \end{cases}$$

δ_0 denoting the Dirac measure on \mathbb{R}^n giving unit mass to the point 0.

Infinite propagation speed. Notice that if g is bounded, continuous, $g \geq 0$, $g \not\equiv 0$, then

$$u(x, t) = \frac{1}{(4\pi t)^{n/2}} \int_{\mathbb{R}^n} e^{-\frac{|x-y|^2}{4t}} g(y) dy$$

is in fact positive for *all* points $x \in \mathbb{R}^n$ and times $t > 0$. We interpret this observation by saying the heat equation forces *infinite propagation speed* for disturbances. If the initial temperature is nonnegative and is positive somewhere, the temperature at any later time (no matter how small) is everywhere positive. (We will learn in §2.4.3 that the wave equation in contrast supports finite propagation speed for disturbances.)

c. Nonhomogeneous problem. Now let us turn our attention to the *nonhomogeneous* initial-value problem

$$(12) \quad \begin{cases} u_t - \Delta u = f & \text{in } \mathbb{R}^n \times (0, \infty) \\ u = 0 & \text{on } \mathbb{R}^n \times \{t = 0\}. \end{cases}$$

How can we produce a formula for the solution? If we recall the motivation leading up to (9), we should note further that the mapping $(x, t) \mapsto \Phi(x - y, t - s)$ is a solution of the heat equation (for given $y \in \mathbb{R}^n$, $0 < s < t$). Now for fixed s , the function

$$u = u(x, t; s) = \int_{\mathbb{R}^n} \Phi(x - y, t - s) f(y, s) dy$$

solves

$$(12_s) \quad \begin{cases} u_t(\cdot; s) - \Delta u(\cdot; s) = 0 & \text{in } \mathbb{R}^n \times (s, \infty) \\ u(\cdot; s) = f(\cdot, s) & \text{on } \mathbb{R}^n \times \{t = s\}, \end{cases}$$

which is just an initial-value problem of the form (8), with the starting time $t = 0$ replaced by $t = s$ and g replaced by $f(\cdot, s)$. Thus $u(\cdot; s)$ is certainly not a solution of (12).

However *Duhamel's principle** asserts that we can build a solution of (12) out of the solutions of (12_s) , by integrating with respect to s . The idea is to consider

$$u(x, t) = \int_0^t u(x, t; s) ds \quad (x \in \mathbb{R}^n, t \geq 0).$$

Rewriting, we have

$$(13) \quad \begin{aligned} u(x, t) &= \int_0^t \int_{\mathbb{R}^n} \Phi(x - y, t - s) f(y, s) dy ds \\ &= \int_0^t \frac{1}{(4\pi(t - s))^{n/2}} \int_{\mathbb{R}^n} e^{-\frac{|x-y|^2}{4(t-s)}} f(y, s) dy ds, \end{aligned}$$

for $x \in \mathbb{R}^n$, $t > 0$.

To confirm that formula (13) works, let us for simplicity assume $f \in C_1^2(\mathbb{R}^n \times [0, \infty))$ and f has compact support.

*Duhamel's principle has wide applicability to linear ODE and PDE and does not depend on the specific structure of the heat equation. It yields, for example, the solution of the nonhomogeneous transport equation, obtained by different means in §2.1.2. We will invoke Duhamel's principle for the wave equation in §2.4.2.

THEOREM 2 (Solution of nonhomogeneous problem). *Define u by (13).*

Then

$$(i) \quad u \in C_1^2(\mathbb{R}^n \times (0, \infty)),$$

$$(ii) \quad u_t(x, t) - \Delta u(x, t) = f(x, t) \quad (x \in \mathbb{R}^n, t > 0),$$

and

$$(iii) \quad \lim_{\substack{(x,t) \rightarrow (x^0,0) \\ x \in \mathbb{R}^n, t > 0}} u(x, t) = 0 \quad \text{for each point } x^0 \in \mathbb{R}^n.$$

Proof. 1. Since Φ has a singularity at $(0, 0)$, we cannot directly justify differentiating under the integral sign. We instead proceed somewhat as in the proof of Theorem 1 in §2.2.1.

First we change variables, to write

$$u(x, t) = \int_0^t \int_{\mathbb{R}^n} \Phi(y, s) f(x - y, t - s) dy ds.$$

As $f \in C_1^2(\mathbb{R}^n \times [0, \infty))$ has compact support and $\Phi = \Phi(y, s)$ is smooth near $s = t > 0$, we compute

$$\begin{aligned} u_t(x, t) &= \int_0^t \int_{\mathbb{R}^n} \Phi(y, s) f_t(x - y, t - s) dy ds \\ &\quad + \int_{\mathbb{R}^n} \Phi(y, t) f(x - y, 0) dy \end{aligned}$$

and

$$u_{x_i x_j}(x, t) = \int_0^t \int_{\mathbb{R}^n} \Phi(y, s) f_{x_i x_j}(x - y, t - s) dy ds \quad (i, j = 1, \dots, n).$$

Thus $u_t, D_x^2 u$, and likewise $u, D_x u$, belong to $C(\mathbb{R}^n \times (0, \infty))$.

2. We then calculate

$$\begin{aligned} (14) \quad u_t(x, t) - \Delta u(x, t) &= \int_0^t \int_{\mathbb{R}^n} \Phi(y, s) \left[\left(\frac{\partial}{\partial t} - \Delta_x \right) f(x - y, t - s) \right] dy ds \\ &\quad + \int_{\mathbb{R}^n} \Phi(y, t) f(x - y, 0) dy \\ &= \int_\epsilon^t \int_{\mathbb{R}^n} \Phi(y, s) \left[\left(-\frac{\partial}{\partial s} - \Delta_y \right) f(x - y, t - s) \right] dy ds \\ &\quad + \int_0^\epsilon \int_{\mathbb{R}^n} \Phi(y, s) \left[\left(-\frac{\partial}{\partial s} - \Delta_y \right) f(x - y, t - s) \right] dy ds \\ &\quad + \int_{\mathbb{R}^n} \Phi(y, t) f(x - y, 0) dy \\ &=: I_\epsilon + J_\epsilon + K. \end{aligned}$$

Now

$$(15) \quad |J_\varepsilon| \leq (\|f_t\|_{L^\infty} + \|D^2 f\|_{L^\infty}) \int_0^\varepsilon \int_{\mathbb{R}^n} \Phi(y, s) dy ds \leq \varepsilon C,$$

by the lemma. Integrating by parts, we also find

$$(16) \quad \begin{aligned} I_\varepsilon &= \int_\varepsilon^t \int_{\mathbb{R}^n} \left[\left(\frac{\partial}{\partial s} - \Delta_y \right) \Phi(y, s) \right] f(x - y, t - s) dy ds \\ &\quad + \int_{\mathbb{R}^n} \Phi(y, \varepsilon) f(x - y, t - \varepsilon) dy \\ &\quad - \int_{\mathbb{R}^n} \Phi(y, t) f(x - y, 0) dy \\ &= \int_{\mathbb{R}^n} \Phi(y, \varepsilon) f(x - y, t - \varepsilon) dy - K, \end{aligned}$$

since Φ solves the heat equation. Combining (14)–(16), we ascertain

$$\begin{aligned} u_t(x, t) - \Delta u(x, t) &= \lim_{\varepsilon \rightarrow 0} \int_{\mathbb{R}^n} \Phi(y, \varepsilon) f(x - y, t - \varepsilon) dy \\ &= f(x, t) \quad (x \in \mathbb{R}^n, t > 0), \end{aligned}$$

the limit as $\varepsilon \rightarrow 0$ being computed as in the proof of Theorem 1. Finally note $\|u(\cdot, t)\|_{L^\infty} \leq t\|f\|_{L^\infty} \rightarrow 0$. \square

Solution of homogeneous problem with general initial data. We can of course combine Theorems 1 and 2 to discover that


$$(17) \quad u(x, t) = \int_{\mathbb{R}^n} \Phi(x - y, t) g(y) dy + \int_0^t \int_{\mathbb{R}^n} \Phi(x - y, t - s) f(y, s) dy ds$$

is, under the hypotheses on g and f as above, a solution of

$$(18) \quad \begin{cases} u_t - \Delta u = f & \text{in } \mathbb{R}^n \times (0, \infty) \\ u = g & \text{on } \mathbb{R}^n \times \{t = 0\}. \end{cases}$$

2.3.2. Mean-value formula.

First we recall some useful notation from §A.2. Assume $U \subset \mathbb{R}^n$ is open and bounded, and fix a time $T > 0$.



The region U_T

DEFINITIONS.

(i) We define the parabolic cylinder

$$U_T := U \times (0, T].$$

(ii) The parabolic boundary of U_T is


$$\Gamma_T := \bar{U}_T - U_T.$$

We interpret U_T as being the *parabolic interior* of $\bar{U} \times [0, T]$: note carefully that U_T includes the top $U \times \{t = T\}$. The parabolic boundary Γ_T comprises the bottom and vertical sides of $U \times [0, T]$, but not the top.

We want next to derive a kind of analogue to the mean-value property for harmonic functions, as discussed in §2.2.2. There is no such simple formula. However let us observe that for fixed x the spheres $\partial B(x, r)$ are level sets of the fundamental solution $\Phi(x - y)$ for Laplace's equation. This suggests that perhaps for fixed (x, t) the level sets of fundamental solution $\Phi(x - y, t - s)$ for the heat equation may be relevant.

DEFINITION. For fixed $x \in \mathbb{R}^n$, $t \in \mathbb{R}$, $r > 0$, we define

$$E(x, t; r) := \left\{ (y, s) \in \mathbb{R}^{n+1} \mid s \leq t, \Phi(x - y, t - s) \geq \frac{1}{r^n} \right\}.$$



A “heat ball”

This is a region in space-time, the boundary of which is a level set of $\Phi(x-y, t-s)$. Note that the point (x, t) is at the center of the top. $E(x, t; r)$ is sometimes called a “heat ball”.

THEOREM 3 (A mean-value property for the heat equation). *Let $u \in C_1^2(U_T)$ solve the heat equation. Then*

$$(19) \quad u(x, t) = \frac{1}{4r^n} \iint_{E(x, t; r)} u(y, s) \frac{|x-y|^2}{(t-s)^2} dy ds$$

for each $E(x, t; r) \subset U_T$.

Formula (19) is a sort of analogue for the heat equation of the mean-value formulas for Laplace’s equation. Observe that the right-hand side involves only $u(y, s)$ for times $s \leq t$. This is reasonable, as the value $u(x, t)$ should not depend upon future times.

Proof. Shift the space and time coordinates so that $x = 0$ and $t = 0$. Upon mollifying if necessary, we may assume u is smooth. Write $E(r) = E(0, 0; r)$ and set

$$(20) \quad \begin{aligned} \phi(r) &:= \frac{1}{r^n} \iint_{E(r)} u(y, s) \frac{|y|^2}{s^2} dy ds \\ &= \iint_{E(1)} u(ry, r^2s) \frac{|y|^2}{s^2} dy ds. \end{aligned}$$

We compute

$$\begin{aligned} \phi'(r) &= \iint_{E(1)} \sum_{i=1}^n u_{y_i} y_i \frac{|y|^2}{s^2} + 2ru_s \frac{|y|^2}{s} dy ds \\ &= \frac{1}{r^{n+1}} \iint_{E(r)} \sum_{i=1}^n u_{y_i} y_i \frac{|y|^2}{s^2} + 2u_s \frac{|y|^2}{s} dy ds \\ &=: A + B. \end{aligned}$$

Also, let us introduce the useful function

$$(21) \quad \psi := -\frac{n}{2} \log(-4\pi s) + \frac{|y|^2}{4s} + n \log r$$

and observe $\psi = 0$ on $\partial E(r)$, since $\Phi(y, -s) = r^{-n}$ on $\partial E(r)$. We utilize (21) to write

$$\begin{aligned} B &= \frac{1}{r^{n+1}} \iint_{E(r)} 4u_s \sum_{i=1}^n y_i \psi_{y_i} dy ds \\ &= -\frac{1}{r^{n+1}} \iint_{E(r)} 4nu_s \psi + 4 \sum_{i=1}^n u_{s y_i} y_i \psi dy ds; \end{aligned}$$

there is no boundary term since $\psi = 0$ on $\partial E(r)$. Integrating by parts with respect to s , we discover

$$\begin{aligned} B &= \frac{1}{r^{n+1}} \iint_{E(r)} -4nu_s \psi + 4 \sum_{i=1}^n u_{y_i} y_i \psi_s dy ds \\ &= \frac{1}{r^{n+1}} \iint_{E(r)} -4nu_s \psi + 4 \sum_{i=1}^n u_{y_i} y_i \left(-\frac{n}{2s} - \frac{|y|^2}{4s^2} \right) dy ds \\ &= \frac{1}{r^{n+1}} \iint_{E(r)} -4nu_s \psi - \frac{2n}{s} \sum_{i=1}^n u_{y_i} y_i dy ds - A. \end{aligned}$$

Consequently, since u solves the heat equation,

$$\begin{aligned} \phi'(r) &= A + B \\ &= \frac{1}{r^{n+1}} \iint_{E(r)} -4n\Delta u \psi - \frac{2n}{s} \sum_{i=1}^n u_{y_i} y_i dy ds \\ &= \sum_{i=1}^n \frac{1}{r^{n+1}} \iint_{E(r)} 4nu_{y_i} \psi_{y_i} - \frac{2n}{s} u_{y_i} y_i dy ds \\ &= 0, \quad \text{according to (21)}. \end{aligned}$$


Thus ϕ is constant, and therefore

$$\phi(r) = \lim_{t \rightarrow 0} \phi(t) = u(0, 0) \left(\lim_{t \rightarrow 0} \frac{1}{t^n} \iint_{E(t)} \frac{|y|^2}{s^2} dy ds \right) = 4u(0, 0),$$

as

$$\frac{1}{t^n} \iint_{E(t)} \frac{|y|^2}{s^2} dy ds = \iint_{E(1)} \frac{|y|^2}{s^2} dy ds = 4.$$

We omit the details of this last computation. □



Strong maximum principle for the heat equation

2.3.3. Properties of solutions.

a. Strong maximum principle, uniqueness. First we employ the mean-value property to give a quick proof of the strong maximum principle.

THEOREM 4 (Strong maximum principle for the heat equation). *Assume $u \in C_1^2(U_T) \cap C(\bar{U}_T)$ solves the heat equation in U_T .*

(i) *Then*

$$\max_{\bar{U}_T} u = \max_{\Gamma_T} u.$$

(ii) *Furthermore, if U is connected and there exists a point $(x_0, t_0) \in U_T$ such that*

$$u(x_0, t_0) = \max_{\bar{U}_T} u,$$

then

$$u \text{ is constant in } \bar{U}_{t_0}.$$

Assertion (i) is the *maximum principle* for the heat equation and (ii) is the *strong maximum principle*. Similar assertions are valid with “min” replacing “max”.

Interpretation. *So if u attains its maximum (or minimum) at an interior point, then u is constant at all earlier times.* This accords with our strong intuitive understanding of the variable t as denoting time: the solution will

be constant on the time interval $[0, t_0]$ provided the initial and boundary conditions are constant. However, the solution may change at times $t > t_0$, provided the boundary conditions alter after t_0 . The solution will however not respond to changes in boundary conditions until these changes happen.

Take note that whereas all this is obvious on intuitive, physical grounds, such insights do not constitute a proof. The task is to *deduce* such behavior from the PDE.

Proof. 1. Suppose there exists a point $(x_0, t_0) \in U_T$ with $u(x_0, t_0) = M := \max_{\bar{U}_T} u$. Then for all sufficiently small $r > 0$, $E(x_0, t_0; r) \subset U_T$; and we employ the mean-value property to deduce

$$M = u(x_0, t_0) = \frac{1}{4r^n} \iint_{E(x_0, t_0; r)} u(y, s) \frac{|x_0 - y|^2}{(t_0 - s)^2} dy ds \leq M,$$

since

$$1 = \frac{1}{4r^n} \iint_{E(x_0, t_0; r)} \frac{|x_0 - y|^2}{(t_0 - s)^2} dy ds.$$

Equality holds only if u is identically equal to M within $E(x_0, t_0; r)$. Consequently

$$u(y, s) = M \quad \text{for all } (y, s) \in E(x_0, t_0; r).$$

Draw any line segment L in U_T connecting (x_0, t_0) with some other point $(y_0, s_0) \in U_T$, with $s_0 < t_0$. Consider

$$r_0 := \min\{s \geq s_0 \mid u(x, t) = M \text{ for all points } (x, t) \in L, s \leq t \leq t_0\}.$$

Since u is continuous, the minimum is attained. Assume $r_0 > s_0$. Then $u(z_0, r_0) = M$ for some point (z_0, r_0) on $L \cap U_T$ and so $u \equiv M$ on $E(z_0, r_0; r)$ for all sufficiently small $r > 0$. Since $E(z_0, r_0; r)$ contains $L \cap \{r_0 - \sigma \leq t \leq r_0\}$ for some small $\sigma > 0$, we have a contradiction. Thus $r_0 = s_0$, and hence $u \equiv M$ on L .

2. Now fix any point $x \in U$ and any time $0 \leq t < t_0$. There exist points $\{x_0, x_1, \dots, x_m = x\}$ such that the line segments in \mathbb{R}^n connecting x_{i-1} to x_i lie in U for $i = 1, \dots, m$. (This follows since the set of points in U which can be so connected to x_0 by a polygonal path is nonempty, open and relatively closed in U .) Select times $t_0 > t_1 > \dots > t_m = t$. Then the line segments in \mathbb{R}^{n+1} connecting (x_{i-1}, t_{i-1}) to (x_i, t_i) ($i = 1, \dots, m$) lie in U_T . According to step 1, $u \equiv M$ on each such segment and so $u(x, t) = M$. \square

Infinite propagation speed again. The strong maximum principle implies that if U is connected and $u \in C_1^2(U_T) \cap C(\bar{U}_T)$ satisfies

$$\begin{cases} u_t - \Delta u = 0 & \text{in } U_T \\ u = 0 & \text{on } \partial U \times [0, T] \\ u = g & \text{on } U \times \{t = 0\} \end{cases}$$

where $g \geq 0$, then u is positive *everywhere* within U_T if g is positive *some-where* on U . This is another illustration of infinite propagation speed for disturbances.

An important application of the maximum principle is the following uniqueness assertion.

THEOREM 5 (Uniqueness on bounded domains). *Let $g \in C(\Gamma_T)$, $f \in C(U_T)$. Then there exists at most one solution $u \in C_1^2(U_T) \cap C(\bar{U}_T)$ of the initial/boundary-value problem*

$$(22) \quad \begin{cases} u_t - \Delta u = f & \text{in } U_T \\ u = g & \text{on } \Gamma_T. \end{cases}$$

Proof. If u and \tilde{u} are two solutions of (22), apply Theorem 4 to $w := \pm(u - \tilde{u})$. □

We next extend our uniqueness assertion to the *Cauchy problem*, that is, the initial-value problem for $U = \mathbb{R}^n$. As we are no longer on a bounded region, we must introduce some control on the behavior of solutions for large $|x|$.

THEOREM 6 (Maximum principle for the Cauchy problem). *Suppose $u \in C_1^2(\mathbb{R}^n \times (0, T]) \cap C(\mathbb{R}^n \times [0, T])$ solves*

$$(23) \quad \begin{cases} u_t - \Delta u = 0 & \text{in } \mathbb{R}^n \times (0, T) \\ u = g & \text{on } \mathbb{R}^n \times \{t = 0\} \end{cases}$$

and satisfies the growth estimate

$$(24) \quad u(x, t) \leq Ae^{a|x|^2} \quad (x \in \mathbb{R}^n, 0 \leq t \leq T)$$

for constants $A, a > 0$. Then

$$\sup_{\mathbb{R}^n \times [0, T]} u = \sup_{\mathbb{R}^n} g.$$

Proof. 1. First assume

$$(25) \quad 4aT < 1,$$

in which case

$$(26) \quad 4a(T + \varepsilon) < 1$$

for some $\varepsilon > 0$. Fix $y \in \mathbb{R}^n$, $\mu > 0$, and define

$$v(x, t) := u(x, t) - \frac{\mu}{(T + \varepsilon - t)^{n/2}} e^{\frac{|x-y|^2}{4(T+\varepsilon-t)}} \quad (x \in \mathbb{R}^n, t > 0).$$

A direct calculation (cf. §2.3.1) shows

$$v_t - \Delta v = 0 \quad \text{in } \mathbb{R}^n \times (0, T].$$

Fix $r > 0$ and set $U := B^0(y, r)$, $U_T = B^0(y, r) \times (0, T]$. Then according to Theorem 4,

$$(27) \quad \max_{\bar{U}_T} v = \max_{\Gamma_T} v.$$

2. Now if $x \in \mathbb{R}^n$,

$$(28) \quad \begin{aligned} v(x, 0) &= u(x, 0) - \frac{\mu}{(T + \varepsilon)^{n/2}} e^{\frac{|x-y|^2}{4(T+\varepsilon)}} \\ &\leq u(x, 0) = g(x); \end{aligned}$$

and if $|x - y| = r$, $0 \leq t \leq T$, then

$$\begin{aligned} v(x, t) &= u(x, t) - \frac{\mu}{(T + \varepsilon - t)^{n/2}} e^{\frac{r^2}{4(T+\varepsilon-t)}} \\ &\leq Ae^{a|x|^2} - \frac{\mu}{(T + \varepsilon - t)^{n/2}} e^{\frac{r^2}{4(T+\varepsilon-t)}} \quad \text{by (24)} \\ &\leq Ae^{a(|y|+r)^2} - \frac{\mu}{(T + \varepsilon)^{n/2}} e^{\frac{r^2}{4(T+\varepsilon)}}. \end{aligned}$$

Now according to (26), $\frac{1}{4(T+\varepsilon)} = a + \gamma$ for some $\gamma > 0$. Thus we may continue the calculation above to find

$$(29) \quad v(x, t) \leq Ae^{a(|y|+r)^2} - \mu(4(a + \gamma))^{n/2} e^{(a+\gamma)r^2} \leq \sup_{\mathbb{R}^n} g,$$

for r selected sufficiently large. Thus (27)–(29) imply

$$v(y, t) \leq \sup_{\mathbb{R}^n} g$$

for all $y \in \mathbb{R}^n$, $0 \leq t \leq T$, provided (25) is valid. Let $\mu \rightarrow 0$.

3. In the general case that (25) fails, we repeatedly apply the result above on the time intervals $[0, T_1]$, $[T_1, 2T_1,]$, etc., for $T_1 = \frac{1}{8a}$. \square

THEOREM 7 (Uniqueness for Cauchy problem). *Let $g \in C(\mathbb{R}^n)$, $f \in C(\mathbb{R}^n \times [0, T])$. Then there exists at most one solution $u \in C_1^2(\mathbb{R}^n \times (0, T]) \cap C(\mathbb{R}^n \times [0, T])$ of the initial-value problem*

$$(30) \quad \begin{cases} u_t - \Delta u = f & \text{in } \mathbb{R}^n \times (0, T) \\ u = g & \text{on } \mathbb{R}^n \times \{t = 0\} \end{cases}$$

satisfying the growth estimate

$$(31) \quad |u(x, t)| \leq Ae^{a|x|^2} \quad (x \in \mathbb{R}^n, 0 \leq t \leq T)$$

for constants $A, a > 0$.

Proof. If u and \tilde{u} both satisfy (30), (31), we apply Theorem 6 to $w := \pm(u - \tilde{u})$. \square

Nonphysical solutions. There are in fact infinitely many solutions of

$$(32) \quad \begin{cases} u_t - \Delta u = 0 & \text{in } \mathbb{R}^n \times (0, T) \\ u = 0 & \text{on } \mathbb{R}^n \times \{t = 0\}; \end{cases}$$

see for instance John [J2, Chapter 7]. Each of these solutions besides $u \equiv 0$ grows very rapidly as $|x| \rightarrow \infty$.

There is an interesting point here: although $u \equiv 0$ is certainly the “physically correct” solution of (32), this initial-value problem in fact admits other, “nonphysical”, solutions. Theorem 7 provides a criterion which excludes the “wrong” solutions. We will encounter somewhat analogous situations in our study of Hamilton–Jacobi equations and conservation laws, in Chapters 3, 10 and 11.

b. Regularity. We next demonstrate that solutions of the heat equation are automatically smooth.


THEOREM 8 (Smoothness). *Suppose $u \in C_1^2(U_T)$ solves the heat equation in U_T . Then*

$$u \in C^\infty(U_T).$$

This regularity assertion is valid even if u attains nonsmooth boundary values on Γ_T .

Proof. 1. Recall from §A.2 that we write

$$C(x, t; r) = \{(y, s) \mid |x - y| \leq r, t - r^2 \leq s \leq t\}$$



to denote the closed circular cylinder of radius r , height r^2 , and top center point (x, t) .

Fix $(x_0, t_0) \in U_T$ and choose $r > 0$ so small that $C := C(x_0, t_0; r) \subset U_T$. Define also the smaller cylinders $C' := C(x_0, t_0; \frac{3}{4}r)$, $C'' := C(x_0, t_0; \frac{1}{2}r)$, which have the same top center point (x_0, t_0) .

Choose a smooth cutoff function $\zeta = \zeta(x, t)$ such that

$$\begin{cases} 0 \leq \zeta \leq 1, \quad \zeta \equiv 1 \text{ on } C', \\ \zeta \equiv 0 \text{ near the parabolic boundary of } C. \end{cases}$$

Extend $\zeta \equiv 0$ in $(\mathbb{R}^n \times [0, t_0]) - C$.

2. Assume temporarily that $u \in C^\infty(U_T)$ and set

$$v(x, t) := \zeta(x, t)u(x, t) \quad (x \in \mathbb{R}^n, 0 \leq t \leq t_0).$$

Then

$$v_t = \zeta u_t + \zeta_t u, \quad \Delta v = \zeta \Delta u + 2D\zeta \cdot Du + u \Delta \zeta.$$

Consequently

$$(33) \quad v = 0 \quad \text{on } \mathbb{R}^n \times \{t = 0\},$$

and

$$(34) \quad v_t - \Delta v = \zeta_t u - 2D\zeta \cdot Du - u \Delta \zeta =: \tilde{f}$$

in $\mathbb{R}^n \times (0, t_0)$. Now set

$$\tilde{v}(x, t) := \int_0^t \int_{\mathbb{R}^n} \Phi(x - y, t - s) \tilde{f}(y, s) dy ds.$$

According to Theorem 2

$$(35) \quad \begin{cases} \tilde{v}_t - \Delta \tilde{v} = \tilde{f} & \text{in } \mathbb{R}^n \times (0, t_0) \\ \tilde{v} = 0 & \text{on } \mathbb{R}^n \times \{t = 0\}. \end{cases}$$

Since $|v|, |\tilde{v}| \leq A$ for some constant A , Theorem 7 implies $v \equiv \tilde{v}$; that is,

$$(36) \quad v(x, t) = \int_0^t \int_{\mathbb{R}^n} \Phi(x - y, t - s) \tilde{f}(y, s) dy ds.$$

Now suppose $(x, t) \in C''$. As $\zeta \equiv 0$ off the cylinder C , (34) and (36) imply

$$u(x, t) = \iint_C \Phi(x - y, t - s) [(\zeta_s(y, s) - \Delta\zeta(y, s))u(y, s) - 2D\zeta(y, s) \cdot Du(y, s)] dy ds.$$

Note in this equation that the expression in the square brackets vanishes in some region *near* the singularity of Φ . Integrate the last term by parts:

$$(37) \quad u(x, t) = \iint_C [\Phi(x - y, t - s)(\zeta_s(y, s) + \Delta\zeta(y, s)) + 2D_y\Phi(x - y, t - s) \cdot D\zeta(y, s)] u(y, s) dy ds.$$

We have proved this formula assuming $u \in C^\infty$. If u satisfies only the hypotheses of the theorem, we derive (37) with $u^\varepsilon = \eta_\varepsilon * u$ replacing u , η_ε being the standard mollifier in the variables x and t , and let $\varepsilon \rightarrow 0$.

3. Formula (37) has the form

$$(38) \quad u(x, t) = \iint_C K(x, t, y, s) u(y, s) dy ds \quad ((x, t) \in C''),$$

where

$$K(x, t, y, s) = 0 \quad \text{for all points } (y, s) \in C',$$

since $\zeta \equiv 1$ on C' . Note also K is smooth on $C - C'$. In view of expression (38), we see u is C^∞ within $C'' = C(x_0, t_0; \frac{1}{2}r)$. \square

c. Local estimates for solutions of the heat equation. Let us now record some estimates on the derivatives of solutions to the heat equation, paying attention to the differences between derivatives with respect to x_i ($i = 1, \dots, n$) and with respect to t .

THEOREM 9 (Estimates on derivatives). *There exists for each pair of integers $k, l = 0, 1, \dots$ a constant $C_{k,l}$ such that*

$$\max_{C(x,t;r/2)} |D_x^k D_t^l u| \leq \frac{C_{kl}}{r^{k+2l+n+2}} \|u\|_{L^1(C(x,t;r))}$$

for all cylinders $C(x, t; r/2) \subset C(x, t; r) \subset U_T$ and all solutions u of the heat equation in U_T .

Proof. 1. Fix some point in U_T . Upon shifting the coordinates, we may as well assume the point is $(0, 0)$. Suppose first that the cylinder $C(1) := C(0, 0; 1)$ lies in U_T . Let $C(\frac{1}{2}) := C(0, 0; \frac{1}{2})$. Then, as in the proof of Theorem 8,

$$u(x, t) = \iint_{C(1)} K(x, t, y, s) u(y, s) dy ds \quad ((x, t) \in C(\frac{1}{2}))$$

for some smooth function K . Consequently

$$(39) \quad \begin{aligned} |D_x^k D_t^l u(x, t)| &\leq \iint_{C(1)} |D_t^l D_x^k K(x, t, y, s)| |u(y, s)| dy ds \\ &\leq C_{kl} \|u\|_{L^1(C(1))} \end{aligned}$$

for some constant C_{kl} .

2. Now suppose the cylinder $C(r) := C(0, 0; r)$ lies in U_T . Let $C(r/2) = C(0, 0; r/2)$. We rescale by defining

$$v(x, t) := u(rx, r^2 t).$$

Then $v_t - \Delta v = 0$ in the cylinder $C(1)$. According to (39),

$$|D_x^k D_t^l v(x, t)| \leq C_{kl} \|v\|_{L^1(C(1))} \quad ((x, t) \in C(\frac{1}{2})).$$

But $D_x^k D_t^l v(x, t) = r^{2l+k} D_x^k D_t^l u(rx, r^2 t)$ and $\|v\|_{L^1(C(1))} = \frac{1}{r^{n+2}} \|u\|_{L^1(C(r))}$. Therefore

$$\max_{C(r/2)} |D_x^k D_t^l u| \leq \frac{C_{kl}}{r^{2l+k+n+2}} \|u\|_{L^1(C(r))}. \quad \square$$

If u solves the heat equation within U_T , then for each time $0 < t \leq T$, the mapping $x \mapsto u(x, t)$ is analytic. (See Mikhailov [M].) However the mapping $t \mapsto u(x, t)$ is not in general analytic.

2.3.4. Energy methods.

a. Uniqueness. We investigate again the initial/boundary-value problem

$$(40) \quad \begin{cases} u_t - \Delta u = f & \text{in } U_T \\ u = g & \text{on } \Gamma_T. \end{cases}$$

We earlier invoked the maximum principle to show uniqueness and now—by analogy with §2.2.5—provide an alternative argument based upon integration by parts. We assume as usual that $U \subset \mathbb{R}^n$ is open and bounded and that ∂U is C^1 . The terminal time $T > 0$ is given.

THEOREM 10 (Uniqueness). *There exists only one solution $u \in C_1^2(\bar{U}_T)$ of the initial/boundary-value problem (40).*

Proof. 1. If \tilde{u} is another solution, $w := u - \tilde{u}$ solves

$$(41) \quad \begin{cases} w_t - \Delta w = 0 & \text{in } U_T \\ w = 0 & \text{on } \Gamma_T. \end{cases}$$

2. Set

$$e(t) := \int_U w^2(x, t) dx \quad (0 \leq t \leq T).$$

Then

$$\begin{aligned} \dot{e}(t) &= 2 \int_U w w_t dx \quad \left(\dot{} = \frac{d}{dt} \right) \\ &= 2 \int_U w \Delta w dx \\ &= -2 \int_U |Dw|^2 dx \leq 0, \end{aligned}$$

and so

$$e(t) \leq e(0) = 0 \quad (0 \leq t \leq T).$$

Consequently $w = u - \tilde{u} \equiv 0$ in U_T . □

Observe that the foregoing is a time-dependent variant of the proof of Theorem 16 in §2.2.5.

b. Backwards uniqueness. A rather more subtle question asks about uniqueness *backwards in time* for the heat equation. For this, suppose u and \tilde{u} are both smooth solutions of the heat equation in U_T , with the same boundary conditions on ∂U :

$$(42) \quad \begin{cases} u_t - \Delta u = 0 & \text{in } U_T \\ u = g & \text{on } \partial U \times [0, T], \end{cases}$$

$$(43) \quad \begin{cases} \tilde{u}_t - \Delta \tilde{u} = 0 & \text{in } U_T \\ \tilde{u} = g & \text{on } \partial U \times [0, T], \end{cases}$$

for some function g . Note carefully that we are *not* supposing $u = \tilde{u}$ at time $t = 0$.

THEOREM 11 (Backwards uniqueness). *Suppose $u, \tilde{u} \in C^2(\bar{U}_T)$ solve (42), (43). If*

$$u(x, T) = \tilde{u}(x, T) \quad (x \in U),$$

then

$$u \equiv \tilde{u} \quad \text{within } U_T.$$

In other words, if two temperature distributions on U agree at some time $T > 0$ and have had the same boundary values for times $0 \leq t \leq T$, then these temperatures must have been identically equal within U at all earlier times. This is not at all obvious.

Proof. 1. Write $w := u - \tilde{u}$ and, as in the proof of Theorem 10, set

$$e(t) := \int_U w^2(x, t) dx \quad (0 \leq t \leq T).$$

As before

$$(44) \quad \dot{e}(t) = -2 \int_U |Dw|^2 dx \quad \left(\dot{\cdot} = \frac{d}{dt} \right).$$

Furthermore

$$(45) \quad \begin{aligned} \ddot{e}(t) &= -4 \int_U Dw \cdot Dw_t dx \\ &= 4 \int_U \Delta w w_t dx \\ &= 4 \int_U (\Delta w)^2 dx \quad \text{by (41)}. \end{aligned}$$

Now since $w = 0$ on ∂U ,

$$\begin{aligned} \int_U |Dw|^2 dx &= - \int_U w \Delta w dx \\ &\leq \left(\int_U w^2 dx \right)^{1/2} \left(\int_U (\Delta w)^2 dx \right)^{1/2}. \end{aligned}$$

Thus (44) and (45) imply

$$\begin{aligned} (\dot{e}(t))^2 &= 4 \left(\int_U |Dw|^2 dx \right)^2 \\ &\leq \left(\int_U w^2 dx \right) \left(4 \int_U (\Delta w)^2 dx \right) \\ &= e(t) \ddot{e}(t). \end{aligned}$$

Hence

$$(46) \quad \ddot{e}(t)e(t) \geq (\dot{e}(t))^2 \quad (0 \leq t \leq T).$$

2. Now if $e(t) = 0$ for all $0 \leq t \leq T$, we are done. Otherwise there exists an interval $[t_1, t_2] \subset [0, T]$, with

$$(47) \quad e(t) > 0 \quad \text{for } t_1 \leq t < t_2, \quad e(t_2) = 0.$$

3. Now write

$$(48) \quad f(t) := \log e(t) \quad (t_1 \leq t < t_2).$$

Then

$$\ddot{f}(t) = \frac{\ddot{e}(t)}{e(t)} - \frac{\dot{e}(t)^2}{e(t)^2} \geq 0 \quad \text{by (46),}$$

and so f is convex on the interval (t_1, t_2) . Consequently if $0 < \tau < 1$, $t_1 < t < t_2$, we have

$$f((1 - \tau)t_1 + \tau t) \leq (1 - \tau)f(t_1) + \tau f(t).$$

Recalling (48), we deduce

$$e((1 - \tau)t_1 + \tau t) \leq e(t_1)^{1-\tau} e(t)^\tau,$$

and so

$$0 \leq e((1 - \tau)t_1 + \tau t_2) \leq e(t_1)^{1-\tau} e(t_2)^\tau \quad (0 < \tau < 1).$$

But in view of (47) this inequality implies $e(t) = 0$ for all times $t_1 \leq t \leq t_2$, a contradiction. \square

2.4. WAVE EQUATION

In this section we investigate the *wave equation*

$$(1) \quad u_{tt} - \Delta u = 0$$

and the *nonhomogeneous wave equation*

$$(2) \quad u_{tt} - \Delta u = f,$$

subject to appropriate initial and boundary conditions. Here $t > 0$ and $x \in U$, where $U \subset \mathbb{R}^n$ is open. The unknown is $u : \bar{U} \times [0, \infty) \rightarrow \mathbb{R}$, $u = u(x, t)$, and the Laplacian Δ is taken with respect to the spatial variables

$x = (x_1, \dots, x_n)$. In (2) the function $f : U \times [0, \infty) \rightarrow \mathbb{R}$ is given. A common abbreviation is to write

$$\square u := u_{tt} - \Delta u.$$

We shall discover that solutions of the wave equation behave quite differently than solutions of Laplace's equation or the heat equation. For example, these solutions are generally not C^∞ , exhibit finite speed of propagation, etc.

Physical interpretation. The wave equation is a simplified model for a vibrating string ($n = 1$), membrane ($n = 2$), or elastic solid ($n = 3$). In these physical interpretations $u(x, t)$ represents the displacement in some direction of the point x at time $t \geq 0$.

Let V represent any smooth subregion of U . The acceleration within V is then

$$\frac{d^2}{dt^2} \int_V u \, dx = \int_V u_{tt} \, dx$$

and the net contact force is

$$- \int_{\partial V} \mathbf{F} \cdot \nu \, dS,$$

where \mathbf{F} denotes the force acting on V through ∂V and the mass density is taken to be unity. Newton's law asserts that the mass times the acceleration equals the net force:

$$\int_V u_{tt} \, dx = - \int_{\partial V} \mathbf{F} \cdot \nu \, dS.$$

This identity obtains for each subregion V and so

$$u_{tt} = - \operatorname{div} \mathbf{F}.$$

For elastic bodies, \mathbf{F} is a function of the displacement gradient Du , whence

$$u_{tt} + \operatorname{div} \mathbf{F}(Du) = 0.$$

For small Du , the linearization $\mathbf{F}(Du) \approx -aDu$ is often appropriate; and so

$$u_{tt} - a\Delta u = 0.$$

This is the wave equation if $a = 1$.

This physical interpretation strongly suggests it will be mathematically appropriate to specify *two* initial conditions, on the *displacement* u and the *velocity* u_t , at time $t = 0$.

2.4.1. Solution by spherical means.

We began §§2.2.1 and 2.3.1 by searching for certain scaling invariant solutions of Laplace's equation and the heat equation. For the wave equation however we will instead present the (reasonably) elegant method of solving (1) first for $n = 1$ directly and then for $n \geq 2$ by the method of spherical means.

a. Solution for $n = 1$, d'Alembert's formula. We first focus our attention on the initial-value problem for the one-dimensional wave equation in all of \mathbb{R} :

$$(3) \quad \begin{cases} u_{tt} - u_{xx} = 0 & \text{in } \mathbb{R} \times (0, \infty) \\ u = g, \quad u_t = h & \text{on } \mathbb{R} \times \{t = 0\}, \end{cases}$$

where g, h are given. We desire to derive a formula for u in terms of g and h .

Let us first note that the PDE in (3) can be “factored”, to read

$$(4) \quad \left(\frac{\partial}{\partial t} + \frac{\partial}{\partial x} \right) \left(\frac{\partial}{\partial t} - \frac{\partial}{\partial x} \right) u = u_{tt} - u_{xx} = 0.$$

Write

$$(5) \quad v(x, t) := \left(\frac{\partial}{\partial t} - \frac{\partial}{\partial x} \right) u(x, t).$$

Then (4) says

$$v_t(x, t) + v_x(x, t) = 0 \quad (x \in \mathbb{R}, t > 0).$$

This is a transport equation with constant coefficients. Applying formula (3) from §2.1.1 (with $n = 1, b = 1$), we find

$$(6) \quad v(x, t) = a(x - t)$$

for $a(x) := v(x, 0)$. Combining now (4)–(6), we obtain

$$u_t(x, t) - u_x(x, t) = a(x - t) \quad \text{in } \mathbb{R} \times (0, \infty).$$

This is a nonhomogeneous transport equation; and so formula (5) from §2.1.2 (with $n = 1, b = -1, f(x, t) = a(x - t)$) implies for $b(x) := u(x, 0)$ that

$$(7) \quad \begin{aligned} u(x, t) &= \int_0^t a(x + (t - s) - s) ds + b(x + t) \\ &= \frac{1}{2} \int_{x-t}^{x+t} a(y) dy + b(x + t). \end{aligned}$$

We lastly invoke the initial conditions in (3) to compute a and b . The first initial condition in (3) gives

$$b(x) = g(x) \quad (x \in \mathbb{R}),$$

whereas the second initial condition and (5) imply

$$a(x) = v(x, 0) = u_t(x, 0) - u_x(x, 0) = h(x) - g'(x) \quad (x \in \mathbb{R}).$$

Our substituting into (7) now yields

$$u(x, t) = \frac{1}{2} \int_{x-t}^{x+t} h(y) - g'(y) dy + g(x+t).$$

Hence

$$(8) \quad u(x, t) = \frac{1}{2} [g(x+t) + g(x-t)] + \frac{1}{2} \int_{x-t}^{x+t} h(y) dy \quad (x \in \mathbb{R}, t \geq 0).$$

This is *d'Alembert's formula*.

We have derived formula (8) assuming u is a (sufficiently smooth) solution of (3). We need to check that this really is a solution.

THEOREM 1 (Solution of wave equation, $n = 1$). *Assume $g \in C^2(\mathbb{R})$, $h \in C^1(\mathbb{R})$, and define u by d'Alembert's formula (8). Then*

- (i) $u \in C^2(\mathbb{R} \times [0, \infty))$,
- (ii) $u_{tt} - u_{xx} = 0$ in $\mathbb{R} \times (0, \infty)$,

and

- (iii) $\lim_{\substack{(x,t) \rightarrow (x^0,0) \\ t > 0}} u(x, t) = g(x^0), \quad \lim_{\substack{(x,t) \rightarrow (x^0,0) \\ t > 0}} u_t(x, t) = h(x^0)$
for each point $x^0 \in \mathbb{R}$.

The proof is a straightforward calculation.

Remarks. (i) In view of (8), our solution u has the form

$$u(x, t) = F(x+t) + G(x-t)$$

for appropriate functions F and G . Conversely any function of this form solves $u_{tt} - u_{xx} = 0$. Hence the general solution of the one-dimensional wave equation is a sum of the general solution of $u_t - u_x = 0$ and the general solution of $u_t + u_x = 0$. This is a consequence of the factorization (4). See Problem 19.

(ii) We see from (8) that if $g \in C^k$ and $h \in C^{k-1}$, then $u \in C^k$ but is not in general smoother. Thus the wave equation does *not* cause instantaneous smoothing of the initial data, as does the heat equation.

A reflection method. To illustrate a further application of d'Alembert's formula, let us next consider this initial/boundary-value problem on the half-line $\mathbb{R}_+ = \{x > 0\}$:

$$(9) \quad \begin{cases} u_{tt} - u_{xx} = 0 & \text{in } \mathbb{R}_+ \times (0, \infty) \\ u = g, \quad u_t = h & \text{on } \mathbb{R}_+ \times \{t = 0\} \\ u = 0 & \text{on } \{x = 0\} \times (0, \infty), \end{cases}$$

where g, h are given, with $g(0) = h(0) = 0$.

We convert (9) into the form (3) by extending u, g, h to all of \mathbb{R} by *odd reflection*. That is, we set

$$\begin{aligned} \tilde{u}(x, t) &:= \begin{cases} u(x, t) & (x \geq 0, t \geq 0) \\ -u(-x, t) & (x \leq 0, t \geq 0), \end{cases} \\ \tilde{g}(x) &:= \begin{cases} g(x) & (x \geq 0) \\ -g(-x) & (x \leq 0), \end{cases} \\ \tilde{h}(x) &:= \begin{cases} h(x) & (x \geq 0) \\ -h(-x) & (x \leq 0). \end{cases} \end{aligned}$$

Then (9) becomes

$$\begin{cases} \tilde{u}_{tt} = \tilde{u}_{xx} & \text{in } \mathbb{R} \times (0, \infty) \\ \tilde{u} = \tilde{g}, \quad \tilde{u}_t = \tilde{h} & \text{on } \mathbb{R} \times \{t = 0\}. \end{cases}$$

Hence d'Alembert's formula (8) implies

$$\tilde{u}(x, t) = \frac{1}{2}[\tilde{g}(x+t) + \tilde{g}(x-t)] + \frac{1}{2} \int_{x-t}^{x+t} \tilde{h}(y) dy.$$

Recalling the definitions of $\tilde{u}, \tilde{g}, \tilde{h}$ above, we can transform this expression to read for $x \geq 0, t \geq 0$:

$$(10) \quad u(x, t) = \begin{cases} \frac{1}{2}[g(x+t) + g(x-t)] + \frac{1}{2} \int_{x-t}^{x+t} h(y) dy & \text{if } x \geq t \geq 0 \\ \frac{1}{2}[g(x+t) - g(t-x)] + \frac{1}{2} \int_{-x+t}^{x+t} h(y) dy & \text{if } 0 \leq x \leq t. \end{cases}$$

If $h \equiv 0$, we can understand formula (10) as saying that an initial displacement g splits into two parts, one moving to the right with speed one and the other to the left with speed one. The latter then reflects off the point $x = 0$, where the vibrating string is held fixed.

Note that our solution does not belong to C^2 , unless $g''(0) = 0$. □

b. Spherical means. Now suppose $n \geq 2$, $m \geq 2$, and $u \in C^m(\mathbb{R}^n \times [0, \infty))$ solves the initial-value problem

$$(11) \quad \begin{cases} u_{tt} - \Delta u = 0 & \text{in } \mathbb{R}^n \times (0, \infty) \\ u = g, \quad u_t = h & \text{on } \mathbb{R}^n \times \{t = 0\}. \end{cases}$$

We intend to derive an explicit formula for u in terms of g, h . The plan will be to study first the average of u over certain spheres. These averages, taken as functions of the time t and the radius r , turn out to solve the Euler–Poisson–Darboux equation, a PDE which we can for odd n convert into the ordinary one-dimensional wave equation. Applying d’Alembert’s formula, or more precisely its variant (10), eventually leads us to a formula for the solution.

NOTATION. (i) Let $x \in \mathbb{R}^n$, $t > 0$, $r > 0$. Define

$$(12) \quad U(x; r, t) := \int_{\partial B(x, r)} u(y, t) dS(y),$$

the average of $u(\cdot, t)$ over the sphere $\partial B(x, r)$.

(ii) Similarly,

$$(13) \quad \begin{cases} G(x; r) := \int_{\partial B(x, r)} g(y) dS(y) \\ H(x; r) := \int_{\partial B(x, r)} h(y) dS(y). \end{cases}$$

For fixed x , we hereafter regard U as a function of r and t and discover a partial differential equation that U solves:

LEMMA 1 (Euler–Poisson–Darboux equation). *Fix $x \in \mathbb{R}^n$, and let u satisfy (11). Then $U \in C^m(\bar{\mathbb{R}}_+ \times [0, \infty))$ and*

$$(14) \quad \begin{cases} U_{tt} - U_{rr} - \frac{n-1}{r}U_r = 0 & \text{in } \mathbb{R}_+ \times (0, \infty) \\ U = G, \quad U_t = H & \text{on } \mathbb{R}_+ \times \{t = 0\}. \end{cases}$$

The partial differential equation in (14) is the *Euler–Poisson–Darboux equation*. (Note that the term $U_{rr} + \frac{n-1}{r}U_r$ is the radial part of the Laplacian Δ in polar coordinates.)

Proof. 1. As in the proof of Theorem 2 in §2.2.2 we compute for $r > 0$

$$(15) \quad U_r(x; r, t) = \frac{r}{n} \int_{B(x, r)} \Delta u(y, t) dy.$$

From this equality we deduce $\lim_{r \rightarrow 0^+} U_r(x; r, t) = 0$. We next differentiate (15), to discover after some computations that

$$(16) \quad U_{rr}(x; r, t) = \int_{\partial B(x,r)} \Delta u \, dS + \left(\frac{1}{n} - 1 \right) \int_{B(x,r)} \Delta u \, dy.$$

Thus $\lim_{r \rightarrow 0^+} U_{rr}(x; r, t) = \frac{1}{n} \Delta u(x, t)$. Using formula (16), we can similarly compute U_{rrr} , etc., and so verify that $U \in C^m(\mathbb{R}_+ \times [0, \infty))$.

2. Continuing the calculation above, we see from (15) that

$$\begin{aligned} U_r &= \frac{r}{n} \int_{B(x,r)} u_{tt} \, dy \quad \text{by (11)} \\ &= \frac{1}{n\alpha(n)} \frac{1}{r^{n-1}} \int_{B(x,r)} u_{tt} \, dy. \end{aligned}$$

Thus

$$r^{n-1} U_r = \frac{1}{n\alpha(n)} \int_{B(x,r)} u_{tt} \, dy,$$

and so

$$\begin{aligned} (r^{n-1} U_r)_r &= \frac{1}{n\alpha(n)} \int_{\partial B(x,r)} u_{tt} \, dS \\ &= r^{n-1} \int_{\partial B(x,r)} u_{tt} \, dS = r^{n-1} U_{tt}. \end{aligned} \quad \square$$

c. Solution for $n = 3, 2$, Kirchhoff's and Poisson's formulas. The plan in the ensuing subsections will be to transform the Euler–Poisson–Darboux equation (14) into the usual one-dimensional wave equation. As the full procedure is rather complicated, we pause here to handle the simpler cases $n = 3, 2$, in that order.

Solution for $n = 3$. Let us therefore hereafter take $n = 3$, and suppose $u \in C^2(\mathbb{R}^3 \times [0, \infty))$ solves the initial-value problem (11). We recall the definitions (12), (13) of U, G, H and then set

$$(17) \quad \tilde{U} := rU,$$

$$(18) \quad \tilde{G} := rG, \quad \tilde{H} := rH.$$

We now assert that \tilde{U} solves

$$(19) \quad \begin{cases} \tilde{U}_{tt} - \tilde{U}_{rr} = 0 & \text{in } \mathbb{R}_+ \times (0, \infty) \\ \tilde{U} = \tilde{G}, \quad \tilde{U}_t = \tilde{H} & \text{on } \mathbb{R}_+ \times \{t = 0\} \\ \tilde{U} = 0 & \text{on } \{r = 0\} \times (0, \infty). \end{cases}$$

Indeed

$$\begin{aligned}\tilde{U}_{tt} &= rU_{tt} \\ &= r \left[U_{rr} + \frac{2}{r}U_r \right] \quad \text{by (14), with } n = 3 \\ &= rU_{rr} + 2U_r = (U + rU_r)_r = \tilde{U}_{rr}.\end{aligned}$$

Notice also that $\tilde{G}_{rr}(0) = 0$. Applying formula (10) to (19), we find for $0 \leq r \leq t$

$$(20) \quad \tilde{U}(x; r, t) = \frac{1}{2}[\tilde{G}(r+t) - \tilde{G}(t-r)] + \frac{1}{2} \int_{-r+t}^{r+t} \tilde{H}(y) dy.$$

Since (12) implies $u(x, t) = \lim_{r \rightarrow 0^+} U(x; r, t)$, we conclude from (17), (18), (20) that

$$\begin{aligned}u(x, t) &= \lim_{r \rightarrow 0^+} \frac{\tilde{U}(x; r, t)}{r} \\ &= \lim_{r \rightarrow 0^+} \left[\frac{\tilde{G}(t+r) - \tilde{G}(t-r)}{2r} + \frac{1}{2r} \int_{t-r}^{t+r} \tilde{H}(y) dy \right] \\ &= \tilde{G}'(t) + \tilde{H}(t).\end{aligned}$$

Owing then to (13), we deduce

$$(21) \quad u(x, t) = \frac{\partial}{\partial t} \left(t \int_{\partial B(x,t)} g dS \right) + t \int_{\partial B(x,t)} h dS.$$

But

$$\int_{\partial B(x,t)} g(y) dS(y) = \int_{\partial B(0,1)} g(x + tz) dS(z);$$

and so

$$\begin{aligned}\frac{\partial}{\partial t} \left(\int_{\partial B(x,t)} g dS \right) &= \int_{\partial B(0,1)} Dg(x + tz) \cdot z dS(z) \\ &= \int_{\partial B(x,t)} Dg(y) \cdot \left(\frac{y-x}{t} \right) dS(y).\end{aligned}$$

Returning to (21), we therefore conclude

$$(22) \quad u(x, t) = \int_{\partial B(x,t)} th(y) + g(y) + Dg(y) \cdot (y-x) dS(y) \quad (x \in \mathbb{R}^3, t > 0).$$

This is *Kirchhoff's formula* for the solution of the initial-value problem (11) in three dimensions.

Solution for $n = 2$. No transformation like (17) works to convert the Euler–Poisson–Darboux equation into the one-dimensional wave equation when $n = 2$. Instead we will take the initial-value problem (11) for $n = 2$ and simply regard it as a problem for $n = 3$, in which the third spatial variable x_3 does not appear.

Indeed, assuming $u \in C^2(\mathbb{R}^2 \times [0, \infty))$ solves (11) for $n = 2$, let us write

$$(23) \quad \bar{u}(x_1, x_2, x_3, t) := u(x_1, x_2, t).$$

Then (11) implies

$$(24) \quad \begin{cases} \bar{u}_{tt} - \Delta \bar{u} = 0 & \text{in } \mathbb{R}^3 \times (0, \infty) \\ \bar{u} = \bar{g}, \quad \bar{u}_t = \bar{h} & \text{on } \mathbb{R}^3 \times \{t = 0\}, \end{cases}$$

for

$$\bar{g}(x_1, x_2, x_3) := g(x_1, x_2), \quad \bar{h}(x_1, x_2, x_3) := h(x_1, x_2).$$

If we write $x = (x_1, x_2) \in \mathbb{R}^2$ and $\bar{x} = (x_1, x_2, 0) \in \mathbb{R}^3$, then (24) and Kirchhoff's formula (in the form (21)) imply

$$(25) \quad \begin{aligned} u(x, t) &= \bar{u}(\bar{x}, t) \\ &= \frac{\partial}{\partial t} \left(t \int_{\partial \bar{B}(\bar{x}, t)} \bar{g} \, d\bar{S} \right) + t \int_{\partial \bar{B}(\bar{x}, t)} \bar{h} \, d\bar{S}, \end{aligned}$$

where $\bar{B}(\bar{x}, t)$ denotes the ball in \mathbb{R}^3 with center \bar{x} , radius $t > 0$ and where $d\bar{S}$ denotes two-dimensional surface measure on $\partial \bar{B}(\bar{x}, t)$. We simplify (25) by observing

$$\begin{aligned} \int_{\partial \bar{B}(\bar{x}, t)} \bar{g} \, d\bar{S} &= \frac{1}{4\pi t^2} \int_{\partial \bar{B}(\bar{x}, t)} \bar{g} \, d\bar{S} \\ &= \frac{2}{4\pi t^2} \int_{B(x, t)} g(y) (1 + |D\gamma(y)|^2)^{1/2} \, dy, \end{aligned}$$

where $\gamma(y) = (t^2 - |y - x|^2)^{1/2}$ for $y \in B(x, t)$. The factor “2” enters since $\partial \bar{B}(\bar{x}, t)$ consists of two hemispheres. Observe that $(1 + |D\gamma|^2)^{1/2} = t(t^2 - |y - x|^2)^{-1/2}$. Therefore

$$\begin{aligned} \int_{\partial \bar{B}(\bar{x}, t)} \bar{g} \, d\bar{S} &= \frac{1}{2\pi t} \int_{B(x, t)} \frac{g(y)}{(t^2 - |y - x|^2)^{1/2}} \, dy \\ &= \frac{t}{2} \int_{B(x, t)} \frac{g(y)}{(t^2 - |y - x|^2)^{1/2}} \, dy. \end{aligned}$$

Consequently formula (25) becomes

$$(26) \quad u(x, t) = \frac{1}{2} \frac{\partial}{\partial t} \left(t^2 \int_{B(x, t)} \frac{g(y)}{(t^2 - |y - x|^2)^{1/2}} dy \right) + \frac{t^2}{2} \int_{B(x, t)} \frac{h(y)}{(t^2 - |y - x|^2)^{1/2}} dy.$$

But

$$t^2 \int_{B(x, t)} \frac{g(y)}{(t^2 - |y - x|^2)^{1/2}} dy = t \int_{B(0, 1)} \frac{g(x + tz)}{(1 - |z|^2)^{1/2}} dz,$$

and so

$$\begin{aligned} & \frac{\partial}{\partial t} \left(t^2 \int_{B(x, t)} \frac{g(y)}{(t^2 - |y - x|^2)^{1/2}} dy \right) \\ &= \int_{B(0, 1)} \frac{g(x + tz)}{(1 - |z|^2)^{1/2}} dz + t \int_{B(0, 1)} \frac{Dg(x + tz) \cdot z}{(1 - |z|^2)^{1/2}} dz \\ &= t \int_{B(x, t)} \frac{g(y)}{(t^2 - |y - x|^2)^{1/2}} dy + t \int_{B(x, t)} \frac{Dg(y) \cdot (y - x)}{(t^2 - |y - x|^2)^{1/2}} dy. \end{aligned}$$

Hence we can rewrite (26) and obtain the relation

$$(27) \quad u(x, t) = \frac{1}{2} \int_{B(x, t)} \frac{tg(y) + t^2 h(y) + t Dg(y) \cdot (y - x)}{(t^2 - |y - x|^2)^{1/2}} dy$$

for $x \in \mathbb{R}^2$, $t > 0$. This is *Poisson's formula* for the solution of the initial-value problem (11) in two dimensions.

The trick of solving the problem for $n = 3$ first and then dropping to $n = 2$ is the *method of descent*.

d. Solution for odd n . In this subsection we solve the Euler–Poisson–Darboux PDE for odd $n \geq 3$. We first record some technical facts.

LEMMA 2 (Some useful identities). *Let $\phi : \mathbb{R} \rightarrow \mathbb{R}$ be C^{k+1} . Then for $k = 1, 2, \dots$*

- (i) $\left(\frac{d^2}{dr^2} \right) \left(\frac{1}{r} \frac{d}{dr} \right)^{k-1} (r^{2k-1} \phi(r)) = \left(\frac{1}{r} \frac{d}{dr} \right)^k \left(r^{2k} \frac{d\phi}{dr}(r) \right),$
- (ii) $\left(\frac{1}{r} \frac{d}{dr} \right)^{k-1} (r^{2k-1} \phi(r)) = \sum_{j=0}^{k-1} \beta_j^k r^{j+1} \frac{d^j \phi}{dr^j}(r),$
where the constants β_j^k ($j = 0, \dots, k-1$) are independent of ϕ .

Furthermore,

$$(iii) \quad \beta_0^k = 1 \cdot 3 \cdot 5 \cdots (2k - 1).$$

The proof by induction is left as an exercise.

Now assume

$$n \geq 3 \text{ is an odd integer}$$

and set

$$n = 2k + 1 \quad (k \geq 1).$$

Henceforth suppose $u \in C^{k+1}(\mathbb{R}^n \times [0, \infty))$ solves the initial-value problem (11). Then the function U defined by (12) is C^{k+1} .

NOTATION. We write

$$(28) \quad \begin{cases} \tilde{U}(r, t) := \left(\frac{1}{r} \frac{\partial}{\partial r}\right)^{k-1} (r^{2k-1} U(x; r, t)) \\ \tilde{G}(r) := \left(\frac{1}{r} \frac{\partial}{\partial r}\right)^{k-1} (r^{2k-1} G(x; r)) \\ \tilde{H}(r) := \left(\frac{1}{r} \frac{\partial}{\partial r}\right)^{k-1} (r^{2k-1} H(x; r)) \end{cases} \quad (r > 0, t \geq 0).$$

Then

$$(29) \quad \tilde{U}(r, 0) = \tilde{G}(r), \quad \tilde{U}_t(r, 0) = \tilde{H}(r).$$

Next we combine Lemma 1 and the identities provided by Lemma 2 to demonstrate that the transformation (28) of U into \tilde{U} in effect converts the Euler–Poisson–Darboux equation into the wave equation.

LEMMA 3 (\tilde{U} solves the one-dimensional wave equation). *We have*

$$\begin{cases} \tilde{U}_{tt} - \tilde{U}_{rr} = 0 & \text{in } \mathbb{R}_+ \times (0, \infty) \\ \tilde{U} = \tilde{G}, \quad \tilde{U}_t = \tilde{H} & \text{on } \mathbb{R}_+ \times \{t = 0\} \\ \tilde{U} = 0 & \text{on } \{r = 0\} \times (0, \infty). \end{cases}$$

Proof. If $r > 0$,

$$\begin{aligned} \tilde{U}_{rr} &= \left(\frac{\partial^2}{\partial r^2}\right) \left(\frac{1}{r} \frac{\partial}{\partial r}\right)^{k-1} (r^{2k-1} U) \\ &= \left(\frac{1}{r} \frac{\partial}{\partial r}\right)^k (r^{2k} U_r) \quad \text{by Lemma 2(i)} \\ &= \left(\frac{1}{r} \frac{\partial}{\partial r}\right)^{k-1} [r^{2k-1} U_{rr} + 2kr^{2k-2} U_r] \\ &= \left(\frac{1}{r} \frac{\partial}{\partial r}\right)^{k-1} \left[r^{2k-1} \left(U_{rr} + \frac{n-1}{r} U_r \right) \right] \quad (n = 2k + 1) \\ &= \left(\frac{1}{r} \frac{\partial}{\partial r}\right)^{k-1} (r^{2k-1} U_{tt}) = \tilde{U}_{tt}, \end{aligned}$$

the next-to-last equality holding according to (14). Using Lemma 2(ii) we conclude as well that $\tilde{U} = 0$ on $\{r = 0\}$. \square

In view of Lemma 3, (29), and formula (10), we conclude for $0 \leq r \leq t$ that

$$(30) \quad \tilde{U}(r, t) = \frac{1}{2}[\tilde{G}(r+t) - \tilde{G}(t-r)] + \frac{1}{2} \int_{t-r}^{t+r} \tilde{H}(y) dy$$

for all $r \in \mathbb{R}$, $t \geq 0$. But recall $u(x, t) = \lim_{r \rightarrow 0} U(x; r, t)$. Furthermore Lemma 2(ii) asserts

$$\begin{aligned} \tilde{U}(r, t) &= \left(\frac{1}{r} \frac{\partial}{\partial r} \right)^{k-1} (r^{2k-1} U(x; r, t)) \\ &= \sum_{j=0}^{k-1} \beta_j^k r^{j+1} \frac{\partial^j}{\partial r^j} U(x; r, t), \end{aligned}$$

and so

$$\lim_{r \rightarrow 0} \frac{\tilde{U}(r, t)}{\beta_0^k r} = \lim_{r \rightarrow 0} U(x; r, t) = u(x, t).$$

Thus (30) implies

$$\begin{aligned} u(x, t) &= \frac{1}{\beta_0^k} \lim_{r \rightarrow 0} \left[\frac{\tilde{G}(t+r) - \tilde{G}(t-r)}{2r} + \frac{1}{2r} \int_{t-r}^{t+r} \tilde{H}(y) dy \right] \\ &= \frac{1}{\beta_0^k} [\tilde{G}'(t) + \tilde{H}(t)]. \end{aligned}$$

Finally then, since $n = 2k + 1$, (30) and Lemma 2(iii) yield this representation formula:

$$(31) \quad \left\{ \begin{aligned} u(x, t) &= \frac{1}{\gamma_n} \left[\left(\frac{\partial}{\partial t} \right) \left(\frac{1}{t} \frac{\partial}{\partial t} \right)^{\frac{n-3}{2}} \left(t^{n-2} \int_{\partial B(x,t)} g dS \right) \right. \\ &\quad \left. + \left(\frac{1}{t} \frac{\partial}{\partial t} \right)^{\frac{n-3}{2}} \left(t^{n-2} \int_{\partial B(x,t)} h dS \right) \right] \\ &\text{where } n \text{ is odd and } \gamma_n = 1 \cdot 3 \cdot 5 \cdots (n-2), \end{aligned} \right.$$

for $x \in \mathbb{R}^n$, $t > 0$.

We note that $\gamma_3 = 1$, and so (31) agrees for $n = 3$ with (21) and thus with Kirchhoff's formula (22).

It remains to check that formula (31) really provides a solution of (11).

THEOREM 2 (Solution of wave equation in odd dimensions). *Assume n is an odd integer, $n \geq 3$, and suppose also $g \in C^{m+1}(\mathbb{R}^n)$, $h \in C^m(\mathbb{R}^n)$, for $m = \frac{n+1}{2}$. Define u by (31). Then*

$$(i) \quad u \in C^2(\mathbb{R}^n \times [0, \infty)),$$

$$(ii) \quad u_{tt} - \Delta u = 0 \quad \text{in } \mathbb{R}^n \times (0, \infty),$$

and

$$(iii) \quad \lim_{\substack{(x,t) \rightarrow (x^0,0) \\ x \in \mathbb{R}^n, t > 0}} u(x, t) = g(x^0), \quad \lim_{\substack{(x,t) \rightarrow (x^0,0) \\ x \in \mathbb{R}^n, t > 0}} u_t(x, t) = h(x^0)$$

for each point $x^0 \in \mathbb{R}^n$.

Proof. 1. Suppose first $g \equiv 0$, so that

$$(32) \quad u(x, t) = \frac{1}{\gamma_n} \left(\frac{1}{t} \frac{\partial}{\partial t} \right)^{\frac{n-3}{2}} (t^{n-2} H(x; t)).$$

Then Lemma 2(i) lets us compute

$$u_{tt} = \frac{1}{\gamma_n} \left(\frac{1}{t} \frac{\partial}{\partial t} \right)^{\frac{n-1}{2}} (t^{n-1} H_t).$$

From the calculation in the proof of Theorem 2 in §2.2.2, we see as well that

$$H_t = \frac{t}{n} \int_{B(x,t)} \Delta h \, dy.$$

Consequently

$$\begin{aligned} u_{tt} &= \frac{1}{n\alpha(n)\gamma_n} \left(\frac{1}{t} \frac{\partial}{\partial t} \right)^{\frac{n-1}{2}} \left(\int_{B(x,t)} \Delta h \, dy \right) \\ &= \frac{1}{n\alpha(n)\gamma_n} \left(\frac{1}{t} \frac{\partial}{\partial t} \right)^{\frac{n-3}{2}} \left(\frac{1}{t} \int_{\partial B(x,t)} \Delta h \, dS \right). \end{aligned}$$

On the other hand,

$$\Delta H(x; t) = \Delta_x \int_{\partial B(0,t)} h(x+y) \, dS(y) = \int_{\partial B(x,t)} \Delta h \, dS.$$

Consequently (32) and the calculations above imply $u_{tt} = \Delta u$ in $\mathbb{R}^n \times (0, \infty)$.

A similar computation works if $h \equiv 0$.

2. We leave it as an exercise to confirm, using Lemma 2(ii)–(iii), that u takes on the correct initial conditions. \square

Remarks. (i) Notice that to compute $u(x, t)$ we need only have information on g, h and their derivatives on the sphere $\partial B(x, t)$, and not on the entire ball $B(x, t)$.

(ii) Comparing formula (31) with d'Alembert's formula (8) ($n = 1$), we observe that the latter does not involve the derivatives of g . This suggests that for $n > 1$, a solution of the wave equation (11) need not for times $t > 0$ be as smooth as its initial value g : irregularities in g may focus at times $t > 0$, thereby causing u to be less regular. (We will see later in §2.4.3 that the "energy norm" of u does *not* deteriorate for $t > 0$.)

(iii) Once again (as in the case $n = 1$) we see the phenomenon of finite propagation speed of the initial disturbance.

(iv) A completely different derivation of formula (31) (using the heat equation!) is in §4.3.3. \square

e. Solution for even n . Assume now

n is an even integer.

Suppose u is a C^m solution of (11), $m = \frac{n+2}{2}$. We want to fashion a representation formula like (31) for u . The trick, as above for $n = 2$, is to note that

$$(33) \quad \bar{u}(x_1, \dots, x_{n+1}, t) := u(x_1, \dots, x_n, t)$$

solves the wave equation in $\mathbb{R}^{n+1} \times (0, \infty)$, with the initial conditions

$$\bar{u} = \bar{g}, \quad \bar{u}_t = \bar{h} \quad \text{on } \mathbb{R}^{n+1} \times \{t = 0\},$$

where

$$(34) \quad \begin{cases} \bar{g}(x_1, \dots, x_{n+1}) := g(x_1, \dots, x_n) \\ \bar{h}(x_1, \dots, x_{n+1}) := h(x_1, \dots, x_n). \end{cases}$$

As $n + 1$ is odd, we may employ (31) (with $n + 1$ replacing n) to secure a representation formula for \bar{u} in terms of \bar{g}, \bar{h} . But then (33) and (34) yield at once a formula for u in terms of g, h . This is again the method of descent.

To carry out the details, let us fix $x \in \mathbb{R}^n$, $t > 0$, and write $\bar{x} = (x_1, \dots, x_n, 0) \in \mathbb{R}^{n+1}$. Then (31), with $n + 1$ replacing n , gives

$$(35) \quad u(x, t) = \frac{1}{\gamma_{n+1}} \left[\frac{\partial}{\partial t} \left(\frac{1}{t} \frac{\partial}{\partial t} \right)^{\frac{n-2}{2}} \left(t^{n-1} \int_{\partial \bar{B}(\bar{x}, t)} \bar{g} d\bar{S} \right) + \left(\frac{1}{t} \frac{\partial}{\partial t} \right)^{\frac{n-2}{2}} \left(t^{n-1} \int_{\partial \bar{B}(\bar{x}, t)} \bar{h} d\bar{S} \right) \right],$$

$\bar{B}(\bar{x}, t)$ denoting the ball in \mathbb{R}^{n+1} with center \bar{x} and radius t and $d\bar{S}$ denoting n -dimensional surface measure on $\partial\bar{B}(\bar{x}, t)$. Now

$$(36) \quad \int_{\partial\bar{B}(\bar{x}, t)} \bar{g} d\bar{S} = \frac{1}{(n+1)\alpha(n+1)t^n} \int_{\partial\bar{B}(\bar{x}, t)} \bar{g} d\bar{S}.$$

Note that $\partial\bar{B}(x, t) \cap \{y_{n+1} \geq 0\}$ is the graph of the function $\gamma(y) := (t^2 - |y - x|^2)^{1/2}$ for $y \in B(x, t) \subset \mathbb{R}^n$. Likewise $\partial\bar{B}(x, t) \cap \{y_{n+1} \leq 0\}$ is the graph of $-\gamma$. Thus (36) implies

$$(37) \quad \int_{\partial\bar{B}(\bar{x}, t)} \bar{g} d\bar{S} = \frac{2}{(n+1)\alpha(n+1)t^n} \int_{B(x, t)} g(y)(1 + |D\gamma(y)|^2)^{1/2} dy,$$

the factor “2” entering because $\partial\bar{B}(\bar{x}, t)$ comprises two hemispheres. Note that $(1 + |D\gamma(y)|^2)^{1/2} = t(t^2 - |y - x|^2)^{-1/2}$. Our substituting this into (37) yields

$$\begin{aligned} \int_{\partial\bar{B}(\bar{x}, t)} \bar{g} d\bar{S} &= \frac{2}{(n+1)\alpha(n+1)t^{n-1}} \int_{B(x, t)} \frac{g(y)}{(t^2 - |y - x|^2)^{1/2}} dy \\ &= \frac{2t\alpha(n)}{(n+1)\alpha(n+1)} \int_{B(x, t)} \frac{g(y)}{(t^2 - |y - x|^2)^{1/2}} dy. \end{aligned}$$

We insert this formula and the similar one with h in place of g into (35) and find

$$\begin{aligned} u(x, t) = & \frac{1}{\gamma_{n+1}} \frac{2\alpha(n)}{(n+1)\alpha(n+1)} \left[\frac{\partial}{\partial t} \left(\frac{1}{t} \frac{\partial}{\partial t} \right)^{\frac{n-2}{2}} \left(t^n \int_{B(x, t)} \frac{g(y)}{(t^2 - |y - x|^2)^{1/2}} dy \right) \right. \\ & \left. + \left(\frac{1}{t} \frac{\partial}{\partial t} \right)^{\frac{n-2}{2}} \left(t^n \int_{B(x, t)} \frac{h(y)}{(t^2 - |y - x|^2)^{1/2}} dy \right) \right]. \end{aligned}$$

Since $\gamma_{n+1} = 1 \cdot 3 \cdot 5 \cdots (n-1)$ and $\alpha(n) = \frac{\pi^{n/2}}{\Gamma(\frac{n+2}{2})}$, we may compute $\gamma_n = 2 \cdot 4 \cdots (n-2) \cdot n$.

Hence the resulting representation formula for even n is

$$(38) \quad \left\{ \begin{aligned} u(x, t) &= \frac{1}{\gamma_n} \left[\left(\frac{\partial}{\partial t} \right) \left(\frac{1}{t} \frac{\partial}{\partial t} \right)^{\frac{n-2}{2}} \left(t^n \int_{B(x, t)} \frac{g(y)}{(t^2 - |y - x|^2)^{1/2}} dy \right) \right. \\ & \quad \left. + \left(\frac{1}{t} \frac{\partial}{\partial t} \right)^{\frac{n-2}{2}} \left(t^n \int_{B(x, t)} \frac{h(y)}{(t^2 - |y - x|^2)^{1/2}} dy \right) \right], \\ & \text{where } n \text{ is even and } \gamma_n = 2 \cdot 4 \cdots (n-2) \cdot n, \end{aligned} \right.$$

for $x \in \mathbb{R}^n$, $t > 0$.

Since $\gamma_2 = 2$, this agrees with Poisson's formula (27) if $n = 2$.

THEOREM 3 (Solution of wave equation in even dimensions). *Assume n is an even integer, $n \geq 2$, and suppose also $g \in C^{m+1}(\mathbb{R}^n)$, $h \in C^m(\mathbb{R}^n)$, for $m = \frac{n+2}{2}$. Define u by (38). Then*

$$(i) \quad u \in C^2(\mathbb{R}^n \times [0, \infty)),$$

$$(ii) \quad u_{tt} - \Delta u = 0 \quad \text{in } \mathbb{R}^n \times (0, \infty),$$

and

$$(iii) \quad \lim_{\substack{(x,t) \rightarrow (x^0,0) \\ x \in \mathbb{R}^n, t > 0}} u(x,t) = g(x^0), \quad \lim_{\substack{(x,t) \rightarrow (x^0,0) \\ x \in \mathbb{R}^n, t > 0}} u_t(x,t) = h(x^0)$$

for each point $x^0 \in \mathbb{R}^n$.

This follows from Theorem 2. Observe, in contrast to formula (31), that to compute $u(x, t)$ for even n we need information on $u = g$, $u_t = h$ on all of $B(x, t)$ and not just on $\partial B(x, t)$.

Huygens' principle. Comparing (31) and (38), we observe that if n is odd and $n \geq 3$, the data g and h at a given point $x \in \mathbb{R}^n$ affect the solution u only on the boundary $\{(y, t) \mid t > 0, |x - y| = t\}$ of the cone $C = \{(y, t) \mid t > 0, |x - y| < t\}$. On the other hand, if n is even, the data g and h affect u within all of C . In other words, a "disturbance" originating at x propagates along a sharp wavefront in odd dimensions, but in even dimensions it continues to have effects even after the leading edge of the wavefront passes. This is Huygens' principle.

2.4.2. Nonhomogeneous problem.

We next investigate the initial-value problem for the nonhomogeneous wave equation

$$(39) \quad \begin{cases} u_{tt} - \Delta u = f & \text{in } \mathbb{R}^n \times (0, \infty) \\ u = 0, \quad u_t = 0 & \text{on } \mathbb{R}^n \times \{t = 0\}. \end{cases}$$

Motivated by Duhamel's principle (introduced earlier in §2.3.1), we define $u = u(x, t; s)$ to be the solution of

$$(40_s) \quad \begin{cases} u_{tt}(\cdot; s) - \Delta u(\cdot; s) = 0 & \text{in } \mathbb{R}^n \times (s, \infty) \\ u(\cdot; s) = 0, \quad u_t(\cdot; s) = f(\cdot, s) & \text{on } \mathbb{R}^n \times \{t = s\}. \end{cases}$$

Now set

$$(41) \quad u(x, t) := \int_0^t u(x, t; s) ds \quad (x \in \mathbb{R}^n, t \geq 0).$$

Duhamel's principle asserts this is a solution of

$$(42) \quad \begin{cases} u_{tt} - \Delta u = f & \text{in } \mathbb{R}^n \times (0, \infty) \\ u = 0, \quad u_t = 0 & \text{on } \mathbb{R}^n \times \{t = 0\}. \end{cases}$$

THEOREM 4 (Solution of nonhomogeneous wave equation). *Assume that $n \geq 2$ and $f \in C^{[n/2]+1}(\mathbb{R}^n \times [0, \infty))$. Define u by (41). Then*

(i) $u \in C^2(\mathbb{R}^n \times [0, \infty))$,

(ii) $u_{tt} - \Delta u = f$ in $\mathbb{R}^n \times (0, \infty)$,

and

(iii) $\lim_{\substack{(x,t) \rightarrow (x^0,0) \\ x \in \mathbb{R}^n, t > 0}} u(x,t) = 0, \quad \lim_{\substack{(x,t) \rightarrow (x^0,0) \\ x \in \mathbb{R}^n, t > 0}} u_t(x,t) = 0$ for each point $x^0 \in \mathbb{R}^n$.

Proof. 1. If n is odd, $[\frac{n}{2}] + 1 = \frac{n+1}{2}$. According to Theorem 2, we have $u(\cdot, \cdot; s) \in C^2(\mathbb{R}^n \times [s, \infty))$ for each $s \geq 0$, and so $u \in C^2(\mathbb{R}^n \times [0, \infty))$. If n is even, $[\frac{n}{2}] + 1 = \frac{n+2}{2}$. Hence $u \in C^2(\mathbb{R}^n \times [0, \infty))$, according to Theorem 3.

2. We then compute

$$u_t(x, t) = u(x, t; t) + \int_0^t u_t(x, t; s) ds = \int_0^t u_t(x, t; s) ds,$$

$$u_{tt}(x, t) = u_t(x, t; t) + \int_0^t u_{tt}(x, t; s) ds = f(x, t) + \int_0^t u_{tt}(x, t; s) ds.$$

Furthermore

$$\Delta u(x, t) = \int_0^t \Delta u(x, t; s) ds = \int_0^t u_{tt}(x, t; s) ds.$$

Thus

$$u_{tt}(x, t) - \Delta u(x, t) = f(x, t) \quad (x \in \mathbb{R}^n, t > 0),$$

and clearly $u(x, 0) = u_t(x, 0) = 0$ for $x \in \mathbb{R}^n$. □

The solution of the general nonhomogeneous problem is consequently the sum of the solution of (11) (given by formulas (8), (31) or (38)) and the solution of (42) (given by (41)).

Examples. (i) Let us work out explicitly how to solve (42) for $n = 1$. In this case d'Alembert's formula (8) gives

$$u(x, t; s) = \frac{1}{2} \int_{x-t+s}^{x+t-s} f(y, s) dy, \quad u(x, t) = \frac{1}{2} \int_0^t \int_{x-t+s}^{x+t-s} f(y, s) dy ds.$$

That is,

$$(43) \quad u(x, t) = \frac{1}{2} \int_0^t \int_{x-s}^{x+s} f(y, t-s) dy ds \quad (x \in \mathbb{R}, t \geq 0).$$

(ii) For $n = 3$, Kirchhoff's formula (22) implies

$$u(x, t; s) = (t - s) \int_{\partial B(x, t-s)} f(y, s) dS,$$

so that

$$\begin{aligned} u(x, t) &= \int_0^t (t - s) \left(\int_{\partial B(x, t-s)} f(y, s) dS \right) ds \\ &= \frac{1}{4\pi} \int_0^t \int_{\partial B(x, t-s)} \frac{f(y, s)}{(t - s)} dS ds \\ &= \frac{1}{4\pi} \int_0^t \int_{\partial B(x, r)} \frac{f(y, t - r)}{r} dS dr. \end{aligned}$$

Therefore

$$(44) \quad u(x, t) = \frac{1}{4\pi} \int_{B(x, t)} \frac{f(y, t - |y - x|)}{|y - x|} dy \quad (x \in \mathbb{R}^3, t \geq 0)$$

solves (42) for $n = 3$. The integrand on the right is called a *retarded potential*. \square

2.4.3. Energy methods.


The explicit formulas (31) and (38) demonstrate the necessity of making more and more smoothness assumptions upon the data g and h to ensure the existence of a C^2 solution of the wave equation for larger and larger n . This suggests that perhaps some other way of measuring the size and smoothness of functions may be more appropriate. Indeed we will see in this subsection that the wave equation is nicely behaved (for all n) with respect to certain integral “energy” norms.

a. Uniqueness. Let $U \subset \mathbb{R}^n$ be a bounded, open set with a smooth boundary ∂U , and as usual set $U_T = U \times (0, T]$, $\Gamma_T = \bar{U}_T - U_T$, where $T > 0$.

We are interested in the initial/boundary-value problem

$$(45) \quad \begin{cases} u_{tt} - \Delta u = f & \text{in } U_T \\ u = g & \text{on } \Gamma_T \\ u_t = h & \text{on } U \times \{t = 0\}. \end{cases}$$

THEOREM 5 (Uniqueness for wave equation). *There exists at most one function $u \in C^2(\bar{U}_T)$ solving (45).*



Cone of dependence

Proof. If \tilde{u} is another such solution, then $w := u - \tilde{u}$ solves

$$\begin{cases} w_{tt} - \Delta w = 0 & \text{in } U_T \\ w = 0 & \text{on } \Gamma_T \\ w_t = 0 & \text{on } U \times \{t = 0\}. \end{cases}$$

Define the “energy”

$$E(t) := \frac{1}{2} \int_U w_t^2(x, t) + |Dw(x, t)|^2 dx \quad (0 \leq t \leq T).$$

We compute

$$\begin{aligned} \dot{E}(t) &= \int_U w_t w_{tt} + Dw \cdot Dw_t dx \quad \left(\dot{\cdot} = \frac{d}{dt} \right) \\ &= \int_U w_t (w_{tt} - \Delta w) dx = 0. \end{aligned}$$

There is no boundary term since $w = 0$, and hence $w_t = 0$, on $\partial U \times [0, T]$. Thus for all $0 \leq t \leq T$, $E(t) = E(0) = 0$, and so $w_t, Dw \equiv 0$ within U_T . Since $w \equiv 0$ on $U \times \{t = 0\}$, we conclude $w = u - \tilde{u} \equiv 0$ in U_T . \square

b. Domain of dependence. As another illustration of energy methods, let us examine again the domain of dependence of solutions to the wave equation in all of space. For this, suppose $u \in C^2$ solves

$$u_{tt} - \Delta u = 0 \quad \text{in } \mathbb{R}^n \times (0, \infty).$$

Fix $x_0 \in \mathbb{R}^n$, $t_0 > 0$ and consider the *backwards wave cone* with apex (x_0, t_0)

$$K(x_0, t_0) := \{(x, t) \mid 0 \leq t \leq t_0, |x - x_0| \leq t_0 - t\}.$$

THEOREM 6 (Finite propagation speed). *If $u \equiv u_t \equiv 0$ on $B(x_0, t_0) \times \{t = 0\}$, then $u \equiv 0$ within the cone $K(x_0, t_0)$.*

In particular, we see that any “disturbance” originating outside $B(x_0, t_0)$ has no effect on the solution within $K(x_0, t_0)$ and consequently has finite propagation speed. We already know this from the representation formulas (31) and (38), at least assuming $g = u$ and $h = u_t$ on $\mathbb{R}^n \times \{t = 0\}$ are sufficiently smooth. The point is that energy methods provide a *much* simpler proof.

Proof. Define the local energy

$$e(t) := \frac{1}{2} \int_{B(x_0, t_0-t)} u_t^2(x, t) + |Du(x, t)|^2 dx \quad (0 \leq t \leq t_0).$$

Then

$$\begin{aligned} \dot{e}(t) &= \int_{B(x_0, t_0-t)} u_t u_{tt} + Du \cdot Du_t dx - \frac{1}{2} \int_{\partial B(x_0, t_0-t)} u_t^2 + |Du|^2 dS \\ &= \int_{B(x_0, t_0-t)} u_t (u_{tt} - \Delta u) dx \\ (46) \quad &+ \int_{\partial B(x_0, t_0-t)} \frac{\partial u}{\partial \nu} u_t dS - \frac{1}{2} \int_{\partial B(x_0, t_0-t)} u_t^2 + |Du|^2 dS \\ &= \int_{\partial B(x_0, t_0-t)} \frac{\partial u}{\partial \nu} u_t - \frac{1}{2} u_t^2 - \frac{1}{2} |Du|^2 dS. \end{aligned}$$

Now

$$(47) \quad \left| \frac{\partial u}{\partial \nu} u_t \right| \leq |u_t| |Du| \leq \frac{1}{2} u_t^2 + \frac{1}{2} |Du|^2,$$

by the Cauchy–Schwarz and Cauchy inequalities (§B.2). Inserting (47) into (46), we find $\dot{e}(t) \leq 0$; and so $e(t) \leq e(0) = 0$ for all $0 \leq t \leq t_0$. Thus $u_t, Du \equiv 0$, and consequently $u \equiv 0$ within the cone $K(x_0, t_0)$. \square

A generalization of this proof to more complicated geometry appears later, in §7.2.4. See also §12.1 for a similar calculation for a nonlinear wave equation.

2.5. PROBLEMS

In the following exercises, all given functions are assumed smooth, unless otherwise stated.

1. Write down an explicit formula for a function u solving the initial-value problem

$$\begin{cases} u_t + b \cdot Du + cu = 0 & \text{in } \mathbb{R}^n \times (0, \infty) \\ u = g & \text{on } \mathbb{R}^n \times \{t = 0\}. \end{cases}$$

Here $c \in \mathbb{R}$ and $b \in \mathbb{R}^n$ are constants.

2. Prove that Laplace's equation $\Delta u = 0$ is rotation invariant; that is, if O is an orthogonal $n \times n$ matrix and we define

$$v(x) := u(Ox) \quad (x \in \mathbb{R}^n),$$

then $\Delta v = 0$.

3. Modify the proof of the mean-value formulas to show for $n \geq 3$ that

$$u(0) = \int_{\partial B(0,r)} g dS + \frac{1}{n(n-2)\alpha(n)} \int_{B(0,r)} \left(\frac{1}{|x|^{n-2}} - \frac{1}{r^{n-2}} \right) f dx,$$

provided

$$\begin{cases} -\Delta u = f & \text{in } B^0(0,r) \\ u = g & \text{on } \partial B(0,r). \end{cases}$$

4. Give a direct proof that if $u \in C^2(U) \cap C(\bar{U})$ is harmonic within a bounded open set U , then

$$\max_{\bar{U}} u = \max_{\partial U} u.$$

(Hint: Define $u_\varepsilon := u + \varepsilon|x|^2$ for $\varepsilon > 0$, and show u_ε cannot attain its maximum over \bar{U} at an interior point.)

5. We say $v \in C^2(\bar{U})$ is *subharmonic* if

$$-\Delta v \leq 0 \quad \text{in } U.$$

- (a) Prove for subharmonic v that

$$v(x) \leq \int_{B(x,r)} v dy \quad \text{for all } B(x,r) \subset U.$$

- (b) Prove that therefore $\max_{\bar{U}} v = \max_{\partial U} v$.
- (c) Let $\phi : \mathbb{R} \rightarrow \mathbb{R}$ be smooth and convex. Assume u is harmonic and $v := \phi(u)$. Prove v is subharmonic.
- (d) Prove $v := |Du|^2$ is subharmonic, whenever u is harmonic.

6. Let U be a bounded, open subset of \mathbb{R}^n . Prove that there exists a constant C , depending only on U , such that

$$\max_{\bar{U}} |u| \leq C(\max_{\partial U} |g| + \max_{\bar{U}} |f|)$$

whenever u is a smooth solution of

$$\begin{cases} -\Delta u = f & \text{in } U \\ u = g & \text{on } \partial U. \end{cases}$$

(Hint: $-\Delta(u + \frac{|x|^2}{2n}\lambda) \leq 0$, for $\lambda := \max_{\bar{U}} |f|$.)

7. Use Poisson's formula for the ball to prove

$$r^{n-2} \frac{r - |x|}{(r + |x|)^{n-1}} u(0) \leq u(x) \leq r^{n-2} \frac{r + |x|}{(r - |x|)^{n-1}} u(0)$$

whenever u is positive and harmonic in $B^0(0, r)$. This is an explicit form of Harnack's inequality.

8. Prove Theorem 15 in §2.2.4. (Hint: Since $u \equiv 1$ solves (44) for $g \equiv 1$, the theory automatically implies

$$\int_{\partial B(0,1)} K(x, y) dS(y) = 1$$

for each $x \in B^0(0, 1)$.)

9. Let u be the solution of

$$\begin{cases} \Delta u = 0 & \text{in } \mathbb{R}_+^n \\ u = g & \text{on } \partial \mathbb{R}_+^n \end{cases}$$

given by Poisson's formula for the half-space. Assume g is bounded and $g(x) = |x|$ for $x \in \partial \mathbb{R}_+^n$, $|x| \leq 1$. Show Du is *not* bounded near $x = 0$. (Hint: Estimate $\frac{u(\lambda e_n) - u(0)}{\lambda}$.)

10. (Reflection principle)

(a) Let U^+ denote the open half-ball $\{x \in \mathbb{R}^n \mid |x| < 1, x_n > 0\}$. Assume $u \in C^2(\bar{U}^+)$ is harmonic in U^+ , with $u = 0$ on $\partial U^+ \cap \{x_n = 0\}$. Set

$$v(x) := \begin{cases} u(x) & \text{if } x_n \geq 0 \\ -u(x_1, \dots, x_{n-1}, -x_n) & \text{if } x_n < 0 \end{cases}$$

for $x \in U = B^0(0, 1)$. Prove $v \in C^2(U)$ and thus v is harmonic within U .

- (b) Now assume only that $u \in C^2(U^+) \cap C(\overline{U^+})$. Show that v is harmonic within U . (Hint: Use Poisson's formula for the ball.)
11. (Kelvin transform for Laplace's equation) The *Kelvin transform* $\mathcal{K}u = \bar{u}$ of a function $u : \mathbb{R}^n \rightarrow \mathbb{R}$ is

$$\bar{u}(x) := u(\bar{x})|\bar{x}|^{n-2} = u(x/|x|)|x|^{2-n} \quad (x \neq 0),$$

where $\bar{x} = x/|x|^2$. Show that if u is harmonic, then so is \bar{u} .

(Hint: First show that $D_x \bar{x}(D_x \bar{x})^T = |\bar{x}|^4 I$. The mapping $x \rightarrow \bar{x}$ is *conformal*, meaning angle preserving.)

12. Suppose u is smooth and solves $u_t - \Delta u = 0$ in $\mathbb{R}^n \times (0, \infty)$.
- (a) Show $u_\lambda(x, t) := u(\lambda x, \lambda^2 t)$ also solves the heat equation for each $\lambda \in \mathbb{R}$.
- (b) Use (a) to show $v(x, t) := x \cdot Du(x, t) + 2tu_t(x, t)$ solves the heat equation as well.
13. Assume $n = 1$ and $u(x, t) = v(\frac{x}{\sqrt{t}})$.

- (a) Show

$$u_t = u_{xx}$$

if and only if

$$(*) \quad v'' + \frac{z}{2}v' = 0.$$

Show that the general solution of (*) is

$$v(z) = c \int_0^z e^{-s^2/4} ds + d.$$

- (b) Differentiate $u(x, t) = v(\frac{x}{\sqrt{t}})$ with respect to x and select the constant c properly, to obtain the fundamental solution Φ for $n = 1$. Explain why this procedure produces the fundamental solution. (Hint: What is the initial condition for u ?)
14. Write down an explicit formula for a solution of

$$\begin{cases} u_t - \Delta u + cu = f & \text{in } \mathbb{R}^n \times (0, \infty) \\ u = g & \text{on } \mathbb{R}^n \times \{t = 0\}, \end{cases}$$

where $c \in \mathbb{R}$.

15. Given $g : [0, \infty) \rightarrow \mathbb{R}$, with $g(0) = 0$, derive the formula

$$u(x, t) = \frac{x}{\sqrt{4\pi}} \int_0^t \frac{1}{(t-s)^{3/2}} e^{\frac{-x^2}{4(t-s)}} g(s) ds$$

for a solution of the initial/boundary-value problem

$$\begin{cases} u_t - u_{xx} = 0 & \text{in } \mathbb{R}_+ \times (0, \infty) \\ u = 0 & \text{on } \mathbb{R}_+ \times \{t = 0\} \\ u = g & \text{on } \{x = 0\} \times [0, \infty). \end{cases}$$

(Hint: Let $v(x, t) := u(x, t) - g(t)$ and extend v to $\{x < 0\}$ by odd reflection.)

16. Give a direct proof that if U is bounded and $u \in C_1^2(U_T) \cap C(\bar{U}_T)$ solves the heat equation, then

$$\max_{\bar{U}_T} u = \max_{\Gamma_T} u.$$

(Hint: Define $u_\varepsilon := u - \varepsilon t$ for $\varepsilon > 0$, and show u_ε cannot attain its maximum over \bar{U}_T at a point in U_T .)

17. We say $v \in C_1^2(U_T)$ is a *subsolution* of the heat equation if

$$v_t - \Delta v \leq 0 \quad \text{in } U_T.$$

- (a) Prove for a subsolution v that

$$v(x, t) \leq \frac{1}{4r^n} \iint_{E(x, t; r)} v(y, s) \frac{|x - y|^2}{(t - s)^2} dy ds$$

for all $E(x, t; r) \subset U_T$.

- (b) Prove that therefore $\max_{\bar{U}_T} v = \max_{\Gamma_T} v$.
 (c) Let $\phi : \mathbb{R} \rightarrow \mathbb{R}$ be smooth and convex. Assume u solves the heat equation and $v := \phi(u)$. Prove v is a subsolution.
 (d) Prove $v := |Du|^2 + u_t^2$ is a subsolution, whenever u solves the heat equation.
18. (Stokes' rule) Assume u solves the initial-value problem

$$\begin{cases} u_{tt} - \Delta u = 0 & \text{in } \mathbb{R}^n \times (0, \infty) \\ u = 0, \quad u_t = h & \text{on } \mathbb{R}^n \times \{t = 0\}. \end{cases}$$

Show that $v := u_t$ solves

$$\begin{cases} v_{tt} - \Delta v = 0 & \text{in } \mathbb{R}^n \times (0, \infty) \\ v = h, \quad v_t = 0 & \text{on } \mathbb{R}^n \times \{t = 0\}. \end{cases}$$

This is *Stokes' rule*.

19. (a) Show the general solution of the PDE $u_{xy} = 0$ is

$$u(x, y) = F(x) + G(y)$$

for arbitrary functions F, G .

- (b) Using the change of variables $\xi = x + t$, $\eta = x - t$, show $u_{tt} - u_{xx} = 0$ if and only if $u_{\xi\eta} = 0$.
- (c) Use (a) and (b) to rederive d'Alembert's formula.
- (d) Under what conditions on the initial data g, h is the solution u a right-moving wave? A left-moving wave?
20. Assume that for some attenuation function $\alpha = \alpha(r)$ and delay function $\beta = \beta(r) \geq 0$, there exist for *all* profiles ϕ solutions of the wave equation in $(\mathbb{R}^n - \{0\}) \times \mathbb{R}$ having the form

$$u(x, t) = \alpha(r)\phi(t - \beta(r)).$$

Here $r = |x|$ and we assume $\beta(0) = 0$.

Show that this is possible only if $n = 1$ or 3 , and compute the form of the functions α, β .

(T. Morley, SIAM Review 27 (1985), 69–71)

21. (a) Assume $\mathbf{E} = (E^1, E^2, E^3)$ and $\mathbf{B} = (B^1, B^2, B^3)$ solve Maxwell's equations

$$\begin{cases} \mathbf{E}_t = \text{curl } \mathbf{B}, & \mathbf{B}_t = -\text{curl } \mathbf{E} \\ \text{div } \mathbf{B} = \text{div } \mathbf{E} = 0. \end{cases}$$

Show

$$\mathbf{E}_{tt} - \Delta \mathbf{E} = 0, \quad \mathbf{B}_{tt} - \Delta \mathbf{B} = 0.$$

- (b) Assume that $\mathbf{u} = (u^1, u^2, u^3)$ solves the evolution equations of linear elasticity

$$\mathbf{u}_{tt} - \mu \Delta \mathbf{u} - (\lambda + \mu) D(\text{div } \mathbf{u}) = \mathbf{0} \quad \text{in } \mathbb{R}^3 \times (0, \infty).$$

Show $w := \text{div } \mathbf{u}$ and $\mathbf{w} := \text{curl } \mathbf{u}$ each solve wave equations, but with differing speeds of propagation.

22. Let u denote the density of particles moving to the right with speed one along the real line and let v denote the density of particles moving to the left with speed one. If at rate $d > 0$ right-moving particles randomly become left-moving, and vice versa, we have the system of PDE

$$\begin{cases} u_t + u_x = d(v - u) \\ v_t - v_x = d(u - v). \end{cases}$$

Show that both $w := u$ and $w := v$ solve the telegraph equation

$$w_{tt} + 2dw_t - w_{xx} = 0.$$

23. Let S denote the square lying in $\mathbb{R} \times (0, \infty)$ with corners at the points $(0, 1), (1, 2), (0, 3), (-1, 2)$. Define

$$f(x, t) := \begin{cases} -1 & \text{for } (x, t) \in S \cap \{t > x + 2\} \\ 1 & \text{for } (x, t) \in S \cap \{t < x + 2\} \\ 0 & \text{otherwise.} \end{cases}$$

Assume u solves

$$\begin{cases} u_{tt} - u_{xx} = f & \text{in } \mathbb{R} \times (0, \infty) \\ u = 0, u_t = 0 & \text{on } \mathbb{R} \times \{t = 0\}. \end{cases}$$

Describe the shape of u for times $t > 3$.

(J. G. Kingston, SIAM Review 30 (1988), 645–649)

24. (Equipartition of energy) Let u solve the initial-value problem for the wave equation in one dimension:

$$\begin{cases} u_{tt} - u_{xx} = 0 & \text{in } \mathbb{R} \times (0, \infty) \\ u = g, u_t = h & \text{on } \mathbb{R} \times \{t = 0\}. \end{cases}$$

Suppose g, h have compact support. The *kinetic energy* is $k(t) := \frac{1}{2} \int_{-\infty}^{\infty} u_t^2(x, t) dx$ and the *potential energy* is $p(t) := \frac{1}{2} \int_{-\infty}^{\infty} u_x^2(x, t) dx$. Prove

- (a) $k(t) + p(t)$ is constant in t ,
 (b) $k(t) = p(t)$ for all large enough times t .

2.6. REFERENCES

- Section 2.2 A good source for more on Laplace's and Poisson's equations is Gilbarg–Trudinger [G-T, Chapters 2-4]. The proof of analyticity is from Mikhailov [M]. J. Cooper helped me with Green's functions.
- Section 2.3 See John [J2, Chapter 7] or Friedman [Fr1] for further information concerning the heat equation. Theorem 3 is due to N. Watson (Proc. London Math. Society 26 (1973), 385–417), as is the proof of Theorem 4. Theorem 6 is taken from John [J2], and Theorem 8 follows Mikhailov [M]. Theorem 11 is from Payne [Pa, §2.3].
- Section 2.4 See Antman (Amer. Math. Monthly 87 (1980), 359–370) for a careful derivation of the one-dimensional wave equation as a model for a vibrating string. The solution of the wave equation presented here follows Folland [F1], Strauss [St2].
- Section 2.5 J. Goldstein contributed Problem 24.

NONLINEAR FIRST-ORDER PDE

- 3.1 Complete integrals, envelopes
- 3.2 Characteristics
- 3.3 Introduction to Hamilton–Jacobi equations
- 3.4 Introduction to conservation laws
- 3.5 Problems
- 3.6 References

In this chapter we investigate general nonlinear first-order partial differential equations of the form

$$F(Du, u, x) = 0,$$

where $x \in U$ and U is an open subset of \mathbb{R}^n . Here

$$F : \mathbb{R}^n \times \mathbb{R} \times \bar{U} \rightarrow \mathbb{R}$$

is given, and $u : \bar{U} \rightarrow \mathbb{R}$ is the unknown, $u = u(x)$.

NOTATION. Let us write

$$F = F(p, z, x) = F(p_1, \dots, p_n, z, x_1, \dots, x_n)$$

for $p \in \mathbb{R}^n$, $z \in \mathbb{R}$, $x \in U$. Thus “ p ” is the name of the variable for which we substitute the gradient $Du(x)$, and “ z ” is the variable for which we substitute $u(x)$. We also assume hereafter that F is smooth and set

$$\begin{cases} D_p F = (F_{p_1}, \dots, F_{p_n}) \\ D_z F = F_z \\ D_x F = (F_{x_1}, \dots, F_{x_n}). \end{cases}$$

□

We are concerned with discovering solutions u of the PDE $F(Du, u, x) = 0$ in U , usually subject to the boundary condition

$$u = g \quad \text{on } \Gamma,$$

where Γ is some given subset of ∂U and $g : \Gamma \rightarrow \mathbb{R}$ is prescribed.

Nonlinear first-order partial differential equations arise in a variety of physical theories, primarily in dynamics (to generate canonical transformations), continuum mechanics (to record conservation of mass, momentum, energy, etc.) and optics (to describe wavefronts). Although the strong nonlinearity generally precludes our deriving any simple formulas for solutions, we can, remarkably, often employ calculus to glean fairly detailed information about solutions. Such techniques, discussed in §§3.1 and 3.2, are typically only local. In §§3.3 and 3.4 we will for the important cases of Hamilton–Jacobi equations and conservation laws derive certain global representation formulas for appropriately defined weak solutions.

3.1. COMPLETE INTEGRALS, ENVELOPES

3.1.1. Complete integrals.

We begin our analysis of the nonlinear first-order PDE

$$(1) \quad F(Du, u, x) = 0$$

by describing some simple classes of solutions and then learning how to build from them more complicated solutions.

Suppose first $A \subset \mathbb{R}^n$ is an open set. Assume for each parameter $a = (a_1, \dots, a_n) \in A$ we have a C^2 solution $u = u(x; a)$ of the PDE (1).

NOTATION. We write

$$(2) \quad (D_a u, D_{x_a}^2 u) := \begin{pmatrix} u_{a_1} & u_{x_1 a_1} & \cdots & u_{x_n a_1} \\ \vdots & \vdots & \ddots & \vdots \\ u_{a_n} & u_{x_1 a_n} & \cdots & u_{x_n a_n} \end{pmatrix}_{n \times (n+1)}.$$

DEFINITION. A C^2 function $u = u(x; a)$ is called a complete integral in $U \times A$ provided

$$(i) \quad u(x; a) \text{ solves the PDE (1) for each } a \in A$$

and

$$(ii) \quad \text{rank}(D_a u, D_{x_a}^2 u) = n \quad (x \in U, a \in A).$$

Interpretation. Condition (ii) ensures $u(x; a)$ “depends on all the n independent parameters a_1, \dots, a_n ”. To see this, suppose $B \subset \mathbb{R}^{n-1}$ is open, and for each $b \in B$ assume $v = v(x; b)$ ($x \in U$) is a solution of (1). Suppose also there exists a C^1 mapping $\psi : A \rightarrow B$, $\psi = (\psi^1, \dots, \psi^{n-1})$, such that

$$(3) \quad u(x; a) = v(x; \psi(a)) \quad (x \in U, a \in A).$$

That is, we are supposing the function $u(x; a)$ “really depends only on the $n - 1$ parameters b_1, \dots, b_{n-1} ”. But then

$$u_{x_i a_j}(x; a) = \sum_{k=1}^{n-1} v_{x_i b_k}(x; \psi(a)) \psi_{a_j}^k(a) \quad (i, j = 1, \dots, n).$$

Consequently

$$\det(D_{x a}^2 u) = \sum_{k_1, \dots, k_n=1}^{n-1} v_{x_1 b_{k_1}} \dots v_{x_n b_{k_n}} \det \begin{pmatrix} \psi_{a_1}^{k_1} & \dots & \psi_{a_n}^{k_1} \\ & \ddots & \\ \psi_{a_1}^{k_n} & \dots & \psi_{a_n}^{k_n} \end{pmatrix} = 0,$$

since for each choice of $k_1, \dots, k_n \in \{1, \dots, n-1\}$, at least two rows in the corresponding matrix are equal. As

$$u_{a_j}(x; a) = \sum_{k=1}^{n-1} v_{b_k}(x; \psi(a)) \psi_{a_j}^k(a) \quad (j = 1, \dots, n),$$

a similar argument shows that the determinant of each $n \times n$ submatrix of $(D_a u, D_{x a}^2 u)$ equals zero, and thus this matrix has rank strictly less than n .

Example 1. *Clairaut's equation* from differential geometry is the PDE

$$(4) \quad x \cdot Du + f(Du) = u,$$

where $f : \mathbb{R}^n \rightarrow \mathbb{R}$ is given. A complete integral is

$$(5) \quad u(x; a) = a \cdot x + f(a) \quad (x \in U)$$

for $a \in \mathbb{R}^n$. □

Example 2. The *eikonal** equation from geometric optics is the PDE

$$(6) \quad |Du| = 1.$$

A complete integral is

$$(7) \quad u(x; a, b) = a \cdot x + b \quad (x \in U)$$

for $x \in U$, $a \in \partial B(0, 1)$, $b \in \mathbb{R}$. □

* $\epsilon\iota\kappa\acute{\omega}\nu$ = image (Greek).

Example 3. The *Hamilton–Jacobi equation* from mechanics is in its simplest form the partial differential equation

$$(8) \quad u_t + H(Du) = 0,$$

where $H : \mathbb{R}^n \rightarrow \mathbb{R}$. Here u depends on $x = (x_1, \dots, x_n) \in \mathbb{R}^n$ and $t \in \mathbb{R}$. As before we have set $t = x_{n+1}$ and written $Du = D_x u = (u_{x_1}, \dots, u_{x_n})$. A complete integral is

$$(9) \quad u(x, t; a, b) = a \cdot x - tH(a) + b \quad (x \in \mathbb{R}^n, t \geq 0)$$

where $a \in \mathbb{R}^n$, $b \in \mathbb{R}$. □

3.1.2. New solutions from envelopes.

We next demonstrate how to build more complicated solutions of our nonlinear first-order PDE (1), solutions which depend on an arbitrary *function* of $n - 1$ variables and not just on n parameters. We will construct these new solutions as envelopes of complete integrals or, more generally, of other m -parameter families of solutions.

DEFINITION. Let $u = u(x; a)$ be a C^1 function of $x \in U$, $a \in A$, where $U \subset \mathbb{R}^n$ and $A \subset \mathbb{R}^m$ are open sets. Consider the vector equation

$$(10) \quad D_a u(x; a) = 0 \quad (x \in U, a \in A).$$

Suppose that we can solve (10) for the parameter a as a C^1 function of x ,

$$(11) \quad a = \phi(x);$$

thus

$$(12) \quad D_a u(x; \phi(x)) = 0 \quad (x \in U).$$

We then call

$$(13) \quad v(x) := u(x; \phi(x)) \quad (x \in U)$$

the envelope of the functions $\{u(\cdot; a)\}_{a \in A}$.

By forming envelopes, we can build new solutions of our nonlinear first-order partial differential equation:

THEOREM 1 (Construction of new solutions). Suppose for each $a \in A$ as above that $u = u(\cdot; a)$ solves the partial differential equation (1). Assume further that the envelope v , defined by (12) and (13) above, exists and is a C^1 function. Then v solves (1) as well.

The envelope v defined above is sometimes called a *singular integral* of (1).

Proof. We have $v(x) = u(x; \phi(x))$; and so for $i = 1, \dots, n$

$$\begin{aligned} v_{x_i}(x) &= u_{x_i}(x; \phi(x)) + \sum_{j=1}^m u_{a_j}(x, \phi(x)) \phi_{x_i}^j(x) \\ &= u_{x_i}(x; \phi(x)), \text{ according to (12)}. \end{aligned}$$

Hence for each $x \in U$,

$$F(Dv(x), v(x), x) = F(Du(x; \phi(x)), u(x; \phi(x)), x) = 0. \quad \square$$

The geometric meaning is that for each $x \in U$, the graph of v is tangent to the graph of $u(\cdot; a)$ for $a = \phi(x)$. Thus $Dv = D_x u(\cdot; a)$ at x , for $a = \phi(x)$.

Example 4. Consider the PDE

$$(14) \quad u^2(1 + |Du|^2) = 1.$$

A complete integral is

$$u(x, a) = \pm(1 - |x - a|^2)^{1/2} \quad (|x - a| < 1).$$

We compute

$$D_a u = \frac{\mp(x - a)}{(1 - |x - a|^2)^{1/2}} = 0$$

provided $a = \phi(x) = x$. Thus $v \equiv \pm 1$ are singular integrals of (14). \square

To generate still more solutions of the PDE (1) from a complete integral, we vary the above construction. Choose any open set $A' \subset \mathbb{R}^{n-1}$ and any C^1 function $h : A' \rightarrow \mathbb{R}$, so that the graph of h lies within A . Let us write

$$a = (a_1, \dots, a_n) = (a', a_n) \quad \text{for } a' = (a_1, \dots, a_{n-1}) \in \mathbb{R}^{n-1}.$$

DEFINITION. The general integral (depending on h) is the envelope $v' = v'(x)$ of the functions

$$u'(x; a') = u(x; a', h(a')) \quad (x \in U, a' \in A'),$$

provided this envelope exists and is C^1 .

In other words, in computing the envelope we are now restricting only to parameters a of the form $a = (a', h(a'))$, for some explicit choice of the function h . Thus from a complete integral, which depends upon n arbitrary constants a_1, \dots, a_n , we build (whenever the foregoing construction works) a solution depending on an arbitrary function h of $n - 1$ variables.

Example 5. Let $H(p) = |p|^2$, $h \equiv 0$ in Example 3 above. Then

$$u'(x, t; a) = x \cdot a - t|a|^2.$$

We calculate the envelope by setting $D_a u' = x - 2ta = 0$. Hence $a = \frac{x}{2t}$, and so

$$v'(x, t) = x \cdot \frac{x}{2t} - t \left| \frac{x}{2t} \right|^2 = \frac{|x|^2}{4t} \quad (x \in \mathbb{R}^n, t > 0)$$

solves the Hamilton–Jacobi equation $v'_t + |Dv'|^2 = 0$. \square

Remark. It is tempting to believe that once we can find as above a solution of (1) depending on an arbitrary function h , we have found all the solutions of (1). But this need not be so. Suppose our PDE has the structure

$$F(Du, u, x) = F_1(Du, u, x)F_2(Du, u, x) = 0.$$

If $u_1(x, a)$ is a complete integral of the PDE $F_1(Du, u, x) = 0$ and we succeed in finding a general integral corresponding to any function h , we will still have missed all the solutions of the PDE $F_2(Du, u, x) = 0$.

3.2. CHARACTERISTICS

3.2.1. Derivation of characteristic ODE.

We return to our basic nonlinear first-order PDE

$$(1) \quad F(Du, u, x) = 0 \quad \text{in } U,$$

subject now to the boundary condition

$$(2) \quad u = g \quad \text{on } \Gamma,$$

where $\Gamma \subseteq \partial U$ and $g : \Gamma \rightarrow \mathbb{R}$ are given. We hereafter suppose that F, g are smooth functions.

We develop next the method of *characteristics*, which solves (1), (2) by converting the PDE into an appropriate system of ODE. This is the plan. Suppose u solves (1), (2) and fix any point $x \in U$. We would like to calculate $u(x)$ by finding some curve lying within U , connecting x with a point $x^0 \in \Gamma$ and along which we can compute u . Since (2) says $u = g$ on Γ , we know the value of u at the one end x^0 . We hope then to be able to calculate u all along the curve, and so in particular at x .

Finding the characteristic ODE. How can we choose a path in U so all this will work? Let us suppose the curve is described parametrically by the function $\mathbf{x}(s) = (x^1(s), \dots, x^n(s))$, the parameter s lying in some subinterval $I \subseteq \mathbb{R}$. Assuming u is a C^2 solution of (1), we define also

$$(3) \quad z(s) := u(\mathbf{x}(s)).$$

In addition, set

$$(4) \quad \mathbf{p}(s) := Du(\mathbf{x}(s));$$

that is, $\mathbf{p}(s) = (p^1(s), \dots, p^n(s))$, where

$$(5) \quad p^i(s) = u_{x_i}(\mathbf{x}(s)) \quad (i = 1, \dots, n).$$

So $z(\cdot)$ gives the values of u along the curve and $\mathbf{p}(\cdot)$ records the values of the gradient Du . We must choose the function $\mathbf{x}(\cdot)$ in such a way that we can compute $z(\cdot)$ and $\mathbf{p}(\cdot)$.

For this, first differentiate (5):

$$(6) \quad \dot{p}^i(s) = \sum_{j=1}^n u_{x_i x_j}(\mathbf{x}(s)) \dot{x}^j(s) \quad \left(\dot{\cdot} = \frac{d}{ds} \right).$$

This expression is not too promising, since it involves the second derivatives of u . On the other hand, we can also differentiate the PDE (1) with respect to x_i :

$$(7) \quad \sum_{j=1}^n F_{p_j}(Du, u, x) u_{x_j x_i} + F_z(Du, u, x) u_{x_i} + F_{x_i}(Du, u, x) = 0.$$

We are able to employ this identity to get rid of the second derivative terms in (6), provided we first set

$$(8) \quad \dot{x}^j(s) = F_{p_j}(\mathbf{p}(s), z(s), \mathbf{x}(s)) \quad (j = 1, \dots, n).$$

Assuming now (8) holds, we evaluate (7) at $x = \mathbf{x}(s)$, obtaining thereby from (3), (4) the identity:

$$\begin{aligned} & \sum_{j=1}^n F_{p_j}(\mathbf{p}(s), z(s), \mathbf{x}(s)) u_{x_i x_j}(\mathbf{x}(s)) \\ & + F_z(\mathbf{p}(s), z(s), \mathbf{x}(s)) p^i(s) + F_{x_i}(\mathbf{p}(s), z(s), \mathbf{x}(s)) = 0. \end{aligned}$$

Substitute this expression and (8) into (6):

$$(9) \quad \begin{aligned} \dot{p}^i(s) &= -F_{x_i}(\mathbf{p}(s), z(s), \mathbf{x}(s)) \\ &\quad - F_z(\mathbf{p}(s), z(s), \mathbf{x}(s))p^i(s) \quad (i = 1, \dots, n). \end{aligned}$$

Finally we differentiate (3):

$$(10) \quad \dot{z}(s) = \sum_{j=1}^n u_{x_j}(\mathbf{x}(s))\dot{x}^j(s) = \sum_{j=1}^n p^j(s)F_{p_j}(\mathbf{p}(s), z(s), \mathbf{x}(s)),$$

the second equality holding by (5) and (8).

The characteristic equations. We summarize by rewriting equations (8)–(10) in vector notation:

$$(11) \quad \begin{cases} (a) \ \dot{\mathbf{p}}(s) = -D_x F(\mathbf{p}(s), z(s), \mathbf{x}(s)) - D_z F(\mathbf{p}(s), z(s), \mathbf{x}(s))\mathbf{p}(s) \\ (b) \ \dot{z}(s) = D_p F(\mathbf{p}(s), z(s), \mathbf{x}(s)) \cdot \mathbf{p}(s) \\ (c) \ \dot{\mathbf{x}}(s) = D_p F(\mathbf{p}(s), z(s), \mathbf{x}(s)). \end{cases}$$

Furthermore,

$$(12) \quad F(\mathbf{p}(s), z(s), \mathbf{x}(s)) \equiv 0.$$

These identities hold for $s \in I$.

The important system (11) of $2n + 1$ first-order ODE comprises the *characteristic equations* of the nonlinear first-order PDE (1). The functions $\mathbf{p}(\cdot) = (p^1(\cdot), \dots, p^n(\cdot))$, $z(\cdot)$, $\mathbf{x}(\cdot) = (x^1(\cdot), \dots, x^n(\cdot))$ are called the *characteristics*. We will sometimes refer to $\mathbf{x}(\cdot)$ as the *projected characteristic*: it is the projection of the full characteristics $(\mathbf{p}(\cdot), z(\cdot), \mathbf{x}(\cdot)) \subset \mathbb{R}^{2n+1}$ onto the physical region $U \subset \mathbb{R}^n$.

We have proved:

THEOREM 1 (Structure of characteristic ODE). *Let $u \in C^2(U)$ solve the nonlinear, first-order partial differential equation (1) in U . Assume $\mathbf{x}(\cdot)$ solves the ODE (11)(c), where $\mathbf{p}(\cdot) = Du(\mathbf{x}(\cdot))$, $z(\cdot) = u(\mathbf{x}(\cdot))$. Then $\mathbf{p}(\cdot)$ solves the ODE (11)(a) and $z(\cdot)$ solves the ODE (11)(b), for those s such that $\mathbf{x}(s) \in U$.*

We still need to discover appropriate initial conditions for the system of ODE (11), in order that this theorem be useful. We accomplish this in §3.2.3 below.

Remark. The characteristic ODE are truly remarkable in that they form an exact system of equations for $\mathbf{x}(\cdot)$, $z(\cdot) = u(\mathbf{x}(\cdot))$, and $\mathbf{p}(\cdot) = Du(\mathbf{x}(\cdot))$, whenever u is a smooth solution of the general nonlinear PDE (1). The key step in the derivation is our setting $\dot{\mathbf{x}} = D_p F$, so that—as explained above—the terms involving second derivatives drop out. We thereby obtain *closure* and in particular are not forced to introduce ODE for the second and higher derivatives of u .

3.2.2. Examples.

Before continuing our investigation of the characteristic equations (11), we pause to consider some special cases for which the structure of these equations is especially simple. We illustrate as well how we can sometimes actually solve the characteristic ODE and thereby explicitly compute solutions of certain first-order PDE, subject to appropriate boundary conditions.

a. F linear. Consider first the situation that our PDE (1) is linear and homogeneous and thus has the form

$$(13) \quad F(Du, u, x) = \mathbf{b}(x) \cdot Du(x) + c(x)u(x) = 0 \quad (x \in U).$$

Then $F(p, z, x) = \mathbf{b}(x) \cdot p + c(x)z$, and so

$$D_p F = \mathbf{b}(x).$$

In this circumstance equation (11)(c) becomes

$$(14) \quad \dot{\mathbf{x}}(s) = \mathbf{b}(\mathbf{x}(s)),$$

an ODE involving only the function $\mathbf{x}(\cdot)$. Furthermore equation (11)(b) becomes

$$(15) \quad \dot{z}(s) = \mathbf{b}(\mathbf{x}(s)) \cdot \mathbf{p}(s).$$

Then equation (12) simplifies (15), yielding

$$(16) \quad \dot{z}(s) = -c(\mathbf{x}(s))z(s).$$

This ODE is linear in $z(\cdot)$, once we know the function $\mathbf{x}(\cdot)$ by solving (14). In summary,

$$(17) \quad \begin{cases} \text{(a)} & \dot{\mathbf{x}}(s) = \mathbf{b}(\mathbf{x}(s)) \\ \text{(b)} & \dot{z}(s) = -c(\mathbf{x}(s))z(s) \end{cases}$$

comprise the characteristic equations for the linear, first-order PDE (13). (We will see later that the equation for $\mathbf{p}(\cdot)$ is not needed.) \square

Example 1. We demonstrate the utility of equations (17) by explicitly solving the problem

$$(18) \quad \begin{cases} x_1 u_{x_2} - x_2 u_{x_1} = u & \text{in } U \\ u = g & \text{on } \Gamma, \end{cases}$$

where U is the quadrant $\{x_1 > 0, x_2 > 0\}$ and $\Gamma = \{x_1 > 0, x_2 = 0\} \subseteq \partial U$. The PDE in (18) is of the form (12), for $\mathbf{b} = (-x_2, x_1)$ and $c = -1$. Thus the equations (17) read

$$(19) \quad \begin{cases} \dot{x}^1 = -x^2, & \dot{x}^2 = x^1 \\ \dot{z} = z. \end{cases}$$

Accordingly we have

$$\begin{cases} x^1(s) = x^0 \cos s, & x^2(s) = x^0 \sin s \\ z(s) = z^0 e^s = g(x^0) e^s, \end{cases}$$

where $x^0 \geq 0$, $0 \leq s \leq \frac{\pi}{2}$. Fix a point $(x_1, x_2) \in U$. We select $s > 0$, $x^0 > 0$ so that $(x_1, x_2) = (x^1(s), x^2(s)) = (x^0 \cos s, x^0 \sin s)$. That is, $x^0 = (x_1^2 + x_2^2)^{1/2}$, $s = \arctan\left(\frac{x_2}{x_1}\right)$. Therefore

$$u(x) = u(x^1(s), x^2(s)) = z(s) = g(x^0) e^s = g((x_1^2 + x_2^2)^{1/2}) e^{\arctan\left(\frac{x_2}{x_1}\right)}.$$

□

b. F quasilinear. The partial differential equation (1) is quasilinear should it have the form

$$(20) \quad F(Du, u, x) = \mathbf{b}(x, u(x)) \cdot Du(x) + c(x, u(x)) = 0.$$

In this circumstance $F(p, z, x) = \mathbf{b}(x, z) \cdot p + c(x, z)$, whence

$$D_p F = \mathbf{b}(x, z).$$

Hence equation (11)(c) reads

$$\dot{\mathbf{x}}(s) = \mathbf{b}(\mathbf{x}(s), z(s)),$$

and (11)(b) becomes

$$\dot{z}(s) = \mathbf{b}(\mathbf{x}(s), z(s)) \cdot \mathbf{p}(s) = -c(\mathbf{x}(s), z(s)), \quad \text{by (12).}$$

Consequently

$$(21) \quad \begin{cases} \text{(a)} & \dot{\mathbf{x}}(s) = \mathbf{b}(\mathbf{x}(s), z(s)) \\ \text{(b)} & \dot{z}(s) = -c(\mathbf{x}(s), z(s)) \end{cases}$$

are the characteristic equations for the quasilinear first-order PDE (20). (Once again we do not require the equation for $\mathbf{p}(\cdot)$) □

Example 2. The characteristic ODE (21) are in general difficult to solve, and so we work out in this example the simpler case of a boundary-value problem for a semilinear PDE:

$$(22) \quad \begin{cases} u_{x_1} + u_{x_2} = u^2 & \text{in } U \\ u = g & \text{on } \Gamma. \end{cases}$$

Now U is the half-space $\{x_2 > 0\}$ and $\Gamma = \{x_2 = 0\} = \partial U$. Here $\mathbf{b} = (1, 1)$ and $c = -z^2$. Then (21) becomes

$$\begin{cases} \dot{x}^1 = 1, \dot{x}^2 = 1 \\ \dot{z} = z^2. \end{cases}$$

Consequently

$$\begin{cases} x^1(s) = x^0 + s, x^2(s) = s \\ z(s) = \frac{z^0}{1-sz^0} = \frac{g(x^0)}{1-sg(x^0)}, \end{cases}$$

where $x^0 \in \mathbb{R}$, $s \geq 0$, provided the denominator is not zero.

Fix a point $(x_1, x_2) \in U$. We select $s > 0$ and $x^0 \in \mathbb{R}$ so that $(x_1, x_2) = (x^1(s), x^2(s)) = (x^0 + s, s)$; that is, $x^0 = x_1 - x_2$, $s = x_2$. Then

$$u(x) = u(x^1(s), x^2(s)) = z(s) = \frac{g(x^0)}{1-sg(x^0)} = \frac{g(x_1 - x_2)}{1 - x_2g(x_1 - x_2)}.$$

This solution of course makes sense only if $1 - x_2g(x_1 - x_2) \neq 0$. \square

c. F fully nonlinear. In the general case, we must integrate the full characteristic equations (11), if possible.

Example 3. Consider the fully nonlinear problem

$$(23) \quad \begin{cases} u_{x_1}u_{x_2} = u & \text{in } U \\ u = x_2^2 & \text{on } \Gamma, \end{cases}$$

where $U = \{x_1 > 0\}$, $\Gamma = \{x_1 = 0\} = \partial U$. Here $F(p, z, x) = p_1p_2 - z$, and hence the characteristic ODE (11) become

$$\begin{cases} \dot{p}^1 = p^1, \dot{p}^2 = p^2 \\ \dot{z} = 2p^1p^2 \\ \dot{x}^1 = p^2, \dot{x}^2 = p^1. \end{cases}$$

We integrate these equations to find

$$\begin{cases} x^1(s) = p_2^0(e^s - 1), x^2(s) = x^0 + p_1^0(e^s - 1) \\ z(s) = z^0 + p_1^0p_2^0(e^{2s} - 1) \\ p^1(s) = p_1^0e^s, p^2(s) = p_2^0e^s, \end{cases}$$

where $x^0 \in \mathbb{R}$, $s \in \mathbb{R}$, and $z^0 = (x^0)^2$.

We must determine $p^0 = (p_1^0, p_2^0)$. Since $u = x_2^2$ on Γ , $p_2^0 = u_{x_2}(0, x^0) = 2x^0$. Furthermore the PDE $u_{x_1}u_{x_2} = u$ itself implies $p_1^0p_2^0 = z^0 = (x^0)^2$, and so $p_1^0 = \frac{x^0}{2}$. Consequently the formulas above become

$$\begin{cases} x^1(s) = 2x^0(e^s - 1), x^2(s) = \frac{x^0}{2}(e^s + 1) \\ z(s) = (x^0)^2 e^{2s} \\ p^1(s) = \frac{x^0}{2} e^s, p^2(s) = 2x^0 e^s. \end{cases}$$

Fix a point $(x_1, x_2) \in U$. Select s and x^0 so that $(x_1, x_2) = (x^1(s), x^2(s)) = (2x^0(e^s - 1), \frac{x^0}{2}(e^s + 1))$. This equality implies $x^0 = \frac{4x_2 - x_1}{4}$, $e^s = \frac{x_1 + 4x_2}{4x_2 - x_1}$, and so

$$u(x) = u(x^1(s), x^2(s)) = z(s) = (x^0)^2 e^{2s} = \frac{(x_1 + 4x_2)^2}{16}.$$

□

3.2.3. Boundary conditions.

We return now to developing the general theory and intend in the section following to invoke the characteristic ODE (11) actually to solve the boundary-value problem (1), (2), at least in a small region near an appropriate portion Γ of ∂U .

a. Straightening the boundary. To simplify subsequent calculations, it is convenient first to change variables, so as to “flatten out” part of the boundary ∂U . To accomplish this, we first fix any point $x^0 \in \partial U$. Then utilizing the notation from §C.1, we find smooth mappings $\Phi, \Psi : \mathbb{R}^n \rightarrow \mathbb{R}^n$ such that $\Psi = \Phi^{-1}$ and Φ straightens out ∂U near x^0 . (See the illustration in §C.1.)

Given any function $u : U \rightarrow \mathbb{R}$, let us write $V := \Phi(U)$ and set

$$(24) \quad v(y) := u(\Psi(y)) \quad (y \in V).$$

Then

$$(25) \quad u(x) = v(\Phi(x)) \quad (x \in U).$$

Now suppose that u is a C^1 solution of our boundary-value problem (1), (2) in U . What PDE does v then satisfy in V ?

According to (25), we see

$$u_{x_i}(x) = \sum_{k=1}^n v_{y_k}(\Phi(x)) \Phi_{x_i}^k(x) \quad (i = 1, \dots, n);$$

that is,

$$Du(x) = Dv(y)D\Phi(x).$$

Thus (1) implies

$$(26) \quad F(Dv(y)D\Phi(\Psi(y)), v(y), \Psi(y)) = F(Du(x), u(x), x) = 0.$$

This is an expression having the form

$$G(Dv(y), v(y), y) = 0 \quad \text{in } V.$$

In addition $v = h$ on Δ , where $\Delta := \Phi(\Gamma)$ and $h(y) := g(\Psi(y))$.

In summary, our problem (1), (2) transforms to read

$$(27) \quad \begin{cases} G(Dv, v, y) = 0 & \text{in } V \\ v = h & \text{on } \Delta, \end{cases}$$

for G, h as above. The point is that if we change variables to straighten out the boundary near x^0 , the boundary-value problem (1), (2) converts into a problem having the same form.

b. Compatibility conditions on boundary data. In view of the foregoing computations, if we are given a point $x^0 \in \Gamma$, we may as well assume from the outset that Γ is flat near x^0 , lying in the plane $\{x_n = 0\}$.

We intend now to utilize the characteristic ODE to construct a solution (1), (2), at least near x^0 , and for this we must discover appropriate initial conditions

$$(28) \quad \mathbf{p}(0) = p^0, \quad z(0) = z^0, \quad \mathbf{x}(0) = x^0.$$

Now clearly if the curve $\mathbf{x}(\cdot)$ passes through x^0 , we should insist that

$$(29) \quad z^0 = g(x^0).$$

What should we require concerning $\mathbf{p}(0) = p^0$? Since (2) implies $u(x_1, \dots, x_{n-1}, 0) = g(x_1, \dots, x_{n-1})$ near x^0 , we may differentiate to find

$$u_{x_i}(x^0) = g_{x_i}(x^0) \quad (i = 1, \dots, n-1).$$

As we also want the PDE (1) to hold, we should therefore insist $p^0 = (p_1^0, \dots, p_n^0)$ satisfies these relations:

$$(30) \quad \begin{cases} p_i^0 = g_{x_i}(x^0) & (i = 1, \dots, n-1) \\ F(p^0, z^0, x^0) = 0. \end{cases}$$

These identities provide n equations for the n quantities $p^0 = (p_1^0, \dots, p_n^0)$.

We call (29) and (30) the *compatibility conditions*. A triple $(p^0, z^0, x^0) \in \mathbb{R}^{2n+1}$ verifying (29), (30) is *admissible*. Note z^0 is uniquely determined by the boundary condition and our choice of the point x^0 , but a vector p^0 satisfying (30) may not exist or may not be unique.

c. Noncharacteristic boundary data. So now assume as above that $x^0 \in \Gamma$, that Γ near x^0 lies in the plane $\{x_n = 0\}$, and that the triple (p^0, z^0, x^0) is admissible. We are planning to construct a solution u of (1), (2) in U near x^0 by integrating the characteristic ODE (11). So far we have ascertained $\mathbf{x}(0) = x^0$, $z(0) = z^0$, $\mathbf{p}(0) = p^0$ are appropriate boundary conditions for the characteristic ODE, with $\mathbf{x}(\cdot)$ intersecting Γ at x^0 . But we will need in fact to solve these ODE for *nearby* initial points as well and must consequently now ask if we can somehow appropriately perturb (p^0, z^0, x^0) , keeping the compatibility conditions.

In other words, given a point $y = (y_1, \dots, y_{n-1}, 0) \in \Gamma$, with y close to x^0 , we intend to solve the characteristic ODE

$$(31) \quad \begin{cases} \text{(a) } \dot{\mathbf{p}}(s) = -D_x F(\mathbf{p}(s), z(s), \mathbf{x}(s)) - D_z F(\mathbf{p}(s), z(s), \mathbf{x}(s))\mathbf{p}(s) \\ \text{(b) } \dot{z}(s) = D_p F(\mathbf{p}(s), z(s), \mathbf{x}(s)) \cdot \mathbf{p}(s) \\ \text{(c) } \dot{\mathbf{x}}(s) = D_p F(\mathbf{p}(s), z(s), \mathbf{x}(s)), \end{cases}$$

with the initial conditions

$$(32) \quad \mathbf{p}(0) = \mathbf{q}(y), \quad z(0) = g(y), \quad \mathbf{x}(0) = y.$$

Our task then is to find a function $\mathbf{q}(\cdot) = (q^1(\cdot), \dots, q^n(\cdot))$, so that

$$(33) \quad \mathbf{q}(x^0) = p^0$$

and $(\mathbf{q}(y), g(y), y)$ is admissible; that is, the compatibility conditions

$$(34) \quad \begin{cases} q^i(y) = g_{x_i}(y) & (i = 1, \dots, n-1) \\ F(\mathbf{q}(y), g(y), y) = 0 \end{cases}$$

hold for all $y \in \Gamma$ close to x^0 .

LEMMA 1 (Noncharacteristic boundary conditions). *There exists a unique solution $\mathbf{q}(\cdot)$ of (33), (34) for all $y \in \Gamma$ sufficiently close to x^0 , provided*

$$(35) \quad F_{p_n}(p^0, z^0, x^0) \neq 0.$$

We say the admissible triple (p^0, z^0, x^0) is *noncharacteristic* if (35) holds. We henceforth assume this condition.

Proof. Our problem is to find $q^n(y)$ so that

$$F(\mathbf{q}(y), g(y), y) = 0,$$

where $q^i(y) = g_{x_i}(y)$ for $i = 1, \dots, n-1$. Since $F(p^0, z^0, x^0) = 0$, the Implicit Function Theorem (§C.7) implies we can indeed locally and uniquely solve for $q^n(y)$, provided that the noncharacteristic condition (35) is valid. \square

General noncharacteristic condition. If Γ is not flat near x^0 , the condition that Γ be noncharacteristic reads

$$(36) \quad D_p F(p^0, z^0, x^0) \cdot \nu(x^0) \neq 0,$$

$\nu(x^0)$ denoting the outward unit normal to ∂U at x^0 . See Problem 7.

3.2.4. Local solution.

Remember that our aim is to use the characteristic ODE to build a solution u of (1), (2), at least near Γ . So as before we select a point $x^0 \in \Gamma$ and, as shown in §3.2.3, may as well assume that near x^0 the surface Γ is flat, lying in the plane $\{x_n = 0\}$. Suppose further that (p^0, z^0, x^0) is an admissible triple of boundary data, which is noncharacteristic. According to Lemma 1 there is a function $\mathbf{q}(\cdot)$ so that $p^0 = \mathbf{q}(x^0)$ and the triple $(\mathbf{q}(y), g(y), y)$ is admissible, for all y sufficiently close to x^0 .

Given any such point $y = (y_1, \dots, y_{n-1}, 0)$, we solve the characteristic ODE (31), subject to initial conditions (32).

NOTATION. Let us write

$$\begin{cases} \mathbf{p}(s) = \mathbf{p}(y, s) = \mathbf{p}(y_1, \dots, y_{n-1}, s) \\ z(s) = z(y, s) = z(y_1, \dots, y_{n-1}, s) \\ \mathbf{x}(s) = \mathbf{x}(y, s) = \mathbf{x}(y_1, \dots, y_{n-1}, s) \end{cases}$$

to display the dependence of the solution of (31), (32) on s and y . Also, we will henceforth when convenient regard x^0 as lying in \mathbb{R}^{n-1} . \square

LEMMA 2 (Local invertibility). *Assume we have the noncharacteristic condition $F_{p_n}(p^0, z^0, x^0) \neq 0$. Then there exist an open interval $I \subseteq \mathbb{R}$ containing 0, a neighborhood W of x^0 in $\Gamma \subset \mathbb{R}^{n-1}$, and a neighborhood V of x^0 in \mathbb{R}^n , such that for each $x \in V$ there exist unique $s \in I$, $y \in W$ such that*

$$x = \mathbf{x}(y, s).$$

The mappings $x \mapsto s, y$ are C^2 .

Proof. We have $\mathbf{x}(x^0, 0) = x^0$. Consequently the Inverse Function Theorem (§C.6) gives the result, provided $\det D\mathbf{x}(x^0, 0) \neq 0$. Now

$$\mathbf{x}(y, 0) = (y, 0) \quad (y \in \Gamma);$$

and so if $i = 1, \dots, n-1$,

$$x_{y_i}^j(x^0, 0) = \begin{cases} \delta_{ij} & (j = 1, \dots, n-1) \\ 0 & (j = n). \end{cases}$$

Furthermore equation (31)(c) implies

$$x_s^j(x^0, 0) = F_{p_j}(p^0, z^0, x^0).$$

Thus

$$D\mathbf{x}(x^0, 0) = \begin{pmatrix} 1 & 0 & F_{p_1}(p^0, z^0, x^0) \\ \vdots & \vdots & \vdots \\ 0 & 1 & \vdots \\ 0 & \cdots & 0 & F_{p_n}(p^0, z^0, x^0) \end{pmatrix}_{n \times n},$$

whence $\det D\mathbf{x}(x^0, 0) \neq 0$ follows from the noncharacteristic condition (35). \square

In view of Lemma 2 for each $x \in V$, we can locally uniquely solve the equation

$$(37) \quad \begin{cases} x = \mathbf{x}(y, s), \\ \text{for } y = \mathbf{y}(x), s = s(x). \end{cases}$$

Finally, let us define

$$(38) \quad \begin{cases} u(x) := z(\mathbf{y}(x), s(x)) \\ \mathbf{p}(x) := \mathbf{p}(\mathbf{y}(x), s(x)) \end{cases}$$

for $x \in V$ and s, y as in (37).

We come finally to our principal assertion, namely, that we can locally weave together the solutions of the characteristic ODE into a solution of the PDE.

THEOREM 2 (Local Existence Theorem). *The function u defined above is C^2 and solves the PDE*

$$F(Du(x), u(x), x) = 0 \quad (x \in V),$$

with the boundary condition

$$u(x) = g(x) \quad (x \in \Gamma \cap V).$$

Proof. 1. First of all, fix $y \in \Gamma$ close to x^0 and, as above, solve the characteristic ODE (31), (32) for $\mathbf{p}(s) = \mathbf{p}(y, s)$, $z(s) = z(y, s)$, and $\mathbf{x}(s) = \mathbf{x}(y, s)$.

2. We assert that if $y \in \Gamma$ is sufficiently close to x^0 , then

$$(39) \quad f(y, s) := F(\mathbf{p}(y, s), z(y, s), \mathbf{x}(y, s)) = 0 \quad (s \in I).$$

To see this, note

$$(40) \quad f(y, 0) = F(\mathbf{p}(y, 0), z(y, 0), \mathbf{x}(y, 0)) = F(\mathbf{q}(y), g(y), y) = 0,$$

by the compatibility condition (34). Furthermore

$$\begin{aligned} f_s(y, s) &= \sum_{j=1}^n F_{p_j} \dot{p}^j + F_z \dot{z} + \sum_{j=1}^n F_{x_j} \dot{x}^j \\ &= \sum_{j=1}^n F_{p_j} (-F_{x_j} - F_z p^j) + F_z \left(\sum_{j=1}^n F_{p_j} p^j \right) \\ &\quad + \sum_{j=1}^n F_{x_j} (F p_j) \quad \text{according to (31)} \\ &= 0. \end{aligned}$$

This calculation and (40) prove (39).

3. In view of Lemma 2 and (37)–(39), we have

$$F(\mathbf{p}(x), u(x), x) = 0 \quad (x \in V).$$

To conclude, we must therefore show

$$(41) \quad \mathbf{p}(x) = Du(x) \quad (x \in V).$$

In order to prove (41), let us first demonstrate for $s \in I$, $y \in W$ that

$$(42) \quad z_s(y, s) = \sum_{j=1}^n p^j(y, s) x_s^j(y, s)$$

and

$$(43) \quad z_{y_i}(y, s) = \sum_{j=1}^n p^j(y, s) x_{y_i}^j(y, s) \quad (i = 1, \dots, n-1).$$

These formulas are obviously consistent with the equality (41) and will later help us prove it. The identity (42) results at once from the characteristic ODE (31)(b),(c). To establish (43), fix $y \in \Gamma$, $i \in \{1, \dots, n-1\}$, and set

$$(44) \quad r^i(s) := z_{y_i}(y, s) - \sum_{j=1}^n p^j(y, s) x_{y_i}^j(y, s).$$

We first note $r^i(0) = g_{x_i}(y) - q^i(y) = 0$ according to the compatibility condition (34). In addition, we can compute

$$(45) \quad \dot{r}^i(s) = z_{y_i s} - \sum_{j=1}^n p_s^j x_{y_i}^j + p^j x_{y_i s}^j.$$

To simplify this expression, let us first differentiate the identity (42) with respect to y_i :

$$(46) \quad z_{s y_i} = \sum_{j=1}^n p_{y_i}^j x_s^j + p^j x_{s y_i}^j.$$

Substituting (46) into (45), we discover

$$(47) \quad \dot{r}^i(s) = \sum_{j=1}^n p_{y_i}^j x_s^j - p_s^j x_{y_i}^j = \sum_{j=1}^n p_{y_i}^j F_{p_j} - (-F_{x_j} - F_z p^j) x_{y_i}^j,$$

according to (31)(a). Now differentiate (39) with respect to y_i :

$$\sum_{j=1}^n F_{p_j} p_{y_i}^j + F_z z_{y_i} + \sum_{j=1}^n F_{x_j} x_{y_i}^j = 0.$$

We employ this identity in (47), thereby obtaining

$$(48) \quad \dot{r}^i(s) = F_z \left(\sum_{j=1}^n p^j x_{y_i}^j - z_{y_i} \right) = -F_z r^i(s).$$

Hence $r^i(\cdot)$ solves the linear ODE (48), with the initial condition $r^i(0) = 0$. Consequently $r^i(s) = 0$ ($s \in I$, $i = 1, \dots, n-1$), and so identity (43) is verified.

4. We finally employ (42), (43) in proving (41). Indeed, if $j = 1, \dots, n$,

$$\begin{aligned} u_{x_j} &= z_s s_{x_j} + \sum_{i=1}^{n-1} z_{y_i} y_{x_j}^i \quad \text{by (38)} \\ &= \sum_{k=1}^n p^k x_s^k s_{x_j} + \sum_{i=1}^{n-1} \sum_{k=1}^n p^k x_{y_i}^k y_{x_j}^i \quad \text{by (42), (43)} \\ &= \sum_{k=1}^n p^k \left(x_s^k s_{x_j} + \sum_{i=1}^{n-1} x_{y_i}^k y_{x_j}^i \right) \\ &= \sum_{k=1}^n p^k x_{x_j}^k = \sum_{k \neq 1}^n p^k \delta_{jk} = p^j. \end{aligned}$$

This assertion at last establishes (41) and so finishes up the proof. \square

3.2.5. Applications.

We turn now to various special cases, to see how the local existence theory simplifies in these circumstances.

a. F linear. Recall that a linear, homogeneous, first-order PDE has the form

$$(49) \quad F(Du, u, x) = \mathbf{b}(x) \cdot Du(x) + c(x)u(x) = 0 \quad (x \in U).$$

Our noncharacteristic assumption (36) at a point $x^0 \in \Gamma$ as above becomes

$$(50) \quad \mathbf{b}(x^0) \cdot \nu(x^0) \neq 0$$

and thus does not involve z^0 or p^0 at all. Furthermore if we specify the boundary condition


$$(51) \quad u = g \quad \text{on } \Gamma,$$

we can uniquely solve equation (34) for $\mathbf{q}(y)$ if $y \in \Gamma$ is near x^0 . Thus we can apply the Local Existence Theorem 2 to construct a unique solution of (49), (51) in some neighborhood V containing x^0 . Note carefully that although we have utilized the full characteristic equations (31) in the proof of Theorem 2, once we know the solution exists, we can use the reduced equations (17) (which do not involve $\mathbf{p}(\cdot)$) to compute the solution. Observe also that the projected characteristics $\mathbf{x}(\cdot)$ emanating from distinct points on Γ cannot cross, owing to uniqueness of solutions of the initial-value problem for the ODE (17)(a).

Example 4. Suppose the trajectories of the ODE

$$(52) \quad \dot{\mathbf{x}}(s) = \mathbf{b}(\mathbf{x}(s))$$

are as drawn for Case 1. We are thus assuming the vector field \mathbf{b} vanishes within U only at one point, which we will take to be the origin 0, and $\mathbf{b} \cdot \nu < 0$ on $\Gamma := \partial U$.




Case 1: flow to an attracting point

Can we solve the linear boundary-value problem

$$(53) \quad \begin{cases} \mathbf{b} \cdot Du = 0 & \text{in } U \\ u = g & \text{on } \Gamma ? \end{cases}$$

Invoking Theorem 2, we see that there exists a unique solution u defined near Γ and indeed that $u(\mathbf{x}(s)) \equiv u(\mathbf{x}(0)) = g(x^0)$ for each solution of the ODE (52), with the initial condition $\mathbf{x}(0) = x^0 \in \Gamma$. However, this solution cannot be smoothly continued to all of U (unless g is constant): any smooth solution of (53) is constant on trajectories of (52) and thus takes on different values near $x = 0$.




Case 2: flow across a domain

But now suppose the trajectories of the ODE (52) look like the illustration for Case 2. We are assuming that each trajectory of the ODE (except those through the characteristic points A, B) enters U precisely once, somewhere through the set

$$\Gamma := \{x \in \partial U \mid \mathbf{b}(x) \cdot \nu(x) < 0\},$$

and exits U precisely once. In this circumstance we can find a smooth solution of (53) by setting u to be constant along each flow line.

Assume finally the flow looks like Case 3. We can now define u to be constant along trajectories, but then u will be discontinuous (unless $g(B) = g(D)$). Note that the point D is characteristic and that the local existence theory fails near D . \square



Case 3: flow with characteristic points

b. F quasilinear. Should F be quasilinear, the PDE (1) is

$$(54) \quad F(Du, u, x) = \mathbf{b}(x, u) \cdot Du + c(x, u) = 0.$$

The noncharacteristic assumption (36) at a point $x^0 \in \Gamma$ reads $\mathbf{b}(x^0, z^0) \cdot \nu(x^0) \neq 0$, where $z^0 = g(x^0)$. As in the preceding example, if we specify the boundary condition

$$(55) \quad u = g \quad \text{on } \Gamma,$$

we can uniquely solve the equations (34) for $\mathbf{q}(y)$ if $y \in \Gamma$ near x^0 . Thus Theorem 2 yields the existence of a unique solution of (54), (55) in some neighborhood V of x^0 . We can compute this solution in V using the reduced characteristic equations (21), which do not explicitly involve $\mathbf{p}(\cdot)$.

In contrast to the linear case, however, *it is possible that the projected characteristics emanating from distinct points in Γ may intersect outside V* ; such an occurrence usually signals the failure of our local solution to exist within all of U .

Example 5 (Characteristics for conservation laws). As an instance of a quasilinear first-order PDE, we turn now to the *scalar conservation law*

$$(56) \quad \begin{aligned} G(Du, u_t, u, x, t) &= u_t + \operatorname{div} \mathbf{F}(u) \\ &= u_t + \mathbf{F}'(u) \cdot Du = 0 \end{aligned}$$

in $U = \mathbb{R}^n \times (0, \infty)$, subject to the initial condition

$$(57) \quad u = g \quad \text{on } \Gamma = \mathbb{R}^n \times \{t = 0\}.$$

Here $\mathbf{F} : \mathbb{R} \rightarrow \mathbb{R}^n$, $\mathbf{F} = (F^1, \dots, F^n)$, and, as usual, we have set $t = x_{n+1}$. Also, “div” denotes the divergence with respect to the spatial variables (x_1, \dots, x_n) , and $Du = D_x u = (u_{x_1}, \dots, u_{x_n})$.

Since the direction $t = x_{n+1}$ plays a special role, we appropriately modify our notation. Writing now $q = (p, p_{n+1})$ and $y = (x, t)$, we have

$$G(q, z, y) = p_{n+1} + \mathbf{F}'(z) \cdot p,$$

and consequently

$$D_q G = (\mathbf{F}'(z), 1), \quad D_y G = 0, \quad D_z G = \mathbf{F}''(z) \cdot p.$$

Clearly the noncharacteristic condition (35) is satisfied at each point $y^0 = (x^0, 0) \in \Gamma$. Furthermore equation (21)(a) becomes

$$(58) \quad \begin{cases} \dot{x}^i(s) = F^{i'}(z(s)) & (i = 1, \dots, n) \\ \dot{x}^{n+1}(s) = 1. \end{cases}$$

Hence $x^{n+1}(s) = s$, in agreement with our having written $x_{n+1} = t$ above. In other words, we can identify the parameter s with the time t .

Equation (21)(b) reads $\dot{z}(s) = 0$. Consequently

$$(59) \quad z(s) = z^0 = g(x^0);$$

and (58) implies

$$(60) \quad \mathbf{x}(s) = \mathbf{F}'(g(x^0))s + x^0.$$

Thus the projected characteristic $\mathbf{y}(s) = (\mathbf{x}(s), s) = (\mathbf{F}'(g(x^0))s + x^0, s)$ ($s \geq 0$) is a straight line, along which u is constant.

Crossing characteristics. But suppose now we apply the same reasoning to a different initial point $z^0 \in \Gamma$, where $g(x^0) \neq g(z^0)$. *The projected characteristics may possibly then intersect at some time $t > 0$.* Since Theorem 1 tells us $u \equiv g(x^0)$ on the projected characteristic through x^0 and $u \equiv g(z^0)$ on the projected characteristic through z^0 , an apparent contradiction arises. The resolution is that *the initial-value problem (56), (57) does not in general have a smooth solution, existing for all times $t > 0$.* \square

We will discuss in §3.4 the interesting possibility of extending the local solution (guaranteed to exist for short times by Theorem 2) to all times $t > 0$, as a kind of “weak” or “generalized” solution.

An implicit formula. We can eliminate s from equations (59), (60) to derive an implicit formula for u . Indeed given $x \in \mathbb{R}^n$ and $t > 0$, we see that since $s = t$,

$$u(\mathbf{x}(t), t) = z(t) = g(\mathbf{x}(t) - t\mathbf{F}'(z^0)) = g(\mathbf{x}(t) - t\mathbf{F}'(u(\mathbf{x}(t), t))).$$

Hence

$$(61) \quad u = g(x - t\mathbf{F}'(u)).$$

This implicit formula for u as a function of x and t is a nonlinear analogue of equation (3) in §2.1. It is easy to check that (61) does indeed give a solution, provided

$$1 + tDg(x - t\mathbf{F}'(u)) \cdot \mathbf{F}''(u) \neq 0.$$

In particular if $n = 1$, we require

$$1 + tg'(x - tF'(u))F''(u) \neq 0.$$

Note that if $F'' > 0$, but $g' < 0$, then this will definitely be false at some time $t > 0$. This failure of the implicit formula (61) reflects also the failure of the characteristic method. \square

c. \mathbf{F} fully nonlinear. The form of the full characteristic equations can be quite complicated for fully nonlinear first-order PDE, but sometimes a remarkable mathematical structure emerges.

Example 6 (Characteristics for the Hamilton–Jacobi equation). We look now at the general Hamilton–Jacobi PDE

$$(62) \quad G(Du, u_t, u, x, t) = u_t + H(Du, x) = 0,$$

where $Du = D_x u = (u_{x_1}, \dots, u_{x_n})$. Then writing $q = (p, p_{n+1})$, $y = (x, t)$, we have

$$G(q, z, y) = p_{n+1} + H(p, x);$$

and so

$$D_q G = (D_p H(p, x), 1), \quad D_y G = (D_x H(p, x), 0), \quad D_z G = 0.$$

Thus equation (11)(c) becomes

$$(63) \quad \begin{cases} \dot{x}^i(s) = H_{p_i}(\mathbf{p}(s), \mathbf{x}(s)) & (i = 1, \dots, n) \\ \dot{x}^{n+1}(s) = 1. \end{cases}$$

In particular we can identify the parameter s with the time t . Equation (11)(a) for the case at hand reads

$$\begin{cases} \dot{p}^i(s) = -H_{x_i}(\mathbf{p}(s), \mathbf{x}(s)) & (i = 1, \dots, n) \\ \dot{p}^{n+1}(s) = 0; \end{cases}$$

the equation (11)(b) is

$$\begin{aligned} \dot{z}(s) &= D_p H(\mathbf{p}(s), \mathbf{x}(s)) \cdot \mathbf{p}(s) + p^{n+1}(s) \\ &= D_p H(\mathbf{p}(s), \mathbf{x}(s)) \cdot \mathbf{p}(s) - H(\mathbf{p}(s), \mathbf{x}(s)). \end{aligned}$$

In summary, the characteristic equations for the Hamilton–Jacobi equation are

$$(64) \quad \begin{cases} \text{(a)} & \dot{\mathbf{p}}(s) = -D_x H(\mathbf{p}(s), \mathbf{x}(s)) \\ \text{(b)} & \dot{z}(s) = D_p H(\mathbf{p}(s), \mathbf{x}(s)) \cdot \mathbf{p}(s) - H(\mathbf{p}(s), \mathbf{x}(s)) \\ \text{(c)} & \dot{\mathbf{x}}(s) = D_p H(\mathbf{p}(s), \mathbf{x}(s)) \end{cases}$$

for $\mathbf{p}(\cdot) = (p^1(\cdot), \dots, p^n(\cdot))$, $z(\cdot)$, and $\mathbf{x}(\cdot) = (x^1(\cdot), \dots, x^n(\cdot))$.

The first and third of these equalities,

$$(65) \quad \begin{cases} \dot{\mathbf{x}} = D_p H(\mathbf{p}, \mathbf{x}) \\ \dot{\mathbf{p}} = -D_x H(\mathbf{p}, \mathbf{x}), \end{cases}$$

are called *Hamilton's equations*. We will discuss these ODE and their relationship to the Hamilton–Jacobi equation in much more detail, just below in §3.3. Observe that the equation for $z(\cdot)$ is trivial, once $\mathbf{x}(\cdot)$ and $\mathbf{p}(\cdot)$ have been found by solving Hamilton's equations. \square

As for conservation laws (Example 5), the initial-value problem for the Hamilton–Jacobi equation does not in general have a smooth solution u lasting for all times $t > 0$.

3.3. INTRODUCTION TO HAMILTON–JACOBI EQUATIONS

In this section we study in some detail the initial-value problem for the Hamilton–Jacobi equation:

$$(1) \quad \begin{cases} u_t + H(Du) = 0 & \text{in } \mathbb{R}^n \times (0, \infty) \\ u = g & \text{on } \mathbb{R}^n \times \{t = 0\}. \end{cases}$$

Here $u : \mathbb{R}^n \times [0, \infty) \rightarrow \mathbb{R}$ is the unknown, $u = u(x, t)$, and $Du = D_x u = (u_{x_1}, \dots, u_{x_n})$. We are given the *Hamiltonian* $H : \mathbb{R}^n \rightarrow \mathbb{R}$ and the initial function $g : \mathbb{R}^n \rightarrow \mathbb{R}$.

Our goal is to find a formula for an appropriate weak or generalized solution, existing for all times $t > 0$, even after the method of characteristics has failed.

3.3.1. Calculus of variations, Hamilton’s ODE.

Remember from §3.2.5 that two of the characteristic equations associated with the Hamilton–Jacobi PDE

$$u_t + H(Du, x) = 0$$

are Hamilton’s ODE

$$\begin{cases} \dot{\mathbf{x}} = D_p H(\mathbf{p}, \mathbf{x}) \\ \dot{\mathbf{p}} = -D_x H(\mathbf{p}, \mathbf{x}), \end{cases}$$

which arise in the classical calculus of variations and in mechanics. (Note the x -dependence in H here.) In this section we recall the derivation of these ODE from a variational principle. We will then discover in §3.3.2 that this discussion contains a clue as to how to build a weak solution of the initial-value problem (1).

a. The calculus of variations. Assume that $L : \mathbb{R}^n \times \mathbb{R}^n \rightarrow \mathbb{R}$ is a given smooth function, hereafter called the *Lagrangian*.

NOTATION. We write

$$L = L(v, x) = L(v_1, \dots, v_n, x_1, \dots, x_n) \quad (v, x \in \mathbb{R}^n)$$

and

$$\begin{cases} D_v L = (L_{v_1} \cdots L_{v_n}) \\ D_x L = (L_{x_1} \cdots L_{x_n}). \end{cases}$$


Thus in the formula (2) below “ v ” is the name of the variable for which we substitute $\dot{\mathbf{w}}(s)$, and “ x ” is the variable for which we substitute $\mathbf{w}(s)$. □

Now fix two points $x, y \in \mathbb{R}^n$ and a time $t > 0$. We introduce then the *action functional*

$$(2) \quad I[\mathbf{w}(\cdot)] := \int_0^t L(\dot{\mathbf{w}}(s), \mathbf{w}(s)) ds \quad \left(\dot{\cdot} = \frac{d}{ds} \right),$$

defined for functions $\mathbf{w}(\cdot) = (w^1(\cdot), w^2(\cdot), \dots, w^n(\cdot))$ belonging to the *admissible class*

$$\mathcal{A} := \{ \mathbf{w}(\cdot) \in C^2([0, t]; \mathbb{R}^n) \mid \mathbf{w}(0) = y, \mathbf{w}(t) = x \}.$$



A problem in the calculus of variations

Thus a C^2 curve $\mathbf{w}(\cdot)$ lies in \mathcal{A} if it starts at the point y at time 0 and reaches the point x at time t .

A basic problem in the *calculus of variations* is to find a curve $\mathbf{x}(\cdot) \in \mathcal{A}$ satisfying

$$(3) \quad I[\mathbf{x}(\cdot)] = \min_{\mathbf{w}(\cdot) \in \mathcal{A}} I[\mathbf{w}(\cdot)].$$

That is, we are asking for a function $\mathbf{x}(\cdot)$ which minimizes the functional $I[\cdot]$ among all admissible candidates $\mathbf{w}(\cdot) \in \mathcal{A}$.

We assume next that there in fact exists a function $\mathbf{x}(\cdot) \in \mathcal{A}$ satisfying our calculus of variations problem and will deduce some of its properties.

THEOREM 1 (Euler–Lagrange equations). *The function $\mathbf{x}(\cdot)$ solves the system of Euler–Lagrange equations*

$$(4) \quad -\frac{d}{ds} (D_v L(\dot{\mathbf{x}}(s), \mathbf{x}(s))) + D_x L(\dot{\mathbf{x}}(s), \mathbf{x}(s)) = 0 \quad (0 \leq s \leq t).$$

This is a vector equation, consisting of n coupled second-order equations.

Proof. 1. Choose a smooth function $\mathbf{y} : [0, t] \rightarrow \mathbb{R}^n$, $\mathbf{y}(\cdot) = (y^1(\cdot), \dots, y^n(\cdot))$, satisfying

$$(5) \quad \mathbf{y}(0) = \mathbf{y}(t) = 0,$$

and define for $\tau \in \mathbb{R}$

$$(6) \quad \mathbf{w}(\cdot) := \mathbf{x}(\cdot) + \tau \mathbf{y}(\cdot).$$

Then $\mathbf{w}(\cdot) \in \mathcal{A}$ and so

$$I[\mathbf{x}(\cdot)] \leq I[\mathbf{w}(\cdot)].$$

Thus the real-valued function

$$i(\tau) := I[\mathbf{x}(\cdot) + \tau\mathbf{y}(\cdot)]$$

has a minimum at $\tau = 0$, and consequently

$$(7) \quad i'(0) = 0 \quad \left(' = \frac{d}{d\tau} \right),$$

provided $i'(0)$ exists.

2. We explicitly compute this derivative. Observe

$$i(\tau) = \int_0^t L(\dot{\mathbf{x}}(s) + \tau\dot{\mathbf{y}}(s), \mathbf{x}(s) + \tau\mathbf{y}(s)) ds,$$

and so

$$i'(\tau) = \int_0^t \sum_{i=1}^n L_{v_i}(\dot{\mathbf{x}} + \tau\dot{\mathbf{y}}, \mathbf{x} + \tau\mathbf{y})\dot{y}^i + L_{x_i}(\dot{\mathbf{x}} + \tau\dot{\mathbf{y}}, \mathbf{x} + \tau\mathbf{y})y^i ds.$$

Set $\tau = 0$ and remember (7):

$$0 = i'(0) = \int_0^t \sum_{i=1}^n L_{v_i}(\dot{\mathbf{x}}, \mathbf{x})\dot{y}^i + L_{x_i}(\dot{\mathbf{x}}, \mathbf{x})y^i ds.$$

We recall (5) and then integrate by parts in the first term inside the integral, to discover

$$0 = \sum_{i=1}^n \int_0^t \left[-\frac{d}{ds} (L_{v_i}(\dot{\mathbf{x}}, \mathbf{x})) + L_{x_i}(\dot{\mathbf{x}}, \mathbf{x}) \right] y^i ds.$$

This identity is valid for all smooth functions $\mathbf{y}(\cdot) = (y^1(\cdot), \dots, y^n(\cdot))$ satisfying the boundary conditions (5), and so for $0 \leq s \leq t$

$$-\frac{d}{ds} (L_{v_i}(\dot{\mathbf{x}}, \mathbf{x})) + L_{x_i}(\dot{\mathbf{x}}, \mathbf{x}) = 0 \quad (i = 1, \dots, n). \quad \square$$

Critical points. We have just demonstrated that any minimizer $\mathbf{x}(\cdot) \in \mathcal{A}$ of $I[\cdot]$ solves the Euler–Lagrange system of ODE. It is of course possible that a curve $\mathbf{x}(\cdot) \in \mathcal{A}$ may solve the Euler–Lagrange equations without necessarily being a minimizer: in this case we say $\mathbf{x}(\cdot)$ is a *critical point* of $I[\cdot]$. So every minimizer is a critical point, but a critical point need not be a minimizer.

Example. If $L(v, x) = \frac{1}{2}m|v|^2 - \phi(x)$, where $m > 0$, the corresponding Euler–Lagrange equation is

$$m\ddot{\mathbf{x}}(s) = \mathbf{f}(\mathbf{x}(s))$$

for $\mathbf{f} := -D\phi$. This is Newton’s law for the motion of a particle of mass m moving in the force field \mathbf{f} generated by the potential ϕ . (See Feynman–Leighton–Sands [F-L-S, Chapter 19].) \square

b. Hamilton’s equations. We now transform the Euler–Lagrange equations, a system of n second-order ODE, into Hamilton’s equations, a system of $2n$ first-order ODE. We hereafter assume the C^2 function $\mathbf{x}(\cdot)$ is a critical point of the action functional and thus solves the Euler–Lagrange equations (4).

First we set

$$(8) \quad \mathbf{p}(s) := D_v L(\dot{\mathbf{x}}(s), \mathbf{x}(s)) \quad (0 \leq s \leq t);$$

$\mathbf{p}(\cdot)$ is called the *generalized momentum* corresponding to the *position* $\mathbf{x}(\cdot)$ and *velocity* $\dot{\mathbf{x}}(\cdot)$. We next make this important hypothesis:

$$(9) \quad \left\{ \begin{array}{l} \text{Suppose for all } x, p \in \mathbb{R}^n \text{ that the equation} \\ \quad p = D_v L(v, x) \\ \text{can be uniquely solved for } v \text{ as a smooth} \\ \text{function of } p \text{ and } x, v = \mathbf{v}(p, x). \end{array} \right.$$

We will examine this assumption in more detail later: see §3.3.2.

DEFINITION. The Hamiltonian H associated with the Lagrangian L is

$$H(p, x) := p \cdot \mathbf{v}(p, x) - L(\mathbf{v}(p, x), x) \quad (p, x \in \mathbb{R}^n),$$

where the function $\mathbf{v}(\cdot)$ is defined implicitly by (9).

Example (continued). The Hamiltonian corresponding to the Lagrangian $L(v, x) = \frac{1}{2}m|v|^2 - \phi(x)$ is

$$H(p, x) = \frac{1}{2m}|p|^2 + \phi(x).$$

The Hamiltonian is thus the total energy, the sum of the kinetic and potential energies (whereas the Lagrangian is the difference between the kinetic and potential energies). \square

Next we rewrite the Euler–Lagrange equations in terms of $\mathbf{p}(\cdot), \mathbf{x}(\cdot)$:

THEOREM 2 (Derivation of Hamilton’s ODE). *The functions $\mathbf{x}(\cdot)$ and $\mathbf{p}(\cdot)$ satisfy Hamilton’s equations:*

$$(10) \quad \begin{cases} \dot{\mathbf{x}}(s) = D_{\mathbf{p}}H(\mathbf{p}(s), \mathbf{x}(s)) \\ \dot{\mathbf{p}}(s) = -D_{\mathbf{x}}H(\mathbf{p}(s), \mathbf{x}(s)) \end{cases}$$

for $0 \leq s \leq t$. Furthermore,

the mapping $s \mapsto H(\mathbf{p}(s), \mathbf{x}(s))$ is constant.

The equations (10) comprise a coupled system of $2n$ first-order ODE for $\mathbf{x}(\cdot) = (x^1(\cdot), \dots, x^n(\cdot))$ and $\mathbf{p}(\cdot) = (p^1(\cdot), \dots, p^n(\cdot))$ (defined by (8)).

Proof. First note from (8) and (9) that $\dot{\mathbf{x}}(s) = \mathbf{v}(\mathbf{p}(s), \mathbf{x}(s))$.

Let us hereafter write $\mathbf{v}(\cdot) = (v^1(\cdot), \dots, v^n(\cdot))$. We compute for $i = 1, \dots, n$ that

$$\begin{aligned} H_{x_i}(p, x) &= \sum_{k=1}^n p_k v_{x_i}^k(p, x) - L_{v_k}(\mathbf{v}(p, x), x) v_{x_i}^k(p, x) - L_{x_i}(\mathbf{v}(p, x), x) \\ &= -L_{x_i}(q, x) \quad \text{according to (9)} \end{aligned}$$

and

$$\begin{aligned} H_{p_i}(p, x) &= v^i(p, x) + \sum_{k=1}^n p_k v_{p_i}^k(p, x) - L_{v_k}(\mathbf{v}(p, x), x) v_{p_i}^k(p, x) \\ &= v^i(p, x), \quad \text{again by (9)}. \end{aligned}$$

Thus

$$H_{p_i}(\mathbf{p}(s), \mathbf{x}(s)) = v^i(\mathbf{p}(s), \mathbf{x}(s)) = \dot{x}^i(s),$$

and likewise

$$\begin{aligned} H_{x_i}(\mathbf{p}(s), \mathbf{x}(s)) &= -L_{x_i}(\mathbf{v}(\mathbf{p}(s), \mathbf{x}(s)), \mathbf{x}(s)) = -L_{x_i}(\dot{\mathbf{x}}(s), \mathbf{x}(s)) \\ &= -\frac{d}{ds}(L_{v_i}(\dot{\mathbf{x}}(s), \mathbf{x}(s))) \quad \text{according to (4)} \\ &= -\dot{p}^i(s). \end{aligned}$$

Finally, observe

$$\begin{aligned} \frac{d}{ds}H(\mathbf{p}, \mathbf{x}) &= \sum_{i=1}^n H_{p_i}(\mathbf{p}, \mathbf{x})\dot{p}^i + H_{x_i}(\mathbf{p}, \mathbf{x})\dot{x}^i \\ &= \sum_{i=1}^n H_{p_i}(\mathbf{p}, \mathbf{x})(-H_{x_i}(\mathbf{p}, \mathbf{x})) + H_{x_i}(\mathbf{p}, \mathbf{x})H_{p_i}(\mathbf{p}, \mathbf{x}) = 0. \end{aligned}$$

□

See Arnold [Ar1, Chapter 9] for more on Hamilton’s ODE and Hamilton–Jacobi PDE in classical mechanics. We are employing here different notation than is customary in mechanics: our notation is better overall for PDE theory.

3.3.2. Legendre transform, Hopf–Lax formula.

Now let us try to find a connection between the Hamilton–Jacobi PDE and the calculus of variations problem (2)–(4). To simplify further, we also drop the x -dependence in the Hamiltonian, so that afterwards $H = H(p)$. We start by reexamining the definition of the Hamiltonian in §3.3.1.

a. Legendre transform. We hereafter suppose the Lagrangian $L : \mathbb{R}^n \rightarrow \mathbb{R}$ satisfies these conditions:

$$(11) \quad \text{the mapping } v \mapsto L(v) \text{ is convex}$$

and

$$(12) \quad \lim_{|v| \rightarrow \infty} \frac{L(v)}{|v|} = +\infty.$$

The convexity implies L is continuous.

DEFINITION. *The Legendre transform of L is*

$$(13) \quad L^*(p) := \sup_{q \in \mathbb{R}^n} \{p \cdot v - L(v)\} \quad (p \in \mathbb{R}^n).$$

This is also referred to as the *Fenchel transform*.

Motivation for Legendre transform. Why do we make this definition? For some insight let us note in view of (12) that the “sup” in (13) is really a “max”; that is, there exists some $v^* \in \mathbb{R}^n$ for which

$$L^*(p) = p \cdot v^* - L(v^*)$$

and the mapping $v \mapsto p \cdot v - L(v)$ has a maximum at $v = v^*$. But then $p = DL(v^*)$, provided L is differentiable at v^* . Hence the equation $p = DL(v)$ is solvable (although perhaps not uniquely) for v in terms of p , $v^* = \mathbf{v}(p)$. Therefore

$$L^*(p) = p \cdot \mathbf{v}(p) - L(\mathbf{v}(p)).$$

However, this is almost exactly the definition of the Hamiltonian H associated with L in §3.3.1 (where, recall, we are now assuming the variable x does not appear). We consequently henceforth write

$$(14) \quad H = L^*.$$

Thus (13) tells us how to obtain the Hamiltonian H from the Lagrangian L .

Now we ask the converse question: given H , how do we compute L ?

THEOREM 3 (Convex duality of Hamiltonian and Lagrangian). *Assume L satisfies (11), (12) and define H by (13), (14).*

(i) *Then*

the mapping $p \mapsto H(p)$ is convex

and

$$\lim_{|p| \rightarrow \infty} \frac{H(p)}{|p|} = +\infty.$$

(ii) *Furthermore*

$$(15) \quad L = H^*.$$

Thus H is the Legendre transform of L , and vice versa:

$$L = H^*, \quad H = L^*.$$

We say H and L are *dual* convex functions. The identity (15) implies that the three statements

$$(16) \quad \begin{cases} p \cdot v = L(v) + H(p) \\ p = DL(v) \\ v = DH(p) \end{cases}$$

are equivalent provided H is differentiable at p and L is differentiable at v : see Problem 11.

Proof. 1. For each fixed v , the function $p \mapsto p \cdot v - L(v)$ is linear; and consequently the mapping

$$p \mapsto H(p) = L^*(p) = \sup_{v \in \mathbb{R}^n} \{p \cdot v - L(v)\}$$

is convex. Indeed, if $0 \leq \tau \leq 1$, $p, \hat{p} \in \mathbb{R}^n$, we have

$$\begin{aligned} H(\tau p + (1 - \tau)\hat{p}) &= \sup_v \{(\tau p + (1 - \tau)\hat{p}) \cdot v - L(v)\} \\ &\leq \tau \sup_v \{p \cdot v - L(v)\} \\ &\quad + (1 - \tau) \sup_v \{\hat{p} \cdot v - L(v)\} \\ &= \tau H(p) + (1 - \tau)H(\hat{p}). \end{aligned}$$

2. Fix any $\lambda > 0$, $p \neq 0$. Then

$$\begin{aligned} H(p) &= \sup_{v \in \mathbb{R}^n} \{p \cdot v - L(v)\} \\ &\geq \lambda |p| - L\left(\lambda \frac{p}{|p|}\right) \quad (v = \lambda \frac{p}{|p|}) \\ &\geq \lambda |p| - \max_{B(0, \lambda)} L. \end{aligned}$$

Thus $\liminf_{|p| \rightarrow \infty} \frac{H(p)}{|p|} \geq \lambda$ for all $\lambda > 0$.

3. In view of (14)

$$H(p) + L(v) \geq p \cdot v$$

for all $p, v \in \mathbb{R}^n$, and consequently

$$L(v) \geq \sup_{p \in \mathbb{R}^n} \{p \cdot v - H(p)\} = H^*(v).$$

On the other hand

$$\begin{aligned} H^*(v) &= \sup_{p \in \mathbb{R}^n} \{p \cdot v - \sup_{r \in \mathbb{R}^n} \{p \cdot r - L(r)\}\} \\ &= \sup_{p \in \mathbb{R}^n} \inf_{r \in \mathbb{R}^n} \{p \cdot (v - r) + L(r)\}. \end{aligned}$$

Now since $v \mapsto L(v)$ is convex, according to §B.1 there exists $s \in \mathbb{R}^n$ such that

$$L(r) \geq L(v) + s \cdot (r - v) \quad (r \in \mathbb{R}^n).$$

(If L is differentiable at q , take $s = DL(v)$.) Putting $p = s$ above, we compute

$$H^*(v) \geq \inf_{r \in \mathbb{R}^n} \{s \cdot (v - r) + L(r)\} = L(v). \quad \square$$

b. Hopf–Lax formula. Let us now return to the initial-value problem (1) for the Hamilton–Jacobi equation and conclude from (64) in §3.2.5 that the corresponding characteristic equations are

$$\begin{cases} \dot{\mathbf{p}} = 0 \\ \dot{z} = DH(\mathbf{p}) \cdot \mathbf{p} - H(\mathbf{p}) \\ \dot{\mathbf{x}} = DH(\mathbf{p}). \end{cases}$$

The first and third of these are Hamilton’s ODE, which we in §3.3.1 derived from a minimization problem for associated Lagrangian $L = H^*$. Remembering (16), we can therefore understand the second of the characteristic equations as asserting

$$\dot{z} = DH(\mathbf{p}) \cdot \mathbf{p} - H(\mathbf{p}) = L(\dot{\mathbf{x}}).$$

But at least for such short times that (1) has a smooth solution u , we have $z(t) = u(\mathbf{x}(t), t)$ and consequently

$$u(x, t) = \int_0^t L(\dot{\mathbf{x}}(s)) ds + g(\mathbf{x}(0)).$$

Our intention is to modify this expression, to make sense even for large times $t > 0$ when (1) does not have a smooth solution. The variational principle for the action discussed in §3.3.1 provides the clue. Given $x \in \mathbb{R}^n$ and $t > 0$, we therefore propose to *minimize* among curves $\mathbf{w}(\cdot)$ satisfying $\mathbf{w}(t) = x$ the expression

$$\int_0^t L(\dot{\mathbf{w}}(s)) ds + g(\mathbf{w}(0)),$$

which is the action augmented with the value of the initial data. We accordingly now *define*

$$(17) \quad u(x, t) := \inf \left\{ \int_0^t L(\dot{\mathbf{w}}(s)) ds + g(\mathbf{w}(0)) \mid \mathbf{w}(t) = x \right\},$$

the infimum taken over all C^1 functions $\mathbf{w}(\cdot)$. (Better justification for this guess will be provided much later, in Chapter 10.)

We must investigate the sense in which the function u given by (17) actually solves the initial-value problem for the Hamilton–Jacobi PDE:

$$(18) \quad \begin{cases} u_t + H(Du) = 0 & \text{in } \mathbb{R}^n \times (0, \infty) \\ u = g & \text{on } \mathbb{R}^n \times \{t = 0\}. \end{cases}$$

Recall we are assuming H is smooth,

$$(19) \quad \begin{cases} H \text{ is convex and} \\ \lim_{|p| \rightarrow \infty} \frac{H(p)}{|p|} = +\infty. \end{cases}$$

We henceforth suppose also

$$(20) \quad g : \mathbb{R}^n \rightarrow \mathbb{R} \quad \text{is Lipschitz continuous;}$$

this means $\text{Lip}(g) := \sup_{\substack{x, y \in \mathbb{R}^n \\ x \neq y}} \left\{ \frac{|g(x) - g(y)|}{|x - y|} \right\} < \infty$.

First we note that formula (17) can be simplified:

THEOREM 4 (Hopf–Lax formula). *If $x \in \mathbb{R}^n$ and $t > 0$, then the solution $u = u(x, t)$ of the minimization problem (17) is*

$$(21) \quad u(x, t) = \min_{y \in \mathbb{R}^n} \left\{ tL \left(\frac{x - y}{t} \right) + g(y) \right\}.$$

DEFINITION. *We call the expression on the right-hand side of (21) the Hopf–Lax formula.*

Proof. 1. Fix any $y \in \mathbb{R}^n$ and define $\mathbf{w}(s) := y + \frac{s}{t}(x - y)$ ($0 \leq s \leq t$). Then the definition (17) of u implies

$$u(x, t) \leq \int_0^t L(\dot{\mathbf{w}}(s)) ds + g(y) = tL\left(\frac{x - y}{t}\right) + g(y),$$

and so

$$u(x, t) \leq \inf_{y \in \mathbb{R}^n} \left\{ tL\left(\frac{x - y}{t}\right) + g(y) \right\}.$$

2. On the other hand, if $\mathbf{w}(\cdot)$ is any C^1 function satisfying $\mathbf{w}(t) = x$, we have

$$L\left(\frac{1}{t} \int_0^t \dot{\mathbf{w}}(s) ds\right) \leq \frac{1}{t} \int_0^t L(\dot{\mathbf{w}}(s)) ds$$

by Jensen's inequality (§B.1). Thus if we write $y = \mathbf{w}(0)$, we find

$$tL\left(\frac{x - y}{t}\right) + g(y) \leq \int_0^t L(\dot{\mathbf{w}}(s)) ds + g(y);$$

and consequently

$$\inf_{y \in \mathbb{R}^n} \left\{ tL\left(\frac{x - y}{t}\right) + g(y) \right\} \leq u(x, t).$$

3. We have so far shown

$$u(x, t) = \inf_{y \in \mathbb{R}^n} \left\{ tL\left(\frac{x - y}{t}\right) + g(y) \right\},$$

and leave it as an exercise to prove that the infimum above is really a minimum. \square

We now commence a study of various properties of the function u defined by the Hopf–Lax formula (21). Our ultimate goal is showing this formula provides a reasonable weak solution of the initial-value problem (18) for the Hamilton–Jacobi equation.

First, we record some preliminary observations.

LEMMA 1 (A functional identity). *For each $x \in \mathbb{R}^n$ and $0 \leq s < t$, we have*

$$(22) \quad u(x, t) = \min_{y \in \mathbb{R}^n} \left\{ (t - s)L\left(\frac{x - y}{t - s}\right) + u(y, s) \right\}.$$

In other words, to compute $u(\cdot, t)$, we can calculate u at time s and then use $u(\cdot, s)$ as the initial condition on the remaining time interval $[s, t]$.

Proof. 1. Fix $y \in \mathbb{R}^n$, $0 < s < t$ and choose $z \in \mathbb{R}^n$ so that

$$(23) \quad u(y, s) = sL\left(\frac{y-z}{s}\right) + g(z).$$

Now since L is convex and $\frac{x-z}{t} = (1 - \frac{s}{t})\frac{x-y}{t-s} + \frac{s}{t}\frac{y-z}{s}$, we have

$$L\left(\frac{x-z}{t}\right) \leq \left(1 - \frac{s}{t}\right)L\left(\frac{x-y}{t-s}\right) + \frac{s}{t}L\left(\frac{y-z}{s}\right).$$

Thus

$$\begin{aligned} u(x, t) &\leq tL\left(\frac{x-z}{t}\right) + g(z) \leq (t-s)L\left(\frac{x-y}{t-s}\right) + sL\left(\frac{y-z}{s}\right) + g(z) \\ &= (t-s)L\left(\frac{x-y}{t-s}\right) + u(y, s), \end{aligned}$$

by (23). This inequality is true for each $y \in \mathbb{R}^n$. Therefore, since $y \mapsto u(y, s)$ is continuous (according to the first part of the proof Lemma 2 below), we have

$$(24) \quad u(x, t) \leq \min_{y \in \mathbb{R}^n} \left\{ (t-s)L\left(\frac{x-y}{t-s}\right) + u(y, s) \right\}.$$

2. Now choose w such that

$$(25) \quad u(x, t) = tL\left(\frac{x-w}{t}\right) + g(w),$$

and set $y := \frac{s}{t}x + (1 - \frac{s}{t})w$. Then $\frac{x-y}{t-s} = \frac{x-w}{t} = \frac{y-w}{s}$. Consequently

$$\begin{aligned} (t-s)L\left(\frac{x-y}{t-s}\right) + u(y, s) &\leq (t-s)L\left(\frac{x-w}{t}\right) + sL\left(\frac{y-w}{s}\right) + g(w) \\ &= tL\left(\frac{x-w}{t}\right) + g(w) = u(x, t), \end{aligned}$$

by (25). Hence

$$(26) \quad \min_{y \in \mathbb{R}^n} \left\{ (t-s)L\left(\frac{x-y}{t-s}\right) + u(y, s) \right\} \leq u(x, t).$$

□

LEMMA 2 (Lipschitz continuity). *The function u is Lipschitz continuous in $\mathbb{R}^n \times [0, \infty)$, and*

$$u = g \quad \text{on } \mathbb{R}^n \times \{t = 0\}.$$

Proof. 1. Fix $t > 0$, $x, \hat{x} \in \mathbb{R}^n$. Choose $y \in \mathbb{R}^n$ such that

$$(27) \quad tL\left(\frac{x-y}{t}\right) + g(y) = u(x, t).$$

Then

$$\begin{aligned} u(\hat{x}, t) - u(x, t) &= \min_z \left\{ tL\left(\frac{\hat{x}-z}{t}\right) + g(z) \right\} - tL\left(\frac{x-y}{t}\right) - g(y) \\ &\leq g(\hat{x} - x + y) - g(y) \leq \text{Lip}(g)|\hat{x} - x|. \end{aligned}$$

Hence

$$u(\hat{x}, t) - u(x, t) \leq \text{Lip}(g)|\hat{x} - x|;$$

and, interchanging the roles of \hat{x} and x , we find

$$(28) \quad |u(x, t) - u(\hat{x}, t)| \leq \text{Lip}(g)|x - \hat{x}|.$$

2. Now select $x \in \mathbb{R}^n$, $t > 0$. Choosing $y = x$ in (21), we discover

$$(29) \quad u(x, t) \leq tL(0) + g(x).$$

Furthermore,

$$\begin{aligned} u(x, t) &= \min_{y \in \mathbb{R}^n} \left\{ tL\left(\frac{x-y}{t}\right) + g(y) \right\} \\ &\geq g(x) + \min_{y \in \mathbb{R}^n} \left\{ -\text{Lip}(g)|x-y| + tL\left(\frac{x-y}{t}\right) \right\} \\ &= g(x) - t \max_{z \in \mathbb{R}^n} \{ \text{Lip}(g)|z| - L(z) \} \quad (z = \frac{x-y}{t}) \\ &= g(x) - t \max_{w \in B(0, \text{Lip}(g))} \max_{z \in \mathbb{R}^n} \{ w \cdot z - L(z) \} \\ &= g(x) - t \max_{B(0, \text{Lip}(g))} H. \end{aligned}$$

This inequality and (29) imply

$$|u(x, t) - g(x)| \leq Ct$$

for

$$(30) \quad C := \max(|L(0)|, \max_{B(0, \text{Lip}(g))} |H|).$$

3. Finally select $x \in \mathbb{R}^n$, $0 < \hat{t} < t$. Then $\text{Lip}(u(\cdot, t)) \leq \text{Lip}(g)$ by (28) above. Consequently Lemma 1 and calculations like those employed in step 2 above imply

$$|u(x, t) - u(x, \hat{t})| \leq C|t - \hat{t}|$$

for the constant C defined by (30). \square

Now Rademacher's Theorem (which we will prove later, in §5.8.3) asserts that a Lipschitz function is differentiable almost everywhere. Consequently in view of Lemma 2 our function u defined by the Hopf–Lax formula (21) is differentiable for a.e. $(x, t) \in \mathbb{R}^n \times (0, \infty)$. The next theorem asserts u in fact solves the Hamilton–Jacobi PDE wherever u is differentiable.

THEOREM 5 (Solving the Hamilton–Jacobi equation). *Suppose $x \in \mathbb{R}^n$, $t > 0$, and u defined by the Hopf–Lax formula (21) is differentiable at a point $(x, t) \in \mathbb{R}^n \times (0, \infty)$. Then*

$$u_t(x, t) + H(Du(x, t)) = 0.$$

Proof. 1. Fix $v \in \mathbb{R}^n$, $h > 0$. Owing to Lemma 1,

$$\begin{aligned} u(x + hv, t + h) &= \min_{y \in \mathbb{R}^n} \left\{ hL \left(\frac{x + hv - y}{h} \right) + u(y, t) \right\} \\ &\leq hL(v) + u(x, t). \end{aligned}$$

Hence

$$\frac{u(x + hv, t + h) - u(x, t)}{h} \leq L(v).$$

Let $h \rightarrow 0^+$, to compute

$$v \cdot Du(x, t) + u_t(x, t) \leq L(v).$$

This inequality is valid for all $v \in \mathbb{R}^n$, and so

$$(31) \quad u_t(x, t) + H(Du(x, t)) = u_t(x, t) + \max_{v \in \mathbb{R}^n} \{v \cdot Du(x, t) - L(v)\} \leq 0.$$

The first equality holds since $H = L^*$.

2. Now choose z such that $u(x, t) = tL\left(\frac{x-z}{t}\right) + g(z)$. Fix $h > 0$ and set $s = t - h$, $y = \frac{s}{t}x + (1 - \frac{s}{t})z$. Then $\frac{x-z}{t} = \frac{y-z}{s}$, and thus

$$\begin{aligned} u(x, t) - u(y, s) &\geq tL\left(\frac{x-z}{t}\right) + g(z) - \left[sL\left(\frac{y-z}{s}\right) + g(z) \right] \\ &= (t-s)L\left(\frac{x-z}{t}\right). \end{aligned}$$

That is,

$$\frac{u(x, t) - u\left(\left(1 - \frac{h}{t}\right)x + \frac{h}{t}z, t - h\right)}{h} \geq L\left(\frac{x - z}{t}\right).$$

Let $h \rightarrow 0^+$, to see that

$$\frac{x - z}{t} \cdot Du(x, t) + u_t(x, t) \geq L\left(\frac{x - z}{t}\right).$$

Consequently

$$\begin{aligned} u_t(x, t) + H(Du(x, t)) &= u_t(x, t) + \max_{v \in \mathbb{R}^n} \{v \cdot Du(x, t) - L(v)\} \\ &\geq u_t(x, t) + \frac{x - z}{t} \cdot Du(x, t) - L\left(\frac{x - z}{t}\right) \\ &\geq 0. \end{aligned}$$

This inequality and (31) complete the proof. \square

We summarize:

THEOREM 6 (Hopf–Lax formula as solution). *The function u defined by the Hopf–Lax formula (21) is Lipschitz continuous, is differentiable a.e. in $\mathbb{R}^n \times (0, \infty)$, and solves the initial-value problem*

$$(32) \quad \begin{cases} u_t + H(Du) = 0 & \text{a.e. in } \mathbb{R}^n \times (0, \infty) \\ u = g & \text{on } \mathbb{R}^n \times \{t = 0\}. \end{cases}$$

3.3.3. Weak solutions, uniqueness.

a. Semiconcavity. In view of Theorem 6 above it may seem reasonable to define a weak solution of the initial-value problem (18) to be a Lipschitz function which agrees with g on $\mathbb{R}^n \times \{t = 0\}$ and solves the PDE a.e. on $\mathbb{R}^n \times (0, \infty)$. However this turns out to be an inadequate definition, *as such weak solutions would not in general be unique.*

Example. Consider the initial-value problem

$$(33) \quad \begin{cases} u_t + |u_x|^2 = 0 & \text{in } \mathbb{R} \times (0, \infty) \\ u = 0 & \text{on } \mathbb{R} \times \{t = 0\}. \end{cases}$$

One obvious solution is

$$u_1(x, t) \equiv 0.$$

However the function

$$u_2(x, t) := \begin{cases} 0 & \text{if } |x| \geq t \\ x - t & \text{if } 0 \leq x \leq t \\ -x - t & \text{if } -t \leq x \leq 0 \end{cases}$$

is Lipschitz continuous and also solves the PDE a.e. (everywhere, in fact, except on the lines $x = 0, \pm t$). It is easy to see that actually there are infinitely many Lipschitz functions satisfying (33). \square

This example shows we must presumably require more of a weak solution than merely that it satisfy the PDE a.e. We will look to the Hopf–Lax formula (21) for a further clue as to what is needed to ensure uniqueness. The following lemma demonstrates that u inherits a kind of “one-sided” second-derivative estimate from the initial function g .

LEMMA 3 (Semiconcavity). *Suppose there exists a constant C such that*

$$(34) \quad g(x+z) - 2g(x) + g(x-z) \leq C|z|^2$$

for all $x, z \in \mathbb{R}^n$. Define u by the Hopf–Lax formula (21). Then

$$u(x+z, t) - 2u(x, t) + u(x-z, t) \leq C|z|^2$$

for all $x, z \in \mathbb{R}^n$, $t > 0$.

We say g is *semiconcave* provided (34) holds. It is easy to check that (34) is valid if g is C^2 and $\sup_{\mathbb{R}^n} |D^2g| < \infty$. Note that g is semiconcave if and only if the mapping $x \mapsto g(x) - \frac{C}{2}|x|^2$ is concave for some constant C .

Proof. Choose $y \in \mathbb{R}^n$ so that $u(x, t) = tL\left(\frac{x-y}{t}\right) + g(y)$. Then, putting $y+z$ and $y-z$ in the Hopf–Lax formulas for $u(x+z, t)$ and $u(x-z, t)$, we find

$$\begin{aligned} & u(x+z, t) - 2u(x, t) + u(x-z, t) \\ & \leq \left[tL\left(\frac{x-y}{t}\right) + g(y+z) \right] - 2 \left[tL\left(\frac{x-y}{t}\right) + g(y) \right] \\ & \quad + \left[tL\left(\frac{x-y}{t}\right) + g(y-z) \right] \\ & = g(y+z) - 2g(y) + g(y-z) \\ & \leq C|z|^2, \quad \text{by (34).} \end{aligned}$$

\square

As a semiconcavity condition for u will turn out to be important, we pause to identify some other circumstances under which it is valid. We will no longer assume g to be semiconcave but will suppose the Hamiltonian H to be uniformly convex.

DEFINITION. A C^2 convex function $H : \mathbb{R}^n \rightarrow \mathbb{R}$ is called uniformly convex (with constant $\theta > 0$) if

$$(35) \quad \sum_{i,j=1}^n H_{p_i p_j}(p) \xi_i \xi_j \geq \theta |\xi|^2 \quad \text{for all } p, \xi \in \mathbb{R}^n.$$

We now prove that even if g is not semiconcave, the uniform convexity of H forces u to become semiconcave for times $t > 0$: this is a kind of mild regularizing effect for the Hopf–Lax solution of the initial-value problem (18).

LEMMA 4 (Semiconcavity again). *Suppose that H is uniformly convex (with constant θ) and u is defined by the Hopf–Lax formula (21). Then*

$$u(x+z, t) - 2u(x, t) + u(x-z, t) \leq \frac{1}{\theta t} |z|^2$$

for all $x, z \in \mathbb{R}^n$, $t > 0$.

Proof. 1. We note first using Taylor’s formula that (35) implies

$$(36) \quad H\left(\frac{p_1 + p_2}{2}\right) \leq \frac{1}{2}H(p_1) + \frac{1}{2}H(p_2) - \frac{\theta}{8}|p_1 - p_2|^2.$$

Next we claim that for the Lagrangian L we have the estimate

$$(37) \quad \frac{1}{2}L(v_1) + \frac{1}{2}L(v_2) \leq L\left(\frac{v_1 + v_2}{2}\right) + \frac{1}{8\theta}|v_1 - v_2|^2$$

for all $v_1, v_2 \in \mathbb{R}^n$. Verification is left as an exercise.

2. Now choose y so that $u(x, t) = tL\left(\frac{x-y}{t}\right) + g(y)$. Then using the same value of y in the Hopf–Lax formulas for $u(x+z, t)$ and $u(x-z, t)$, we calculate

$$\begin{aligned} & u(x+z, t) - 2u(x, t) + u(x-z, t) \\ & \leq \left[tL\left(\frac{x+z-y}{t}\right) + g(y) \right] - 2 \left[tL\left(\frac{x-y}{t}\right) + g(y) \right] \\ & \quad + \left[tL\left(\frac{x-z-y}{t}\right) + g(y) \right] \\ & = 2t \left[\frac{1}{2}L\left(\frac{x+z-y}{t}\right) + \frac{1}{2}L\left(\frac{x-z-y}{t}\right) - L\left(\frac{x-y}{t}\right) \right] \\ & \leq 2t \frac{1}{8\theta} \left| \frac{2z}{t} \right|^2 \leq \frac{1}{\theta t} |z|^2, \end{aligned}$$

the next-to-last inequality following from (37). □

b. Weak solutions, uniqueness. In this section we show that semi-concavity conditions of the sorts discovered for the Hopf–Lax solution u in Lemmas 3 and 4 can be utilized as uniqueness criteria.

DEFINITION. We say that a Lipschitz continuous function $u : \mathbb{R}^n \times [0, \infty) \rightarrow \mathbb{R}$ is a weak solution of the initial-value problem:

$$(38) \quad \begin{cases} u_t + H(Du) = 0 & \text{in } \mathbb{R}^n \times (0, \infty) \\ u = g & \text{on } \mathbb{R}^n \times \{t = 0\} \end{cases}$$

provided

$$(a) \quad u(x, 0) = g(x) \quad (x \in \mathbb{R}^n),$$

$$(b) \quad u_t(x, t) + H(Du(x, t)) = 0 \quad \text{for a.e. } (x, t) \in \mathbb{R}^n \times (0, \infty),$$

and

$$(c) \quad u(x + z, t) - 2u(x, t) + u(x - z, t) \leq C \left(1 + \frac{1}{t}\right) |z|^2$$

for some constant $C \geq 0$ and all $x, z \in \mathbb{R}^n$, $t > 0$.

Next we prove that a weak solution of (38) is unique, the key point being that this uniqueness assertion follows from the *inequality* condition (c).

THEOREM 7 (Uniqueness of weak solutions). *Assume H is C^2 and satisfies (19) and g satisfies (20). Then there exists at most one weak solution of the initial-value problem (38).*

Proof*. 1. Suppose that u and \tilde{u} are two weak solutions of (38) and write $w := u - \tilde{u}$.

Observe now that at any point (y, s) where both u and \tilde{u} are differentiable and solve our PDE, we have

$$\begin{aligned} w_t(y, s) &= u_t(y, s) - \tilde{u}_t(y, s) \\ &= -H(Du(y, s)) + H(D\tilde{u}(y, s)) \\ &= -\int_0^1 \frac{d}{dr} H(rDu(y, s) + (1-r)D\tilde{u}(y, s)) dr \\ &= -\int_0^1 DH(rDu(y, s) + (1-r)D\tilde{u}(y, s)) dr \cdot (Du(y, s) - D\tilde{u}(y, s)) \\ &=: -\mathbf{b}(y, s) \cdot Dw(y, s). \end{aligned}$$

Consequently

$$(39) \quad w_t + \mathbf{b} \cdot Dw = 0 \quad \text{a.e.}$$

*Omit on first reading.

2. Write $v := \phi(w) \geq 0$, where $\phi : \mathbb{R} \rightarrow [0, \infty)$ is a smooth function to be selected later. We multiply (39) by $\phi'(w)$ to discover

$$(40) \quad v_t + \mathbf{b} \cdot Dv = 0 \quad \text{a.e.}$$

3. Now choose $\varepsilon > 0$ and define $u^\varepsilon := \eta_\varepsilon * u$, $\tilde{u}^\varepsilon := \eta_\varepsilon * \tilde{u}$, where η_ε is the standard mollifier in the x and t variables. Then according to §C.4

$$(41) \quad |Du^\varepsilon| \leq \text{Lip}(u), \quad |D\tilde{u}^\varepsilon| \leq \text{Lip}(\tilde{u}),$$

and

$$(42) \quad Du^\varepsilon \rightarrow Du, \quad D\tilde{u}^\varepsilon \rightarrow D\tilde{u} \quad \text{a.e., as } \varepsilon \rightarrow 0.$$

Furthermore inequality (c) in the definition of weak solution implies

$$(43) \quad D^2u^\varepsilon, D^2\tilde{u}^\varepsilon \leq C \left(1 + \frac{1}{s}\right) I$$

for an appropriate constant C and all $\varepsilon > 0$, $y \in \mathbb{R}^n$, $s > 2\varepsilon$. Verification is left as an exercise.

4. Write

$$(44) \quad \mathbf{b}_\varepsilon(y, s) := \int_0^1 DH(rDu^\varepsilon(y, s) + (1-r)D\tilde{u}^\varepsilon(y, s)) dr.$$

Then (40) becomes

$$v_t + \mathbf{b}_\varepsilon \cdot Dv = (\mathbf{b}_\varepsilon - \mathbf{b}) \cdot Dv \quad \text{a.e.};$$

hence

$$(45) \quad v_t + \text{div}(v\mathbf{b}_\varepsilon) = (\text{div } \mathbf{b}_\varepsilon)v + (\mathbf{b}_\varepsilon - \mathbf{b}) \cdot Dv \quad \text{a.e.}$$

5. Now

$$(46) \quad \begin{aligned} \text{div } \mathbf{b}_\varepsilon &= \int_0^1 \sum_{k,l=1}^n H_{p_k p_l}(rDu^\varepsilon + (1-r)D\tilde{u}^\varepsilon)(ru^\varepsilon_{x_l x_k} + (1-r)\tilde{u}^\varepsilon_{x_l x_k}) dr \\ &\leq C \left(1 + \frac{1}{s}\right) \end{aligned}$$

for some constant C , in view of (41), (43). Here we note that H convex implies $D^2H \geq 0$.

6. Fix $x_0 \in \mathbb{R}^n$, $t_0 > 0$, and set

$$(47) \quad R := \max\{|DH(p)| \mid |p| \leq \max(\text{Lip}(u), \text{Lip}(\tilde{u}))\}.$$

Define also the cone

$$C := \{(x, t) \mid 0 \leq t \leq t_0, |x - x_0| \leq R(t_0 - t)\}.$$

Next write

$$e(t) = \int_{B(x_0, R(t_0-t))} v(x, t) \, dx$$

and compute for a.e. $t > 0$:

$$\begin{aligned} \dot{e}(t) &= \int_{B(x_0, R(t_0-t))} v_t \, dx - R \int_{\partial B(x_0, R(t_0-t))} v \, dS \\ &= \int_{B(x_0, R(t_0-t))} -\text{div}(v \mathbf{b}_\varepsilon) + (\text{div} \mathbf{b}_\varepsilon)v + (\mathbf{b}_\varepsilon - \mathbf{b}) \cdot Dv \, dx \\ &\quad - R \int_{\partial B(x_0, R(t_0-t))} v \, dS \quad \text{by (45)} \\ &= - \int_{\partial B(x_0, R(t_0-t))} v(\mathbf{b}_\varepsilon \cdot \nu + R) \, dS \\ &\quad + \int_{B(x_0, R(t_0-t))} (\text{div} \mathbf{b}_\varepsilon)v + (\mathbf{b}_\varepsilon - \mathbf{b}) \cdot Dv \, dx \\ &\leq \int_{B(x_0, R(t_0-t))} (\text{div} \mathbf{b}_\varepsilon)v + (\mathbf{b}_\varepsilon - \mathbf{b}) \cdot Dv \, dx \quad \text{by (41), (44)} \\ &\leq C \left(1 + \frac{1}{t}\right) e(t) + \int_{B(x_0, R(t_0-t))} (\mathbf{b}_\varepsilon - \mathbf{b}) \cdot Dv \, dx \end{aligned}$$

by (46). The last term on the right-hand side goes to zero as $\varepsilon \rightarrow 0$, for a.e. $t > 0$, according to (41), (42) and the Dominated Convergence Theorem. Thus

$$(48) \quad \dot{e}(t) \leq C \left(1 + \frac{1}{t}\right) e(t) \quad \text{for a.e. } 0 < t < t_0.$$

7. Fix $0 < \varepsilon < r < t_0$ and choose the function $\phi(z)$ to equal zero if

$$|z| \leq \varepsilon[\text{Lip}(u) + \text{Lip}(\tilde{u})]$$

and to be positive otherwise. Since $u \equiv \tilde{u}$ on $\mathbb{R}^n \times \{t = 0\}$,

$$v = \phi(w) = \phi(u - \tilde{u}) = 0 \quad \text{at } \{t = \varepsilon\}.$$

Thus $e(\varepsilon) = 0$. Consequently Gronwall's inequality (§B.2) and (48) imply

$$e(r) \leq e(\varepsilon)e^{\int_\varepsilon^r C(1+\frac{1}{s})ds} = 0.$$

Hence

$$|u - \tilde{u}| \leq \varepsilon[\text{Lip}(u) + \text{Lip}(\tilde{u})] \quad \text{on } B(x_0, R(t_0 - r)).$$

This inequality is valid for all $\varepsilon > 0$, and so $u \equiv \tilde{u}$ in $B(x_0, R(t_0 - r))$. Therefore, in particular, $u(x_0, t_0) = \tilde{u}(x_0, t_0)$. \square

In light of Lemmas 3, 4 and Theorem 7, we have

THEOREM 8 (Hopf–Lax formula as weak solution). *Suppose H is C^2 and satisfies (19) and g satisfies (20). If either g is semiconcave or H is uniformly convex, then*

$$u(x, t) = \min_{y \in \mathbb{R}^n} \left\{ tL\left(\frac{x-y}{t}\right) + g(y) \right\}$$

is the unique weak solution of the initial-value problem (38) for the Hamilton–Jacobi equation.

Examples. (i) Consider the initial-value problem:

$$(49) \quad \begin{cases} u_t + \frac{1}{2}|Du|^2 = 0 & \text{in } \mathbb{R}^n \times (0, \infty) \\ u = |x| & \text{on } \mathbb{R}^n \times \{t = 0\}. \end{cases}$$

Here $H(p) = \frac{1}{2}|p|^2$ and so $L(v) = \frac{1}{2}|v|^2$. The Hopf–Lax formula for the unique, weak solution of (49) is

$$(50) \quad u(x, t) = \min_{y \in \mathbb{R}^n} \left\{ \frac{|x-y|^2}{2t} + |y| \right\}.$$

Assume $|x| > t$. Then

$$D_y \left(\frac{|x-y|^2}{2t} + |y| \right) = \frac{y-x}{t} + \frac{y}{|y|} \quad (y \neq 0);$$

and this expression equals zero if $x = y + \frac{y}{|y|}t$, $y = (|x| - t)\frac{x}{|x|} \neq 0$. Thus $u(x, t) = |x| - \frac{t}{2}$ if $|x| > t$. If $|x| \leq t$, the minimum in (50) is attained at $y = 0$. Consequently

$$u(x, t) = \begin{cases} |x| - t/2 & \text{if } |x| \geq t \\ \frac{|x|^2}{2t} & \text{if } |x| \leq t. \end{cases}$$

Observe that the solution becomes semiconcave at times $t > 0$, even though the initial function $g(x) = |x|$ is not semiconcave. This accords with Lemma 4.

(ii) We next examine the problem with reversed initial conditions:

$$(51) \quad \begin{cases} u_t + \frac{1}{2}|Du|^2 = 0 & \text{in } \mathbb{R}^n \times (0, \infty) \\ u = -|x| & \text{on } \mathbb{R}^n \times \{t = 0\}. \end{cases}$$

Then

$$u(x, t) = \min_{y \in \mathbb{R}^n} \left\{ \frac{|x - y|^2}{2t} - |y| \right\}.$$

Now

$$D_y \left(\frac{|x - y|^2}{2t} - |y| \right) = \frac{y - x}{t} - \frac{y}{|y|} \quad (y \neq 0),$$

and this equals zero if $x = y - \frac{y}{|y|}t$, $y = (|x| + t)\frac{x}{|x|}$. Thus

$$u(x, t) = -|x| - \frac{t}{2} \quad (x \in \mathbb{R}^n, t \geq 0).$$

The initial function $g(x) = -|x|$ is semiconcave, and the solution remains so for times $t > 0$. \square

In Chapter 10 we will again study Hamilton–Jacobi PDE and discover another and better notion of weak solution, applicable even if H is not convex.

3.4. INTRODUCTION TO CONSERVATION LAWS

In this section we investigate the initial-value problem for scalar conservation laws in one space dimension:

$$(1) \quad \begin{cases} u_t + F(u)_x = 0 & \text{in } \mathbb{R} \times (0, \infty) \\ u = g & \text{on } \mathbb{R} \times \{t = 0\}. \end{cases}$$

Here $F : \mathbb{R} \rightarrow \mathbb{R}$ and $g : \mathbb{R} \rightarrow \mathbb{R}$ are given and $u : \mathbb{R} \times [0, \infty) \rightarrow \mathbb{R}$ is the unknown, $u = u(x, t)$. As noted in §3.2, the method of characteristics demonstrates that there does not in general exist a smooth solution of (1), existing for all times $t > 0$. By analogy with the developments in §3.3.3, we therefore look for some sort of weak or generalized solution.

3.4.1. Shocks, entropy condition.

a. Integral solutions; Rankine–Hugoniot condition. We open our discussion by noting that since we cannot in general find a smooth solution of (1), we must devise some way to interpret a less regular function u as somehow “solving” this initial-value problem. But as it stands, the PDE does not even make sense unless u is differentiable. However, observe that if we temporarily assume u is smooth, we can as follows rewrite, so that the resulting expression does not directly involve the derivatives of u . The idea is to multiply the PDE in (1) by a smooth function v and then to integrate by parts, thereby transferring the derivatives onto v .

More precisely, assume

$$(2) \quad v : \mathbb{R} \times [0, \infty) \rightarrow \mathbb{R} \text{ is smooth, with compact support.}$$

We call v a *test function*. Now multiply the PDE $u_t + F(u)_x = 0$ by v and integrate by parts:

$$(3) \quad \begin{aligned} 0 &= \int_0^\infty \int_{-\infty}^\infty (u_t + F(u)_x) v \, dx dt \\ &= - \int_0^\infty \int_{-\infty}^\infty uv_t \, dx dt - \int_{-\infty}^\infty uv \, dx|_{t=0} - \int_0^\infty \int_{-\infty}^\infty F(u)v_x \, dx dt. \end{aligned}$$

In view of the initial condition $u = g$ on $\mathbb{R} \times \{t = 0\}$, we thereby obtain the identity

$$(4) \quad \int_0^\infty \int_{-\infty}^\infty uv_t + F(u)v_x \, dx dt + \int_{-\infty}^\infty gv \, dx|_{t=0} = 0.$$

We derived this equality supposing u to be a smooth solution of (1), but the resulting formula has meaning even if u is only bounded.

DEFINITION. We say that $u \in L^\infty(\mathbb{R} \times (0, \infty))$ is an integral solution of (1), provided equality (4) holds for each test function v satisfying (2).

Suppose then that we have an integral solution of (1). What can we deduce about this solution from the identities (4)?

We partially answer this question by looking at a situation for which u , although not continuous, has a particularly simple structure. Let us in fact suppose in some open region $V \subset \mathbb{R} \times (0, \infty)$ that u is smooth on either side of a smooth curve C . Let V_l be that part of V on the left of the curve and let V_r be that part on the right. We assume that u is an integral solution of

(1), and that u and its first derivatives are uniformly continuous in V_l and in V_r .

First of all, choose a test function v with compact support in V_l . Then (4) becomes

$$(5) \quad 0 = \int_0^\infty \int_{-\infty}^\infty uv_t + F(u)v_x dxdt = - \int_0^\infty \int_{-\infty}^\infty [u_t + F(u)_x]v dxdt,$$

the integration by parts being justified since u is C^1 in V_l and v vanishes near the boundary of V_l . The identity (5) holds for all test functions v with compact support in V_l , and so

$$(6) \quad u_t + F(u)_x = 0 \quad \text{in } V_l.$$

Likewise,

$$(7) \quad u_t + F(u)_x = 0 \quad \text{in } V_r.$$

Jump conditions along shocks. Now select a test function v with compact support in V , but which does not necessarily vanish along the curve C . Again employing (4), we deduce


$$(8) \quad \begin{aligned} 0 &= \int_0^\infty \int_{-\infty}^\infty uv_t + F(u)v_x dxdt \\ &= \iint_{V_l} uv_t + F(u)v_x dxdt + \iint_{V_r} uv_t + F(u)v_x dxdt. \end{aligned}$$

Now since v has compact support within V , we have

$$(9) \quad \begin{aligned} \iint_{V_l} uv_t + F(u)v_x dxdt &= - \iint_{V_l} [u_t + F(u)_x]v dxdt \\ &\quad + \int_C (u_l \nu^2 + F(u_l) \nu^1) v dl \\ &= \int_C (u_l \nu^2 + F(u_l) \nu^1) v dl \end{aligned}$$

in view of (6). Here $\nu = (\nu^1, \nu^2)$ is the unit normal to the curve C , pointing from V_l into V_r , and the subscript “ l ” denotes the limit from the left. Similarly, (7) implies

$$\iint_{V_r} uv_t + F(u)v_x dxdt = - \int_C (u_r \nu^2 + F(u_r) \nu^1) v dl,$$



Rankine-Hugoniot condition

the subscript “ r ” denoting the limit from the right. Adding this identity to (9) and recalling (8) gives us

$$\int_C [(F(u_l) - F(u_r))\nu^1 + (u_l - u_r)\nu^2]v \, dl = 0.$$

This equality holds for all test functions v as above, and so

$$(10) \quad (F(u_l) - F(u_r))\nu^1 + (u_l - u_r)\nu^2 = 0 \quad \text{along } C.$$

Now suppose C is represented parametrically as $\{(x, t) \mid x = s(t)\}$ for some smooth function $s(\cdot) : [0, \infty) \rightarrow \mathbb{R}$. We can then take $\nu = (\nu^1, \nu^2) = (1 + \dot{s}^2)^{-1/2}(1, -\dot{s})$. Consequently (10) implies

$$(11) \quad F(u_l) - F(u_r) = \dot{s}(u_l - u_r)$$

in V , along the curve C .

NOTATION.

$$\begin{cases} [[u]] = u_l - u_r = \text{jump in } u \text{ across the curve } C \\ [[F(u)]] = F(u_l) - F(u_r) = \text{jump in } F(u) \\ \sigma = \dot{s} = \text{speed of the curve } C. \end{cases}$$

Let us then rewrite (11) as the identity

$$(12) \quad [[F(u)]] = \sigma[[u]]$$

along the discontinuity curve. This is the *Rankine–Hugoniot condition* along the *shock curve* C . Observe that the speed σ and the values $u_l, u_r, F(u_l)$ and $F(u_r)$ will generally vary along the curve C . The point is that even though these quantities may change, the expressions $[[F(u)]] = F(u_l) - F(u_r)$ and $\sigma[[u]] = \dot{s}(u_l - u_r)$ must always exactly balance.

Example 1 (Shock waves). Let us consider the initial-value problem for *Burgers' equation*:

$$(13) \quad \begin{cases} u_t + \left(\frac{u^2}{2}\right)_x = 0 & \text{in } \mathbb{R} \times (0, \infty) \\ u = g & \text{on } \mathbb{R} \times \{t = 0\}, \end{cases}$$

with the initial data

$$(14) \quad g(x) = \begin{cases} 1 & \text{if } x \leq 0 \\ 1 - x & \text{if } 0 \leq x \leq 1 \\ 0 & \text{if } x \geq 1. \end{cases}$$

According to the characteristic equations (cf. §3.2.5) any smooth solution u of (13), (14) takes the constant value $z^0 = g(x^0)$ along the projected characteristic

$$\mathbf{y}(s) = (g(x^0)s + x^0, s) \quad (s \geq 0)$$

for each $x^0 \in \mathbb{R}$. Thus


$$u(x, t) := \begin{cases} 1 & \text{if } x \leq t, 0 \leq t \leq 1 \\ \frac{1-x}{1-t} & \text{if } t \leq x \leq 1, 0 \leq t \leq 1 \\ 0 & \text{if } x \geq 1, 0 \leq t \leq 1. \end{cases}$$

Observe that for $t \geq 1$ this method breaks down, since the projected characteristics then cross. So how should we define u for $t \geq 1$?

Let us set $s(t) = \frac{1+t}{2}$ and write

$$u(x, t) := \begin{cases} 1 & \text{if } x < s(t) \\ 0 & \text{if } s(t) < x \end{cases}$$

if $t \geq 1$. Now along the curve parameterized by $s(\cdot)$, $u_l = 1$, we have $u_r = 0$, $F(u_l) = \frac{1}{2}(u_l)^2 = \frac{1}{2}$, $F(u_r) = 0$. Thus $[[F(u)]] = \frac{1}{2} = \sigma[[u]]$, as required by the Rankine–Hugoniot condition (12). \square



Formation of a shock

b. Shocks, entropy condition.

We try now to solve a similar problem by the same techniques.

Example 2 (Rarefaction waves and nonphysical shocks). Again consider the initial-value problem (13), for which now we take

$$(15) \quad g(x) = \begin{cases} 0 & \text{if } x < 0 \\ 1 & \text{if } x > 0. \end{cases}$$


The method of characteristics this time does not lead to any ambiguity in defining u but does fail to provide any information within the wedge $\{0 < x < t\}$. To illustrate this lack of knowledge, let us first set

$$u_1(x, t) := \begin{cases} 0 & \text{if } x < \frac{t}{2} \\ 1 & \text{if } x > \frac{t}{2}. \end{cases}$$


It is easy to check that the Rankine–Hugoniot condition holds and, indeed, that u is an integral solution of (13), (15). However, we can create another such solution by writing

$$u_2(x, t) := \begin{cases} 1 & \text{if } x > t \\ \frac{x}{t} & \text{if } 0 < x < t \\ 0 & \text{if } x < 0. \end{cases}$$

The function u_2 , called a *rarefaction wave*, is also a continuous integral solution of (13), (15). \square



A “nonphysical” shock



Rarefaction wave

Thus we see that *integral solutions are not in general unique*. Presumably the class of integral solutions includes various “nonphysical” solutions, which we want somehow to exclude. Can we find some further criterion which ensures uniqueness?

Entropy condition. Let us recall from §3.2.5 that for the general scalar conservation law of the form

$$u_t + F(u)_x = 0,$$

the solution u , whenever smooth, takes the constant value $z^0 = g(x^0)$ along the projected characteristic

$$(16) \quad \mathbf{y}(s) = (F'(g(x^0))s + x^0, s) \quad (s \geq 0).$$

Now we know that typically we will encounter the crossing of characteristics, and resultant discontinuities in the solution, if we move *forward* in time. However, we can hope that if we start at some point in $\mathbb{R} \times (0, \infty)$ and

go *backwards* in time along a characteristic, we will not cross any others. In other words, let us consider the class of, say, piecewise-smooth integral solutions of (1) with the property that if we move backwards in t along any characteristic, we will not encounter any lines of discontinuity for u .

So now suppose at some point on a curve C of discontinuities that u has distinct left and right limits, u_l and u_r , and that a characteristic from the left and a characteristic from the right hit C at this point. Then in view of (16) we deduce

$$(17) \quad F'(u_l) > \sigma > F'(u_r).$$

These inequalities are called the *entropy condition* (from a rough analogy with the thermodynamic principle that physical entropy cannot decrease as time goes forward). A curve of discontinuity for u is called a *shock* provided both the Rankine–Hugoniot identity (12) and the entropy inequalities (17) hold.

Let us further interpret the entropy condition under the additional assumption that

$$(18) \quad F \text{ is uniformly convex.}$$

This means $F'' \geq \theta > 0$ for some constant θ . Thus in particular F' is strictly increasing. Then (17) is equivalent to our requiring the inequality

$$(19) \quad u_l > u_r$$


along any shock curve. □

Example 3. We again return to Burgers' equation (13), now for the initial function

$$(20) \quad g(x) = \begin{cases} 0 & \text{if } x < 0 \\ 1 & \text{if } 0 \leq x \leq 1 \\ 0 & \text{if } x > 1. \end{cases}$$

For $0 \leq t \leq 2$, we may combine the analysis in Examples 1 and 2 above to find

$$(21) \quad u(x, t) := \begin{cases} 0 & \text{if } x < 0 \\ \frac{x}{t} & \text{if } 0 < x < t \\ 1 & \text{if } t < x < 1 + \frac{t}{2} \\ 0 & \text{if } x > 1 + \frac{t}{2} \end{cases} \quad (0 \leq t \leq 2).$$



For times $t \geq 2$, we expect the shock wave parameterized by $s(\cdot)$ to continue, with $u = x/t$ to the left of $s(\cdot)$, $u = 0$ to the right. This is compatible with the entropy condition (19). We calculate the behavior of the shock curve by applying the Rankine–Hugoniot jump condition (12). Now

$$[[u]] = \frac{s(t)}{t}, \quad [[F(u)]] = \frac{1}{2} \left(\frac{s(t)}{t} \right)^2, \quad \sigma = \dot{s}(t)$$

along the shock curve for $t \geq 0$. Thus (12) implies

$$\dot{s}(t) = \frac{s(t)}{2t} \quad (t \geq 2).$$

Additionally $s(2) = 2$, and so we can solve this ODE to find $s(t) = (2t)^{1/2}$ ($t \geq 2$). Hence we may augment (21) by setting

$$u(x, t) = \begin{cases} 0 & \text{if } x < 0 \\ \frac{x}{t} & \text{if } 0 < x < (2t)^{1/2} \\ 0 & \text{if } x > (2t)^{1/2} \end{cases} \quad (t \geq 2).$$

See the illustration. □

3.4.2. Lax–Oleinik formula.

We now try to obtain a formula for an appropriate weak solution of the initial-value problem (1), assuming as above that the flux function F is uniformly convex. With no loss of generality we may as well also take

$$(22) \quad F(0) = 0.$$

As motivation, suppose now $g \in L^\infty(\mathbb{R})$ and define

$$(23) \quad h(x) := \int_0^x g(y) dy \quad (x \in \mathbb{R}).$$

Recall the Hopf–Lax formula from §3.3 and set

$$(24) \quad w(x, t) := \min_{y \in \mathbb{R}} \left\{ tL \left(\frac{x-y}{t} \right) + h(y) \right\} \quad (x \in \mathbb{R}, t > 0),$$

where

$$(25) \quad L = F^*.$$

Thus w is the unique, weak solution of this initial-value problem for the Hamilton–Jacobi equation:

$$(26) \quad \begin{cases} w_t + F(w_x) = 0 & \text{in } \mathbb{R} \times (0, \infty) \\ w = h & \text{on } \mathbb{R} \times \{t = 0\}. \end{cases}$$

For the moment assume w is smooth. We now differentiate the PDE and its initial condition with respect to x , to deduce

$$\begin{cases} w_{xt} + F(w_x)_x = 0 & \text{in } \mathbb{R} \times (0, \infty) \\ w_x = g & \text{on } \mathbb{R} \times \{t = 0\}. \end{cases}$$

Hence if we set $u = w_x$, we discover u solves problem (1).

The foregoing computation is only formal, as we know that w defined by (24) is not in general smooth. But recall from §3.3 that w is in fact differentiable a.e. Consequently

$$(27) \quad u(x, t) := \frac{\partial}{\partial x} \left[\min_{y \in \mathbb{R}} \left\{ tL \left(\frac{x-y}{t} \right) + h(y) \right\} \right]$$

is defined for a.e. (x, t) and is presumably a leading candidate for some sort of weak solution of the initial-value problem (1). Our intention henceforth is to justify this expectation.

First, we will need to rewrite the expression (27) into a more useful form.

NOTATION. Since F is uniformly convex, F' is strictly increasing and onto. Write

$$(28) \quad G := (F')^{-1}$$

for the inverse of F' .

THEOREM 1 (Lax–Oleinik formula). *Assume $F : \mathbb{R} \rightarrow \mathbb{R}$ is smooth and uniformly convex and $g \in L^\infty(\mathbb{R})$.*

(i) *For each time $t > 0$, there exists for all but at most countably many values of $x \in \mathbb{R}$ a unique point $y(x, t)$ such that*

$$\min_{y \in \mathbb{R}} \left\{ tL \left(\frac{x - y}{t} \right) + h(y) \right\} = tL \left(\frac{x - y(x, t)}{t} \right) + h(y(x, t)).$$

(ii) *The mapping $x \mapsto y(x, t)$ is nondecreasing.*

(iii) *For each time $t > 0$, the function u defined by (27) is*

$$(29) \quad u(x, t) = G \left(\frac{x - y(x, t)}{t} \right)$$

for a.e. x . In particular, formula (29) holds for a.e. $(x, t) \in \mathbb{R} \times (0, \infty)$.

DEFINITION. *We call equation (29) the Lax–Oleinik formula for the solution (1), where h is defined by (23) and L by (25).*

Proof. 1. First, we note

$$L(v) = \max_{p \in \mathbb{R}} (vp - F(p)) = vp^* - F(p^*),$$

where $F'(p^*) = v$. But then $p^* = G(v)$ according to (28), and so

$$L(v) = vG(v) - F(G(v)) \quad (v \in \mathbb{R})$$

(cf. §3.3.1). In particular, L is C^2 . Furthermore

$$(30) \quad L'(v) = G(v) + vG'(v) - F'(G(v))G'(v) = G(v)$$

by (28), and $L''(v) = G'(v) > 0$. This and (22) imply L is nonnegative and strictly convex.

2. Fix $t > 0$, $x_1 < x_2$. As in §3.3 there exists at least one point $y_1 \in \mathbb{R}$ such that

$$(31) \quad \left\{ tL \left(\frac{x_1 - y_1}{t} \right) + h(y_1) \right\} = \min_{y \in \mathbb{R}} \left\{ tL \left(\frac{x_1 - y}{t} \right) + h(y) \right\}.$$

We next claim

$$(32) \quad tL \left(\frac{x_2 - y_1}{t} \right) + h(y_1) < tL \left(\frac{x_2 - y}{t} \right) + h(y) \quad \text{if } y < y_1.$$

To see this, we calculate $x_2 - y_1 = \tau(x_1 - y_1) + (1 - \tau)(x_2 - y)$ and $x_1 - y = (1 - \tau)(x_1 - y_1) + \tau(x_2 - y)$ for

$$0 < \tau := \frac{y_1 - y}{x_2 - x_1 + y_1 - y} < 1.$$

Since $L'' > 0$, we thus have

$$\begin{aligned} L\left(\frac{x_2 - y_1}{t}\right) &< \tau L\left(\frac{x_1 - y_1}{t}\right) + (1 - \tau)L\left(\frac{x_2 - y}{t}\right), \\ L\left(\frac{x_1 - y}{t}\right) &< (1 - \tau)L\left(\frac{x_1 - y_1}{t}\right) + \tau L\left(\frac{x_2 - y}{t}\right); \end{aligned}$$

and hence

$$(33) \quad L\left(\frac{x_2 - y_1}{t}\right) + L\left(\frac{x_1 - y}{t}\right) < L\left(\frac{x_1 - y_1}{t}\right) + L\left(\frac{x_2 - y}{t}\right).$$

Now notice from (31) that

$$tL\left(\frac{x_1 - y_1}{t}\right) + h(y_1) \leq tL\left(\frac{x_1 - y}{t}\right) + h(y).$$

We multiply (33) by t , add $h(y_1) + h(y)$ to both sides, and add the resulting expression to the above inequality to obtain (32).

3. In view of (31), in computing the minimum of $tL\left(\frac{x_2 - y}{t}\right) + h(y)$ we need only consider those $y \geq y_1$, where y_1 satisfies (31). Now for each $x \in \mathbb{R}$ and $t > 0$, define the point $y(x, t)$ to equal the smallest of those points y giving the minimum of $tL\left(\frac{x - y}{t}\right) + h(y)$. Then the mapping $x \mapsto y(x, t)$ is nondecreasing and is thus continuous for all but at most countably many x . At a point x of continuity of $y(\cdot, t)$, $y(x, t)$ is the unique value of y yielding the minimum.

4. According to the theory developed in §3.3 for each fixed $t > 0$, the mapping

$$\begin{aligned} x \mapsto w(x, t) &:= \min_{y \in \mathbb{R}} \left\{ tL\left(\frac{x - y}{t}\right) + h(y) \right\} \\ &= tL\left(\frac{x - y(x, t)}{t}\right) + h(y(x, t)) \end{aligned}$$

is differentiable a.e. Furthermore the mapping $x \mapsto y(x, t)$ is monotone and consequently differentiable a.e. as well. Thus given $t > 0$, for a.e. x the

mappings $x \mapsto L\left(\frac{x-y(x,t)}{t}\right)$ and so also $x \mapsto h(y(x,t))$ are differentiable as well.

Consequently formula (27) becomes

$$\begin{aligned} u(x,t) &= \frac{\partial}{\partial x} \left[tL\left(\frac{x-y(x,t)}{t}\right) + h(y(x,t)) \right] \\ &= L'\left(\frac{x-y(x,t)}{t}\right) (1 - y_x(x,t)) + \frac{\partial}{\partial x} h(y(x,t)). \end{aligned}$$

But since $y \mapsto tL\left(\frac{x-y}{t}\right) + h(y)$ has a minimum at $y = y(x,t)$, the mapping $z \mapsto tL\left(\frac{x-y(z,t)}{t}\right) + h(y(z,t))$ has a minimum at $z = x$. Therefore

$$-L'\left(\frac{x-y(x,t)}{t}\right) y_x(x,t) + h_x(y(x,t)) = 0,$$

and hence

$$u(x,t) = L'\left(\frac{x-y(x,t)}{t}\right) = G\left(\frac{x-y(x,t)}{t}\right),$$

according to (30). □

We now investigate the precise sense in which formula (29) provides us with a solution of the initial-value problem (1).

THEOREM 2 (Lax–Oleinik formula as integral solution). *Under the assumptions of Theorem 1, the function u defined by (29) is an integral solution of the initial-value problem (1).*

Proof. As above, define

$$w(x,t) = \min_{y \in \mathbb{R}} \left\{ tL\left(\frac{x-y}{t}\right) + h(y) \right\} \quad (x \in \mathbb{R}, t > 0).$$

Then Theorem 6 in §3.3.2 tells us w is Lipschitz continuous, is differentiable for a.e. (x,t) , and solves

$$(34) \quad \begin{cases} w_t + F(w_x) = 0 & \text{a.e. in } \mathbb{R} \times (0, \infty) \\ w = h & \text{on } \mathbb{R} \times \{t = 0\}. \end{cases}$$

Choose any test function v satisfying (2). Multiply the PDE $w_t + F(w_x) = 0$ by v_x and integrate over $\mathbb{R} \times (0, \infty)$:

$$(35) \quad 0 = \int_0^\infty \int_{-\infty}^\infty [w_t + F(w_x)] v_x \, dx dt.$$

Observe

$$\begin{aligned} \int_0^\infty \int_{-\infty}^\infty w_t v_x \, dx dt &= - \int_0^\infty \int_{-\infty}^\infty w v_{tx} \, dx dt - \int_{-\infty}^\infty w v_x \, dx|_{t=0} \\ &= \int_0^\infty \int_{-\infty}^\infty w_x v_t \, dx dt + \int_{-\infty}^\infty w_x v \, dx|_{t=0}. \end{aligned}$$

These integrations by parts are valid since the mapping $x \mapsto w(x, t)$ is Lipschitz continuous, and thus absolutely continuous, for each time $t > 0$. Likewise $t \mapsto w(x, t)$ is absolutely continuous for each $x \in \mathbb{R}$. Now $w(x, 0) = h(x) = \int_0^x g(y) \, dy$, and so $w_x(x, 0) = g(x)$ for a.e. x . Consequently

$$\int_0^\infty \int_{-\infty}^\infty w_t v_x \, dx dt = \int_0^\infty \int_{-\infty}^\infty w_x v_t \, dx dt + \int_{-\infty}^\infty g v \, dx|_{t=0}.$$

Substitute this identity into (35) and recall $u = w_x$ a.e., to derive the integral identity (4). \square

3.4.3. Weak solutions, uniqueness.

a. Entropy condition revisited. We have already seen in §3.4.1 that integral solutions of (1) are not generally unique. Since we believe the Lax–Oleinik formula does in fact provide the “correct” solution of this initial-value problem, we must see if it satisfies some appropriate form of the entropy condition discussed in §3.4.1. This is not straightforward, however, since it is not usually the case that the function u defined by the Lax–Oleinik formula is smooth, or even piecewise smooth.

We identify now a kind of “one-sided” derivative estimate for the function u defined by the Lax–Oleinik formula (27). This estimate—which is an analogue for conservation laws of the semiconcavity estimate from Lemmas 3, 4 in §3.3.3 for Hamilton–Jacobi equations—will turn out to be a uniqueness criterion.

LEMMA (A one-sided jump estimate). *Under the assumptions of Theorem 1, there exists a constant C such that the function u defined by the Lax–Oleinik formula (29) satisfies the inequality*

$$(36) \quad u(x + z, t) - u(x, t) \leq \frac{C}{t} z$$

for all $t > 0$ and $x, z \in \mathbb{R}$, $z > 0$.

DEFINITION. We call inequality (36) the entropy condition.

It follows from (36) that for $t > 0$ the function $x \mapsto u(x, t) - \frac{C}{t}x$ is nonincreasing and consequently has left- and right-hand limits at each point. Thus also $x \mapsto u(x, t)$ has left- and right-hand limits at each point, with $u_l(x, t) \geq u_r(x, t)$. In particular, the original form of the entropy condition (19) holds at any point of discontinuity.

Proof. We know from §3.3 that in computing the minimum in (29) we need only consider those y such that $|\frac{x-y}{t}| \leq C$ for some constant C ; verification is left to the reader. Consequently we may assume, upon redefining G if necessary off some bounded interval, that G is Lipschitz continuous.

As $G = (F')^{-1}$ and $y(\cdot, t)$ are nondecreasing, we have

$$\begin{aligned} u(x, t) &= G\left(\frac{x - y(x, t)}{t}\right) \\ &\geq G\left(\frac{x - y(x + z, t)}{t}\right) \quad \text{for } z > 0 \\ &\geq G\left(\frac{x + z - y(x + z, t)}{t}\right) - \frac{\text{Lip}(G)z}{t} \\ &= u(x + z, t) - \frac{\text{Lip}(G)z}{t}. \end{aligned} \quad \square$$

b. Weak solutions, uniqueness. We now establish the important assertion that an integral solution which satisfies the entropy condition is unique.

DEFINITION. We say that a function $u \in L^\infty(\mathbb{R} \times (0, \infty))$ is an entropy solution of the initial-value problem

$$(37) \quad \begin{cases} u_t + F(u)_x = 0 & \text{in } \mathbb{R} \times (0, \infty) \\ u = g & \text{on } \mathbb{R} \times \{t = 0\} \end{cases}$$

provided

$$(i) \quad \int_0^\infty \int_{-\infty}^\infty uv_t + F(u)v_x \, dxdt + \int_{-\infty}^\infty gv \, dx|_{t=0} = 0$$

for all test functions $v : \mathbb{R} \times [0, \infty) \rightarrow \mathbb{R}$ with compact support and

$$(ii) \quad u(x + z, t) - u(x, t) \leq C\left(1 + \frac{1}{t}\right)z$$

for some constant $C \geq 0$ and a.e. $x, z \in \mathbb{R}$, $t > 0$, with $z > 0$.

THEOREM 3 (Uniqueness of entropy solutions). Assume F is convex and smooth. Then there exists—up to a set of measure zero—at most one entropy solution of (37).

Proof*. 1. Assume that u and \tilde{u} are two entropy solutions of (37), and write $w := u - \tilde{u}$. Observe for any point (x, t) that

$$\begin{aligned} F(u(x, t)) - F(\tilde{u}(x, t)) &= \int_0^1 \frac{d}{dr} F(ru(x, t) + (1-r)\tilde{u}(x, t)) dr \\ &= \int_0^1 F'(ru(x, t) + (1-r)\tilde{u}(x, t)) dr (u(x, t) - \tilde{u}(x, t)) \\ &=: b(x, t)w(x, t). \end{aligned}$$

Consequently if v is a test function as above,

$$\begin{aligned} (38) \quad 0 &= \int_0^\infty \int_{-\infty}^\infty (u - \tilde{u})v_t + [F(u) - F(\tilde{u})]v_x dxdt \\ &= \int_0^\infty \int_{-\infty}^\infty w[v_t + bv_x] dxdt. \end{aligned}$$

2. Now take $\varepsilon > 0$ and define $u^\varepsilon = \eta_\varepsilon * u$, $\tilde{u}^\varepsilon = \eta_\varepsilon * \tilde{u}$, where η_ε is the standard mollifier in the x and t variables. Then according to §C.4

$$(39) \quad \|u^\varepsilon\|_{L^\infty} \leq \|u\|_{L^\infty}, \quad \|\tilde{u}^\varepsilon\|_{L^\infty} \leq \|\tilde{u}\|_{L^\infty},$$

$$(40) \quad u^\varepsilon \rightarrow u, \quad \tilde{u}^\varepsilon \rightarrow \tilde{u} \quad \text{a.e., as } \varepsilon \rightarrow 0.$$

Furthermore the entropy inequality (ii) implies

$$(41) \quad u_x^\varepsilon(x, t), \quad \tilde{u}_x^\varepsilon(x, t) \leq C \left(1 + \frac{1}{t}\right)$$

for an appropriate constant C and all $\varepsilon > 0$, $x \in \mathbb{R}$, $t > 0$.

3. Write

$$b_\varepsilon(x, t) := \int_0^1 F'(ru^\varepsilon(x, t) + (1-r)\tilde{u}^\varepsilon(x, t)) dr.$$

Then (38) becomes

$$(42) \quad 0 = \int_0^\infty \int_{-\infty}^\infty w[v_t + b_\varepsilon v_x] dxdt + \int_0^\infty \int_{-\infty}^\infty w[b - b_\varepsilon]v_x dxdt.$$

4. Now select $T > 0$ and any smooth function $\psi : \mathbb{R} \times (0, T) \rightarrow \mathbb{R}$ with compact support. We choose v^ε to be the solution of the following terminal-value problem for a linear transport equation:

$$(43) \quad \begin{cases} v_t^\varepsilon + b_\varepsilon v_x^\varepsilon = \psi & \text{in } \mathbb{R} \times (0, T) \\ v^\varepsilon = 0 & \text{on } \mathbb{R} \times \{t = T\}. \end{cases}$$

*Omit on first reading.

Let us solve (43) by the method of characteristics. For this, fix $x \in \mathbb{R}$, $0 \leq t \leq T$, and denote by $x_\varepsilon(\cdot)$ the solution of the ODE

$$(44) \quad \begin{cases} \dot{x}_\varepsilon(s) = b_\varepsilon(x_\varepsilon(s), s) & (s \geq t) \\ x_\varepsilon(t) = x, \end{cases}$$

and set

$$(45) \quad v^\varepsilon(x, t) := - \int_t^T \psi(x_\varepsilon(s), s) ds \quad (x \in \mathbb{R}, 0 \leq t \leq T).$$

Then v^ε is smooth and is the unique solution of (43). Since $|b_\varepsilon|$ is bounded and ψ has compact support, v^ε has compact support in $\mathbb{R} \times [0, T]$.

5. We now claim that for each $s > 0$, there exists a constant C_s such that

$$(46) \quad |v_x^\varepsilon| \leq C_s \quad \text{on } \mathbb{R} \times (s, T).$$

To prove this, first note that if $0 < s \leq t \leq T$, then

$$(47) \quad b_{\varepsilon,x}(x, t) = \int_0^1 F''(ru^\varepsilon + (1-r)\tilde{u}^\varepsilon)(ru_x^\varepsilon + (1-r)\tilde{u}_x^\varepsilon) dr \leq \frac{C}{t} \leq \frac{C}{s}$$

by (41), since F is convex.

Next, differentiate the PDE in (43) with respect to x :

$$(48) \quad v_{tx}^\varepsilon + b_\varepsilon v_{xx}^\varepsilon + b_{\varepsilon,x} v_x^\varepsilon = \psi_x.$$

Now set $a(x, t) := e^{\lambda t} v_x^\varepsilon(x, t)$, for

$$(49) \quad \lambda = \frac{C}{s} + 1.$$

Then

$$(50) \quad \begin{aligned} a_t + b_\varepsilon a_x &= \lambda a + e^{\lambda t} [v_{xt}^\varepsilon + b_\varepsilon v_{xx}^\varepsilon] \\ &= \lambda a + e^{\lambda t} [-b_{\varepsilon,x} v_x^\varepsilon + \psi_x] \quad \text{by (48)} \\ &= [\lambda - b_{\varepsilon,x}] a + e^{\lambda t} \psi_x. \end{aligned}$$

Since v^ε has compact support, a attains a nonnegative maximum over $\mathbb{R} \times [s, T]$ at some finite point (x_0, t_0) . If $t_0 = T$, then $v_x = 0$. If $0 \leq t_0 < T$, then

$$a_t(x_0, t_0) \leq 0, \quad a_x(x_0, t_0) = 0.$$

Consequently equation (50) gives

$$(51) \quad [\lambda - b_{\varepsilon, x}]a + e^{\lambda t_0} \psi_x \leq 0 \quad \text{at } (x_0, t_0).$$

But since $b_{\varepsilon, x} \leq \frac{C}{s}$ and λ is given by (49), inequality (51) implies

$$a(x_0, t_0) \leq -e^{\lambda t_0} \psi_x \leq e^{\lambda T} \|\psi_x\|_{L^\infty}.$$

A similar argument shows

$$a(x_1, t_1) \geq -e^{\lambda T} \|\psi_x\|_{L^\infty}$$

at any point (x_1, t_1) where a attains a nonpositive minimum. These two estimates and the definition of a imply (46).

6. We will need one more inequality, namely

$$(52) \quad \int_{-\infty}^{\infty} |v_x^\varepsilon(x, t)| dx \leq D$$

for all $0 \leq t \leq \tau$ and some constant D , provided τ is small enough.

To prove this, choose $\tau > 0$ so small that $\psi = 0$ on $\mathbb{R} \times (0, \tau)$. Then if $0 \leq t \leq \tau$, we see from (45) that v^ε is constant along the characteristic curve $x_\varepsilon(\cdot)$ (solving (44)) for $t \leq s \leq \tau$. Select any partition $x_0 < x_1 < \cdots < x_N$. Then $y_0 < y_1 < \cdots < y_N$, where $y_i := x_i^\varepsilon(\tau)$ ($i = 1, \dots, N$) for

$$\begin{cases} \dot{x}_i^\varepsilon(s) = b_\varepsilon(x_i^\varepsilon(s), s) & (t \leq s \leq \tau) \\ x_i^\varepsilon(t) = x_i. \end{cases}$$

As v^ε is constant along each characteristic curve $x_i^\varepsilon(\cdot)$, we have

$$\sum_{i=1}^N |v^\varepsilon(x_i, t) - v^\varepsilon(x_{i-1}, t)| = \sum_{i=1}^N |v^\varepsilon(y_i, \tau) - v^\varepsilon(y_{i-1}, \tau)| \leq \text{var } v^\varepsilon(\cdot, \tau),$$

“var” denoting variation with respect to x . Taking the supremum over all such partitions, we find

$$\int_{-\infty}^{\infty} |v_x^\varepsilon(x, t)| dx = \text{var } v^\varepsilon(\cdot, t) \leq \text{var } v^\varepsilon(\cdot, \tau) = \int_{-\infty}^{\infty} |v_x^\varepsilon(x, \tau)| dx \leq C,$$

since v^ε has compact support and estimate (46) is valid for $s = \tau$.

7. Now, at last, we complete the proof by setting $v = v^\varepsilon$ in (42) and substituting, using (43):

$$\begin{aligned} \int_0^\infty \int_{-\infty}^\infty w\psi \, dxdt &= \int_0^\infty \int_{-\infty}^\infty w[b_\varepsilon - b]v_x^\varepsilon \, dxdt \\ &= \int_\tau^T \int_{-\infty}^\infty w[b_\varepsilon - b]v_x^\varepsilon \, dxdt \\ &\quad + \int_0^\tau \int_{-\infty}^\infty w[b_\varepsilon - b]v_x^\varepsilon \, dxdt \\ &=: I_\tau^\varepsilon + J_\tau^\varepsilon. \end{aligned}$$

Then in view of (40), (46), and the Dominated Convergence Theorem,

$$I_\tau^\varepsilon \rightarrow 0 \quad \text{as } \varepsilon \rightarrow 0$$

for each $\tau > 0$. On the other hand, if $0 < \tau < T$, we see

$$|J_\tau^\varepsilon| \leq \tau C \max_{0 \leq t \leq \tau} \int_{-\infty}^\infty |v_x^\varepsilon| \, dx \leq \tau C, \quad \text{by (52).}$$

Thus

$$\int_0^\infty \int_{-\infty}^\infty w\psi \, dxdt = 0$$

for all smooth functions ψ as above, and so $w = u - \tilde{u} = 0$ a.e. □

3.4.4. Riemann's problem.

The initial-value problem (1) with the piecewise-constant initial function

$$(53) \quad g(x) = \begin{cases} u_l & \text{if } x < 0 \\ u_r & \text{if } x > 0 \end{cases}$$


is called *Riemann's problem* for the scalar conservation law (1). Here the constants u_l, u_r are the left and right *initial states*, $u_l \neq u_r$.

We continue to assume F is uniformly convex and C^2 , and as before we write $G = (F')^{-1}$.

THEOREM 4 (Solution of Riemann's problem).

(i) If $u_l > u_r$, the unique entropy solution of the Riemann problem (1), (53) is

$$(54) \quad u(x, t) := \begin{cases} u_l & \text{if } \frac{x}{t} < \sigma \\ u_r & \text{if } \frac{x}{t} > \sigma \end{cases} \quad (x \in \mathbb{R}, t > 0),$$



Shock wave solving Riemann's problem for $u_l > u_r$

where

$$(55) \quad \sigma := \frac{F(u_l) - F(u_r)}{u_l - u_r}.$$

(ii) If $u_l < u_r$, the unique entropy solution of the Riemann problem (1), (53) is

$$(56) \quad u(x, t) := \begin{cases} u_l & \text{if } \frac{x}{t} < F'(u_l) \\ G\left(\frac{x}{t}\right) & \text{if } F'(u_l) < \frac{x}{t} < F'(u_r) \\ u_r & \text{if } \frac{x}{t} > F'(u_r) \end{cases} \quad (x \in \mathbb{R}, t > 0).$$


Shocks and rarefactions. In the first case the states u_l and u_r are separated by a *shock wave* with constant speed σ . In the second case the states u_l and u_r are separated by a *rarefaction wave*.

We know from the theory set forth in §§3.4.2–3.4.3 that the Lax–Oleinik formula must generate these solutions, and it is an interesting exercise to verify this directly. We will instead construct the functions (54), (56) from first principles and verify they are in fact entropy solutions. By uniqueness, then, they must agree with Lax–Oleinik formulas. This is a nice illustration of the power of the uniqueness assertion, Theorem 3.

Proof. 1. Assume $u_l > u_r$. Clearly u defined by (54), (55) is then an integral solution of our PDE. In particular since $\sigma = \frac{[F(u)]}{[u]}$, the Rankine–Hugoniot condition holds. Furthermore note

$$F'(u_r) < \sigma = \frac{F(u_l) - F(u_r)}{u_l - u_r} = \int_{u_r}^{u_l} F'(r) dr < F'(u_l)$$

in accordance with (17). Since $u_l > u_r$, the entropy condition holds as well. Uniqueness follows from Theorem 3.



Rarefaction wave solving Riemann's problem for $u_l < u_r$

2. Assume now that $u_l < u_r$. We must first check that u defined by (56) solves the conservation law in the region $\{F'(u_l) < \frac{x}{t} < F'(u_r)\}$. To verify this, let us ask the general question as to when a function u of the form

$$u(x, t) = v\left(\frac{x}{t}\right)$$

solves (1). We compute

$$\begin{aligned} u_t + F(u)_x &= u_t + F'(u)u_x \\ &= -v'\left(\frac{x}{t}\right) \frac{x}{t^2} + F'(v)v'\left(\frac{x}{t}\right) \frac{1}{t} \\ &= v'\left(\frac{x}{t}\right) \frac{1}{t} \left[F'(v) - \frac{x}{t} \right]. \end{aligned}$$

Thus, assuming v' never vanishes, we find $F'(v(\frac{x}{t})) = \frac{x}{t}$. Hence

$$u(x, t) = v\left(\frac{x}{t}\right) = G\left(\frac{x}{t}\right)$$

solves the conservation law. Now $v(\frac{x}{t}) = u_l$ provided $\frac{x}{t} = F'(u_l)$, and similarly $v(\frac{x}{t}) = u_r$ if $\frac{x}{t} = F'(u_r)$.

As a consequence we see that the rarefaction wave u defined by (56) is continuous in $\mathbb{R} \times (0, \infty)$ and is a solution of the PDE $u_t + F(u)_x = 0$ in each of its regions of definition. It is easy to check that u is thus an integral solution of (1), (53). Furthermore, since as noted in §3.4.3 we may as well assume G is Lipschitz continuous, we have

$$u(x+z, t) - u(x, t) = G\left(\frac{x+z}{t}\right) - G\left(\frac{x}{t}\right) \leq \frac{\text{Lip}(G)z}{t}$$

if $F'(u_l)t < x < x+z < F'(u_r)t$. This inequality implies that u also satisfies the entropy condition. Uniqueness is once more a consequence of Theorem 3.

□

3.4.5. Long time behavior.

a. Decay in sup-norm. We now employ the Lax–Oleinik formula (29) to study the behavior of our entropy solution u of (1) as $t \rightarrow \infty$. We assume below that F is smooth, uniformly convex, $F(0) = 0$, and g is bounded and summable.

THEOREM 5 (Asymptotics in L^∞ -norm). *There exists a constant C such that*

$$(57) \quad |u(x, t)| \leq \frac{C}{t^{1/2}}$$

for all $x \in \mathbb{R}$, $t > 0$.

Proof. 1. Set

$$(58) \quad \sigma := F'(0);$$

then

$$(59) \quad G(\sigma) = 0,$$

and therefore

$$(60) \quad L(\sigma) = \sigma G(\sigma) - F(G(\sigma)) = 0, \quad L'(\sigma) = 0.$$

2. In view of (60) and the uniform convexity of L ,

$$(61) \quad \begin{aligned} tL\left(\frac{x-y}{t}\right) &= tL\left(\frac{x-y-\sigma t}{t} + \sigma\right) \\ &\geq t\left[L(\sigma) + L'(\sigma)\left(\frac{x-y-\sigma t}{t}\right) + \theta\left(\frac{x-y-\sigma t}{t}\right)^2\right] \\ &= \theta\frac{|x-y-\sigma t|^2}{t} \end{aligned}$$

for some constant $\theta > 0$. Since $h = \int_0^x g \, dy$ is bounded by $M := \|g\|_{L^1}$, we see from (61) that

$$tL\left(\frac{x-y}{t}\right) + h(y) \geq \theta\frac{|x-y-\sigma t|^2}{t} - M.$$

On the other hand,

$$tL\left(\frac{x-(x-\sigma t)}{t}\right) + h(x-\sigma t) \leq M.$$

Thus at the minimizing point $y(x, t)$ we have

$$\theta \frac{|x - y(x, t) - \sigma t|^2}{t} \leq 2M,$$

and so

$$(62) \quad \left| \frac{x - y(x, t)}{t} - \sigma \right| \leq \frac{C}{t^{1/2}}$$

for some constant C .

3. But since $G(\sigma) = 0$, for any $x \in \mathbb{R}$, $t > 0$ we have

$$\begin{aligned} |u(x, t)| &= \left| G\left(\frac{x - y(x, t)}{t}\right) \right| \\ &= \left| G\left(\frac{x - y(x, t)}{t} - \sigma + \sigma\right) - G(\sigma) \right| \\ &\leq \text{Lip}(G) \left| \frac{x - y(x, t)}{t} - \sigma \right| \leq \frac{C}{t^{1/2}}, \end{aligned}$$

according to (62). □

Example 3 in §3.4.1 shows this $t^{-1/2}$ decay rate to be optimal.

b. Decay to N-wave. Estimate (57) asserts that the L^∞ -norm of u goes to zero as $t \rightarrow \infty$. On the other hand we note from Example 3 in §3.4.1 that the L^1 -norm of u need not go to zero; indeed, the integral of u over \mathbb{R} is conserved (Problem 19). We instead show here that u evolves in L^1 into a simple shape, assuming now that

g has compact support.


Given constants p, q, d, σ , with $p, q \geq 0$, $d > 0$, we define the corresponding *N-wave* to be the function

$$(63) \quad N(x, t) := \begin{cases} \frac{1}{d} \left(\frac{x}{t} - \sigma\right) & \text{if } -(pdt)^{1/2} < x - \sigma t < (qdt)^{1/2} \\ 0 & \text{otherwise.} \end{cases}$$

The constant σ is the *velocity* of the *N-wave*.

Now define σ by (58), set

$$(64) \quad d := F''(0) > 0,$$



N-wave

and also write

$$(65) \quad p := -2 \min_{y \in \mathbb{R}} \int_{-\infty}^y g \, dx, \quad q := 2 \max_{y \in \mathbb{R}} \int_y^{\infty} g \, dx.$$

Note $p, q \geq 0$ and

$$(66) \quad G'(\sigma) = \frac{1}{d}.$$

THEOREM 6 (Asymptotics in L^1 -norm). *Assume that $p, q > 0$. Then there exists a constant C such that*

$$(67) \quad \int_{-\infty}^{\infty} |u(\cdot, t) - N(\cdot, t)| \, dx \leq \frac{C}{t^{1/2}}$$

for all $t > 0$.

Proof. 1. From estimate (62) in the proof of Theorem 5 we have

$$(68) \quad \left| \frac{(x - \sigma t) - y(x, t)}{t} \right| \leq \frac{C}{t^{1/2}}.$$

Now

$$\begin{aligned} u(x, t) &= G\left(\frac{x - y(x, t)}{t}\right) \\ &= G\left(\frac{(x - \sigma t) - y(x, t)}{t} + \sigma\right) \\ &= G(\sigma) + G'(\sigma) \left(\frac{(x - \sigma t) - y(x, t)}{t}\right) \\ &\quad + O\left(\left|\frac{(x - \sigma t) - y(x, t)}{t}\right|^2\right). \end{aligned}$$

Consequently (59), (66) and (68) imply

$$(69) \quad \left| u(x, t) - \frac{1}{d} \frac{(x - \sigma t) - y(x, t)}{t} \right| \leq \frac{C}{t}.$$

2. Since g has compact support, we may assume for some constant $R > 0$ that $g \equiv 0$ on $\mathbb{R} \cap \{|x| \geq R\}$. Therefore

$$h(x) = \begin{cases} h_- & \text{if } x \leq -R \\ h_+ & \text{if } x \geq R, \end{cases}$$

for constants h_{\pm} . A calculation shows

$$(70) \quad \min_{\mathbb{R}} h = -\frac{p}{2} + h_- = -\frac{q}{2} + h_+.$$

We next set

$$(71) \quad \varepsilon = \varepsilon(t) := \frac{A}{t^{1/2}} \quad (t > 0),$$

the constant A to be selected later.

3. We now claim that if A is sufficiently large, then

$$(72) \quad u(x, t) = 0 \quad \text{for } x - \sigma t < -R - (pd(1 + \varepsilon)t)^{1/2}$$

and

$$(73) \quad u(x, t) = 0 \quad \text{for } x - \sigma t > R + (qd(1 + \varepsilon)t)^{1/2}.$$

In fact, since (64) implies

$$L''(\sigma) = \frac{1}{d},$$

we deduce from (60) and (62) that

$$(74) \quad tL\left(\frac{x - y(x, t)}{t}\right) = \frac{1}{d} \frac{|(x - \sigma t) - y(x, t)|^2}{2t} + O\left(t^{-1/2}\right) \quad \text{as } t \rightarrow \infty.$$

Assume now that

$$(75) \quad x - \sigma t < -R - (pd(1 + \varepsilon)t)^{1/2}.$$

Then $h(x - \sigma t) = h_-$ and so

$$tL\left(\frac{(x - (x - \sigma t))}{t}\right) + h(x - \sigma t) = tL(\sigma) + h_- = h_-.$$

Now if $y(x, t) \leq -R$, then

$$tL \left(\frac{x - y(x, t)}{t} \right) + h(y(x, t)) \geq h_-,$$

since $L \geq 0$. On the other hand if $y(x, t) \geq -R$, we employ (74) and (70) to estimate

$$\begin{aligned} tL \left(\frac{x - y(x, t)}{t} \right) + h(y(x, t)) &\geq \frac{1}{d} \frac{|(x - \sigma t) - y(x, t)|^2}{2t} - \frac{p}{2} + h_- + O(t^{-1/2}) \\ &\geq \frac{pd(1 + \varepsilon)t}{2dt} - \frac{p}{2} + h_- + O(t^{-1/2}) \quad \text{by (75)} \\ &= \frac{p}{2} \frac{A}{t^{1/2}} + h_- + O(t^{-1/2}) \quad \text{by (71)} \\ &\geq h_-, \end{aligned}$$

provided A is large enough.

We conclude that (75) forces $y(x, t) = x - \sigma t$, and so $u(x, t) = G(\sigma) = 0$. This establishes assertion (72), and the proof of (73) is analogous.

4. Next we assert for A and t large enough that

$$(76) \quad y(x, t) \geq -R \quad \text{if } x - \sigma t = R - (pd(1 - \varepsilon)t)^{1/2}.$$

To see this, notice that $y(x, t) \leq -R$ implies as above that

$$tL \left(\frac{x - y(x, t)}{t} \right) + h(y(x, t)) \geq h_-.$$

Select now a point z such that $h(z) = \min h = -\frac{p}{2} + h_-$ and $|z| \leq R$. Then we can as before estimate

$$\begin{aligned} tL \left(\frac{x - z}{t} \right) + h(z) &\leq \frac{1}{d} \frac{|(x - \sigma t) - z|^2}{2t} - \frac{p}{2} + h_- + O(t^{-1/2}) \\ &\leq \frac{pd(1 - \varepsilon)t}{2dt} - \frac{p}{2} + h_- + O(t^{-1/2}) \\ &= -\frac{p}{2} \frac{A}{t^{1/2}} + h_- + O(t^{-1/2}) < h_-, \end{aligned}$$

for A large enough. This proves (76) and a similar argument establishes that

$$(77) \quad y(x, t) \leq R \quad \text{if } x - \sigma t = -R + (qd(1 - \varepsilon)t)^{1/2}.$$

5. Remember from the proof of Theorem 1 in §3.4.2 that the mapping $x \mapsto y(x, t)$ is nondecreasing. Hence (69), (76) and (77) imply for large t that

$$(78) \quad \begin{cases} |u(x, t) - \frac{1}{d} \left(\frac{x}{t} - \sigma \right)| \leq \frac{C}{t} & \text{if} \\ R - (pd(1 - \varepsilon)t)^{1/2} < x - \sigma t < -R + (qd(1 - \varepsilon)t)^{1/2}. \end{cases}$$

According to Theorem 5, we have $|u| = O(t^{-\frac{1}{2}})$ and by definition $|N| = O(t^{-\frac{1}{2}})$. In addition (71) implies $((1 \pm \varepsilon)t)^{\frac{1}{2}} - t^{\frac{1}{2}} = O(1)$. Using these bounds along with (72), (73) and (78), we estimate

$$\int_{-\infty}^{\infty} |u(x, t) - N(x, t)| dx = O\left(t^{-1/2}\right),$$

as asserted. □

Example 3 (continued). Observe that we have $p = 0$, $q = 2$, $\sigma = 0$, $d = 1$ in Example 3 of §3.4.1. In this case

$$N(x, t) = \begin{cases} \frac{x}{t} & \text{if } 0 < x < (2t)^{1/2} \\ 0 & \text{otherwise,} \end{cases}$$

and so in fact $u \equiv N$ for times $t \geq 2$. □

We will study systems of conservation laws in Chapter 11.

3.5. PROBLEMS

In the following exercises, all given functions are assumed smooth, unless otherwise stated.

1. Prove

$$u(x, t, a, b) = a \cdot x - tH(a) + b \quad (a \in \mathbb{R}^n, b \in \mathbb{R})$$

is a complete integral of the Hamilton–Jacobi equation

$$u_t + H(Du) = 0.$$

2. Compute the envelopes of the family of lines

$$x_1 + a^2 x_2 - 2a = 0 \quad (a \in \mathbb{R})$$

in \mathbb{R}^2 and of the family of planes

$$2a_1 x_1 + 2a_2 x_2 - x_3 + a_1^2 + a_2^2 = 0 \quad (a_1, a_2 \in \mathbb{R})$$

in \mathbb{R}^3 . Draw pictures illustrating the geometric meaning of the envelopes.

3. Suppose that the formula $G(x, z, a) = 0$ implicitly defines the function $z = u(x, a)$, where $x, a \in \mathbb{R}^n$. Assume further that we can eliminate the variables a from the identities

$$\begin{cases} G(x, u, a) = 0 \\ G_{x_i}(x, u, a) + G_z(x, u, a)u_{x_i} = 0 \quad (i = 1, \dots, n), \end{cases}$$

to solve for $u = u(x)$.

(a) Find a PDE that u solves if $G = \sum_{i=1}^n a_i x_i^2 + z^3$.

(b) What is the PDE characterizing all spheres in \mathbb{R}^{n+1} with unit radius and center in $\mathbb{R}^n \times \{z = 0\}$?

4. (a) Write down the characteristic equations for the PDE

$$(*) \quad u_t + b \cdot Du = f \quad \text{in } \mathbb{R}^n \times (0, \infty),$$

where $b \in \mathbb{R}^n$, $f = f(x, t)$.

(b) Use the characteristic ODE to solve (*) subject to the initial condition

$$u = g \quad \text{on } \mathbb{R}^n \times \{t = 0\}.$$

Make sure your answer agrees with formula (5) in §2.1.2.

5. Solve using characteristics:

$$(a) \quad x_1 u_{x_1} + x_2 u_{x_2} = 2u, \quad u(x_1, 1) = g(x_1).$$

$$(b) \quad x_1 u_{x_1} + 2x_2 u_{x_2} + u_{x_3} = 3u, \quad u(x_1, x_2, 0) = g(x_1, x_2).$$

$$(c) \quad u u_{x_1} + u_{x_2} = 1, \quad u(x_1, x_1) = \frac{1}{2} x_1.$$

6. Given a smooth vector field \mathbf{b} on \mathbb{R}^n , let $\mathbf{x}(s) = \mathbf{x}(s, x, t)$ solve the ODE

$$\begin{cases} \dot{\mathbf{x}} = \mathbf{b}(\mathbf{x}) & (s \in \mathbb{R}) \\ \mathbf{x}(t) = x. \end{cases}$$

- (a) Define the Jacobian

$$J(s, x, t) := \det D_x \mathbf{x}(s, x, t)$$

and derive *Euler's formula*:

$$J_s = \operatorname{div} \mathbf{b}(\mathbf{x}) J.$$

- (b) Demonstrate that

$$u(x, t) := g(\mathbf{x}(0, x, t)) J(0, x, t)$$

solves

$$\begin{cases} u_t + \operatorname{div}(u\mathbf{b}) = 0 & \text{in } \mathbb{R}^n \times (0, \infty) \\ u = g & \text{on } \mathbb{R}^n \times \{t = 0\}. \end{cases}$$

(Hint: Show $\frac{\partial}{\partial s}(u(\mathbf{x}, s)J) = 0$.)

7. Verify assertion (36) in §3.2.3, that when Γ is not flat near x^0 , the noncharacteristic condition is

$$D_p F(p^0, z^0, x^0) \cdot \nu(x^0) \neq 0.$$

8. Confirm that the formula $u = g(x - t\mathbf{F}'(u))$ from §3.2.5 provides an implicit solution for the conservation law

$$u_t + F(u)_x = 0.$$

9. Consider the problem of minimizing the action $\int_0^t L(\dot{\mathbf{w}}(s), \mathbf{w}(s)) ds$ over the new admissible class

$$\mathcal{A} := \{\mathbf{w}(\cdot) \in C^2([0, t]; \mathbb{R}^n) \mid \mathbf{w}(t) = x\},$$

where we do not require that $\mathbf{w}(0) = y$.

- (a) Show that a minimizer $\mathbf{x}(\cdot) \in \mathcal{A}$ solves the Euler–Lagrange equations

$$-\frac{d}{ds}(D_v L(\dot{\mathbf{x}}(s), \mathbf{x}(s))) + D_x L(\dot{\mathbf{x}}(s), \mathbf{x}(s)) = 0 \quad (0 \leq s \leq t).$$

- (b) Prove that

$$D_v L(\dot{\mathbf{x}}(0), \mathbf{x}(0)) = 0.$$

- (c) Suppose now that $\mathbf{x}(\cdot) \in \mathcal{A}$ minimizes the modified action

$$\int_0^t L(\dot{\mathbf{w}}(s), \mathbf{w}(s)) ds + g(\mathbf{w}(0)).$$

Show that $\mathbf{x}(\cdot)$ solves the usual Euler–Lagrange equations and determine the boundary condition at $s = 0$.

10. If $H : \mathbb{R}^n \rightarrow \mathbb{R}$ is convex, we write $L = H^*$.

- (a) Let $H(p) = \frac{1}{r}|p|^r$, for $1 < r < \infty$. Show

$$L(v) = \frac{1}{s}|v|^s, \quad \text{where } \frac{1}{r} + \frac{1}{s} = 1.$$

- (b) Let $H(p) = \frac{1}{2} \sum_{i,j=1}^n a_{ij} p_i p_j + \sum_{i=1}^n b_i p_i$, where $A = ((a_{ij}))$ is a symmetric, positive definite matrix, $b \in \mathbb{R}^n$. Compute $L(v)$.

11. Let $H : \mathbb{R}^n \rightarrow \mathbb{R}$ be convex. We say v belongs to the *subdifferential* of H at p , written

$$v \in \partial H(p),$$

if

$$H(r) \geq H(p) + v \cdot (r - p) \quad \text{for all } r \in \mathbb{R}^n.$$

Prove $v \in \partial H(p)$ if and only if $p \in \partial L(v)$ if and only if $p \cdot v = H(p) + L(v)$, where $L = H^*$.

12. Assume $L_1, L_2 : \mathbb{R}^n \rightarrow \mathbb{R}$ are convex, smooth and superlinear. Show that

$$\min_{v \in \mathbb{R}^n} (L_1(v) + L_2(v)) = \max_{p \in \mathbb{R}^n} (-H_1(p) - H_2(-p)),$$

where $H_1 = L_1^*, H_2 = L_2^*$.

13. Prove that the Hopf–Lax formula reads

$$\begin{aligned} u(x, t) &= \min_{y \in \mathbb{R}^n} \left\{ tL \left(\frac{x - y}{t} \right) + g(y) \right\} \\ &= \min_{y \in B(x, Rt)} \left\{ tL \left(\frac{x - y}{t} \right) + g(y) \right\} \end{aligned}$$

for $R = \sup_{\mathbb{R}^n} |DH(Dg)|$, $H = L^*$. (This proves *finite propagation speed* for a Hamilton–Jacobi PDE with convex Hamiltonian and Lipschitz continuous initial function g .)

14. Let E be a closed subset of \mathbb{R}^n . Show that if the Hopf–Lax formula could be applied to the initial-value problem

$$\begin{cases} u_t + |Du|^2 = 0 & \text{in } \mathbb{R}^n \times (0, \infty) \\ u = \begin{cases} 0 & x \in E \\ +\infty & x \notin E \end{cases} & \text{on } \mathbb{R}^n \times \{t = 0\}, \end{cases}$$

it would give the solution

$$u(x, t) = \frac{1}{4t} \text{dist}(x, E)^2.$$

15. Provide all details for the proof of Lemma 4 in §3.3.3.
16. Assume u^1, u^2 are two solutions of the initial-value problems

$$\begin{cases} u_t^i + H(Du^i) = 0 & \text{in } \mathbb{R}^n \times (0, \infty) \\ u^i = g^i & \text{on } \mathbb{R}^n \times \{t = 0\} \quad (i = 1, 2), \end{cases}$$

given by the Hopf–Lax formula. Prove the L^∞ -contraction inequality

$$\sup_{\mathbb{R}^n} |u^1(\cdot, t) - u^2(\cdot, t)| \leq \sup_{\mathbb{R}^n} |g^1 - g^2| \quad (t > 0).$$

17. Show that

$$u(x, t) := \begin{cases} -\frac{2}{3} \left(t + \sqrt{3x + t^2} \right) & \text{if } 4x + t^2 > 0 \\ 0 & \text{if } 4x + t^2 < 0 \end{cases}$$

is an (unbounded) entropy solution of $u_t + \left(\frac{u^2}{2}\right)_x = 0$.

18. Assume $u(x+z) - u(x) \leq Ez$ for all $z > 0$. Let $u^\epsilon = \eta_\epsilon * u$, and show

$$u_x^\epsilon \leq E.$$

19. Assume $F(0) = 0$, u is a continuous integral solution of the conservation law

$$\begin{cases} u_t + F(u)_x = 0 & \text{in } \mathbb{R} \times (0, \infty) \\ u = g & \text{on } \mathbb{R} \times \{t = 0\}, \end{cases}$$

and u has compact support in $\mathbb{R} \times [0, T]$ for each time $T > 0$. Prove

$$\int_{-\infty}^{\infty} u(\cdot, t) dx = \int_{-\infty}^{\infty} g dx$$

for all $t > 0$.

20. Compute explicitly the unique entropy solution of

$$\begin{cases} u_t + \left(\frac{u^2}{2}\right)_x = 0 & \text{in } \mathbb{R} \times (0, \infty) \\ u = g & \text{on } \mathbb{R} \times \{t = 0\}, \end{cases}$$

for

$$g(x) = \begin{cases} 1 & \text{if } x < -1 \\ 0 & \text{if } -1 < x < 0 \\ 2 & \text{if } 0 < x < 1 \\ 0 & \text{if } x > 1. \end{cases}$$

Draw a picture documenting your answer, being sure to illustrate what happens for all times $t > 0$.

3.6. REFERENCES

Section 3.1 A nice source for this material is Courant–Hilbert [C-H, Chapter 2].

Section 3.2 This derivation of the characteristic differential equations is found in Carathéodory [C]. The proof of Theorem 2 follows Garabedian [G, Chapter 2], John [J2, Chapter 1], etc. Chester [Ch] and Sneddon [Sn] are also good texts for more

on first-order PDE. Example 3 in §3.2.2 is from Zwillinger [Zw].

Section 3.3 See Lions [Li], Rund [Ru] and Benton [Be] for more on Hamilton–Jacobi PDE. The uniqueness proof, which is due to A. Douglis, is from [Be].

Section 3.4 See Lax [Lx1] and Smoller [S, Chapters 15,16] (from which I took the proof of Theorem 3, due to O. Oleinik). Theorems 5 and 6 are from DiPerna (Indiana U. Math. J. 24 (1975), 1047–1071) and I am indebted to M. Struwe and to E. Wiedemann for help with the proofs. A good overall reference on nonlinear waves is Whitham [Wh].

Section 3.5 Problem 3 is based upon Miller [Mi].

OTHER WAYS TO REPRESENT SOLUTIONS

- 4.1 Separation of variables
- 4.2 Similarity solutions
- 4.3 Transform methods
- 4.4 Converting nonlinear into linear PDE
- 4.5 Asymptotics
- 4.6 Power series
- 4.7 Problems
- 4.8 References

This chapter collects together a wide variety of techniques that are sometimes useful for finding certain more-or-less explicit solutions to various partial differential equations, or at least representation formulas for solutions.

4.1. SEPARATION OF VARIABLES

The method of *separation of variables* tries to construct a solution u to a given partial differential equation as some sort of combination of functions of fewer variables. In other words, the idea is to guess that u can be written as, say, a sum or product of as yet undetermined constituent functions, to plug this guess into the PDE, and finally to choose the simpler functions to ensure u really is a solution.

4.1.1. Examples.

The separation of variables technique is best understood in some examples.

Example 1. Let $U \subset \mathbb{R}^n$ be a bounded, open set with smooth boundary. We consider the initial/boundary-value problem for the heat equation

$$(1) \quad \begin{cases} u_t - \Delta u = 0 & \text{in } U \times (0, \infty) \\ u = 0 & \text{on } \partial U \times [0, \infty) \\ u = g & \text{on } U \times \{t = 0\}, \end{cases}$$

where $g : U \rightarrow \mathbb{R}$ is given. We conjecture there exists a solution having the multiplicative form

$$(2) \quad u(x, t) = v(t)w(x) \quad (x \in U, t \geq 0);$$

that is, we look for a solution of (1) with the variables $x = (x_1, \dots, x_n) \in U$ “separated” from the variable $t \in [0, T]$.

Will this work? To find out, we compute

$$u_t(x, t) = v'(t)w(x), \quad \Delta u(x, t) = v(t)\Delta w(x).$$

Hence

$$0 = u_t(x, t) - \Delta u(x, t) = v'(t)w(x) - v(t)\Delta w(x)$$

if and only if

$$(3) \quad \frac{v'(t)}{v(t)} = \frac{\Delta w(x)}{w(x)}$$

for all $x \in U$ and $t > 0$ such that $w(x), v(t) \neq 0$. Now observe that the left-hand side of (3) depends only on t and the right-hand side depends only on x . This is impossible unless each is constant, say

$$\frac{v'(t)}{v(t)} = \mu = \frac{\Delta w(x)}{w(x)} \quad (t \geq 0, x \in U).$$

Then

$$(4) \quad v' = \mu v,$$

$$(5) \quad \Delta w = \mu w.$$

We must solve these equations for the unknowns w, v and μ .

Notice first that if μ is known, the solution of (4) is $v = de^{\mu t}$ for an arbitrary constant d . Consequently we need only investigate equation (5).

We say that λ is an *eigenvalue* of the operator $-\Delta$ on U (subject to zero boundary conditions) provided there exists a function w , not identically equal to zero, solving

$$\begin{cases} -\Delta w = \lambda w & \text{in } U \\ w = 0 & \text{on } \partial U. \end{cases}$$

The function w is a corresponding *eigenfunction*. (See Chapter 6 for the theory of eigenvalues, eigenfunctions.)

If λ is an eigenvalue and w is a related eigenfunction, we set $\mu = -\lambda$ above, to find

$$(6) \quad u = de^{-\lambda t}w$$

solves

$$(7) \quad \begin{cases} u_t - \Delta u = 0 & \text{in } U \times (0, \infty) \\ u = 0 & \text{on } \partial U \times [0, \infty), \end{cases}$$

with the initial condition $u(\cdot, 0) = dw$. Thus the function u defined by (6) solves problem (1), provided $g = dw$. More generally, if $\lambda_1, \dots, \lambda_m$ are eigenvalues, w_1, \dots, w_m are corresponding eigenfunctions, and d_1, \dots, d_m are constants, then

$$(8) \quad u = \sum_{k=1}^m d_k e^{-\lambda_k t} w_k$$

solves (7), with the initial condition $u(\cdot, 0) = \sum_{k=1}^m d_k w_k$. If we can find m, w_1, \dots , etc. such that $\sum_{k=1}^m d_k w_k = g$, we are done.

We can hope to generalize further by trying to find a countable sequence λ_1, \dots of eigenvalues with corresponding eigenfunctions w_1, \dots , so that

$$(9) \quad \sum_{k=1}^{\infty} d_k w_k = g \quad \text{in } U$$

for appropriate constants d_1, \dots . Then presumably

$$(10) \quad u = \sum_{k=1}^{\infty} d_k e^{-\lambda_k t} w_k$$

will be the solution of the initial-value problem (1).

This is an attractive representation formula for the solution, but depends upon (a) our being able to find eigenvalues, eigenfunctions and constants satisfying (9) and (b) our verifying that the series in (10) converges in some appropriate sense. We will discuss these matters further in Chapters 6, 7, within the context of Galerkin approximations. \square

Take note that only our solution (6) is determined by separation of variables; the more complicated forms (8) and (10) depend upon the linearity of the heat equation.

Example 2. Let us next apply the separation of variables technique to discover a solution of the *porous medium equation*

$$(11) \quad u_t - \Delta(u^\gamma) = 0 \quad \text{in } \mathbb{R}^n \times (0, \infty),$$

where $u \geq 0$ and $\gamma > 1$ is a constant. The expression (11) is a nonlinear diffusion equation, in which the rate of diffusion of some density u depends upon u itself. This PDE describes flow in porous media, thin-film lubrication, and a variety of other phenomena.

As in the previous example, we seek a solution of the form

$$(12) \quad u(x, t) = v(t)w(x) \quad (x \in \mathbb{R}^n, t \geq 0).$$

Inserting into (11), we discover that

$$(13) \quad \frac{v'(t)}{v(t)^\gamma} = \mu = \frac{\Delta w^\gamma(x)}{w(x)}$$

for some constant μ and all $x \in \mathbb{R}^n$, $t \geq 0$, such that $w(x), v(t) \neq 0$.

We solve the ODE for v and find

$$v = ((1 - \gamma)\mu t + \lambda)^{\frac{1}{1-\gamma}},$$

for some constant λ , which we will take to be positive. To discover w , we must then solve the PDE

$$(14) \quad \Delta(w^\gamma) = \mu w.$$

Let us now guess that

$$w = |x|^\alpha,$$

for some constant α that must be determined. Then

$$(15) \quad \mu w - \Delta(w^\gamma) = \mu|x|^\alpha - \alpha\gamma(\alpha\gamma + n - 2)|x|^{\alpha\gamma-2}.$$

So in order that (14) hold in \mathbb{R}^n , we should first require that $\alpha = \alpha\gamma - 2$, and hence

$$(16) \quad \alpha = \frac{2}{\gamma - 1}.$$

Returning to (15), we see that we must further set

$$(17) \quad \mu = \alpha\gamma(\alpha\gamma + n - 2) > 0.$$

In summary then, for each $\lambda > 0$ the function

$$u = ((1 - \gamma)\mu t + \lambda)^{\frac{1}{1-\gamma}} |x|^\alpha$$

solves the porous medium equation (11), the parameters α , μ defined by (16), (17). \square

Remark. Observe that since $\gamma > 1$, this solution blows up for $x \neq 0$ as $t \rightarrow t_*$, for $t_* := \frac{\lambda}{(\gamma-1)\mu}$. Physically, a huge amount of mass “diffuses in from infinity” in finite time. See §4.2.2 for another, better behaved, solution of the porous medium equation, and see §9.4.1 for more on blow-up phenomena for nonlinear diffusion equations.

In the previous example separation of variables worked owing to the homogeneity of the nonlinearity, which is compatible with functions u having the multiplicative form (12). In other circumstances it is profitable to look for a solution in which the variables are separated additively:

Example 3. Let us turn once again to the Hamilton–Jacobi equation

$$(18) \quad u_t + H(Du) = 0 \quad \text{in } \mathbb{R}^n \times (0, \infty)$$

and look for a solution u having the form

$$u(x, t) = w(x) + v(t) \quad (x \in \mathbb{R}^n, t \geq 0).$$

Then

$$0 = u_t(x, t) + H(Du(x, t)) = v'(t) + H(Dw(x))$$

if and only if

$$H(Dw(x)) = \mu = -v'(t) \quad (x \in \mathbb{R}^n, t > 0)$$

for some constant μ . Consequently if

$$H(Dw) = \mu$$

for some $\mu \in \mathbb{R}$, then

$$u(x, t) = w(x) - \mu t + b$$

will for any constant b solve $u_t + H(Du) = 0$. In particular, if we choose $w(x) = a \cdot x$ for some $a \in \mathbb{R}^n$ and set $\mu = H(a)$, we discover the solution

$$u = a \cdot x - H(a)t + b$$

already noted in §3.1. \square

4.1.2. Application: Turing instability.

Separations of variables and eigenfunction expansions, discussed in Example 1 above, are very powerful tools in both pure and applied mathematics. This section discusses an interesting such application.

Assume we are given a smooth vector field $\mathbf{f} = (f^1, f^2)$ on \mathbb{R}^2 for which $\mathbf{0}$ is an equilibrium:

$$\mathbf{f}(\mathbf{0}) = \mathbf{0}.$$

We are interested in comparing the stability of solutions $\mathbf{x} = (x^1, x^2)$ of the system of ODE

$$(19) \quad \dot{\mathbf{x}} = \mathbf{f}(\mathbf{x}) \quad (t \geq 0)$$

with solutions $\mathbf{u} = (u^1, u^2)$ of a corresponding reaction-diffusion system of PDE

$$(20) \quad \begin{cases} \mathbf{u}_t - A\Delta\mathbf{u} = \mathbf{f}(\mathbf{u}) & \text{in } U \times (0, \infty) \\ \mathbf{u} = 0 & \text{on } \partial U \times (0, \infty) \end{cases}$$

in some bounded, smooth region $U \subset \mathbb{R}^2$. The matrix

$$A = \begin{pmatrix} a_1 & 0 \\ 0 & a_2 \end{pmatrix}$$

introduces the diffusion constants $a_1, a_2 \geq 0$. (See §9.2.1 for more on reaction-diffusion equations.)

Linearizations, separation of variables. The linearization of (19) around the equilibrium solution $\mathbf{x} \equiv \mathbf{0}$ is the linear system of ODE

$$(21) \quad \dot{\mathbf{y}} = D\mathbf{f}(\mathbf{0})\mathbf{y} \quad (t \geq 0)$$

where $\mathbf{y} = (y^1, y^2)$. The equilibrium $\mathbf{x} \equiv \mathbf{0}$ is asymptotically stable if each solution \mathbf{y} goes to zero as $t \rightarrow \infty$. This will be so provided the eigenvalues of the matrix $D\mathbf{f}(\mathbf{0})$ have negative real parts.

Similarly, the linearization of (20) around $\mathbf{u} \equiv \mathbf{0}$ is the linear system of PDE

$$(22) \quad \mathbf{v}_t - A\Delta\mathbf{v} = D\mathbf{f}(\mathbf{0})\mathbf{v}$$

for $\mathbf{v} = (v^1, v^2)$. We solve (22) by the separation of variables and subsequent eigenfunction expansion method introduced in §4.1.1. We therefore write

$$(23) \quad \mathbf{v}(x, t) = \sum_{j=1}^{\infty} \mathbf{s}_j(t)w_j(x)$$

for the eigenfunctions $\{w_j\}_{j=1}^{\infty}$ for the Laplacian on U with zero boundary conditions:

$$\begin{cases} -\Delta w_j = \lambda_j w_j & \text{in } U \\ w_j = 0 & \text{on } \partial U. \end{cases}$$

The theory of such eigenvalues and eigenfunctions appears in §6.5, where we will learn in particular that

$$\lambda_j > 0 \quad (j = 1, \dots)$$

and also that we can take $\{w_j\}_{j=1}^{\infty}$ to be orthonormal in $L^2(U)$:

$$\int_U w_i w_j \, dx = \delta_{ij} \quad (i, j = 1, \dots).$$

Plugging (23) into (22), we deduce that for $j = 1, \dots$

$$(24) \quad \mathbf{s}'_j = A_j \mathbf{s}_j$$

for the matrix

$$(25) \quad A_j := D\mathbf{f}(0) - \lambda_j A.$$

The solution $\mathbf{v} \equiv 0$ is stable if and only if each function \mathbf{s}_j decays to 0 as $t \rightarrow \infty$. This occurs provided the eigenvalues of the matrices A_j have negative real parts for $j = 1, \dots$.

We now address the following question: if 0 is an asymptotically stable equilibrium for the system of ODE (19), does it necessarily follow that 0 is an asymptotically stable equilibrium for the system of PDE (20)? The perhaps surprising answer is “no”. The diffusion terms introduced into the PDE (20) can in fact transform a stable point for (19) into an unstable point for (20). This effect is called a *Turing instability*.

Eigenvalues of $D\mathbf{f}(0)$. We investigate this phenomenon by first introducing explicit conditions on $D\mathbf{f}(0)$ that force 0 to be stable for the ODE (19). Let us hereafter write

$$D\mathbf{f}(0) = \begin{pmatrix} f_{z_1}^1(0) & f_{z_2}^1(0) \\ f_{z_1}^2(0) & f_{z_2}^2(0) \end{pmatrix} =: \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix}.$$

Then

$$(26) \quad \det(D\mathbf{f}(0) - \sigma I) = \sigma^2 - \sigma(\alpha + \delta) + \alpha\delta - \gamma\beta.$$

We require that 0 be a stable equilibrium of $Df(0)$ and thus that the roots σ_1, σ_2 of (26) have negative real parts. This means

$$(27) \quad \begin{cases} \alpha + \delta = \sigma_1 + \sigma_2 < 0 \\ \alpha\delta - \gamma\beta = \sigma_1\sigma_2 > 0. \end{cases}$$

Notice that these conditions cover both the cases of real eigenvalues $\sigma_2 \leq \sigma_1 < 0$ and of complex conjugate eigenvalues $\sigma_1 = b + ic, \sigma_2 = b - ic$ with $b < 0$.

Eigenvalues of A_j . We want to see if by adding in diffusion terms $a_1, a_2 \geq 0$ we can force the eigenvalues of A_j to have positive real parts for some j .

We see from (25) that

$$(28) \quad \det(A_j - \sigma I) = \sigma^2 - \sigma(\alpha + \delta - \lambda_j(a_1 + a_2)) + p(\lambda_j)$$

for

$$(29) \quad p(\lambda) := \lambda^2 a_1 a_2 - \lambda(a_1 \delta + a_2 \alpha) + \alpha \delta - \beta \gamma.$$

The roots $\sigma_{1,j}$ and $\sigma_{2,j}$ of the polynomial (28) satisfy

$$\sigma_{1,j} + \sigma_{2,j} = \alpha + \delta - \lambda_j(a_1 + a_2) < 0,$$

since $\alpha + \delta < 0$ according (27), $\lambda_j \geq 0$, $a_1, a_2 > 0$. Consequently for the case of complex conjugate roots $\sigma_{1,j} = b_j + ic_j, \sigma_{2,j} = b_j - ic_j$, the real part b_j is negative. In this circumstance solutions \mathbf{s}_j of the ODE (24) tend to zero as $t \rightarrow \infty$ and we have asymptotic stability: this is not what we are looking for.

Loss of stability. Consequently the only way the PDE system (22) could lose stability is when we have real roots $\sigma_{2,j} \leq \sigma_{1,j}$. We want to try to select $a_1, a_2 \geq 0$ in this case so that $\sigma_{1,j} \geq 0$. Let us imagine starting out with $a_1 = a_2 = 0$ and then increasing these diffusion constants until the system (22) first begins to lose stability, when $\sigma_{1,j} = 0$. This happens provided

$$(30) \quad p(\lambda_j) = 0.$$

We seek algebraic conditions implying (30). We may assume without loss of generality that

$$(31) \quad \delta < 0.$$

Then if $a_2 = 0$, we would have

$$p(\lambda_j) = -\lambda_j a_1 \delta + \alpha \delta - \beta \gamma > 0$$

according to (27). Thus we must have $a_2 > 0$; that is, *we must introduce some diffusion into the second equation of our PDE system (22) for the Turing instability to occur.*

Now if $\alpha \leq 0$, then (27) forces

$$p(\lambda_j) = \lambda_j^2 a_1 a_2 - \lambda_j (a_1 \delta + a_2 \alpha) + (\alpha \delta - \beta \gamma) > 0,$$

and so we cannot achieve (30). Consequently we must require

$$(32) \quad \alpha > 0.$$

We keep (31) and (32) in mind and rewrite the formula $p(\lambda_j) = 0$ to read

$$(33) \quad a_2 = \frac{\alpha \delta - \gamma \beta - \delta \lambda_j a_1}{\lambda_j (\alpha - \lambda_j a_1)}.$$

But then given some $\lambda_j > 0$, we can easily find $a_2 > 0$ and $a_1 \geq 0$ solving (33). Notice that if $\lambda_j \gg 0$, we will need to take a_1 to be small enough to ensure that $\alpha - \lambda_j a_1 > 0$.

Interpretation: activators and inhibitors. The sign conditions

$$\begin{cases} \alpha = f_{z_1}^1(0) > 0, & \beta = f_{z_2}^1(0) < 0 \\ \gamma = f_{z_1}^2(0) > 0, & \delta = f_{z_2}^2(0) < 0 \end{cases}$$

are consistent with our requirements (27), (31) and (32). We may then interpret u^1 as the density of a chemical *activator* and u^2 as the density of an *inhibitor*: since $\alpha > 0$, the activator by itself would increase; but since $\beta < 0$, this growth can be offset by the inhibitor. The signs of γ, δ imply that the inhibitor increases only in response to the presence of the activator. Condition (27) means that activator/inhibitor balance holds for the ODE (19), at least near the origin.

We have discovered that diffusion effects can upset this equilibrium, provided a_2 is sufficiently large and a_1 sufficiently small. The physical interpretation is that the inhibitor u^2 diffuses away from any given point more rapidly than the activator u^1 , and consequently there need not be enough of the inhibitor present to prevent runaway growth of the activator. Such reaction-diffusion instabilities are sometimes proposed as simple models for biological pattern formation: see for example Markowich [Mr].

4.2. SIMILARITY SOLUTIONS

When investigating partial differential equations, it is often profitable to look for specific solutions u , the form of which reflects various symmetries in the structure of the PDE. We have already seen this idea in our derivation of the fundamental solutions for Laplace's equation and the heat equation in §2.2.1 and §2.3.1 and our discovery of rarefaction waves for conservation laws in §3.4.4. Following are some further applications of this important method.

4.2.1. Plane and traveling waves, solitons.

Consider first a partial differential equation involving the two variables $x \in \mathbb{R}$, $t \in \mathbb{R}$. A solution u of the form

$$(1) \quad u(x, t) = v(x - \sigma t) \quad (x \in \mathbb{R}, t \in \mathbb{R})$$

is called a *traveling wave* (with *speed* σ and *profile* v). More generally, a solution u of a PDE in the $n + 1$ variables $x = (x_1, \dots, x_n) \in \mathbb{R}^n$, $t \in \mathbb{R}$ having the form

$$(2) \quad u(x, t) = v(y \cdot x - \sigma t) \quad (x \in \mathbb{R}^n, t \in \mathbb{R})$$

is called a *plane wave* (with *wavefront* normal to $y \in \mathbb{R}^n$, *speed* $\frac{\sigma}{|y|}$, and *profile* v).

a. Exponential solutions. In view of the Fourier transform (discussed later, in §4.3.1), it is particularly enlightening when studying linear partial differential equations to consider complex-valued plane wave solutions of the form

$$(3) \quad u(x, t) = e^{i(y \cdot x - \sigma t)},$$

where $\sigma \in \mathbb{C}$ and $y = (y_1, \dots, y_n) \in \mathbb{R}^n$, σ being the *time frequency* and $\{y_i\}_{i=1}^n$ the *wave numbers*. We will next substitute trial solutions of the form (3) into various linear PDE, paying particular attention to the relationship between y and $\sigma = \sigma(y)$ forced by the structure of the equation.

(i) Heat equation. If u is given by (3), we compute

$$u_t - \Delta u = (-i\sigma + |y|^2)u = 0,$$

provided $\sigma = -i|y|^2$. Hence

$$u = e^{iy \cdot x - |y|^2 t}$$

solves the heat equation for each $y \in \mathbb{R}^n$. Taking real and imaginary parts, we discover further that $e^{-|y|^2 t} \cos(y \cdot x)$ and $e^{-|y|^2 t} \sin(y \cdot x)$ are solutions as well. Notice in this example that since σ is purely imaginary, there results a real, negative exponential term $e^{-|y|^2 t}$ in the formulas, which corresponds to damping or *dissipation*.

(ii) Wave equation, Klein–Gordon equation. Upon our substituting (3) into the wave equation, we discover

$$u_{tt} - \Delta u = (-\sigma^2 + |y|^2)u = 0,$$

provided $\sigma = \pm|y|$. Consequently

$$u = e^{i(y \cdot x \pm |y|t)}$$

solves the wave equation, as do the pair of functions $\cos(y \cdot x \pm |y|t)$ and $\sin(y \cdot x \pm |y|t)$. Since σ is real, there are no dissipation effects in these solutions; and the absolute value of propagation speed $\frac{|\sigma|}{|y|} = 1$ of each such solution is the same.

Turning next to the *Klein–Gordon equation*

$$u_{tt} - \Delta u + m^2 u = 0,$$

our inserting (3) yields

$$u_{tt} - \Delta u + m^2 u = (-\sigma^2 + |y|^2 + m^2)u = 0$$

for $\sigma = \pm(|y|^2 + m^2)^{\frac{1}{2}}$. However notice now that the speed $\frac{\sigma}{|y|}$ of propagation depends nonlinearly upon the frequency of the initial value $e^{iy \cdot x}$, the slower oscillating solutions traveling faster. That waves of different frequencies propagate at different speeds means that the Klein–Gordon equation creates *dispersion*.

(iii) Other dispersive equations. Putting $u = e^{i(y \cdot x - \sigma t)}$ into *Schrödinger's equation*

$$iu_t + \Delta u = 0,$$

we compute

$$iu_t + \Delta u = (\sigma - |y|^2)u = 0$$

when $\sigma = |y|^2$. Therefore

$$u = e^{i(y \cdot x - |y|^2 t)},$$

and so this solution displays dispersion.

For a final example of a dispersive PDE, let $n = 1$ and substitute $u = e^{i(yx - \sigma t)}$ into *Airy's equation*

$$u_t + u_{xxx} = 0.$$

We calculate

$$u_t + u_{xxx} = -i(\sigma + y^3)u = 0,$$

whenever $\sigma = -y^3$.

Phase and group velocities. For a general dispersive linear PDE with constant coefficients, we can in principle compute as above $\sigma = \sigma(y)$. We sometimes refer to $\frac{\sigma(y)}{|y|}$ as the *phase velocity* of the exponential plane-wave solution (3): this is the speed of propagation in the direction of the unit vector $\frac{y}{|y|}$.

However, we will see in §4.3.1 that we can often use the Fourier transform to write more general solutions of our PDE as a linear superposition of such exponential plane-wave solutions:

$$u(x, t) = \int_{\mathbb{R}^n} e^{i(y \cdot x - \sigma(y)t)} a(y) dy$$

for some appropriate function a . To understand the speed of propagation of u , let us consider the limit $t \rightarrow \infty$, while the ratio $v := \frac{x}{t}$ is held fixed. We will learn later in §4.5.3 on stationary phase that the main contribution to the integral

$$\int_{\mathbb{R}^n} e^{i(y \cdot x - \sigma(y)t)} a(y) dy = \int_{\mathbb{R}^n} e^{it(y \cdot v - \sigma(y))} a(y) dy$$

occurs for wave numbers y for which $D\sigma(y) = v$. For this reason, we call $D\sigma(y)$ the *group velocity*.

b. Solitons. We consider next the *Korteweg-de Vries* (KdV) equation in the form

$$(4) \quad u_t + 6uu_x + u_{xxx} = 0 \quad \text{in } \mathbb{R} \times (0, \infty),$$

this nonlinear dispersive equation being a model for surface waves in water. We seek a traveling wave solution having the structure

$$(5) \quad u(x, t) = v(x - \sigma t) \quad (x \in \mathbb{R}, t > 0).$$

Then u solves the KdV equation (4), provided v satisfies the ODE

$$(6) \quad -\sigma v' + 6vv' + v''' = 0 \quad \left(' = \frac{d}{ds} \right).$$

We integrate (6) by first noting

$$(7) \quad -\sigma v + 3v^2 + v'' = a,$$

a denoting some constant. Multiply this equality by v' to obtain

$$-\sigma vv' + 3v^2v' + v''v' = av',$$

and so deduce

$$(8) \quad \frac{(v')^2}{2} = -v^3 + \frac{\sigma}{2}v^2 + av + b$$

where b is another arbitrary constant.

We investigate (8) by looking now only for solutions v which satisfy $v, v', v'' \rightarrow 0$ as $s \rightarrow \pm\infty$ (in which case the function u having the form (5) is called a *solitary* wave). Then (7), (8) imply $a = b = 0$. Equation (8) thereupon simplifies to read

$$\frac{(v')^2}{2} = v^2 \left(-v + \frac{\sigma}{2} \right).$$

Hence $v' = \pm v(\sigma - 2v)^{1/2}$.

We take the minus sign above for computational convenience and obtain then this implicit formula for v :

$$(9) \quad s = - \int_1^{v(s)} \frac{dz}{z(\sigma - 2z)^{1/2}} + c,$$

for some constant c . Now substitute $z = \frac{\sigma}{2} \operatorname{sech}^2 \theta$. It follows that $\frac{dz}{d\theta} = -\sigma \operatorname{sech}^2 \theta \tanh \theta$ and $z(\sigma - 2z)^{1/2} = \frac{\sigma^{3/2}}{2} \operatorname{sech}^2 \theta \tanh \theta$. Hence (9) becomes

$$(10) \quad s = \frac{2}{\sqrt{\sigma}} \theta + c,$$

where θ is implicitly given by the relation

$$(11) \quad \frac{\sigma}{2} \operatorname{sech}^2 \theta = v(s).$$

We lastly combine (10) and (11), to compute

$$v(s) = \frac{\sigma}{2} \operatorname{sech}^2 \left(\frac{\sqrt{\sigma}}{2} (s - c) \right) \quad (s \in \mathbb{R}).$$

Conversely, it is routine to check that v so defined actually solves the ODE (6). The upshot is that

$$u(x, t) = \frac{\sigma}{2} \operatorname{sech}^2\left(\frac{\sqrt{\sigma}}{2}(x - \sigma t - c)\right) \quad (x \in \mathbb{R}, t \geq 0)$$


is a solution of the KdV equation for each $c \in \mathbb{R}$, $\sigma > 0$. A solution of this form is called a *soliton*. Notice that the velocity of the soliton depends upon its height. \square

The KdV equation is in fact utterly remarkable, in that it is *completely integrable*, which means that in principle the exact solution can be computed for arbitrary initial data. The relevant techniques are mostly beyond the scope of this book, but see Problems 11 and 12 and also Drazin [Dr] for more information.

c. Traveling waves for a bistable equation. Consider next the scalar reaction-diffusion equation

$$(12) \quad u_t - u_{xx} = f(u) \quad \text{in } \mathbb{R} \times (0, \infty),$$

where $f : \mathbb{R} \rightarrow \mathbb{R}$ has a “cubic-like” shape.



Graph of the function f

We assume, more precisely, that f is smooth and verifies

$$(13) \quad \begin{cases} (a) & f(0) = f(a) = f(1) = 0 \\ (b) & f < 0 \text{ on } (0, a), f > 0 \text{ on } (a, 1) \\ (c) & f'(0) < 0, f'(1) < 0 \\ (d) & \int_0^1 f(z) dz > 0 \end{cases}$$

for some point $0 < a < 1$.

We look for a traveling wave solution of the form

$$(14) \quad u(x, t) = v(x - \sigma t),$$

the profile v and velocity σ to be determined, such that

$$u \rightarrow 0 \quad \text{as } x \rightarrow -\infty, \quad u \rightarrow 1 \quad \text{as } x \rightarrow +\infty.$$

Now since $f' < 0$ at $z = 0, 1$, the constants 0 and 1 are stable solutions of the PDE (and since $f' \geq 0$ at $z = a$, the constant a is an unstable solution). So we want our traveling wave (14) to interpolate between the two stable states $z = 0, 1$ at $x = \mp\infty$.

Plugging (14) into (12), we see v must satisfy the ordinary differential equation

$$(15) \quad v'' + \sigma v' + f(v) = 0 \quad \left(' = \frac{d}{ds} \right),$$

subject to the conditions

$$(16) \quad \lim_{s \rightarrow +\infty} v(s) = 1, \quad \lim_{s \rightarrow -\infty} v(s) = 0, \quad \lim_{s \rightarrow \pm\infty} v'(s) = 0.$$

We outline now (without complete proofs) a *phase plane analysis* of the ODE problem (15), (16). We begin by setting

$$w := v'.$$

Then (15), (16) transform into the autonomous first-order system:

$$(17) \quad \begin{cases} v' = w \\ w' = -\sigma w - f(v), \end{cases}$$

with


$$(18) \quad \lim_{s \rightarrow \infty} (v, w) = (1, 0), \quad \lim_{s \rightarrow -\infty} (v, w) = (0, 0).$$

Now $(0, 0)$ and $(1, 0)$ are critical points for the system (17), and the eigenvalues of the corresponding linearizations are

$$(19) \quad \lambda_0^\pm = \frac{-\sigma \pm (\sigma^2 - 4f'(0))^{1/2}}{2}, \quad \lambda_1^\pm = \frac{-\sigma \pm (\sigma^2 - 4f'(1))^{1/2}}{2}.$$

In view of (13)(c), $\lambda_0^\pm, \lambda_1^\pm$ are real, with differing sign, and thus $(0, 0)$ and $(1, 0)$ are saddle points for the flow (17). Consequently an “unstable curve” W^u leaves $(0, 0)$ and a “stable curve” W^s approaches $(1, 0)$, as drawn. Furthermore, by calculating eigenvectors corresponding to (19) we see

$$(20) \quad \begin{cases} W^u \text{ is tangent to the line } w = \lambda_0^+ v \text{ at } (0, 0) \\ W^s \text{ is tangent to the line } w = \lambda_1^- (v - 1) \text{ at } (1, 0). \end{cases}$$



Stable and unstable curves

Note that $\lambda_0^\pm, \lambda_1^\pm, W^u$ and W^s depend upon the parameter σ . Our intention is to find $\sigma < 0$ so that

$$(21) \quad W^u = W^s \quad \text{in the region } \{v > 0, w > 0\}.$$

Then we will have a solution of (17), (18), whose path in the phase plane is a heteroclinic orbit connecting $(0, 0)$ to $(1, 0)$.

To establish (21), we fix now a small number $\varepsilon > 0$ and let L denote the vertical line through the point $(a + \varepsilon, 0)$. We claim

$$(22) \quad W^s \cap L \neq \emptyset, \quad W^u \cap L \neq \emptyset$$

if $\sigma < 0$. To check this assertion, define



$$E(v, w) := \frac{w^2}{2} + \int_0^v f(z) dz \quad (v, w \in \mathbb{R})$$

and compute

$$\begin{aligned} \frac{d}{dt} E(v(t), w(t)) &= w(t)w'(t) + f(v(t))v'(t) \\ &= -\sigma w^2(t) \quad \text{by (17)}. \end{aligned}$$

As $\sigma < 0$, we see that E is nondecreasing along trajectories of the ODE (17). Note also that the level sets of E have the shapes illustrated.

Consider next the region R , as drawn below. The unstable curve enters R from $(0, 0)$ and cannot exit through the bottom, top or left-hand side.

Level curves of E The region R


Using (17), we deduce that W^u must exit R through the line L , at a point $(a + \varepsilon, w_0(\sigma))$. Similarly we argue W^s must hit L at a point $(a + \varepsilon, w_1(\sigma))$. This verifies claim (22).

We next observe

$$(23) \quad w_0(0) < w_1(0);$$

this follows since trajectories of (17) for $\sigma = 0$ are contained in level sets of E . We assert further that

$$(24) \quad w_0(\sigma) > w_1(\sigma)$$

The region S

provided $\sigma < 0$ and $|\sigma|$ is large enough. To see this, fix $\beta > 0$ and consider the region S , as drawn.

Now along the line segment $T := \{0 \leq v \leq a + \varepsilon, w = \beta v\}$, we have

$$\frac{w'}{v'} = \frac{-\sigma w - f(v)}{w} = -\sigma - \frac{f(v)}{\beta v}.$$

Since $\left| \frac{f(v)}{v} \right|$ is bounded for $0 \leq v \leq a + \varepsilon$, we see

$$(25) \quad \frac{w'}{v'} \geq -\sigma - \frac{C}{\beta} > \beta \quad \text{on } T,$$

provided $\sigma < 0$ and $|\sigma|$ is large enough.

The calculation (25) shows that W^u cannot exit S through the line segment T , and so $w_0(\sigma) \geq \beta(a + \varepsilon)$ if $\sigma = \sigma(\beta)$ is sufficiently negative. On the other hand, $w_1(\sigma) \leq w_1(0)$ for all $\sigma \leq 0$. Thus we see that (24) will follow once we choose β large enough and then σ sufficiently negative.

Since w_0 and w_1 depend smoothly on σ , we deduce from (23) and (24) that there exists $\sigma < 0$ with

$$(26) \quad w_0(\sigma) = w_1(\sigma).$$

For this velocity σ there consequently exists a solution of the ODE (17), (18). Hence we have found for our reaction-diffusion PDE (12) a traveling wave of the form (14). \square

A more refined analysis demonstrates that the velocity σ verifying (26) is unique. Hence given the nonlinearity f satisfying hypotheses (13), there

exists a *unique* velocity for which there is a corresponding traveling wave. Compare this assertion with the previous example, where we found soliton traveling waves of the KdV equation for each given velocity.

4.2.2. Similarity under scaling.

We next illustrate the possibility of finding other types of “similarity” solutions to PDE.

Example (A scaling invariant solution). Consider again the *porous medium equation*

$$(27) \quad u_t - \Delta(u^\gamma) = 0 \quad \text{in } \mathbb{R}^n \times (0, \infty),$$

where $u \geq 0$ and $\gamma > 1$ is a constant.

As in our earlier derivation of the fundamental solution of the heat equation in §2.3.1, let us look for a solution u having the form

$$(28) \quad u(x, t) = \frac{1}{t^\alpha} v\left(\frac{x}{t^\beta}\right) \quad (x \in \mathbb{R}^n, t > 0),$$

where the constants α, β and the function $v : \mathbb{R}^n \rightarrow \mathbb{R}$ must be determined. Remember that we come upon (28) if we seek a solution u of (27) invariant under the *dilation scaling*

$$u(x, t) \mapsto \lambda^\alpha u(\lambda^\beta x, \lambda t);$$

so that

$$u(x, t) = \lambda^\alpha u(\lambda^\beta x, \lambda t)$$

for all $\lambda > 0$, $x \in \mathbb{R}^n$, $t > 0$. Setting $\lambda = t^{-1}$, we obtain (28) for $v(y) := u(y, 1)$.

We insert (28) into (27) and discover

$$(29) \quad \alpha t^{-(\alpha+1)} v(y) + \beta t^{-(\alpha+1)} y \cdot Dv(y) + t^{-(\alpha\gamma+2\beta)} \Delta(v^\gamma)(y) = 0$$

for $y = t^{-\beta} x$. In order to convert (29) into an expression involving the variable y alone, let us require

$$(30) \quad \alpha + 1 = \alpha\gamma + 2\beta.$$

Then (29) reduces to

$$(31) \quad \alpha v + \beta y \cdot Dv + \Delta(v^\gamma) = 0.$$

At this point we have effected a reduction from $n + 1$ to n variables. We simplify further by supposing v is radial; that is, $v(y) = w(|y|)$ for some $w : \mathbb{R} \rightarrow \mathbb{R}$. Then (31) becomes

$$(32) \quad \alpha w + \beta r w' + (w^\gamma)'' + \frac{n-1}{r} (w^\gamma)' = 0,$$

where $r = |y|$, $' = \frac{d}{dr}$. Now if we set

$$(33) \quad \alpha = n\beta,$$

(32) thereupon simplifies to read

$$(r^{n-1} (w^\gamma)')' + \beta (r^n w)' = 0.$$

Thus

$$r^{n-1} (w^\gamma)' + \beta r^n w = a$$

for some constant a . Assuming $\lim_{r \rightarrow \infty} w, w' = 0$, we conclude $a = 0$; whence

$$(w^\gamma)' = -\beta r w.$$

But then

$$(w^{\gamma-1})' = -\frac{(\gamma-1)}{\gamma} \beta r.$$

Consequently

$$w^{\gamma-1} = b - \frac{\gamma-1}{2\gamma} \beta r^2,$$

b a constant; and so

$$(34) \quad w = \left(b - \frac{\gamma-1}{2\gamma} \beta r^2 \right)^{+\frac{1}{\gamma-1}},$$

where we took the positive part of the right-hand side of (34) to ensure $w \geq 0$. Recalling $v(y) = w(r)$ and (28), we obtain

$$(35) \quad u(x, t) = \frac{1}{t^\alpha} \left(b - \frac{\gamma-1}{2\gamma} \beta \frac{|x|^2}{t^{2\beta}} \right)^{+\frac{1}{\gamma-1}} \quad (x \in \mathbb{R}^n, t > 0),$$

where, from (30), (33),

$$(36) \quad \alpha = \frac{n}{n(\gamma-1)+2}, \quad \beta = \frac{1}{n(\gamma-1)+2}.$$

The formulas (35), (36) are the *Barenblatt–Kompaneetz–Zeldovich solution* of the porous medium equation. \square

Finite propagation speed and degenerate diffusions. Observe that the Barenblatt–Kompaneetz–Zeldovich solution has compact support for each time $t > 0$. This is a general feature for (appropriately defined) weak, nonnegative solutions of the porous medium equation with compactly supported initial data. The nonlinear parabolic PDE (27) becomes degenerate wherever $u = 0$, and the set $\{u > 0\}$ moves with *finite propagation speed*. Consequently the porous medium equation (27) is often regarded as a better model of diffusive spreading than the linear heat equation, which predicts infinite propagation speed.

4.3. TRANSFORM METHODS

In this section we develop some of the theory for the Fourier transform \mathcal{F} , the Radon transform \mathcal{R} and the Laplace transform \mathcal{L} . These provide extremely powerful tools for converting certain linear partial differential equations into either algebraic equations or else differential equations involving fewer variables.

4.3.1. Fourier transform.

In this section all functions are complex-valued, and $\bar{\cdot}$ denotes the complex conjugate.

a. Definitions and properties.

DEFINITION. If $u \in L^1(\mathbb{R}^n)$, we define its *Fourier transform* $\mathcal{F}u = \hat{u}$ by

$$(1) \quad \hat{u}(y) := \frac{1}{(2\pi)^{n/2}} \int_{\mathbb{R}^n} e^{-ix \cdot y} u(x) dx \quad (y \in \mathbb{R}^n)$$

and its *inverse Fourier transform* $\mathcal{F}^{-1}u = \check{u}$ by

$$(2) \quad \check{u}(y) := \frac{1}{(2\pi)^{n/2}} \int_{\mathbb{R}^n} e^{ix \cdot y} u(x) dx \quad (y \in \mathbb{R}^n).$$

Since $|e^{\pm ix \cdot y}| = 1$ and $u \in L^1(\mathbb{R}^n)$, these integrals converge for each $y \in \mathbb{R}^n$.

We intend now to extend definitions (1), (2) to functions $u \in L^2(\mathbb{R}^n)$.

THEOREM 1 (Plancherel's Theorem). *Assume $u \in L^1(\mathbb{R}^n) \cap L^2(\mathbb{R}^n)$. Then $\hat{u}, \check{u} \in L^2(\mathbb{R}^n)$ and*

$$(3) \quad \|\hat{u}\|_{L^2(\mathbb{R}^n)} = \|\check{u}\|_{L^2(\mathbb{R}^n)} = \|u\|_{L^2(\mathbb{R}^n)}.$$

Proof. 1. First we note that if $v, w \in L^1(\mathbb{R}^n)$, then $\hat{v}, \hat{w} \in L^\infty(\mathbb{R}^n)$. Also

$$(4) \quad \int_{\mathbb{R}^n} v(x)\hat{w}(x) dx = \int_{\mathbb{R}^n} \hat{v}(y)w(y) dy,$$

since both expressions equal $\frac{1}{(2\pi)^{n/2}} \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} e^{-ix \cdot y} v(x)w(y) dx dy$. Furthermore, as we will explicitly compute below in Example 1,

$$\int_{\mathbb{R}^n} e^{ix \cdot y - t|x|^2} dx = \left(\frac{\pi}{t}\right)^{n/2} e^{-\frac{|y|^2}{4t}} \quad (t > 0).$$

Consequently if $\varepsilon > 0$ and $v_\varepsilon(x) := e^{-\varepsilon|x|^2}$, we have $\hat{v}_\varepsilon(y) = \frac{e^{-\frac{|y|^2}{4\varepsilon}}}{(2\varepsilon)^{n/2}}$. Thus (4) implies for each $\varepsilon > 0$ that

$$(5) \quad \int_{\mathbb{R}^n} \hat{w}(y)e^{-\varepsilon|y|^2} dy = \frac{1}{(2\varepsilon)^{n/2}} \int_{\mathbb{R}^n} w(x)e^{-\frac{|x|^2}{4\varepsilon}} dx.$$

2. Now take $u \in L^1(\mathbb{R}^n) \cap L^2(\mathbb{R}^n)$ and set $v(x) := \bar{u}(-x)$. Let $w := u * v \in L^1(\mathbb{R}^n) \cap C(\mathbb{R}^n)$ and check (cf. Theorem 2 below) that

$$\hat{w} = (2\pi)^{n/2} \hat{u} \hat{v} \in L^\infty(\mathbb{R}^n).$$

But

$$\hat{v}(y) = \frac{1}{(2\pi)^{n/2}} \int_{\mathbb{R}^n} e^{-ix \cdot y} \bar{u}(-x) dx = \bar{\hat{u}}(y),$$

and so $\hat{w} = (2\pi)^{n/2} |\hat{u}|^2$.

Now w is continuous and thus

$$\lim_{\varepsilon \rightarrow 0} \frac{1}{(2\varepsilon)^{n/2}} \int_{\mathbb{R}^n} w(x)e^{-\frac{|x|^2}{4\varepsilon}} dx = (2\pi)^{n/2} w(0),$$

where we employed the lemma from §2.3.1. Since $\hat{w} = (2\pi)^{n/2} |\hat{u}|^2 \geq 0$, we deduce upon sending $\varepsilon \rightarrow 0^+$ in (5) that \hat{w} is summable, with

$$\int_{\mathbb{R}^n} \hat{w}(y) dy = (2\pi)^{n/2} w(0).$$

Hence

$$\int_{\mathbb{R}^n} |\hat{u}|^2 dy = w(0) = \int_{\mathbb{R}^n} u(x)v(-x) dx = \int_{\mathbb{R}^n} |u|^2 dx.$$

The proof for \tilde{u} is similar. □

Definition of Fourier transform on L^2 . In view of the equality (3) we can define the Fourier transforms of a function $u \in L^2(\mathbb{R}^n)$ as follows. Choose a sequence $\{u_k\}_{k=1}^\infty \subset L^1(\mathbb{R}^n) \cap L^2(\mathbb{R}^n)$ with

$$u_k \rightarrow u \quad \text{in } L^2(\mathbb{R}^n).$$

According to (3), $\|\hat{u}_k - \hat{u}_j\|_{L^2(\mathbb{R}^n)} = \|\widehat{u_k - u_j}\|_{L^2(\mathbb{R}^n)} = \|u_k - u_j\|_{L^2(\mathbb{R}^n)}$, and thus $\{\hat{u}_k\}_{k=1}^\infty$ is a Cauchy sequence in $L^2(\mathbb{R}^n)$. This sequence consequently converges to a limit, which we define to be $\mathcal{F}u = \hat{u}$:

$$\hat{u}_k \rightarrow \hat{u} \quad \text{in } L^2(\mathbb{R}^n).$$

The definition of \hat{u} does not depend upon the choice of approximating sequence $\{\hat{u}_k\}_{k=1}^\infty$. We similarly define \check{u} .

Next we record some useful formulas.

THEOREM 2 (Properties of Fourier transform). *Assume $u, v \in L^2(\mathbb{R}^n)$. Then*

- (i) $\int_{\mathbb{R}^n} u \bar{v} dx = \int_{\mathbb{R}^n} \hat{u} \bar{\hat{v}} dy$.
- (ii) $(D^\alpha u)^\wedge = (iy)^\alpha \hat{u}$ for each multiindex α such that $D^\alpha u \in L^2(\mathbb{R}^n)$.
- (iii) If $u, v \in L^1(\mathbb{R}^n) \cap L^2(\mathbb{R}^n)$, then $(u * v)^\wedge = (2\pi)^{n/2} \hat{u} \hat{v}$.
- (iv) Furthermore, $u = (\hat{u})^\check{}$.

Assertion (iv) is the *Fourier inversion formula*, which represents a function u in terms of the exponential plane waves $e^{ix \cdot y}$, provided $\hat{u} \in L^1(\mathbb{R}^n)$:

$$(6) \quad u(x) = \frac{1}{(2\pi)^{n/2}} \int_{\mathbb{R}^n} e^{ix \cdot y} \hat{u}(y) dy.$$

Proof. 1. Let $u, v \in L^2(\mathbb{R}^n)$ and $\alpha \in \mathbb{C}$. Then

$$\|u + \alpha v\|_{L^2(\mathbb{R}^n)}^2 = \|\hat{u} + \widehat{\alpha v}\|_{L^2(\mathbb{R}^n)}^2.$$

Expanding, we deduce

$$\int_{\mathbb{R}^n} |u|^2 + |\alpha v|^2 + \bar{u}(\alpha v) + u(\bar{\alpha} \bar{v}) dx = \int_{\mathbb{R}^n} |\hat{u}|^2 + |\widehat{\alpha v}|^2 + \hat{u}(\widehat{\alpha v}) + \hat{u}(\bar{\alpha} \bar{\hat{v}}) dy;$$

and so according to Theorem 1,

$$\int_{\mathbb{R}^n} \alpha \bar{u} v + \bar{\alpha} u \bar{v} dx = \int_{\mathbb{R}^n} \alpha \bar{\hat{u}} \hat{v} + \bar{\alpha} \hat{u} \bar{\hat{v}} dy.$$

Take $\alpha = 1, i$ and combine the resulting equalities to deduce

$$\int_{\mathbb{R}^n} u \bar{v} \, dx = \int_{\mathbb{R}^n} \hat{u} \bar{\hat{v}} \, dy.$$

This proves (i).

2. If u is smooth and has compact support, we calculate

$$\begin{aligned} (D^\alpha u)^\wedge(y) &= \frac{1}{(2\pi)^{n/2}} \int_{\mathbb{R}^n} e^{-ix \cdot y} D^\alpha u(x) \, dx \\ &= \frac{(-1)^{|\alpha|}}{(2\pi)^{n/2}} \int_{\mathbb{R}^n} D_x^\alpha (e^{-ix \cdot y}) u(x) \, dx \\ &= \frac{1}{(2\pi)^{n/2}} \int_{\mathbb{R}^n} e^{-ix \cdot y} (iy)^\alpha u(x) \, dx = (iy)^\alpha \hat{u}(y). \end{aligned}$$

By approximation the same formula is true if $D^\alpha u \in L^2(\mathbb{R}^n)$.

3. We compute for $u, v \in L^1(\mathbb{R}^n) \cap L^2(\mathbb{R}^n)$ and $y \in \mathbb{R}^n$ that

$$\begin{aligned} (u * v)^\wedge(y) &= \frac{1}{(2\pi)^{n/2}} \int_{\mathbb{R}^n} e^{-ix \cdot y} \int_{\mathbb{R}^n} u(z) v(x - z) \, dz \, dx \\ &= \frac{1}{(2\pi)^{n/2}} \int_{\mathbb{R}^n} e^{-iz \cdot y} u(z) \left(\int_{\mathbb{R}^n} e^{-i(x-z) \cdot y} v(x - z) \, dx \right) \, dz \\ &= \int_{\mathbb{R}^n} e^{-iz \cdot y} u(z) \, dz \hat{v}(y) = (2\pi)^{n/2} \hat{u}(y) \hat{v}(y). \end{aligned}$$

4. Next we observe that if $u, v \in L^2(\mathbb{R}^n)$, then

$$\int_{\mathbb{R}^n} \check{u} v \, dx = \int_{\mathbb{R}^n} u \check{v} \, dx,$$

since for $u, v \in L^1(\mathbb{R}^n) \cap L^2(\mathbb{R}^n)$ both sides equal

$$\frac{1}{(2\pi)^{n/2}} \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} e^{ix \cdot y} u(y) v(x) \, dx \, dy.$$

We also note that

$$\check{v} = \overline{(\hat{v})^\wedge}.$$

We may therefore employ assertion (i) to compute

$$\int_{\mathbb{R}^n} (\hat{u})^\wedge v \, dx = \int_{\mathbb{R}^n} \hat{u} \check{v} \, dx = \int_{\mathbb{R}^n} \hat{u} \overline{(\hat{v})^\wedge} \, dx = \int_{\mathbb{R}^n} u \bar{v} \, dx = \int_{\mathbb{R}^n} uv \, dx.$$

This holds for all $v \in L^2(\mathbb{R}^n)$, and so statement (iv) follows. \square

b. Applications. The Fourier transform \mathcal{F} is an especially powerful technique for studying linear, constant-coefficient partial differential equations.

Example 1 (Bessel potentials). We investigate first the PDE

$$-\Delta u + u = f \quad \text{in } \mathbb{R}^n,$$

where $f \in L^2(\mathbb{R}^n)$. To find an explicit formula for u , we take the Fourier transform, recalling Theorem 2(ii) to obtain

$$(7) \quad (1 + |y|^2)\hat{u}(y) = \hat{f}(y) \quad (y \in \mathbb{R}^n).$$

The effect of the Fourier transform has been to convert the PDE into the algebraic equation (7), the solution of which is trivial:

$$\hat{u} = \frac{\hat{f}}{1 + |y|^2}.$$

Thus

$$(8) \quad u = \mathcal{F}^{-1} \left(\frac{\hat{f}}{1 + |y|^2} \right),$$

and so the only real problem is to rewrite the right-hand side of (8) into a more explicit form.

Invoking Theorem 2(iii), we see

$$(9) \quad u = \frac{f * B}{(2\pi)^{n/2}},$$

where

$$(10) \quad \hat{B} = \frac{1}{1 + |y|^2}.$$

Although \hat{B} is not necessarily in L^1 or L^2 , we solve formally for B as follows. Since $\frac{1}{a} = \int_0^\infty e^{-ta} dt$ for each $a > 0$, we have $\frac{1}{1+|y|^2} = \int_0^\infty e^{-t(1+|y|^2)} dt$. Thus

$$(11) \quad B = \left(\frac{1}{1 + |y|^2} \right)^\vee = \frac{1}{(2\pi)^{n/2}} \int_0^\infty e^{-t} \left(\int_{\mathbb{R}^n} e^{ix \cdot y - t|y|^2} dy \right) dt.$$

Now if $a, b \in \mathbb{R}$, $b > 0$, and we set $z = b^{1/2}x - \frac{a}{2b^{1/2}}i$, we find

$$\int_{-\infty}^\infty e^{iax - bx^2} dx = \frac{e^{-a^2/4b}}{b^{1/2}} \int_\Gamma e^{-z^2} dz,$$

Γ denoting the contour $\left\{ \operatorname{Im}(z) = -\frac{a}{2b^{1/2}} \right\}$ in the complex plane. Deforming Γ into the real axis, we compute $\int_{\Gamma} e^{-z^2} dz = \int_{-\infty}^{\infty} e^{-x^2} dx = \pi^{1/2}$; and hence

$$(12) \quad \int_{-\infty}^{\infty} e^{iax-bx^2} dx = e^{-a^2/4b} \left(\frac{\pi}{b} \right)^{1/2}.$$

Thus

$$(13) \quad \int_{\mathbb{R}^n} e^{ix \cdot y - t|y|^2} dy = \prod_{j=1}^n \int_{-\infty}^{\infty} e^{ix_j y_j - t y_j^2} dy_j = \left(\frac{\pi}{t} \right)^{n/2} e^{-\frac{|x|^2}{4t}}$$

by (12). Consequently, we conclude from (11), (13) that

$$(14) \quad B(x) = \frac{1}{2^{n/2}} \int_0^{\infty} \frac{e^{-t - \frac{|x|^2}{4t}}}{t^{n/2}} dt \quad (x \in \mathbb{R}^n).$$

B is called a *Bessel potential*. Employing (9), we derive then the formula

$$(15) \quad u(x) = \frac{1}{(4\pi)^{n/2}} \int_0^{\infty} \int_{\mathbb{R}^n} \frac{e^{-t - \frac{|x-y|^2}{4t}}}{t^{n/2}} f(y) dy dt \quad (x \in \mathbb{R}^n). \quad \square$$

Example 2 (Fundamental solution of heat equation). Consider again the initial-value problem for the heat equation

$$(16) \quad \begin{cases} u_t - \Delta u = 0 & \text{in } \mathbb{R}^n \times (0, \infty) \\ u = g & \text{on } \mathbb{R}^n \times \{t = 0\}. \end{cases}$$

We establish a new method for solving (16) by computing \hat{u} , the Fourier transform of u in *the spatial variables x only*. Thus

$$\begin{cases} \hat{u}_t + |y|^2 \hat{u} = 0 & \text{for } t > 0 \\ \hat{u} = \hat{g} & \text{for } t = 0, \end{cases}$$

whence

$$\hat{u} = e^{-t|y|^2} \hat{g}.$$

Consequently $u = \left(e^{-t|y|^2} \hat{g} \right)^{\vee}$, and therefore

$$(17) \quad u = \frac{g * \hat{F}}{(2\pi)^{n/2}},$$

where $\hat{F} = e^{-t|y|^2}$. But then

$$F = \mathcal{F}^{-1} \left(e^{-t|y|^2} \right) = \frac{1}{(2\pi)^{n/2}} \int_{\mathbb{R}^n} e^{ix \cdot y - t|y|^2} dy = \frac{1}{(2t)^{n/2}} e^{-\frac{|x|^2}{4t}}$$

by (13). Invoking (17), we compute

$$(18) \quad u(x, t) = \frac{1}{(4\pi t)^{n/2}} \int_{\mathbb{R}^n} e^{-\frac{|x-y|^2}{4t}} g(y) dy \quad (x \in \mathbb{R}^n, t > 0),$$

in agreement with §2.3.1. The Fourier transform has provided us with a new derivation of the fundamental solution of the heat equation. \square

Example 3 (Fundamental solution of Schrödinger's equation). Let us next look at the initial-value problem for Schrödinger's equation

$$(19) \quad \begin{cases} iu_t + \Delta u = 0 & \text{in } \mathbb{R}^n \times (0, \infty) \\ u = g & \text{on } \mathbb{R}^n \times \{t = 0\}. \end{cases}$$

Here u and g are complex-valued.

If we formally replace t by it on the right-hand side of (18), we obtain the formula

$$(20) \quad u(x, t) = \frac{1}{(4\pi it)^{n/2}} \int_{\mathbb{R}^n} e^{\frac{i|x-y|^2}{4t}} g(y) dy \quad (x \in \mathbb{R}^n, t > 0),$$

where we interpret $i^{\frac{1}{2}}$ as $e^{\frac{i\pi}{4}}$. This expression clearly makes sense for all times $t > 0$, provided $g \in L^1(\mathbb{R}^n)$. Furthermore if $|y|^2 g \in L^1(\mathbb{R}^n)$, we can check by a direct calculation that u solves $iu_t + \Delta u = 0$ in $\mathbb{R}^n \times (0, \infty)$. (We will not discuss here the sense in which $u(\cdot, t) \rightarrow g$ as $t \rightarrow 0^+$, but see §4.5.3 below and Problem 16.)

Let us next rewrite formula (20) as

$$u(x, t) = \frac{e^{\frac{i|x|^2}{4t}}}{(4\pi it)^{n/2}} \int_{\mathbb{R}^n} e^{\frac{-ix \cdot y}{2t}} e^{\frac{i|y|^2}{4t}} g(y) dy.$$

Since $|e^{\frac{i|x|^2}{4t}}, e^{\frac{i|y|^2}{4t}}| = 1$, we can check as in Theorem 1 that if $g \in L^1(\mathbb{R}^n) \cap L^2(\mathbb{R}^n)$, then

$$(21) \quad \|u(\cdot, t)\|_{L^2(\mathbb{R}^n)} = \|g\|_{L^2(\mathbb{R}^n)} \quad (t > 0).$$

Hence the mapping $g \mapsto u(\cdot, t)$ preserves the L^2 -norm. Therefore we can extend formula (20) to functions $g \in L^2(\mathbb{R}^n)$, in the same way that we extended the definition of Fourier transform. \square

We call

$$(22) \quad \Psi(x, t) := \frac{1}{(4\pi it)^{n/2}} e^{\frac{i|x|^2}{4t}} \quad (x \in \mathbb{R}^n, t \neq 0)$$

the *fundamental solution* of Schrödinger's equation. Note that formula (20), $u = g * \Psi$, makes sense for all times $t \neq 0$, even $t < 0$. Thus we in fact have solved

$$(23) \quad \begin{cases} iu_t + \Delta u = 0 & \text{in } \mathbb{R}^n \times (-\infty, \infty) \\ u = g & \text{on } \mathbb{R}^n \times \{t = 0\}. \end{cases}$$

In particular, Schrödinger's equation is *reversible in time*, whereas the heat equation is not (in spite of Theorem 11 in §2.3.4).

Example 4 (Wave equation). We next analyze the initial-value problem for the wave equation

$$(24) \quad \begin{cases} u_{tt} - \Delta u = 0 & \text{in } \mathbb{R}^n \times (0, \infty) \\ u = g, \quad u_t = h & \text{on } \mathbb{R}^n \times \{t = 0\}, \end{cases}$$

where for simplicity we suppose the initial velocity to be zero. Take as before \hat{u} to be the Fourier transform of u in the variable $x \in \mathbb{R}^n$. Then

$$(25) \quad \begin{cases} \hat{u}_{tt} + |y|^2 \hat{u} = 0 & \text{for } t > 0 \\ \hat{u} = \hat{g}, \quad \hat{u}_t = \hat{h} & \text{for } t = 0. \end{cases}$$

This is an ODE for each fixed $y \in \mathbb{R}^n$, the solution of which is

$$(26) \quad \hat{u} = \hat{g} \cos(t|y|) + \frac{\hat{h}}{|y|} \sin(t|y|).$$

Inverting, we find

$$u(x, t) = \left[\hat{g} \cos(t|y|) + \frac{\hat{h}}{|y|} \sin(t|y|) \right]^\vee.$$

In the particular case that $h \equiv 0$, we have

$$(27) \quad u(x, t) = \frac{1}{(2\pi)^{n/2}} \int_{\mathbb{R}^n} \frac{\hat{g}(y)}{2} (e^{i(x \cdot y + t|y|)} + e^{i(x \cdot y - t|y|)}) dy$$

for $x \in \mathbb{R}^n$, $t \geq 0$, a formula we will further analyze in certain asymptotic limits later, in §4.5.3.

Asymptotic equipartition of energy. Assume that $f, g, Dg \in L^2(\mathbb{R}^n)$. Recall from §2.4.3 that the energy of the solution u of the wave equation (24) is

$$E(t) := \frac{1}{2} \int_{\mathbb{R}^n} u_t^2 + |Du|^2 dx \quad (t \geq 0)$$

and that it is constant in time:

$$E(t) = E(0) = \frac{1}{2} \int_{\mathbb{R}^n} h^2 + |Dg|^2 dx.$$

As an application of the representation formula (26) we next show that

$$(28) \quad \lim_{t \rightarrow \infty} \int_{\mathbb{R}^n} |Du|^2 dx = \lim_{t \rightarrow \infty} \int_{\mathbb{R}^n} u_t^2 dx = E(0).$$

This says that asymptotically the total energy splits equally into its potential and kinetic parts (cf. Problem 24 in Chapter 2).

To establish (28), we compute using (26) that

$$\begin{aligned}
 \int_{\mathbb{R}^n} |Du|^2 dx &= \int_{\mathbb{R}^n} |y|^2 |\hat{u}|^2 dy \\
 (29) \qquad &= \int_{\mathbb{R}^n} |y|^2 |\hat{g}|^2 \cos^2(t|y|) + |\hat{h}|^2 \sin^2(t|y|) dy \\
 &\quad + \int_{\mathbb{R}^n} \cos(t|y|) \sin(t|y|) |y| (\hat{h}\bar{\hat{g}} + \hat{g}\bar{\hat{h}}) dy.
 \end{aligned}$$

Now if $f \in C_c^\infty(\mathbb{R}^n)$, then

$$\begin{aligned}
 \int_{\mathbb{R}^n} \cos(t|y|) \sin(t|y|) f dy &= \frac{1}{2} \int_{\mathbb{R}^n} \sin(2t|y|) f dy \\
 &= \frac{1}{2} \int_0^\infty \sin(2tr) \int_{\partial B(0,r)} f dS dr \\
 &= -\frac{1}{4t} \int_0^\infty \frac{d}{dr} (\cos(2tr)) \int_{\partial B(0,r)} f dS dr \\
 &= \frac{1}{4t} \int_0^\infty \cos(2tr) \frac{d}{dr} \left(\int_{\partial B(0,r)} f dS \right) dr \\
 &= O(t^{-1}).
 \end{aligned}$$

Approximating the integrable function $|y|(\hat{h}\bar{\hat{g}} + \hat{g}\bar{\hat{h}})$ by a smooth function with compact support, we see that the last integral in (29) goes to zero as $t \rightarrow \infty$. Using the identity $\cos^2(t|y|) = \frac{1}{2}(\cos(2t|y|) + 1)$, we likewise deduce that

$$\int_{\mathbb{R}^n} |y|^2 |\hat{g}|^2 \cos^2(t|y|) dy \rightarrow \frac{1}{2} \int_{\mathbb{R}^n} |y|^2 |\hat{g}|^2 dy,$$

and similarly

$$\int_{\mathbb{R}^n} |\hat{h}|^2 \sin^2(t|y|) dy \rightarrow \frac{1}{2} \int_{\mathbb{R}^n} |\hat{h}|^2 dy.$$

It follows then from (29) that

$$\lim_{t \rightarrow \infty} \int_{\mathbb{R}^n} |Du|^2 dx = \frac{1}{2} \int_{\mathbb{R}^n} |y|^2 |\hat{g}|^2 + |\hat{h}|^2 dy = \frac{1}{2} \int_{\mathbb{R}^n} |Dg|^2 + |h|^2 dx = E(0).$$

□

Example 5 (Telegraph equation). The initial-value problem for the one-dimensional telegraph equation is

$$\begin{cases} u_{tt} + 2du_t - u_{xx} = 0 & \text{in } \mathbb{R} \times (0, \infty) \\ u = g, \quad u_t = h & \text{on } \mathbb{R} \times \{t = 0\}, \end{cases}$$

for $d > 0$, the term “ $2d\hat{u}_t$ ” representing a physical damping of wave propagation. As before

$$\begin{cases} \hat{u}_{tt} + 2d\hat{u}_t + |y|^2\hat{u} = 0 & \text{for } t > 0 \\ \hat{u} = \hat{g}, \hat{u}_t = \hat{h} & \text{for } t = 0. \end{cases}$$

We again seek a solution of the form $\hat{u} = \beta e^{t\gamma}$ ($\beta, \gamma \in \mathbb{C}$). Plugging in above, we deduce that $\gamma^2 + 2d\gamma + |y|^2 = 0$; whence $\gamma = -d \pm (d^2 - |y|^2)^{1/2}$. Consequently

$$\hat{u}(y, t) = \begin{cases} e^{-dt}(\beta_1(y)e^{\gamma(y)t} + \beta_2(y)e^{-\gamma(y)t}) & \text{if } |y| \leq d \\ e^{-dt}(\beta_1(y)e^{i\delta(y)t} + \beta_2(y)e^{-i\delta(y)t}) & \text{if } |y| \geq d \end{cases}$$

for $\gamma(y) := (d^2 - |y|^2)^{1/2}$ ($|y| \leq d$), $\delta(y) := (|y|^2 - d^2)^{1/2}$ ($|y| \geq d$), where $\beta_1(y)$ and $\beta_2(y)$ are selected so that

$$\hat{g}(y) = \beta_1(y) + \beta_2(y)$$

and

$$\hat{h}(y) = \begin{cases} \beta_1(y)(\gamma(y) - d) + \beta_2(y)(-\gamma(y) - d) & \text{if } |y| \leq d \\ \beta_1(y)(i\delta(y) - d) + \beta_2(y)(-i\delta(y) - d) & \text{if } |y| \geq d. \end{cases}$$

We thereby obtain the representation formula:

$$\begin{aligned} u(x, t) &= \frac{e^{-dt}}{(2\pi)^{1/2}} \int_{\{|y| \leq d\}} \beta_1(y)e^{ixy + \gamma(y)t} + \beta_2(y)e^{ixy - \gamma(y)t} dy \\ &\quad + \frac{e^{-dt}}{(2\pi)^{1/2}} \int_{\{|y| \geq d\}} \beta_1(y)e^{i(xy + \delta(y)t)} + \beta_2(y)e^{i(xy - \delta(y)t)} dy. \end{aligned}$$

Notice the terms e^{-dt} , which causes damping as $t \rightarrow \infty$. □

4.3.2. Radon transform.

The Fourier inversion formula (6) is significant for PDE theory primarily since it represents a function in terms of the exponential plane waves $e^{ix \cdot y}$. We introduce in this subsection the Radon transform \mathcal{R} , which provides for odd dimensions the elegant, alternative decomposition (33), (34) into plane waves. This is sometimes useful because it is often easier to determine information concerning the support of a function from its Radon transform, rather than its Fourier transform.

a. Definitions and properties

NOTATION. We write S^{n-1} for the unit sphere $\partial B(0, 1)$ in \mathbb{R}^n , a typical point of which we denote $\omega = (\omega_1, \dots, \omega_n)$. The plane with unit normal $\omega \in S^{n-1}$ at a distance $s \in \mathbb{R}$ from the origin is

$$\Pi(s, \omega) := \{y \in \mathbb{R}^n \mid y \cdot \omega = s\}.$$

Note that we allow $s < 0$.

DEFINITION. The Radon transform $\mathcal{R}u = \tilde{u}$ of a function $u \in C_c^\infty(\mathbb{R}^n)$ is

$$(30) \quad \tilde{u}(x, \omega) := \int_{\Pi(s, \omega)} u \, dS \quad (s \in \mathbb{R}, \omega \in S^{n-1}).$$

The term on the right is the integral over the plane $\Pi(s, \omega)$ with respect to $(n-1)$ -dimensional surface measure.

THEOREM 3 (Properties of Radon transform). Assume $u \in C_c^\infty(\mathbb{R}^n)$. Then

- (i) $\tilde{u}(-s, -\omega) = \tilde{u}(s, \omega)$.
- (ii) $(D^\alpha u)^\sim = \omega^\alpha \frac{\partial^{|\alpha|}}{\partial s^{|\alpha|}} \tilde{u}$ for each multiindex α .
- (iii) $(\Delta u)^\sim = \frac{\partial^2}{\partial s^2} \tilde{u}$.
- (iv) If $u \equiv 0$ in $\mathbb{R}^n - B(0, R)$, then $\tilde{u}(s, \omega) = 0$ for $|s| \geq R$.

Proof. Assertion (i) is clear, since $\Pi(s, \omega) = \Pi(-s, -\omega)$. To prove (ii), let $\{b_1, \dots, b_{n-1}\}$ be an orthonormal basis of the subspace $\Pi(0, \omega)$. Then $\{b_1, \dots, b_{n-1}, \omega\}$ is an orthonormal basis of \mathbb{R}^n and so

$$Du = \sum_{j=1}^{n-1} (Du \cdot b_j) b_j + (Du \cdot \omega) \omega.$$

Consequently

$$\begin{aligned} \tilde{u}_{x_i} &= \int_{\Pi(s, \omega)} u_{x_i} \, dS \\ &= \sum_{j=1}^{n-1} (b_j \cdot e_i) \int_{\Pi(s, \omega)} Du \cdot b_j \, dS + \omega_i \int_{\Pi(s, \omega)} Du \cdot \omega \, dS \\ &= \omega_i \int_{\Pi(s, \omega)} Du \cdot \omega \, dS. \end{aligned}$$

The integrals of $Du \cdot b_j$ over $\Pi(s, \omega)$ vanish because b_j is tangent to $\Pi(s, \omega)$ and u has compact support. Since

$$\tilde{u}_s = \int_{\Pi(s, \omega)} Du \cdot \omega \, dS,$$

we have proved (ii) for $\alpha = e_i$. The general case follows by induction, and (iii) is immediate since $|\omega| = 1$.

Assertion (iv) is obvious, since $\Pi(s, \omega) \cap B(0, R) = \emptyset$ if $|s| > R$. \square

Next we discover an interesting connection between the Radon and Fourier transforms.

THEOREM 4 (Radon and Fourier transforms). *Assume that $u \in C_c^\infty(\mathbb{R}^n)$. Then*

$$(31) \quad \bar{u}(r, \omega) := \int_{\mathbb{R}} \tilde{u}(s, \omega) e^{-irs} \, ds = (2\pi)^{n/2} \hat{u}(r\omega) \quad (r \in \mathbb{R}, \omega \in S^{n-1}),$$

where $\hat{u} = \mathcal{F}u$ is the Fourier transform.

Proof. As in the previous proof, take $\{b_1, \dots, b_{n-1}\}$ to be an orthonormal basis of $\Pi(0, \omega)$. Then

$$\tilde{u}(s, \omega) = \int_{\mathbb{R}^{n-1}} u \left(\sum_{j=1}^{n-1} y_j b_j + s\omega \right) \, dy,$$

and so

$$\int_{\mathbb{R}} \tilde{u}(s, \omega) e^{-irs} \, ds = \int_{\mathbb{R}} \int_{\mathbb{R}^{n-1}} u \left(\sum_{j=1}^{n-1} y_j b_j + s\omega \right) e^{-irs} \, dy \, ds.$$

We change variables, now writing $x := \sum_{j=1}^{n-1} y_j b_j + s\omega$. Then

$$\int_{\mathbb{R}} \tilde{u}(s, \omega) e^{-irs} \, ds = \int_{\mathbb{R}^n} u(x) e^{-ir(x \cdot \omega)} \, dx = (2\pi)^{n/2} \hat{u}(r\omega). \quad \square$$

Since we know how to invert the Fourier transform, we can likewise invert the Radon transform. The surprise is that we discover a nice formula for odd dimensions.

THEOREM 5 (Inverting the Radon transform).

(i) *We have*

$$(32) \quad u(x) = \frac{1}{2(2\pi)^n} \int_{\mathbb{R}} \int_{S^{n-1}} \bar{u}(r, \omega) r^{n-1} e^{ir\omega \cdot x} dS dr,$$

the function \bar{u} defined by (31).

(ii) *If $n = 2k + 1$ is odd, then*

$$(33) \quad u(x) = \int_{S^{n-1}} r(x \cdot \omega, \omega) dS$$

for

$$(34) \quad r(s, \omega) := \frac{(-1)^k}{2(2\pi)^{2k}} \frac{\partial^{2k}}{\partial s^{2k}} \tilde{u}(s, \omega).$$

Formulas (33) and (34) provide an elegant and useful decomposition of u into plane waves.

Proof. 1. According to (31) we have

$$\begin{aligned} \int_{\mathbb{R}} \int_{S^{n-1}} \bar{u} r^{n-1} e^{ir\omega \cdot x} dS dr &= (2\pi)^{n/2} \int_{\mathbb{R}} \int_{S^{n-1}} \hat{u}(r\omega) r^{n-1} e^{ir\omega \cdot x} dS dr \\ &= 2(2\pi)^{n/2} \int_0^\infty \int_{S^{n-1}} \hat{u}(r\omega) r^{n-1} e^{ir\omega \cdot x} dS dr \\ &= 2(2\pi)^{n/2} \int_{\mathbb{R}^n} \hat{u}(y) e^{iy \cdot x} dy \\ &= 2(2\pi)^n u(x). \end{aligned}$$

For the second equality above, we replaced r by $-r$ and ω by $-\omega$ in computing the integral $\int_{-\infty}^0 \int_{S^{n-1}} \hat{u}(r\omega) r^{n-1} e^{ir\omega \cdot x} dS dr$. The last equality is the Fourier inversion formula (6).

2. The identity (31) also implies that $\bar{u} = (2\pi)^{1/2}(\tilde{u})^\wedge$, the circumflex now denoting the one-dimensional Fourier transform in the variable r (with ω held fixed). Consequently

$$\left(\frac{\partial^{2k}}{\partial s^{2k}} \tilde{u} \right)^\wedge = (ir)^{2k} (\tilde{u})^\wedge = \frac{(-1)^k r^{2k}}{(2\pi)^{1/2}} \bar{u},$$

and hence

$$\frac{\partial^{2k}}{\partial s^{2k}} \tilde{u}(s, \omega) = \frac{(-1)^k}{2\pi} \int_{\mathbb{R}} \bar{u}(r, \omega) r^{2k} e^{irs} dr.$$

Put $s = x \cdot \omega$:

$$\frac{\partial^{2k}}{\partial s^{2k}} \tilde{u}(\omega \cdot x, \omega) = \frac{(-1)^k}{2\pi} \int_{\mathbb{R}} \bar{u} r^{n-1} e^{ir(\omega \cdot x)} dr.$$

Now integrate over the unit sphere, to discover

$$\begin{aligned} \int_{S^{n-1}} \frac{\partial^{2k}}{\partial s^{2k}} \tilde{u}(\omega \cdot x, \omega) dS &= \frac{(-1)^k}{2\pi} \int_{\mathbb{R}} \int_{S^{n-1}} \bar{u} r^{n-1} e^{ir(\omega \cdot x)} dr ds \\ &= \frac{(-1)^k}{2\pi} 2(2\pi)^n u(x) \end{aligned}$$

in light of (32). □

Application (Vanishing of Radon transform). As a quick corollary, we note that

$$(35) \quad \begin{cases} \text{if } n \text{ is odd and } \tilde{u} = 0 \text{ for } |s| \leq R, \\ \text{then } u \equiv 0 \text{ in } B(0, R). \end{cases}$$

Compare this statement with assertion (iv) from Theorem 3. To prove it, observe that for $n = 2k + 1$, $r(s, \omega) = \frac{(-1)^k}{2(2\pi)^{2k}} \frac{\partial^{2k}}{\partial s^{2k}} \tilde{u}(s, \omega) = 0$ if $|s| \leq R$; and consequently formula (33) implies (35). □

b. Applications

Example 6 (Another representation formula for the wave equation). If we fix a unit vector $\omega \in S^{n-1}$, then the plane wave $\gamma(x \cdot \omega - t, \omega)$ solves the wave equation and therefore so does the superposition of plane waves

$$(36) \quad u(x, t) := \int_{S^{n-1}} \gamma(x \cdot \omega - t, \omega) dS.$$

We claim now that if $n = 2k + 1$ is odd and

$$(37) \quad \gamma := \frac{(-1)^k}{2(2\pi)^{2k}} \frac{\partial^{2k}}{\partial s^{2k}} \tilde{g} = \frac{(-1)^k}{2(2\pi)^{2k}} \mathcal{R}(\Delta^k g),$$

then (36) provides a formula for the solution of the initial-value problem for the wave equation:

$$(38) \quad \begin{cases} u_{tt} - \Delta u = 0 & \text{in } \mathbb{R}^n \times (0, \infty) \\ u = g, \quad u_t = 0 & \text{on } \mathbb{R}^n \times \{t = 0\}. \end{cases}$$

To confirm this, note first that $u = g$ on $\mathbb{R}^n \times \{t = 0\}$ in view of Theorem 5(ii). Consequently we need only check the second initial condition, that

$$u_t(x, 0) = - \int_{S^{n-1}} \gamma_s(x \cdot \omega, \omega) dS = 0.$$

In view of (37), it will suffice to show that $\int_{S^{n-1}} \lambda_s(x \cdot \omega, \omega) dS = 0$ when $\lambda := \mathcal{R}l$ for some smooth function l . But

$$\lambda_s = \int_{\Pi(s, \omega)} Dl \cdot \omega dS;$$

and consequently

$$\int_{S^{n-1}} \lambda_s(x \cdot \omega, \omega) dS = \int_{S^{n-1}} \int_{\Pi(x \cdot \omega, \omega)} Dl(y) \cdot \omega dS(y) dS(\omega) = 0,$$

since the integrand is an odd function of ω . □

Example 7 (Huygens' principle for hyperbolic systems). A linear system of first-order PDE

$$\mathbf{u}_t + \sum_{j=1}^n B_j \mathbf{u}_{x_j} = 0$$

for the unknown $\mathbf{u} : \mathbb{R}^n \times [0, \infty) \rightarrow \mathbb{R}^m$, $\mathbf{u} = (u^1, \dots, u^m)$, is called *hyperbolic* if for each $y \in \mathbb{R}^n$ the $m \times m$ matrix

$$\mathbf{B}(y) := \sum_{j=1}^n y_j B_j$$

has m real eigenvalues

$$\lambda_1(y) \leq \lambda_2(y) \leq \dots \leq \lambda_m(y).$$

We suppose further that $\mathbf{B}(y)$ is *diagonalizable*, meaning there exists a smooth, invertible matrix $\mathbf{A}(y)$ such that

$$\mathbf{A}^{-1}(y)\mathbf{B}(y)\mathbf{A}(y) = \mathbf{D}(y) = \text{diag}(\lambda_1(y), \dots, \lambda_m(y)).$$

In §7.3 we will employ the Fourier transform to construct solutions of the initial-value problem

$$(39) \quad \begin{cases} \mathbf{u}_t + \sum_{j=1}^n B_j \mathbf{u}_{x_j} = 0 & \text{in } \mathbb{R}^n \times (0, \infty) \\ \mathbf{u} = \mathbf{g} & \text{on } \mathbb{R}^n \times \{t = 0\} \end{cases}$$

for given data $\mathbf{g} = (g^1, \dots, g^m)$. Here we provide instead a quick application of the Radon transform, to conclude information about the support of the solution when the initial data have compact support.

So let us suppose that \mathbf{u} is a smooth solution of (39) and take the Radon transform in the variables x . We deduce from Theorem 3(ii) that for each fixed $\omega \in S^{n-1}$

$$\begin{cases} \tilde{\mathbf{u}}_t + \mathbf{B}(\omega)\tilde{\mathbf{u}}_s = 0 & \text{in } \mathbb{R} \times (0, \infty) \\ \tilde{\mathbf{u}} = \tilde{\mathbf{g}} & \text{on } \mathbb{R} \times \{t = 0\}. \end{cases}$$

Now put $\mathbf{v} := A^{-1}(\omega)\tilde{\mathbf{u}}$, $\mathbf{h} := A^{-1}(\omega)\tilde{\mathbf{g}}$. Then \mathbf{v} solves the diagonal system

$$\begin{cases} \mathbf{v}_t + \mathbf{D}(\omega)\mathbf{v}_s = 0 & \text{in } \mathbb{R} \times (0, \infty) \\ \mathbf{v} = \mathbf{h} & \text{on } \mathbb{R} \times \{t = 0\}. \end{cases}$$

That is,

$$\begin{cases} v_t^j + \lambda_j(\omega)v_s^j = 0 & \text{in } \mathbb{R} \times (0, \infty) \\ \mathbf{v}^j = \mathbf{h}^j & \text{on } \mathbb{R} \times \{t = 0\}; \end{cases}$$

and so

$$v^j(s, \omega, t) = h^j(s - \lambda_j(\omega)t, \omega)$$

for $j = 1, \dots, m$. It follows that

$$\tilde{\mathbf{u}}(s, \omega, t) = \mathbf{A}(\omega)\mathbf{v}(s, \omega, t) = \sum_{j=1}^m h^j(s - \lambda_j(\omega)t, \omega)\mathbf{a}_j(\omega),$$

$\mathbf{a}_j(\omega)$ denoting the j^{th} column of $\mathbf{A}(\omega)$. The inversion formulas (33), (34) then provide for odd $n = 2k + 1$ the expression

$$(40) \quad \mathbf{u}(x, t) = \int_{S^{n-1}} \sum_{j=1}^m l^j(x \cdot \omega - \lambda_j(\omega)t, \omega)\mathbf{a}_j(\omega) dS$$

for

$$(41) \quad l^j := \frac{(-1)^k}{2(2\pi)^{2k}} \frac{\partial^{2k}}{\partial s^{2k}} h^j(s - \lambda_j(\omega)t, \omega) \quad (j = 1, \dots, m).$$

In particular, if

$$\sigma := \min_{\omega \in S^{n-1}} \min_{1 \leq j \leq m} |\lambda_j(\omega)| > 0$$

and $\text{spt } \mathbf{g} \subset B(0, R)$, then

$$(42) \quad u(x, t) = 0 \quad \text{for } |x| \leq \sigma t - R.$$

This is a form of Huygens' principle for the hyperbolic system (39) in odd dimensions n . □

4.3.3. Laplace transform.

Remember that we write $R_+ = (0, \infty)$.

DEFINITION. If $u \in L^1(\mathbb{R}_+)$, we define its Laplace transform $\mathcal{L}u = u^\#$ to be

$$(43) \quad u^\#(s) := \int_0^\infty e^{-st}u(t) dt \quad (s \geq 0).$$

Whereas the Fourier and Radon transforms are most appropriate for functions defined on all of \mathbb{R} (or \mathbb{R}^n), the Laplace transform is useful for functions defined only on \mathbb{R}_+ . In practice this means that for a partial differential equation involving time, it may be useful to perform a Laplace transform in t , holding the space variables x fixed. (This is the reverse of the technique from Examples 2–5 of §4.3.1.)

Example 8 (Resolvents and Laplace transform). Consider again the heat equation

$$(44) \quad \begin{cases} v_t - \Delta v = 0 & \text{in } U \times (0, \infty) \\ v = f & \text{on } U \times \{t = 0\}, \end{cases}$$

and perform a Laplace transform with respect to time:

$$v^\#(x, s) = \int_0^\infty e^{-st}v(x, t) dt \quad (s > 0).$$

What PDE does $v^\#$ satisfy? We compute

$$\begin{aligned} \Delta v^\#(x, s) &= \int_0^\infty e^{-st} \Delta v(x, t) dt = \int_0^\infty e^{-st} v_t(x, t) dt \\ &= s \int_0^\infty e^{-st} v(x, t) dt + e^{-st} v|_{t=0}^{t=\infty} = s v^\#(x, s) - f(x). \end{aligned}$$

Think now of $s > 0$ being fixed, and write $u(x) := v^\#(x, s)$. Then

$$(45) \quad -\Delta u + su = f \quad \text{in } U.$$

Thus the solution of the resolvent equation (45) with right-hand side f is the Laplace transform of the solution of the heat equation (44) with initial data f . (If $U = \mathbb{R}^n$ and $s = 1$, we could now represent v in terms of the fundamental solution, to rederive formula (15).) \square

The connection between the resolvent equation and the Laplace transform will be made clearer by the discussion in §7.4 of semigroup theory.

Example 9 (Wave equation from the heat equation). Next we employ some Laplace transform ideas to provide a new derivation of the solution for the wave equation (cf. §2.4.1), based—surprisingly—upon the heat equation.

Suppose u is a bounded, smooth solution of the initial-value problem:

$$(46) \quad \begin{cases} u_{tt} - \Delta u = 0 & \text{in } \mathbb{R}^n \times (0, \infty) \\ u = g, \quad u_t = 0 & \text{on } \mathbb{R}^n \times \{t = 0\}, \end{cases}$$

where n is odd and g is smooth, with compact support. We extend u to negative times by writing

$$(47) \quad u(x, t) = u(x, -t) \quad \text{if } x \in \mathbb{R}^n, \quad t < 0.$$

Then

$$u_{tt} - \Delta u = 0 \quad \text{in } \mathbb{R}^n \times \mathbb{R}.$$

Next define

$$(48) \quad v(x, t) := \frac{1}{(4\pi t)^{1/2}} \int_{-\infty}^{\infty} e^{-s^2/4t} u(x, s) ds \quad (x \in \mathbb{R}^n, \quad t > 0).$$

Hence

$$\lim_{t \rightarrow 0} v = g \quad \text{uniformly on } \mathbb{R}^n.$$

In addition

$$\begin{aligned} \Delta v(x, t) &= \frac{1}{(4\pi t)^{1/2}} \int_{-\infty}^{\infty} e^{-s^2/4t} \Delta u(x, s) ds \\ &= \frac{1}{(4\pi t)^{1/2}} \int_{-\infty}^{\infty} e^{-s^2/4t} u_{ss}(x, s) ds \\ &= \frac{1}{(4\pi t)^{1/2}} \int_{-\infty}^{\infty} \frac{s}{2t} e^{-s^2/4t} u_s(x, s) ds \\ &= \frac{1}{(4\pi t)^{1/2}} \int_{-\infty}^{\infty} \left(\frac{s^2}{4t^2} - \frac{1}{2t} \right) e^{-s^2/4t} u(x, s) ds = v_t(x, t). \end{aligned}$$

Consequently v solves this initial-value problem for the heat equation:

$$\begin{cases} v_t - \Delta v = 0 & \text{in } \mathbb{R}^n \times (0, \infty) \\ v = g & \text{on } \mathbb{R}^n \times \{t = 0\}. \end{cases}$$

As v is bounded, we deduce from §2.3 that

$$(49) \quad v(x, t) = \frac{1}{(4\pi t)^{n/2}} \int_{\mathbb{R}^n} e^{-\frac{|x-y|^2}{4t}} g(y) dy.$$

We equate (48) with (49), recall (47), and set $\lambda = \frac{1}{4t}$, thereby obtaining the identity

$$\int_0^\infty u(x, s) e^{-\lambda s^2} ds = \frac{1}{2} \left(\frac{\lambda}{\pi} \right)^{\frac{n-1}{2}} \int_{\mathbb{R}^n} e^{-\lambda|x-y|^2} g(y) dy.$$

Thus

$$(50) \quad \int_0^\infty u(x, s) e^{-\lambda s^2} ds = \frac{n\alpha(n)}{2} \left(\frac{\lambda}{\pi} \right)^{\frac{n-1}{2}} \int_0^\infty e^{-\lambda r^2} r^{n-1} G(x; r) dr,$$

for all $\lambda > 0$, where

$$(51) \quad G(x; r) = \int_{\partial B(x, r)} g(y) dS(y).$$

We will solve (50), (51) for u . To do so, we write $n = 2k + 1$ and note $-\frac{1}{2r} \frac{d}{dr} (e^{-\lambda r^2}) = \lambda e^{-\lambda r^2}$. Hence

$$\begin{aligned} \lambda^{\frac{n-1}{2}} \int_0^\infty e^{-\lambda r^2} r^{n-1} G(x; r) dr &= \int_0^\infty \lambda^k e^{-\lambda r^2} r^{2k} G(x; r) dr \\ &= \frac{(-1)^k}{2^k} \int_0^\infty \left[\left(\frac{1}{r} \frac{d}{dr} \right)^k (e^{-\lambda r^2}) \right] r^{2k} G(x; r) dr \\ &= \frac{1}{2^k} \int_0^\infty r \left[\left(\frac{1}{r} \frac{\partial}{\partial r} \right)^k (r^{2k-1} G(x; r)) \right] e^{-\lambda r^2} dr, \end{aligned}$$

where we integrated by parts k times for the last equality.

Owing to (50) (with r replacing s in the expression on the left), we deduce

$$\int_0^\infty u(x, r) e^{-\lambda r^2} dr = \frac{n\alpha(n)}{\pi^{\frac{n-1}{2}} 2^{k+1}} \int_0^\infty r \left[\left(\frac{1}{r} \frac{\partial}{\partial r} \right)^k (r^{2k-1} G(x; r)) \right] e^{-\lambda r^2} dr.$$

Upon substituting $\tau = r^2$, we see that each side above, taken as a function of λ , is a Laplace transform. As two Laplace transforms agree only if the original functions were identical, we deduce

$$(52) \quad u(x, t) = \frac{n\alpha(n)}{\pi^k 2^{k+1}} t \left(\frac{1}{t} \frac{\partial}{\partial t} \right)^k (t^{2k-1} G(x, t)).$$

Now $n = 2k + 1$ and $\alpha(n) = \frac{\pi^{n/2}}{\Gamma(\frac{n}{2}+1)} = \frac{\pi^{k+\frac{1}{2}}}{\Gamma(\frac{n}{2}+1)}$. Since $\Gamma(\frac{1}{2}) = \pi^{1/2}$ and $\Gamma(x+1) = x\Gamma(x)$ for $x > 0$ (cf. [Rd, Chapter 8]), we can compute

$$\frac{n\alpha(n)}{\pi^k 2^{k+1}} = \frac{n\pi^{1/2}}{2^{k+1}\Gamma(\frac{n}{2}+1)} = \frac{1}{(n-2)(n-4)\cdots 5\cdot 3} = \frac{1}{\gamma_n}.$$

We insert this deduction into (52) and simplify:

$$(53) \quad u(x, t) = \frac{1}{\gamma_n} \frac{\partial}{\partial t} \left(\frac{1}{t} \frac{\partial}{\partial t} \right)^{\frac{n-3}{2}} \left(t^{n-2} \int_{\partial B(x, t)} g \, dS \right) \quad (x \in \mathbb{R}^n, t > 0).$$

This is formula (31) in §2.4.1 (for $h \equiv 0$). □

4.4. CONVERTING NONLINEAR INTO LINEAR PDE

In this section we describe several techniques which are sometime useful for converting certain nonlinear equations into linear equations.

4.4.1. Cole–Hopf transformation.

a. A parabolic PDE with quadratic nonlinearity. We consider first of all an initial-value problem for a quasilinear parabolic equation:

$$(1) \quad \begin{cases} u_t - a\Delta u + b|Du|^2 = 0 & \text{in } \mathbb{R}^n \times (0, \infty) \\ u = g & \text{on } \mathbb{R}^n \times \{t = 0\}, \end{cases}$$

where $a > 0$. This sort of nonlinear PDE arises in stochastic optimal control theory.

Assuming for the moment u is a smooth solution of (1), we set

$$w := \phi(u),$$

where $\phi : \mathbb{R} \rightarrow \mathbb{R}$ is a smooth function, as yet unspecified. We will try to choose ϕ so that w solves a linear equation. We have

$$w_t = \phi'(u)u_t, \quad \Delta w = \phi'(u)\Delta u + \phi''(u)|Du|^2;$$

and consequently (1) implies

$$\begin{aligned} w_t &= \phi'(u)u_t = \phi'(u)[a\Delta u - b|Du|^2] \\ &= a\Delta w - [a\phi''(u) + b\phi'(u)]|Du|^2 \\ &= a\Delta w, \end{aligned}$$

provided we choose ϕ to satisfy $a\phi'' + b\phi' = 0$. We solve this differential equation by setting $\phi = e^{\frac{-bz}{a}}$. Thus we see that if u solves (1), then

$$(2) \quad w = e^{\frac{-bu}{a}}$$

solves this initial-value problem for the heat equation (with conductivity a):

$$(3) \quad \begin{cases} w_t - a\Delta w = 0 & \text{in } \mathbb{R}^n \times (0, \infty) \\ w = e^{\frac{-bg}{a}} & \text{on } \mathbb{R}^n \times \{t = 0\}. \end{cases}$$

Formula (2) is the *Cole-Hopf transformation*.

Now the unique bounded solution of (3) is

$$w(x, t) = \frac{1}{(4\pi at)^{n/2}} \int_{\mathbb{R}^n} e^{\frac{-|x-y|^2}{4at}} e^{\frac{-b}{a}g(y)} dy \quad (x \in \mathbb{R}^n, t > 0);$$

and, since (2) implies

$$u = -\frac{a}{b} \log w,$$

we obtain thereby the explicit formula

$$(4) \quad u(x, t) = -\frac{a}{b} \log \left(\frac{1}{(4\pi at)^{n/2}} \int_{\mathbb{R}^n} e^{\frac{-|x-y|^2}{4at} - \frac{b}{a}g(y)} dy \right) \quad (x \in \mathbb{R}^n, t > 0)$$

for a solution of quasilinear initial-value problem (1).

b. Burgers' equation with viscosity. As a further application, we examine now for $n = 1$ the initial-value problem for the *viscous Burgers' equation*:

$$(5) \quad \begin{cases} u_t - au_{xx} + uu_x = 0 & \text{in } \mathbb{R} \times (0, \infty) \\ u = g & \text{on } \mathbb{R} \times \{t = 0\}. \end{cases}$$

If we set

$$(6) \quad w(x, t) := \int_{-\infty}^x u(y, t) dy$$

and

$$(7) \quad h(x) := \int_{-\infty}^x g(y) dy$$

(cf. §3.4), we have

$$(8) \quad \begin{cases} w_t - aw_{xx} + \frac{1}{2}w_x^2 = 0 & \text{in } \mathbb{R} \times (0, \infty) \\ w = h & \text{on } \mathbb{R} \times \{t = 0\}. \end{cases}$$

This is an equation of the form (1) for $n = 1$, $b = \frac{1}{2}$; and so (4) provides the formula

$$(9) \quad w(x, t) = -2a \log \left(\frac{1}{(4\pi at)^{1/2}} \int_{\mathbb{R}} e^{\frac{-|x-y|^2}{4at} - \frac{h(y)}{2a}} dy \right).$$

But then since $u = w_x$, we find upon differentiating (9) that

$$(10) \quad u(x, t) = \frac{\int_{-\infty}^{\infty} \frac{x-y}{t} e^{\frac{-|x-y|^2}{4at} - \frac{h(y)}{2a}} dy}{\int_{-\infty}^{\infty} e^{\frac{-|x-y|^2}{4at} - \frac{h(y)}{2a}} dy} \quad (x \in \mathbb{R}, t > 0)$$

is a solution of problem (5), where h is defined by (7). We will scrutinize this formula further in §4.5.2.

4.4.2. Potential functions.

Another technique is to utilize a *potential function* to convert a nonlinear system of PDE into a single linear PDE. We consider as an example *Euler's equations* for inviscid, incompressible fluid flow:

$$(11) \quad \begin{cases} \text{(a)} & \mathbf{u}_t + \mathbf{u} \cdot D\mathbf{u} = -Dp + \mathbf{f} & \text{in } \mathbb{R}^3 \times (0, \infty) \\ \text{(b)} & \operatorname{div} \mathbf{u} = 0 & \text{in } \mathbb{R}^3 \times (0, \infty) \\ \text{(c)} & \mathbf{u} = \mathbf{g} & \text{on } \mathbb{R}^3 \times \{t = 0\}. \end{cases}$$

Here the unknowns are the *velocity field* $\mathbf{u} = (u^1, u^2, u^3)$ and the scalar *pressure* p ; the external force $\mathbf{f} = (f^1, f^2, f^3)$ and initial velocity $\mathbf{g} = (g^1, g^2, g^3)$ are given. Here D as usual denotes the gradient in the spatial variables $x = (x_1, x_2, x_3)$. The vector equation 11(a) means

$$u_t^i + \sum_{j=1}^3 u^j u_{x_j}^i = -p_{x_i} + f^i \quad (i = 1, 2, 3).$$

We will assume

$$(12) \quad \operatorname{div} \mathbf{g} = 0.$$

If furthermore there exists a scalar function $h : \mathbb{R}^3 \times (0, \infty) \rightarrow \mathbb{R}$ such that

$$(13) \quad \mathbf{f} = Dh,$$

we say that the external force is derived from the *potential* h .

We will try to find a solution (\mathbf{u}, p) of (11) for which the velocity field \mathbf{u} is also derived from a potential, say

$$(14) \quad \mathbf{u} = Dv.$$

Our flow will then be irrotational, as $\operatorname{curl} \mathbf{u} \equiv 0$. Now equation (11)(b) says

$$(15) \quad 0 = \operatorname{div} \mathbf{u} = \Delta v$$

and so v must be harmonic as a function of x , for each time $t > 0$. Thus if we can find a smooth function v satisfying (15) and $Dv(\cdot, 0) = \mathbf{g}$, we can then recover \mathbf{u} from v by (14).

How do we compute the pressure p ? Let us observe that if $\mathbf{u} = Dv$, then $\mathbf{u} \cdot D\mathbf{u} = \frac{1}{2}D(|Dv|^2)$. Consequently (11)(a) reads $D(v_t + \frac{1}{2}|Dv|^2) = D(-p + h)$, in view of (13). Therefore we may take

$$(16) \quad v_t + \frac{1}{2}|Dv|^2 + p = h.$$

This is *Bernoulli's law*. But now we can employ (16) to calculate p , up to an additive constant, since v and h are already known.

4.4.3. Hodograph and Legendre transforms.

a. Hodograph transform. The *hodograph transform* is a technique for converting certain quasilinear systems of PDE into linear systems, by reversing the roles of the dependent and independent variables. As this method is most easily understood by an example, we investigate here the equations of steady, two-dimensional, irrotational fluid flow:

$$(17) \quad \begin{cases} \text{(a)} & (\sigma^2(\mathbf{u}) - (u^1)^2)u_{x_1}^1 - u^1 u^2 (u_{x_2}^1 + u_{x_1}^2) \\ & + (\sigma^2(\mathbf{u}) - (u^2)^2)u_{x_2}^2 = 0 \\ \text{(b)} & u_{x_2}^1 - u_{x_1}^2 = 0 \end{cases}$$

in \mathbb{R}^2 . The unknown is the *velocity* field $\mathbf{u} = (u^1, u^2)$, and the function $\sigma(\cdot) : \mathbb{R}^2 \rightarrow \mathbb{R}$, the local *sound speed*, is given.

The system (17) is quasilinear. Let us now, however, no longer regard u^1 and u^2 as functions of x_1 and x_2 :

$$(18) \quad u^1 = u^1(x_1, x_2), \quad u^2 = u^2(x_1, x_2),$$

but rather regard x^1 and x^2 as functions of u_1 and u_2 :

$$(19) \quad x^1 = x^1(u_1, u_2), \quad x^2 = x^2(u_1, u_2).$$

We have exchanged sub- and superscripts in the notation to emphasize the interchange between independent and dependent variables.

According to the Inverse Function Theorem (§C.6) we can, locally at least, invert equations (18) to yield (19), provided

$$(20) \quad J = \frac{\partial(u^1, u^2)}{\partial(x_1, x_2)} = u_{x_1}^1 u_{x_2}^2 - u_{x_2}^1 u_{x_1}^2 \neq 0$$

in some region of \mathbb{R}^2 . Assuming now (20) holds, we calculate

$$(21) \quad \begin{cases} u_{x_2}^2 = J x_{u_1}^1, & u_{x_1}^2 = -J x_{u_1}^2 \\ u_{x_2}^1 = -J x_{u_2}^1, & u_{x_1}^1 = J x_{u_2}^2. \end{cases}$$

We insert (21) into (17), to discover

$$(22) \quad \begin{cases} \text{(a)} & (\sigma^2(u) - u_1^2)x_{u_2}^2 + u_1 u_2 (x_{u_2}^1 + x_{u_1}^2) + (\sigma^2(u) - u_2^2)x_{u_1}^1 = 0 \\ \text{(b)} & x_{u_2}^1 - x_{u_1}^2 = 0. \end{cases}$$

This is a *linear* system for $\mathbf{x} = (x^1, x^2)$, as a function of $u = (u_1, u_2)$.

Remark. We can utilize the method of potential functions (§4.4.2) to simplify (22) further. Indeed, equation (22)(b) suggests that we look for a single function $z = z(u)$ such that

$$\begin{cases} x^1 = z_{u_1} \\ x^2 = z_{u_2}. \end{cases}$$

Then (22)(a) transforms into the linear, second-order PDE

$$(23) \quad (\sigma^2(u) - u_1^2)z_{u_2 u_2} + 2u_1 u_2 z_{u_1 u_2} + (\sigma^2(u) - u_2^2)z_{u_1 u_1} = 0.$$

b. Legendre transform. A technique closely related to the hodograph transform is the classical *Legendre transform*, a version of which we have already encountered before, in §3.3. The idea is to regard the components of the gradient of a solution as new independent variables.

Once again an example is instructive. We investigate the *minimal surface equation* (cf. Example 4 in §8.1.2)

$$\operatorname{div} \left(\frac{Du}{(1 + |Du|^2)^{1/2}} \right) = 0,$$

which for $n = 2$ may be rewritten as

$$(24) \quad (1 + u_{x_2}^2)u_{x_1x_1} - 2u_{x_1}u_{x_2}u_{x_1x_2} + (1 + u_{x_1}^2)u_{x_2x_2} = 0.$$

Let us now assume that at least in some region of \mathbb{R}^2 , we can invert the relations

$$(25) \quad p^1 = u_{x_1}(x_1, x_2), \quad p^2 = u_{x_2}(x_1, x_2),$$

to solve for

$$(26) \quad x^1 = x^1(p_1, p_2), \quad x^2 = x^2(p_1, p_2).$$

The Inverse Function Theorem assures us we can do so in a neighborhood of any point where

$$(27) \quad J = \det D^2u \neq 0.$$

Now define

$$(28) \quad v(p) := \mathbf{x}(p) \cdot p - u(\mathbf{x}(p)),$$

where $\mathbf{x} = (x^1, x^2)$ is given by (26), $p = (p_1, p_2)$. We discover after some calculations that

$$(29) \quad \begin{cases} u_{x_1x_1} = Jv_{p_2p_2} \\ u_{x_1x_2} = -Jv_{p_1p_2} \\ u_{x_2x_2} = Jv_{p_1p_1}. \end{cases}$$

Upon substituting the identities (29) into (24), we derive for v the *linear* equation

$$(30) \quad (1 + p_2^2)v_{p_2p_2} + 2p_1p_2v_{p_1p_2} + (1 + p_1^2)v_{p_1p_1} = 0.$$

Remark. The hodograph and Legendre transform techniques for obtaining linear out of nonlinear PDE are in practice tricky to use, as it is usually not possible to transform given boundary conditions very easily.

4.5. ASYMPTOTICS

It is often the case that even when explicit representation formulas can be had for solutions of partial differential equations, these are too complicated to be of much immediate use. In such circumstances it sometimes becomes profitable to study the formulas in various asymptotic limits, whereupon simplifications often appear.

Following are several rather complicated examples, illustrating typical issues involved in asymptotics for PDE. The results in this section are explained only heuristically, mostly without formal proofs.

4.5.1. Singular perturbations.

A *singular perturbation* is a modification of a given PDE by adding a small multiple ε times a higher-order term. In accordance with the informal principle that the behavior of solutions is governed primarily by the highest order terms, a solution u^ε of the perturbed problem will often behave analytically quite differently from a solution u of the original equation.

Example 1 (Transport and small diffusion). We illustrate this idea by studying formally the effects of small diffusion upon the transport of dye within a moving fluid in \mathbb{R}^2 .

Suppose we are given a smooth vector field $\mathbf{b} : \mathbb{R}^2 \rightarrow \mathbb{R}^2$, $\mathbf{b} = (b^1, b^2)$, representing the steady fluid velocity. Assume dye has been continuously injected at unit rate into the fluid at the origin, and let $u(x)$ represent the density of dye at the point $x \in \mathbb{R}^2$, $x = (x_1, x_2)$. Then, formally at least as we shall see,


$$(1) \quad \operatorname{div}(u\mathbf{b}) = \delta_0 \quad \text{in } \mathbb{R}^2,$$

where δ_0 is the Dirac measure on \mathbb{R}^2 giving unit mass to the point 0. This PDE implies that the dye density is transported with the fluid motion at points $x \neq 0$.

Consider now for $\varepsilon > 0$ the singular perturbation:

$$(2) \quad -\varepsilon\Delta u^\varepsilon + \operatorname{div}(u^\varepsilon\mathbf{b}) = \delta_0 \quad \text{in } \mathbb{R}^2.$$

The new term “ $\varepsilon\Delta$ ” represents a small, isotropic diffusion of the dye within the background fluid motion. We are interested in understanding in an approximate way the structure of the solution u^ε of (2) and, in particular, describing if and how u^ε approximates u for small $\varepsilon > 0$.



Flow of dye without diffusion

a. Analysis of problem (1). We turn our attention first to the unperturbed PDE (1). Consider the characteristic ODE

$$(3) \quad \begin{cases} \dot{\mathbf{x}}(t) = \mathbf{b}(\mathbf{x}(t)) & (t \geq 0) \\ \mathbf{x}(0) = 0, \end{cases}$$

the solution $\mathbf{x}(t) = (x^1(t), x^2(t))$ of which we assume to trace out a curve C , as drawn.

Given a point $x \in \mathbb{R}^2$ near C , we write

$$(4) \quad x = \mathbf{x}(t) + y\nu(\mathbf{x}(t)),$$

where $\nu = (\nu^1, \nu^2)$ is the (upward pointing) unit normal to C , $y \in \mathbb{R}$, and t is the time required for the solution of the ODE (3) to reach the point $\mathbf{x}(t)$ along C closest to x . We hereafter regard (t, y) as providing a new coordinate system near the curve C , so that $x = (x^1(y, t), x^2(y, t))$.

Using (3) and (4), we compute

$$\frac{\partial(x^1, x^2)}{\partial(t, y)} = \det \begin{pmatrix} \frac{\partial x^1}{\partial t} & \frac{\partial x^1}{\partial y} \\ \frac{\partial x^2}{\partial t} & \frac{\partial x^2}{\partial y} \end{pmatrix} = \det \begin{pmatrix} b^1 + y\nu^1 & \nu^1 \\ b^2 + y\nu^2 & \nu^2 \end{pmatrix}.$$

Let us write $\sigma = |\mathbf{b}|$, $\nu = (-b^2, b^1)/\sigma$ and $\dot{\nu} = -\sigma\kappa\tau = -\kappa\mathbf{b}$ (where $\sigma =$ speed, $\kappa =$ curvature, $\tau = \frac{\mathbf{b}}{\sigma} =$ unit tangent). We then simplify, to obtain

$$(5) \quad \frac{\partial(x^1, x^2)}{\partial(t, y)} = \sigma(1 - \kappa y).$$

Return now to the PDE (1), which we rewrite to read

$$(6) \quad \mathbf{b} \cdot Du + (\operatorname{div} \mathbf{b})u = \delta_0 \quad \text{in } \mathbb{R}^2.$$

As in §3.2 we see $u \equiv 0$ off the curve C . Let us next guess u has the form

$$(7) \quad u(x) = \rho(t)\delta(y)$$

in the (t, y) -coordinates, δ denoting the Dirac measure on \mathbb{R} giving unit mass to the origin.

What is $\rho(t)$? To compute it, take R to be a small, smooth region in the (x_1, x_2) -plane, with boundary intersecting the curve C at the points $\mathbf{x}(t_1)$ and $\mathbf{x}(t_2)$, $0 < t_1 < t_2$. Let R' denote the corresponding region in the (t, y) -plane. Then using (5), we calculate

$$\int_R u \, dx = \int_{R'} \rho(t) \delta(y) \sigma(t) (1 - \kappa y) \, dy dt = \int_{t_1}^{t_2} \rho(t) \sigma(t) \, dt.$$

Now $\int_R u \, dx$ represents the total amount of dye within the region R , which is to say, the total amount released between times t_1 and t_2 . This is simply $t_2 - t_1$. Thus

$$\int_{t_1}^{t_2} \rho(t) \sigma(t) \, dt = t_2 - t_1.$$

This identity holds for all $0 < t_1 < t_2$, and so $\rho(t) = \sigma(t)^{-1}$. Hence (7) says

$$(8) \quad u(x, t) = \delta(y) / \sigma(t)$$


is a solution of (1), for $\sigma(t) := |\mathbf{b}(\mathbf{x}(t))|$, $t \geq 0$. In other words, u represents the density along the curve C of the dye, whose concentration varies inversely with the speed of the fluid.

We can further confirm this formula as follows. Let $v \in C_c^\infty(\mathbb{R}^2)$. Then (5) lets us compute that

$$\begin{aligned} \int_{\mathbb{R}^2} Dv \cdot \mathbf{b} u \, dx &= \int_{\mathbb{R}^2} Dv \cdot \mathbf{b} \frac{\delta(y)}{\sigma(t)} \sigma(t) (1 - \kappa y) \, dy dt \\ &= \int_0^\infty Dv(\mathbf{x}(t)) \cdot \mathbf{b}(\mathbf{x}(t)) \, dt \\ &= \int_0^\infty \frac{d}{dt} v(\mathbf{x}(t)) \, dt = -v(0). \end{aligned}$$

Hence we may indeed interpret u defined by (8) as a weak solution of the unperturbed PDE (1).

b. Analysis of problem (2) for $0 < \varepsilon \ll 1$. We look now at the perturbed problem (2). We expect that at time $t > 0$, the diffusing dye will fill a ball of radius approximately $O((\varepsilon t)^{1/2})$ about the point $\mathbf{x}(t)$. The dye will thus be mostly concentrated in a plume as drawn, about the curve C .



Flow of dye with diffusion

We wish to understand the structure of the solution u^ϵ of (2) within this plume, as $\epsilon \rightarrow 0$.

Since the width is presumably of order $O(\epsilon^{1/2})$ for times $0 < t_1 \leq t \leq t_2$ and the total mass of dye corresponding to the same time interval is $t_2 - t_1$, we expect u^ϵ to be of order $O(\epsilon^{-1/2})$ along C . This suggests that for us to understand the asymptotics as $\epsilon \rightarrow 0$, we should turn our attention to the *rescaled variables*

$$(9) \quad z := \epsilon^{-1/2}y, \quad v^\epsilon := \epsilon^{1/2}u^\epsilon,$$

the powers of ϵ selected so that $z, v^\epsilon = O(1)$.

We must therefore rewrite the PDE (2) in terms of the new variables t, z and v^ϵ . For this, we need first to study the structure of the velocity field \mathbf{b} along the curve C . Let us therefore write

$$(10) \quad \mathbf{b} = \sigma(t)\boldsymbol{\tau} + \{\alpha(t)\boldsymbol{\tau} + \beta(t)\boldsymbol{\nu}\}y + O(y^2),$$

where, as noted before, $\boldsymbol{\tau} = \mathbf{b}/\sigma$ is a unit tangent vector to C . Now for any smooth function w :

$$w_t = w_{x_1} \frac{\partial x^1}{\partial t} + w_{x_2} \frac{\partial x^2}{\partial t} = w_{x_1} \sigma(1 - \kappa y) \boldsymbol{\tau}^1 + w_{x_2} \sigma(1 - \kappa y) \boldsymbol{\tau}^2$$

and

$$w_y = w_{x_1} \frac{\partial x^1}{\partial y} + w_{x_2} \frac{\partial x^2}{\partial y} = w_{x_1} \boldsymbol{\nu}^1 + w_{x_2} \boldsymbol{\nu}^2.$$

Thus

$$(11) \quad \begin{cases} w_{x_1} = \frac{w_t \boldsymbol{\nu}^2 - w_y \sigma(1 - \kappa y) \boldsymbol{\tau}^2}{\sigma(1 - \kappa y)}, \\ w_{x_2} = \frac{-w_t \boldsymbol{\nu}^1 + w_y \sigma(1 - \kappa y) \boldsymbol{\tau}^1}{\sigma(1 - \kappa y)}. \end{cases}$$

Therefore using (10), (11) (for $w = u^\epsilon$), we can calculate

$$\begin{aligned} \mathbf{b} \cdot Du^\epsilon &= [\sigma \boldsymbol{\tau} + (\alpha \boldsymbol{\tau} + \beta \boldsymbol{\nu})y + O(y^2)] \cdot Du^\epsilon \\ &= \frac{u_t^\epsilon}{(1 - \kappa y)} + \frac{\alpha y u_t^\epsilon}{\sigma(1 - \kappa y)} + \beta y u_y^\epsilon + O(y^2 |Du^\epsilon|). \end{aligned}$$

Since $v^\varepsilon = \varepsilon^{1/2}u^\varepsilon$, $z = \varepsilon^{-1/2}y$, we can rewrite the foregoing as

$$(12) \quad \mathbf{b} \cdot Dv^\varepsilon = v_t^\varepsilon + \beta z v_z^\varepsilon + O(\varepsilon^{1/2}).$$

Similarly, we calculate using (10) and (11) (for $w = b^1, b^2$) that

$$\begin{aligned} \operatorname{div} \mathbf{b} &= \frac{1}{\sigma(1 - \kappa y)} [b_t^1 \nu^2 - b_t^2 \nu^1] + [\tau^1 b_y^2 - \tau^2 b_y^1] \\ &= \frac{1}{\sigma(1 - \kappa y)} [(\sigma \tau^1)_t \nu^2 - (\sigma \tau^2)_t \nu^1] \\ &\quad + [\tau^1(\alpha \tau^2 + \beta \nu^2) - \tau^2(\alpha \tau^1 + \beta \nu^1)] + O(y) \\ &= \left(\frac{\dot{\sigma}}{\sigma} + \beta \right) + O(y). \end{aligned}$$

Here we used the identity $\dot{\tau} = \sigma \kappa \nu$. It follows that

$$(13) \quad (\operatorname{div} \mathbf{b})v^\varepsilon = \left(\frac{\dot{\sigma}}{\sigma} + \beta \right) v^\varepsilon + O(\varepsilon^{1/2}).$$

In addition, a similar heuristic argument, the details of which we omit, suggests that

$$(14) \quad \varepsilon \Delta v^\varepsilon = v_{zz}^\varepsilon + O(\varepsilon^{1/2}).$$

Combining now (12)–(14) and recalling (2), we at last deduce v^ε satisfies

$$(15) \quad v_t^\varepsilon - v_{zz}^\varepsilon + (\beta z v^\varepsilon)_z + \frac{\dot{\sigma}}{\sigma} v^\varepsilon = O(\varepsilon^{1/2}).$$

We suppose now that as $\varepsilon \rightarrow 0$, the functions v^ε converge in some sense to a limit:

$$(16) \quad v^\varepsilon \rightarrow v \quad \text{in } \mathbb{R}^2.$$

Then presumably from (15) we will have

$$(17) \quad v_t - v_{zz} + (\beta z v)_z + \frac{\dot{\sigma}}{\sigma} v = 0 \quad \text{in } \mathbb{R} \times (0, \infty).$$

We therefore expect

$$(18) \quad u^\varepsilon = \varepsilon^{-1/2} v^\varepsilon = \varepsilon^{-1/2} (v + o(1)),$$

with v solving (17). The PDE (17) is consequently a *parabolic approximation* (in the variables $t, z = \varepsilon^{-1/2}y$) to our elliptic equation (2). The proper initial condition should be

$$(19) \quad v = \frac{\delta(z)}{\sigma(0)} \quad \text{on } \mathbb{R} \times \{t = 0\}.$$

We will see in Problem 13 that an explicit solution of (17), (19) can be found, in terms of the solution of an ODE involving β .

4.5.2. Laplace's method.

Laplace's method concerns the asymptotics as $\varepsilon \rightarrow 0$ of integrals involving expressions of the form $e^{-I/\varepsilon}$, I denoting some given function.

Example 2 (Vanishing viscosity method for Burgers' equation). We next investigate the limit as $\varepsilon \rightarrow 0$ of the solution u^ε of the initial-value problem for the viscous Burgers' equation

$$(20) \quad \begin{cases} u_t^\varepsilon + u^\varepsilon u_x^\varepsilon - \varepsilon u_{xx}^\varepsilon = 0 & \text{in } \mathbb{R} \times (0, \infty) \\ u^\varepsilon = g & \text{on } \mathbb{R} \times \{t = 0\}. \end{cases}$$

Remembering formula (10) from §4.4.1, we note

$$(21) \quad u^\varepsilon(x, t) = \frac{\int_{-\infty}^{\infty} \frac{x-y}{t} e^{-\frac{K(x,y,t)}{2\varepsilon}} dy}{\int_{-\infty}^{\infty} e^{-\frac{K(x,y,t)}{2\varepsilon}} dy},$$

for

$$(22) \quad K(x, y, t) := \frac{|x-y|^2}{2t} + h(y) \quad (x, y \in \mathbb{R}, t > 0),$$

where h is an antiderivative of g .

What happens to u^ε as $\varepsilon \rightarrow 0$? Mathematically the term " εu_{xx} " in (20) makes the partial differential equation act somewhat like the heat equation, in that the solution u^ε is infinitely differentiable in $\mathbb{R} \times (0, \infty)$, in spite of the nonlinearity. This follows from the explicit formula (21). On the other hand, an obvious guess is that the solutions u^ε should converge as $\varepsilon \rightarrow 0$ to a solution u of the conservation law

$$(23) \quad \begin{cases} u_t + \left(\frac{u^2}{2}\right)_x = 0 & \text{in } \mathbb{R} \times (0, \infty) \\ u = g & \text{on } \mathbb{R} \times \{t = 0\}. \end{cases}$$

Physically, we regard the term " εu_{xx} " as imposing an "artificial viscosity" effect, which we are now sending to zero. We expect that this *vanishing viscosity* technique should allow us to recover the correct entropy solution u of (23), which may have discontinuities across shock waves, as the limit of the solutions u^ε of (20), which are smooth.

We must understand the limiting behavior of the expression on the right-hand side of (21), as $\varepsilon \rightarrow 0$.

LEMMA (Asymptotics). *Suppose that $k, l : \mathbb{R} \rightarrow \mathbb{R}$ are continuous functions, that l grows at most linearly and that k grows at least quadratically. Assume also there exists a unique point $y_0 \in \mathbb{R}$ such that*

$$k(y_0) = \min_{y \in \mathbb{R}} k(y).$$

Then

$$(24) \quad \lim_{\varepsilon \rightarrow 0} \frac{\int_{-\infty}^{\infty} l(y) e^{-\frac{k(y)}{\varepsilon}} dy}{\int_{-\infty}^{\infty} e^{-\frac{k(y)}{\varepsilon}} dy} = l(y_0).$$

Proof. Write $k_0 = k(y_0)$. Then the function

$$\mu_\varepsilon(y) := \frac{e^{\frac{k_0 - k(y)}{\varepsilon}}}{\int_{-\infty}^{\infty} e^{\frac{k_0 - k(z)}{\varepsilon}} dz} \quad (y \in \mathbb{R})$$

satisfies

$$(25) \quad \begin{cases} \mu_\varepsilon \geq 0, \int_{-\infty}^{\infty} \mu_\varepsilon(y) dy = 1, \\ \mu_\varepsilon(y) \rightarrow 0 \text{ exponentially fast for } y \neq y_0, \text{ as } \varepsilon \rightarrow 0. \end{cases}$$

Consequently

$$\lim_{\varepsilon \rightarrow 0} \frac{\int_{-\infty}^{\infty} l(y) e^{-\frac{k(y)}{\varepsilon}} dy}{\int_{-\infty}^{\infty} e^{-\frac{k(y)}{\varepsilon}} dy} = \lim_{\varepsilon \rightarrow 0} \int_{-\infty}^{\infty} l(y) \mu_\varepsilon(y) dy = l(y_0).$$

□

Return now to (21), (22). We observe $K(x, y, t) = tL\left(\frac{x-y}{t}\right) + h(y)$, where $L = F^*$ for $F(z) = \frac{z^2}{2}$. According to the analysis in §3.4, for each time $t > 0$ the mapping $y \mapsto K(x, y, t)$ attains its minimum at a unique point $y = y(x, t)$ for all but at most countably many points x . But then the lemma implies

$$(26) \quad \lim_{\varepsilon \rightarrow 0} u^\varepsilon(x, t) = \frac{x - y(x, t)}{t} = G\left(\frac{x - y(x, t)}{t}\right) = u(x, t)$$

for $G := (F')^{-1}$.

The final equality in (26) is the Lax–Oleinik formula for the unique entropy solution of the initial-value problem (23). It is a powerful endorsement of the methods from §3.4 that this formula has reappeared in the context of vanishing viscosity. (See also Problem 7.) □

We will later discuss the vanishing viscosity method for symmetric hyperbolic systems in §7.3.2, for Hamilton–Jacobi equations in §10.1, and for systems of conservation laws in §11.4.

4.5.3. Geometric optics, stationary phase.

This section investigates the behavior of certain highly oscillatory solutions of the wave equation. We begin with some crude, but instructive, calculations.

a. Geometric optics.

Example 3 (Oscillating solutions). Let us once more turn our attention to the wave equation

$$(27) \quad u_{tt} - \Delta u = 0 \quad \text{in } \mathbb{R}^n \times (0, \infty),$$

and we now regard the solution u as taking complex values. We fix $\varepsilon > 0$ and seek a solution $u = u^\varepsilon$ of (27) having the form

$$(28) \quad u^\varepsilon(x, t) = e^{\frac{ip^\varepsilon(x, t)}{\varepsilon}} a^\varepsilon(x, t) \quad (x \in \mathbb{R}^n, t \geq 0),$$

the real-valued function p^ε representing the *phase* and the real-valued function a^ε representing the *amplitude*. The proposed form (28) for the solution is called the *geometric optics ansatz**. The idea is that highly oscillatory solutions of the wave equation can be understood by studying a PDE for the phase function in the limit as $\varepsilon \rightarrow 0$. Following is a formal demonstration.

Substituting (28) into (27), we find after some computations that

$$0 = u_{tt}^\varepsilon - \Delta u^\varepsilon = e^{ip^\varepsilon/\varepsilon} \left(\frac{ip_{tt}^\varepsilon}{\varepsilon} a^\varepsilon - \left(\frac{p_t^\varepsilon}{\varepsilon} \right)^2 a^\varepsilon + 2 \frac{ip_t^\varepsilon a_t^\varepsilon}{\varepsilon} + a_{tt}^\varepsilon \right) - e^{ip^\varepsilon/\varepsilon} \left(\frac{i\Delta p^\varepsilon}{\varepsilon} a^\varepsilon - \frac{|Dp^\varepsilon|^2}{\varepsilon^2} a^\varepsilon + \frac{2iDp^\varepsilon \cdot Da^\varepsilon}{\varepsilon} + \Delta a^\varepsilon \right).$$

We cancel the term $e^{ip^\varepsilon/\varepsilon}$ and take the real part of the resulting expression, to find

$$(29) \quad a^\varepsilon((p_t^\varepsilon)^2 - |Dp^\varepsilon|^2) = \varepsilon^2(a_{tt}^\varepsilon - \Delta a^\varepsilon).$$

Now if as $\varepsilon \rightarrow 0$

$$(30) \quad p^\varepsilon \rightarrow p, \quad a^\varepsilon \rightarrow a \neq 0$$

in some sense, then presumably from (29) it follows that

$$(31) \quad p_t \pm |Dp| = 0 \quad \text{in } \mathbb{R}^n \times (0, \infty).$$

* *ansatz* = formulation (German).

We may informally regard the straight line characteristics of these Hamilton–Jacobi PDE as rays along which the solution (28) concentrates in the high-frequency limit as $\varepsilon \rightarrow 0$.

More generally, let us consider the second-order hyperbolic PDE

$$(32) \quad u_{tt} - \sum_{k,l=1}^n a^{kl}(x)u_{x_k x_l} = 0 \quad \text{in } \mathbb{R}^n \times (0, \infty)$$

with $a^{kl} = a^{lk}$ ($k, l = 1, \dots, n$). We again look for a complex-valued solution $u = u^\varepsilon$ of the form (28) and calculate

$$\begin{aligned} 0 &= u_{tt}^\varepsilon - \sum_{k,l=1}^n a^{kl} u_{x_k x_l}^\varepsilon \\ &= e^{ip^\varepsilon/\varepsilon} \left(\frac{ip_{tt}^\varepsilon}{\varepsilon} a^\varepsilon - \left(\frac{p_t^\varepsilon}{\varepsilon} \right)^2 a^\varepsilon + \frac{2ip_t^\varepsilon a_t^\varepsilon}{\varepsilon} + a_{tt}^\varepsilon \right) \\ &\quad - e^{ip^\varepsilon/\varepsilon} \left(\sum_{k,l=1}^n a^{kl} \left(\frac{ip_{x_k x_l}^\varepsilon}{\varepsilon} a^\varepsilon - \frac{p_{x_k}^\varepsilon p_{x_l}^\varepsilon}{\varepsilon^2} a^\varepsilon + \frac{2ip_{x_k}^\varepsilon a_{x_l}^\varepsilon}{\varepsilon} + a_{x_k x_l}^\varepsilon \right) \right). \end{aligned}$$

We once again cancel $e^{ip^\varepsilon/\varepsilon}$ and take real parts to find

$$a^\varepsilon \left((p_t^\varepsilon)^2 - \sum_{k,l=1}^n a^{kl} p_{x_k}^\varepsilon p_{x_l}^\varepsilon \right) = \varepsilon^2 \left(a_{tt}^\varepsilon - \sum_{k,l=1}^n a^{kl} a_{x_k x_l}^\varepsilon \right).$$

Hence if (30) holds in some sense, we may then expect

$$(33) \quad p_t \pm \left(\sum_{k,l=1}^n a^{kl} p_{x_k} p_{x_l} \right)^{1/2} = 0 \quad \text{in } \mathbb{R}^n \times (0, \infty).$$

See below and also §4.6.1, §7.2.4 for further elaboration of these ideas. \square

b. Stationary phase. The foregoing example suggests that the Hamilton–Jacobi PDE (31) somehow “controls the high-frequency asymptotics for the wave equation”. However the range of validity of the geometric optics ansatz is highly uncertain in the preceding strictly formal computations. To understand more clearly the behavior of the solution, we employ next the method of *stationary phase*, which is a variant of Laplace’s method obtained by replacing the -1 in the exponent (cf. §4.5.2) with i .

Example 4 (Stationary phase for the wave equation). Look again at the initial-value problem for the wave equation

$$(34) \quad \begin{cases} u_{tt}^\varepsilon - \Delta u^\varepsilon = 0 & \text{in } \mathbb{R}^n \times (0, \infty) \\ u^\varepsilon = g^\varepsilon, \quad u_t^\varepsilon = 0 & \text{on } \mathbb{R}^n \times \{t = 0\}, \end{cases}$$

where we hereafter assume g^ε to have the rapidly oscillating structure

$$(35) \quad g^\varepsilon(x) = a(x)e^{\frac{ip(x)}{\varepsilon}} \quad (x \in \mathbb{R}^n).$$

Here $\varepsilon > 0$, $a, p \in C_c^\infty(\mathbb{R}^n)$, and we suppose

$$(36) \quad Dp \neq 0 \quad \text{on the support of } a.$$

Utilizing formula (27) from §4.3.1, we can write

$$u^\varepsilon(x, t) = \frac{1}{(2\pi)^{n/2}} \int_{\mathbb{R}^n} \frac{\hat{g}^\varepsilon(y)}{2} (e^{i(x \cdot y + t|y|)} + e^{i(x \cdot y - t|y|)}) dy \quad (x \in \mathbb{R}^n, t > 0).$$

Invoking (35), we see

$$(37) \quad u^\varepsilon(x, t) = \frac{1}{2} (I_+^\varepsilon(x, t) + I_-^\varepsilon(x, t)),$$

where

$$I_\pm^\varepsilon(x, t) = \frac{1}{(2\pi)^n} \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} a(z) e^{i((x-z) \cdot y \pm t|y| + \frac{p(z)}{\varepsilon})} dy dz.$$

Changing variables gives

$$(38) \quad I_\pm^\varepsilon(x, t) = \frac{1}{(2\pi\varepsilon)^n} \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} a(z) e^{\frac{i}{\varepsilon} \phi_\pm(x, y, z, t)} dy dz,$$

for

$$(39) \quad \phi_\pm(x, y, z, t) := (x - z) \cdot y \pm t|y| + p(z).$$

We want to study the asymptotics of I_\pm^ε as $\varepsilon \rightarrow 0$. Let us pause in this example and develop some general machinery, which we will later apply to (38), (39). \square

Example 4 motivates our considering general integral expressions of the form

$$(40) \quad I_\varepsilon := \int_{\mathbb{R}^n} e^{i\frac{\phi(y)}{\varepsilon}} a(y) dy \quad (x \in \mathbb{R}^n),$$

where a, ϕ are smooth functions, a has compact support, and $\varepsilon > 0$. We wish to understand the limiting behavior of I_ε as $\varepsilon \rightarrow 0$.

We first examine the special case that ϕ is linear in y :

LEMMA 1 (Asymptotics for linear terms). *Let $a \in C_c^\infty(\mathbb{R}^n)$ and $p \in \mathbb{R}^n$, $p \neq 0$. Then for $m = 1, 2, \dots$*

$$\int_{\mathbb{R}^n} e^{\frac{i}{\varepsilon} p \cdot y} a(y) dy = O(\varepsilon^m) \quad \text{as } \varepsilon \rightarrow 0.$$

Proof. Without loss we may assume $p = (p_1, \dots, p_n)$, $p_1 \neq 0$. Then for $m = 1, 2, \dots$

$$\begin{aligned} \int_{\mathbb{R}^n} e^{\frac{i}{\varepsilon} p \cdot y} a(y) dy &= \left(\frac{\varepsilon}{ip_1} \right)^m \int_{\mathbb{R}^n} \frac{\partial^m}{\partial y_1^m} (e^{\frac{i}{\varepsilon} p \cdot y}) a(y) dy \\ &= \left(\frac{-\varepsilon}{ip_1} \right)^m \int_{\mathbb{R}^n} e^{\frac{i}{\varepsilon} p \cdot y} \frac{\partial^m}{\partial y_1^m} a(y) dy = O(\varepsilon^m). \quad \square \end{aligned}$$

Next we suppose ϕ is quadratic in y :

LEMMA 2 (Asymptotics for quadratic terms). *Let $a \in C_c^\infty(\mathbb{R}^n)$ and suppose A is a real, nonsingular, symmetric matrix. Then*

$$(41) \quad \frac{1}{(2\pi\varepsilon)^{n/2}} \int_{\mathbb{R}^n} e^{\frac{i}{2\varepsilon} y \cdot A y} a(y) dy = \frac{e^{i\frac{\pi}{4} \operatorname{sgn} A}}{|\det A|^{1/2}} (a(0) + O(\varepsilon)) \quad \text{as } \varepsilon \rightarrow 0.$$

Here $\operatorname{sgn} A$, the *signature* of A , denotes the number of positive eigenvalues of A minus the number of negative eigenvalues.

Proof. 1. First we claim for each $\phi \in C_c^\infty(\mathbb{R}^n)$ that

$$(42) \quad \begin{aligned} \lim_{\delta \rightarrow 0^+} \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} e^{ix \cdot Ax - \delta |x|^2 - ix \cdot y} \phi(y) dx dy \\ = \frac{\pi^{n/2}}{|\det A|^{1/2}} e^{i\frac{\pi}{4} \operatorname{sgn} A} \int_{\mathbb{R}^n} e^{-\frac{i}{4} y \cdot A^{-1} y} \phi(y) dy. \end{aligned}$$

To confirm this, we start by assuming A is diagonal:

$$(43) \quad A = \operatorname{diag}(\lambda_1, \dots, \lambda_n) \quad (\lambda_k \neq 0, k = 1, \dots, n).$$

Now for fixed y , $\lambda \in \mathbb{R}$ and $\delta > 0$, we have

$$\begin{aligned} \int_{\mathbb{R}} e^{i\lambda x^2 - \delta x^2 - ix y} dx &= e^{\frac{y^2}{4(i\lambda - \delta)}} \int_{\mathbb{R}} e^{(i\lambda - \delta) \left(x - \frac{iy}{2(i\lambda - \delta)} \right)^2} dx \\ &= \frac{e^{\frac{y^2}{4(i\lambda - \delta)}}}{(\delta - i\lambda)^{1/2}} \int_{\Gamma} e^{-z^2} dz, \end{aligned}$$

where $\Gamma = \{z = (\delta - i\lambda)^{1/2} \left(x - \frac{iy}{2(i\lambda - \delta)}\right) \mid x \in \mathbb{R}\}$ and we take $\operatorname{Re}(\delta - i\lambda)^{1/2} > 0$. Thus Γ is a line in the complex plane, which intersects the x -axis at an angle less than $\frac{\pi}{4}$. We consequently may deform the integral over Γ into the integral along the real axis: see Problem 15. Hence $\int_{\Gamma} e^{-z^2} dz = \int_{\mathbb{R}} e^{-x^2} dx = \pi^{1/2}$, and thus

$$\int_{\mathbb{R}} e^{i\lambda x^2 - \delta x^2 - ixy} dx = \frac{\pi^{1/2}}{(\delta - i\lambda)^{1/2}} e^{\frac{y^2}{4(i\lambda - \delta)}}.$$

Since A has the diagonal form (43), we consequently deduce

$$\begin{aligned} J_{\delta}(y) &:= \int_{\mathbb{R}^n} e^{ix \cdot Ax - \delta|x|^2 - ix \cdot y} dx \\ &= \prod_{k=1}^n \int_{\mathbb{R}} e^{i\lambda_k x_k^2 - \delta x_k^2 - ix_k y_k} dx_k = \pi^{n/2} \prod_{k=1}^n \frac{e^{\frac{y_k^2}{4(i\lambda_k - \delta)}}}{(\delta - i\lambda_k)^{1/2}}. \end{aligned}$$

2. Now let $\phi \in C_c^{\infty}(\mathbb{R}^n)$. Then

$$\int_{\mathbb{R}^n} \phi(y) J_{\delta}(y) dy = \pi^{n/2} \int_{\mathbb{R}^n} \phi(y) \prod_{k=1}^n \frac{e^{\frac{y_k^2}{4(i\lambda_k - \delta)}}}{(\delta - i\lambda_k)^{1/2}} dy.$$

Applying the Dominated Convergence Theorem, we deduce

$$(44) \quad \lim_{\delta \rightarrow 0} \int_{\mathbb{R}^n} \phi(y) J_{\delta}(y) dy = \pi^{n/2} \int_{\mathbb{R}^n} \phi(y) \prod_{k=1}^n \frac{e^{-\frac{iy_k^2}{4\lambda_k}}}{(-i\lambda_k)^{1/2}} dy.$$

Recall that we are supposing $\operatorname{Re}(-i\lambda_k)^{1/2} > 0$. Thus if $\lambda_k > 0$, $(-i\lambda_k)^{1/2} = |\lambda_k|^{1/2} e^{-\frac{i\pi}{4}}$. If instead $\lambda_k < 0$, then $(-i\lambda_k)^{1/2} = |\lambda_k|^{1/2} e^{\frac{i\pi}{4}}$. Therefore

$$\prod_{k=1}^{\infty} (-i\lambda_k)^{1/2} = |\det A|^{1/2} e^{-\frac{i\pi}{4} \operatorname{sgn} A},$$

and so (44) gives (42), provided A is diagonal.

If A is not diagonal, we rotate to new coordinates to diagonalize A and again verify (42).

3. Let us now write $\alpha_{\varepsilon}(y) := e^{\frac{i}{2\varepsilon} y \cdot Ay}$. Then if $a \in C_c^{\infty}(\mathbb{R}^n)$,

$$\int_{\mathbb{R}^n} a(x) \alpha_{\varepsilon}(x) dx = \int_{\mathbb{R}^n} \hat{a}(y) \hat{\alpha}_{\varepsilon}(-y) dy.$$

According to (42) (with $\frac{1}{2\varepsilon}A$ replacing A):

$$\begin{aligned}\hat{\alpha}_\varepsilon(y) &= \frac{\varepsilon^{n/2}}{|\det A|^{1/2}} e^{\frac{i\pi}{4} \operatorname{sgn} A} e^{-\frac{i\varepsilon}{2} y \cdot A^{-1} y} \\ &= \frac{\varepsilon^{n/2}}{|\det A|^{1/2}} e^{\frac{i\pi}{4} \operatorname{sgn} A} (1 + O(\varepsilon|y|^2)),\end{aligned}$$

$\hat{\alpha}_\varepsilon$ interpreted as in (42). Consequently

$$\frac{1}{\varepsilon^{n/2}} \int_{\mathbb{R}^n} a(x) \alpha_\varepsilon(x) dx = \frac{e^{\frac{i\pi}{4} \operatorname{sgn} A}}{|\det A|^{1/2}} \int_{\mathbb{R}^n} \hat{a}(y) (1 + O(\varepsilon|y|^2)) dy.$$

But $\int_{\mathbb{R}^n} \hat{a}(y) dy = (2\pi)^{\frac{n}{2}} a(0)$ and $\int_{\mathbb{R}^n} \hat{a}(y) |y|^2 dy < \infty$. Formula (41) follows. \square

For a general phase function ϕ , we will employ the following result to change variables and thereby convert locally to one of the earlier cases.

LEMMA 3 (Changing coordinates). *Assume $\phi : \mathbb{R}^n \rightarrow \mathbb{R}$ is smooth.*

(i) *Suppose that*

$$D\phi(0) \neq 0.$$

Then there exists a smooth function $\Phi : \mathbb{R}^n \rightarrow \mathbb{R}^n$ such that

$$(45) \quad \begin{cases} \Phi(0) = 0, & D\Phi(0) = I, \text{ and} \\ \phi(\Phi(x)) = \phi(0) + D\phi(0) \cdot x & \text{for } |x| \text{ small.} \end{cases}$$

(ii) (Morse Lemma) *Suppose instead that*

$$D\phi(0) = 0, \det D^2\phi(0) \neq 0.$$

Then there exists a smooth function $\Phi : \mathbb{R}^n \rightarrow \mathbb{R}^n$ such that

$$(46) \quad \begin{cases} \Phi(0) = 0, & D\Phi(0) = I, \text{ and} \\ \phi(\Phi(x)) = \phi(0) + \frac{1}{2} x \cdot D^2\phi(0)x & \text{for } |x| \text{ small.} \end{cases}$$

In other words, we can change variables near 0 to make ϕ affine in case (i), quadratic in case (ii).

Proof. 1. Assume $\mathbf{r}_n := D\phi(0) \neq 0$. Then there exist vectors $\mathbf{r}_1, \dots, \mathbf{r}_{n-1}$ so that $\{\mathbf{r}_k\}_{k=1}^n$ is an orthogonal basis of \mathbb{R}^n . Define $\mathbf{f} : \mathbb{R}^n \times \mathbb{R}^n \rightarrow \mathbb{R}^n$ by

$$\mathbf{f}(x, y) := (\mathbf{r}_1 \cdot (y - x), \dots, \mathbf{r}_{n-1} \cdot (y - x), \phi(y) - \phi(0) - D\phi(0) \cdot x).$$

Therefore

$$D_y \mathbf{f}(0, 0) = \begin{pmatrix} \mathbf{r}_1 \\ \vdots \\ \mathbf{r}_n \end{pmatrix},$$

the $\{\mathbf{r}_k\}_{k=1}^n$ regarded as row vectors, and so $\det D_y \mathbf{f}(0, 0) \neq 0$. The Implicit Function Theorem (§C.7) implies we can find $\Phi : \mathbb{R}^n \rightarrow \mathbb{R}^n$ such that $\Phi(0) = 0$ and

$$\mathbf{f}(x, \Phi(x)) = 0 \quad \text{for } |x| \text{ small.}$$

In particular,

$$(47) \quad \begin{cases} \phi(\Phi(x)) = \phi(0) + D\phi(0) \cdot x \\ \mathbf{r}_k \cdot (\Phi(x) - x) = 0 \quad (k = 1, \dots, n-1). \end{cases}$$

Differentiating with respect to x , we deduce as well

$$(D\Phi(0) - I)\mathbf{r}_k = 0 \quad (k = 1, \dots, n)$$

and so $D\Phi(0) = I$. This proves assertion (i).

2. Fix $x \in \mathbb{R}^n$. Then $\psi(t) := \phi(tx)$ satisfies

$$\psi(1) = \psi(0) + \psi'(0) + \int_0^1 (1-t)\psi''(t) dt.$$

Thus if $D\phi(0) = 0$, we have

$$(48) \quad \phi(x) = \phi(0) + \frac{1}{2}x \cdot \mathbf{A}(x)x$$

for the symmetric matrix

$$\mathbf{A}(x) := 2 \int_0^1 (1-t)D^2\phi(tx) dt.$$

Observe $\mathbf{A}(0) = D^2\phi(0)$. Let us hereafter suppose $D^2\phi(0)$ is nonsingular, and so the same is true for $\mathbf{A}(x)$, provided $|x|$ is small. Furthermore, we may assume upon rotating to new coordinates if necessary that

$$\mathbf{A}(0) = D^2\phi(0) \quad \text{is diagonal.}$$

3. We now claim that there exists for each $m \in \{0, 1, \dots, n\}$ a smooth mapping $\Phi_m : \mathbb{R}^n \rightarrow \mathbb{R}^n$, such that

$$(49) \quad \begin{cases} \Phi_m(0) = 0, \quad D\Phi_m(0) = I, \quad \text{and} \\ \phi(\Phi_m(x)) = \phi(0) + \frac{1}{2} \sum_{i=1}^m \phi_{x_i x_i}(0)x_i^2 + \frac{1}{2} \sum_{i,j=m+1}^n a_m^{ij}(x)x_i x_j, \end{cases}$$

for $|x|$ small, where $\mathbf{A}_m = ((a_m^{ij}))$ is smooth and symmetric.

Observe in particular that (49) implies

$$(50) \quad a_m^{ij}(0) = \phi_{x_i x_j}(0) \quad (i, j = m+1, \dots, n),$$

and thus $a_m^{m+1, m+1}(x) \neq 0$ for $|x|$ sufficiently small.

4. Assertion (49) for $m = 0$ is (48) with $\mathbf{A}_0 = \mathbf{A}$ and Φ_0 the identity mapping. So assume by induction that (49) holds for some $m \in \{0, \dots, n-1\}$ and write

$$\phi_m(x) := \phi(\Phi_m(x)).$$

Then

$$(51) \quad \phi_m(x) = \phi(0) + \frac{1}{2} \sum_{i=1}^m \phi_{x_i x_i}(0) x_i^2 + \frac{1}{2} \sum_{i,j=m+1}^n a_m^{ij}(x) x_i x_j \quad \text{for } |x| \text{ small.}$$

Define a mapping $\Pi_{m+1} : \mathbb{R}^n \rightarrow \mathbb{R}^n$, $\Pi_{m+1}(y) = x$, by writing

$$\Pi_{m+1}(y) := \left(y_1, \dots, y_m, \left(\frac{a_m^{m+1, m+1}(y)}{\phi_{x_{m+1} x_{m+1}}(0)} \right)^{\frac{1}{2}} \left[y_{m+1} + \sum_{j=m+2}^n \frac{y_j a_m^{m+1, j}(y)}{a_m^{m+1, m+1}(y)} \right], \dots, y_n \right)$$

for small $|y|$. It follows then from (51) that

$$\phi_m(y) = \phi(0) + \frac{1}{2} \sum_{i=1}^{m+1} \phi_{x_i x_i}(0) x_i^2 + \frac{1}{2} \sum_{i,j=m+2}^n b_{m+1}^{ij}(y) x_i x_j,$$

where

$$b_{m+1}^{ij}(y) := \begin{cases} a_m^{ij}(y) - \frac{a_m^{m+1, i}(y) a_m^{m+1, j}(y)}{a_m^{m+1, m+1}(y)} & i, j = m+2, \dots, n \\ 0 & \text{otherwise.} \end{cases}$$

Since $D^2\phi(0)$ is diagonal, (50) implies

$$\Pi_{m+1}(0) = 0, \quad D\Pi_{m+1}(0) = I.$$

Consequently we can define for small $|x|$ the inverse mapping $\Xi_{m+1} := \Pi_{m+1}^{-1}$, $y = \Xi_{m+1}(x)$. Therefore

$$\phi_m(\Xi_{m+1}(x)) = \phi(0) + \frac{1}{2} \sum_{i=1}^{m+1} \phi_{x_i x_i}(0) x_i^2 + \frac{1}{2} \sum_{i,j=m+2}^n a_{m+1}^{ij}(x) x_i x_j,$$

for $\mathbf{A}_{m+1} := \mathbf{B}_{m+1} \circ \Xi_{m+1}$. This is statement (49), with $m + 1$ replacing m and with $\Phi_{m+1} := \Phi_m \circ \Xi_{m+1}$.

The case $m = n$ is assertion (ii) of the lemma. □

The stationary phase method. We can at last combine the information gleaned in Lemmas 1–3 to explain informally the stationary phase technique for deriving the asymptotics of

$$I_\varepsilon = \int_{\mathbb{R}^n} e^{\frac{i\phi(y)}{\varepsilon}} a(y) dy$$

as $\varepsilon \rightarrow 0$. We will assume

$$(52) \quad \begin{cases} D\phi \text{ vanishes within the support of } a \\ \text{only at the points } y_1, \dots, y_N \end{cases}$$

and furthermore

$$(53) \quad D^2\phi(y_k) \text{ is nonsingular} \quad (k = 1, \dots, N).$$

Then for $m = 1, \dots,$

$$\left| \int_{\mathbb{R}^n} e^{\frac{i\phi(y)}{\varepsilon}} \zeta(y) a(y) dy \right| = O(\varepsilon^m),$$

where $\zeta \in C_c^\infty$ vanishes near $\{y_1, \dots, y_N\}$. This follows since we can employ Lemma 3(i) to change variables near any point in the support of ζ to make ϕ affine, with nonvanishing gradient, and apply Lemma 1.

On the other hand if ζ is smooth, ζ vanishes except near y_k , and $\zeta(y_k) = 1$, we can employ Lemma 3(ii) to compute

$$\begin{aligned} \int_{\mathbb{R}^n} e^{i\phi(y)} \zeta(y) a(y) dy &= \int_{\mathbb{R}^n} e^{i\phi(\Phi(x))} \zeta a(\Phi(x)) |\det D\Phi(x)| dx \\ &= e^{\frac{i\phi(y_k)}{\varepsilon}} \int_{\mathbb{R}^n} e^{\frac{i}{2\varepsilon}(x-y_k) \cdot D^2\phi(y_k)(x-y_k)} \zeta a(\Phi(x)) \\ &\hspace{20em} |\det D\Phi(x)| dx \\ &= e^{\frac{i\phi(y_k)}{\varepsilon}} \frac{(2\pi\varepsilon)^{n/2}}{|\det D^2\phi(y_k)|^{1/2}} e^{\frac{i\pi}{4} \operatorname{sgn}(D^2\phi(y_k))} (a(y_k) + O(\varepsilon)), \end{aligned}$$

according to Lemma 2. Using these estimates and a partition of unity, we thereby obtain the asymptotic formula as $\varepsilon \rightarrow 0$:

$$(54) \quad I_\varepsilon = (2\pi\varepsilon)^{n/2} \sum_{k=1}^N \frac{e^{\frac{i\phi(y_k)}{\varepsilon}}}{|\det D^2\phi(y_k)|^{1/2}} e^{\frac{i\pi}{4} \operatorname{sgn}(D^2\phi(y_k))} (a(y_k) + O(\varepsilon)).$$

Example 4 (Stationary phase for the wave equation, continued). We can now apply the foregoing theory to (38), which states

$$I_+^\varepsilon(x, t) = \frac{1}{(2\pi\varepsilon)^n} \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} a(z) e^{i\varepsilon\phi_+(x, y, z, t)} dy dz.$$

This is of the form (40), with (x, t) replacing x and (y, z) replacing y .

Define for fixed $x \in \mathbb{R}^n$, $t > 0$, the set where the mapping $(y, z) \mapsto \phi_+(x, y, z, t)$ is stationary:

$$S^+ := \{(y, z) \mid D_{y,z}\phi_+(x, y, z, t) = 0\}.$$

Recall from (39) that $\phi_+(x, y, z, t) = (x - z) \cdot y + t|y| + p(z)$; and so

$$\begin{cases} D_y\phi_+ = (x - z) + t\frac{y}{|y|} & (y \neq 0), \\ D_z\phi_+ = -y + Dp(z). \end{cases}$$

Consequently

$$(55) \quad S^+ = \left\{ (y, z) \mid x = z - \frac{tDp(z)}{|Dp(z)|}, y = Dp(z) \right\},$$

and we here and henceforth assume that $y = Dp(z) \neq 0$ if $(y, z) \in S^+$.

Now if $y \neq 0$,

$$D_{y,z}^2\phi_+ = \begin{pmatrix} D_y^2\phi_+ & D_{yz}^2\phi_+ \\ D_{yz}^2\phi_+ & D_z^2\phi_+ \end{pmatrix}_{2n \times 2n} = \begin{pmatrix} \frac{t}{|y|}\mathbf{P}(y) & -I \\ -I & D^2p \end{pmatrix},$$

for $\mathbf{P}(y) := I - \frac{y \otimes y}{|y|^2}$. We have $y = Dp(z)$ on the stationary set S^+ , and so

$$\det(D_{y,z}^2\phi_+) = (-1)^n \det\left(I - \frac{t}{|Dp|} D^2p \mathbf{P}(Dp)\right).$$

Now the symmetric matrix $\Sigma(z) := \frac{\mathbf{P}(Dp)}{|Dp|} D^2p \mathbf{P}(Dp)$ has n real eigenvalues $\lambda_1(z), \dots, \lambda_n(z)$. Since $\Sigma(z)Dp = 0$, we may take $\lambda_n(z) = 0$. The other eigenvalues $\lambda_1(z), \dots, \lambda_{n-1}(z)$ turn out to be the principal curvatures $\kappa_1(z), \dots, \kappa_{n-1}(z)$ of the level surface of p passing through z . Since $\mathbf{P}^2 = \mathbf{P}$, the nonzero eigenvalues of $\frac{1}{|Dp|} D^2p \mathbf{P}(Dp)$ are the nonzero principal curvatures. Thus on S^+ ,

$$(56) \quad \det(D_{y,z}^2\phi_+) = (-1)^n \prod_{i=1}^{n-1} (1 - t\kappa_i(z)).$$

We apply the stationary phase estimates for $x_0 \in \mathbb{R}^n$ and small $t_0 > 0$. If t_0 is small enough, we can invoke the Implicit Function Theorem (§C.7) to solve uniquely the expressions

$$x_0 = z - t_0 \frac{Dp(z)}{|Dp(z)|}, \quad y = Dp(z)$$

for $y_0 = y(x_0, t_0)$, $z_0 = z(x_0, t_0)$. Thus the asymptotic formula (54) (with $2n$ replacing n) implies

$$(57) \quad \begin{aligned} I_+^\varepsilon(x_0, t_0) &= \frac{1}{(2\pi\varepsilon)^n} \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} a(z) e^{i\varepsilon \phi_+(x_0, y, z, t_0)} dy dz \\ &= \frac{e^{\frac{i\pi}{4} \operatorname{sgn}(D_{y,z}^2 \phi_+)}}{|\det D_{y,z}^2 \phi_+|^{1/2}} e^{i\varepsilon \phi_+[a(z_0) + O(\varepsilon)]} \quad \text{as } \varepsilon \rightarrow 0, \end{aligned}$$

ϕ_+ and $D_{y,z}^2 \phi_+$ evaluated at (x_0, y_0, z_0, t_0) . Recall further that (56) gives us an explicit function for $\det(D_{y,z}^2 \phi_+)$. A similar asymptotic formula holds for $I_-^\varepsilon(x_0, t_0)$. Since $u^\varepsilon(x_0, t_0) = \frac{1}{2}(I_+^\varepsilon(x_0, t_0) + I_-^\varepsilon(x_0, t_0))$, we derive detailed information concerning the limits as $\varepsilon \rightarrow 0$, at least for small times $t_0 > 0$. \square

Remark (Optics and stationary phase). It remains to discuss briefly the connections between the formal geometric optics and the stationary phase approaches. Recall that the former brought us to the two Hamilton–Jacobi equations

$$(58) \quad p_t \pm |Dp| = 0$$

for the phase function of $u^\varepsilon = a_\varepsilon e^{\frac{ip\varepsilon}{\varepsilon}} = (a + o(1))e^{\frac{ip+o(1)}{\varepsilon}}$. Now the characteristic equations for the PDE $p_t - |Dp| = 0$ are

$$(59) \quad \begin{cases} \dot{\mathbf{x}}(s) = -\frac{\mathbf{p}(s)}{|\mathbf{p}(s)|} \\ \dot{\mathbf{p}}(s) = 0, \end{cases}$$

as previously discussed in §3.2.2. In particular given a point $x \in \mathbb{R}^n$, $t > 0$, where t is small, the projected characteristic $\mathbf{x}(\cdot)$ is a straight line, starting at the unique point z satisfying

$$z = x + t \frac{Dp(z)}{|Dp(z)|}.$$

But this relation is precisely what determines the stationary set S^+ above. Likewise, the characteristics of the partial differential equation $p_t + |Dp| = 0$ determine the stationary set S^- for ϕ_- .

4.5.4. Homogenization.

Homogenization theory studies the effects of high-frequency oscillations in the coefficients upon solutions of PDE. In the simplest setting we are given a partial differential equation with two natural *length scales*, a macroscopic scale of order 1 and a microscopic scale of order ε , the latter measuring the period of the oscillations. For fixed, but small, $\varepsilon > 0$ the solution u^ε of the PDE will in general be complicated, having different behaviors on the two length scales.

Homogenization theory studies the limiting behavior $u^\varepsilon \rightarrow u$ as $\varepsilon \rightarrow 0$. The idea is that in this limit the high-frequency effects will “average out”, and there will be a simpler, effective limiting PDE that u solves. One of the difficulties is even to guess the form of the limiting partial differential equation, and for this, multiscale expansions in ε may be useful.

Example 5 (Periodic homogenization of an elliptic equation). This example assumes some familiarity with the theory of divergence-structure, second-order elliptic PDE, as developed later in Chapter 6.

Let U denote an open, bounded subset of \mathbb{R}^n , with smooth boundary ∂U , and consider this boundary-value problem for a divergence structure PDE:

$$(60) \quad \begin{cases} -\sum_{i,j=1}^n (a^{ij}(\frac{x}{\varepsilon}) u_{x_i}^\varepsilon)_{x_j} = f & \text{in } U \\ u^\varepsilon = 0 & \text{in } \partial U. \end{cases}$$

Here $f : U \rightarrow \mathbb{R}$ is given, as are the coefficients a^{ij} ($i, j = 1, \dots, n$). We will assume the uniform ellipticity condition

$$\sum_{i,j=1}^n a^{ij}(y) \xi_i \xi_j \geq \theta |\xi|^2$$

for some constant $\theta > 0$ and all $y, \xi \in \mathbb{R}^n$. We suppose also

$$(61) \quad \text{the mapping } y \mapsto a^{ij}(y) \text{ is } Q\text{-periodic} \quad (y \in \mathbb{R}^n),$$

Q denoting the unit cube in \mathbb{R}^n . Thus the coefficients $a^{ij}(\frac{x}{\varepsilon})$ in (60) are rapidly oscillating in x for small $\varepsilon > 0$, and we inquire as to the effect this has upon the solution u^ε . (In applications u^ε represents, say, the electric potential within a nonisotropic body having small-scale, periodic structure.)

In the following heuristic discussion let us assume

$$(62) \quad u^\varepsilon \rightarrow u \quad \text{as } \varepsilon \rightarrow 0$$

in some suitable sense and try to determine an equation which u satisfies. The trick is to suppose u^ε admits the following *two-scale* expansion:

$$(63) \quad u^\varepsilon(x) = u_0(x, x/\varepsilon) + \varepsilon u_1(x, x/\varepsilon) + \varepsilon^2 u_2(x, x/\varepsilon) + \dots,$$

where $u_i : U \times Q \rightarrow \mathbb{R}$ ($i = 0, 1, \dots$), $u_i = u_i(x, y)$. We are thus thinking of the terms u_i as being both functions of the macroscopic variable x and periodic functions of the microscopic variable $y = \frac{x}{\varepsilon}$. The plan is to plug (63) into (60) and to determine thereby u_0, u_1 , etc. We are primarily interested in $u = u_0$.

Now if $v(x) = w(x, x/\varepsilon)$ for some function $w = w(x, y)$, then $\frac{\partial}{\partial x_i} v = \left(\frac{\partial}{\partial x_i} + \frac{1}{\varepsilon} \frac{\partial}{\partial y_i} \right) w$, $i = 1, \dots, n$. Thus, writing

$$Lv = - \sum_{i,j=1}^n (a^{ij}(x/\varepsilon) v_{x_i})_{x_j},$$

we have

$$(64) \quad L = \frac{1}{\varepsilon^2} L_1 + \frac{1}{\varepsilon} L_2 + L_3,$$

where

$$(65) \quad \begin{cases} \text{(a)} & L_1 w := - \sum_{i,j=1}^n (a^{ij}(y) w_{y_i})_{y_j}, \\ \text{(b)} & L_2 w := - \sum_{i,j=1}^n (a^{ij}(y) w_{x_i})_{y_j} + (a^{ij}(y) w_{y_i})_{x_j}, \\ \text{(c)} & L_3 w := - \sum_{i,j=1}^n (a^{ij}(y) w_{x_i})_{x_j}. \end{cases}$$

Next plug the expansion (63) into the PDE $Lu^\varepsilon = f$, and utilize the decomposition (64), (65) to find

$$\begin{aligned} \frac{1}{\varepsilon^2} L_1 u_0 + \frac{1}{\varepsilon} (L_1 u_1 + L_2 u_0) + (L_1 u_2 + L_2 u_1 + L_3 u_0) \\ + \{\text{terms involving } \varepsilon, \varepsilon^2, \dots\} = f. \end{aligned}$$

Equating like powers of ε , we deduce

$$(66) \quad \begin{cases} \text{(a)} & L_1 u_0 = 0, \\ \text{(b)} & L_1 u_1 + L_2 u_0 = 0, \\ \text{(c)} & L_1 u_2 + L_2 u_1 + L_3 u_0 = f, \quad \text{etc.} \end{cases}$$

We examine these PDE to deduce information concerning u_0, u_1, u_2 . Now in view of (65)(a), (66)(a) for each fixed x , $u_0(x, y)$ solves $L_1 u_0 = 0$

and is Q -periodic. It turns out that the only such solutions are constant in y . Thus in fact

$$(67) \quad u_0 = u(x) \text{ depends only on } x.$$

Next employ (67), (65)(b), (66)(b) to discover

$$(68) \quad L_1 u_1 = \sum_{i,j=1}^n a^{ij}(y)_{y_j} u_{x_i}.$$

We can as follows separate variables to represent u_1 more simply. For $i = 1, \dots, n$, let $\chi^i = \chi^i(y)$ solve

$$(69) \quad \begin{cases} L_1 \chi^i = -\sum_{j=1}^n a^{ij}(y)_{y_j} & \text{in } Q \\ \chi^i & Q\text{-periodic.} \end{cases}$$

As the right-hand side of the PDE in (69) has integral zero over Q , this problem has a solution χ^i (unique up to an additive constant). Here we are applying the *Fredholm alternative*: see Chapter 6.

Using (69), we obtain

$$(70) \quad u_1(x, y) = -\sum_{i=1}^n \chi^i(y) u_{x_i}(x) + \tilde{u}_1(x),$$

\tilde{u}_1 denoting an arbitrary function of x alone.

Finally let us recall (66)(c):

$$(71) \quad L_1 u_2 = f - L_2 u_1 - L_3 u_0.$$

In view of (65)(a) this PDE will have a Q -periodic solution (in the variable y) only if the integral of the right-hand side over Q is zero. Thus we require

$$(72) \quad \int_Q L_2 u_1 + L_3 u_0 \, dy = \int_Q f \, dy = f(x).$$

Owing to (65)(b) and (70),

$$\begin{aligned} \int_Q L_2 u_1 \, dy &= -\sum_{j,k=1}^n \left(\int_Q a^{jk}(y) u_{1,y_k} \, dy \right)_{x_j} \\ &= \sum_{i,j,k=1}^n \left(\int_Q a^{jk}(y) \chi_{y_k}^i(y) \, dy \right) u_{x_i x_j}. \end{aligned}$$

Since $u = u_0$, this calculation and (72) imply

$$-\sum_{i,j=1}^n \left(\int_Q a^{ij}(y) - \sum_{k=1}^n a^{jk}(y) \chi_{y_k}^i(y) dy \right) u_{x_i x_j}(x) = f(x).$$

That is,

$$(73) \quad \begin{cases} -\sum_{i,j=1}^n \bar{a}^{ij} u_{x_i x_j} = f & \text{in } U \\ u = 0 & \text{on } \partial U, \end{cases}$$

where

$$(74) \quad \bar{a}^{ij} := \int_Q a^{ij}(y) - \sum_{k=1}^n a^{jk}(y) \chi_{y_k}^i(y) dy \quad (i, j = 1, \dots, n)$$

are the *homogenized coefficients* and χ^i solves the *corrector problem* (69) ($i = 1, \dots, n$). Thus we expect $u^\varepsilon \rightarrow u$ as $\varepsilon \rightarrow 0$ and u to solve the limit problem (73). \square

This example clearly illustrates the power of the multiscale expansion method. It is not at all readily apparent that the high-frequency oscillations in the coefficients of (60) lead to a constant coefficient PDE of the precise form (73), (74). We will later introduce some variational principles characterizing $\bar{A} = ((\bar{a}^{ij}))$: see Problem 18.

4.6. POWER SERIES

We discuss in this final section solving boundary-value problems for partial differential equations by looking for solutions expressed as power series.

4.6.1. Noncharacteristic surfaces.

We begin with some fairly general comments concerning the solvability of the k^{th} -order quasilinear PDE

$$(1) \quad \sum_{|\alpha|=k} a_\alpha(D^{k-1}u, \dots, u, x) D^\alpha u + a_0(D^{k-1}u, \dots, u, x) = 0$$

in some open region $U \subset \mathbb{R}^n$. Let us assume that Γ is a smooth, $(n-1)$ -dimensional hypersurface in U , the unit normal to which at any point $x^0 \in \Gamma$ is $\nu(x^0) = \nu = (\nu_1, \dots, \nu_n)$.

NOTATION. The j^{th} normal derivative of u at $x^0 \in \Gamma$ is

$$\frac{\partial^j u}{\partial \nu^j} := \sum_{|\alpha|=j} \binom{j}{\alpha} D^\alpha u \nu^\alpha = \sum_{\alpha_1+\dots+\alpha_n=j} \binom{j}{\alpha} \frac{\partial^j u}{\partial^{\alpha_1} x_1 \dots \partial^{\alpha_n} x_n} \nu_1^{\alpha_1} \dots \nu_n^{\alpha_n}. \quad \square$$

Now let $g_0, \dots, g_{k-1} : \Gamma \rightarrow \mathbb{R}$ be k given functions. The *Cauchy problem* is then to find a function u solving the PDE (1), subject to the boundary conditions

$$(2) \quad u = g_0, \quad \frac{\partial u}{\partial \nu} = g_1, \dots, \quad \frac{\partial^{k-1} u}{\partial \nu^{k-1}} = g_{k-1} \quad \text{on } \Gamma.$$

We say that the equations (2) prescribe the *Cauchy data* g_0, \dots, g_{k-1} on Γ .

We now pose a basic question:

$$(3) \quad \left\{ \begin{array}{l} \text{Assuming } u \text{ is a smooth solution of the PDE (1),} \\ \text{do conditions (2) allow us to compute } \textit{all} \text{ the partial} \\ \text{derivatives of } u \text{ along } \Gamma? \end{array} \right.$$

This must certainly be so, if we are ever going to be able to calculate the terms of a power series representation formula for u .

a. Flat boundaries. We examine first the special circumstance that $U = \mathbb{R}^n$ and Γ is the plane $\{x_n = 0\}$. In this situation we can take $\nu = e_n$, and so the Cauchy conditions (2) read

$$(4) \quad u = g_0, \quad \frac{\partial u}{\partial x_n} = g_1, \dots, \quad \frac{\partial^{k-1} u}{\partial x_n^{k-1}} = g_{k-1} \quad \text{on } \{x_n = 0\}.$$

Which further partial derivatives of u can we compute along the plane $\Gamma = \{x_n = 0\}$? First, notice that since $u = g_0$ on all of Γ , we can differentiate tangentially, that is, with respect to x_i ($i = 1, \dots, n-1$), to find

$$\frac{\partial u}{\partial x_i} = \frac{\partial g_0}{\partial x_i} \quad (i = 1, \dots, n-1).$$

Since we also know from (4) that

$$\frac{\partial u}{\partial x_n} = g_1,$$

we can determine the full gradient Du along $\Gamma = \{x_n = 0\}$. Similarly, we have

$$\left\{ \begin{array}{l} \frac{\partial^2 u}{\partial x_n \partial x_i} = \frac{\partial g_1}{\partial x_i} \quad (i = 1, \dots, n-1) \\ \frac{\partial^2 u}{\partial x_n^2} = g_2, \end{array} \right.$$

and hence we can compute D^2u on Γ . Next, we see

$$\frac{\partial^3 u}{\partial x_i \partial x_j \partial x_m} = \begin{cases} \frac{\partial^3 g_0}{\partial x_i \partial x_j \partial x_m} & \text{if } i, j, m = 1, \dots, n-1 \\ \frac{\partial^2 g_1}{\partial x_i \partial x_j} & \text{if } i, j = 1, \dots, n-1; m = n \\ \frac{\partial g_2}{\partial x_i} & \text{if } i = 1, \dots, n-1; j = m = n \\ g_3 & \text{if } i = j = m = n \end{cases}$$

along Γ , and so we can compute D^3u there. Continuing, it is straightforward to check that employing the Cauchy conditions (4), we can compute $u, Du, \dots, D^{k-1}u$ on Γ .

Difficulties will arise, however, when we try to calculate $D^k u$. In this circumstance it is not hard to verify that we can determine each partial derivative of u of order k along $\Gamma = \{x_n = 0\}$ from the Cauchy data (4), except for the k^{th} -order normal derivative

$$\frac{\partial^k u}{\partial x_n^k}.$$

Here, at last, we turn to PDE (1) for help. We observe from (1) that *if the coefficient $a_{(0, \dots, 0, k)}$ is nonzero*, we can then solve for

$$(5) \quad \frac{\partial^k u}{\partial x_n^k} = -\frac{1}{a_{(0, \dots, 0, k)}} \left[\sum_{\substack{|\alpha|=k \\ \alpha \neq (0, \dots, 0, k)}} a_\alpha D^\alpha u + a_0 \right],$$

with the coefficients a_α ($|\alpha| = k$) and a_0 evaluated at $(D^{k-1}u, \dots, u, x)$ along Γ . Now in view of the remarks above, everything on the right-hand side of equality (5) can be calculated in terms of the Cauchy data along the plane Γ , and thus we have a formula for the missing k^{th} partial derivative. Consequently we can in fact compute all of $D^k u$ on Γ , provided

$$(6) \quad a_{(0, \dots, 0, k)} \neq 0.$$

We say that the plane $\Gamma = \{x_n = 0\}$ is *noncharacteristic* for the PDE (1) if the function $a_{(0, \dots, k)}$ is nonzero for all values of its arguments.

Can we calculate still higher partial derivatives? Assuming the noncharacteristic condition (6), we observe that we can now augment our list (4) of Cauchy data with the new equality

$$(7) \quad \frac{\partial^k u}{\partial x_n^k} = g_k \quad \text{on } \Gamma = \{x_n = 0\},$$

g_k denoting the right-hand side of (5). But then we can, as before, compute all of $D^{k+1}u$ along Γ , except for the term

$$\frac{\partial^{k+1}u}{\partial x_n^{k+1}}.$$

Again we employ the PDE (1). We differentiate (1) with respect to x_n , evaluate the resulting expression on the plane Γ , and rearrange to find

$$\frac{\partial^{k+1}u}{\partial x_n^{k+1}} = \frac{1}{a_{(0,\dots,0,k)}} \{ \dots \},$$

the dots denoting the sum of various expressions, each of which can be computed along Γ in terms of g_0, \dots, g_k . Consequently we can ascertain all of $D^{k+1}u$ on Γ , and an induction verifies that in fact we can compute all the partial derivatives of u on the plane Γ .

b. General surfaces. We now propose to generalize the results obtained above to the general case that Γ is a smooth hypersurface with normal vector field ν .

DEFINITION. We say the surface Γ is noncharacteristic for the partial differential equation (1) provided

$$(8) \quad \sum_{|\alpha|=k} a_\alpha \nu^\alpha \neq 0 \quad \text{on } \Gamma,$$

for all values of the arguments of the coefficients a_α ($|\alpha| = k$).

THEOREM 1 (Cauchy data and noncharacteristic surfaces). Assume that Γ is noncharacteristic for the PDE (1). Then if u is a smooth solution of (1) and u satisfies the Cauchy conditions (2), we can uniquely compute all the partial derivatives of u along Γ in terms of Γ , the functions g_0, \dots, g_{k-1} , and the coefficients a_α ($|\alpha| = k$), a_0 .

Proof. 1. We will reduce to the special case considered above.

For this, let us choose any point $x^0 \in \Gamma$ and recall §C.1 to find smooth maps $\Phi, \Psi : \mathbb{R}^n \rightarrow \mathbb{R}^n$, so that $\Psi = \Phi^{-1}$ and

$$\Phi(\Gamma \cap B(x^0, r)) \subset \{y_n = 0\}$$

for some $r > 0$. Define

$$v(y) := u(\Psi(y)),$$

so that

$$(9) \quad u(x) = v(\Phi(x)).$$

It is relatively easy to check now that v satisfies a quasilinear partial differential equation having the form

$$(10) \quad \sum_{|\alpha|=k} b_\alpha D^\alpha v + b_0 = 0.$$

2. We claim

$$(11) \quad b_{(0,\dots,0,k)} \neq 0 \quad \text{on } \{y_n = 0\}.$$

Indeed from (9) we see that for any multiindex α with $|\alpha| = k$, we have

$$D^\alpha u = \frac{\partial^k v}{\partial y_n^k} (D\Phi^n)^\alpha + \left\{ \text{terms not involving } \frac{\partial^k v}{\partial y_n^k} \right\}.$$

Thus from (1) it follows that

$$\begin{aligned} 0 &= \sum_{|\alpha|=k} a_\alpha D^\alpha u + a_0 \\ &= \sum_{|\alpha|=k} a_\alpha (D\Phi^n)^\alpha \frac{\partial^k v}{\partial y_n^k} + \left\{ \text{terms not involving } \frac{\partial^k v}{\partial y_n^k} \right\}, \end{aligned}$$

and so

$$b_{(0,\dots,0,k)} = \sum_{|\alpha|=k} a_\alpha (D\Phi^n)^\alpha.$$

But $D\Phi^n$ is parallel to ν on Γ . Consequently $b_{(0,\dots,0,k)}$ is a nonzero multiple of the term

$$\sum_{|\alpha|=k} a_\alpha \nu^\alpha \neq 0.$$

This verifies the claim (11).

3. Let us now define the functions $h_0, h_1, \dots, h_{k-1} : \mathbb{R}^{n-1} \rightarrow \mathbb{R}$ by

$$(12) \quad v = h_0, \quad \frac{\partial v}{\partial y_n} = h_1, \dots, \quad \frac{\partial^{k-1} v}{\partial y_n^{k-1}} = h_{k-1} \quad \text{on } \{y_n = 0\}.$$

Thus we can compute h_0, \dots, h_{k-1} near $y = 0$ in terms of Φ and the functions g_0, \dots, g_{k-1} . But then, using (11) and the special case discussed above, we see that we can calculate all of the partial derivatives of v on $\{y_n = 0\}$ near $y = 0$.

And finally, upon recalling (9), we at last observe that we can compute all the partial derivatives of u on Γ near x^0 . \square

Remark. It is sometimes convenient to recast the noncharacteristic condition (8) into a somewhat different form, by representing Γ as the zero set of a function $w : \mathbb{R}^n \rightarrow \mathbb{R}$. So assume that we are given a function w with

$$\Gamma = \{w = 0\}$$

and $Dw \neq 0$ on Γ . Then $\nu = \pm \frac{Dw}{|Dw|}$ on Γ , and so the noncharacteristic condition (8) becomes

$$(13) \quad \sum_{|\alpha|=k} a_\alpha (Dw)^\alpha \neq 0 \quad \text{on } \Gamma.$$

4.6.2. Real analytic functions.

We review in this section the representation of real-valued functions by power series.

DEFINITION. A function $f : \mathbb{R}^n \rightarrow \mathbb{R}$ is called (real) analytic near x_0 if there exist $r > 0$ and constants $\{f_\alpha\}$ such that

$$f(x) = \sum_{\alpha} f_\alpha (x - x_0)^\alpha \quad (|x - x_0| < r),$$

the sum taken over all multiindices α .

Remarks. (i) Remember that we write $x^\alpha = x_1^{\alpha_1} \cdots x_n^{\alpha_n}$, for the multiindex $\alpha = (\alpha_1, \dots, \alpha_n)$.

(ii) If f is analytic near x_0 , then f is C^∞ near x_0 . Furthermore the constants f_α are computed as $f_\alpha = \frac{D^\alpha f(x_0)}{\alpha!}$, where $\alpha! = \alpha_1! \alpha_2! \cdots \alpha_n!$. Thus f equals its Taylor expansion about x_0 :

$$f(x) = \sum_{\alpha} \frac{1}{\alpha!} D^\alpha f(x_0) (x - x_0)^\alpha \quad (|x - x_0| < r).$$

To simplify, we hereafter take $x_0 = 0$.

Example. If $r > 0$, set

$$f(x) := \frac{r}{r - (x_1 + \cdots + x_n)} \quad \text{for } |x| < r/\sqrt{n}.$$

Then

$$\begin{aligned} f(x) &= \frac{1}{1 - \left(\frac{x_1 + \cdots + x_n}{r}\right)} = \sum_{k=0}^{\infty} \left(\frac{x_1 + \cdots + x_n}{r}\right)^k \\ &= \sum_{k=0}^{\infty} \frac{1}{r^k} \sum_{|\alpha|=k} \binom{|\alpha|}{\alpha} x^\alpha = \sum_{\alpha} \frac{|\alpha|!}{r^{|\alpha| \alpha!}} x^\alpha. \end{aligned}$$

We employed the Multinomial Theorem for the third equality above and recalled that $\binom{|\alpha|}{\alpha} = \frac{|\alpha|!}{\alpha!}$. This power series is absolutely convergent for $|x| < r/\sqrt{n}$. Indeed,

$$\sum_{\alpha} \frac{|\alpha|!}{r^{|\alpha|}\alpha!} |x^{\alpha}| = \sum_{k=0}^{\infty} \left(\frac{|x_1| + \cdots + |x_n|}{r} \right)^k < \infty,$$

since $|x_1| + \cdots + |x_n| \leq |x|\sqrt{n} < r$. □

We will see momentarily that the simple power series illustrated in this example is rather important, since we can use it to majorize, and so confirm the convergence of, other power series.

DEFINITION. *Let*

$$f = \sum_{\alpha} f_{\alpha} x^{\alpha}, \quad g = \sum_{\alpha} g_{\alpha} x^{\alpha}$$

be two power series. We say g majorizes f , written

$$g \gg f,$$

provided

$$g_{\alpha} \geq |f_{\alpha}| \quad \text{for all multiindices } \alpha.$$

LEMMA (Majorants).

(i) *If $g \gg f$ and g converges for $|x| < r$, then f also converges for $|x| < r$.*

(ii) *If $f = \sum_{\alpha} f_{\alpha} x^{\alpha}$ converges for $|x| < r$ and $0 < s\sqrt{n} < r$, then f has a majorant for $|x| < s/\sqrt{n}$.*

Proof. 1. To verify assertion (i), we check

$$\sum_{\alpha} |f_{\alpha} x^{\alpha}| \leq \sum_{\alpha} g_{\alpha} |x_1|^{\alpha_1} \cdots |x_n|^{\alpha_n} < \infty \quad \text{if } |x| < r.$$

2. Let $0 < s\sqrt{n} < r$ and set $y := s(1, \dots, 1)$. Then $|y| = s\sqrt{n} < r$ and so $\sum_{\alpha} f_{\alpha} y^{\alpha}$ converges. Thus there exists a constant C such that

$$|f_{\alpha} y^{\alpha}| \leq C \quad \text{for each multiindex } \alpha.$$

In particular,

$$|f_{\alpha}| \leq \frac{C}{y_1^{\alpha_1} \cdots y_n^{\alpha_n}} = \frac{C}{s^{|\alpha|}} \leq C \frac{|\alpha|!}{s^{|\alpha|}\alpha!}.$$

But then

$$g(x) := \frac{Cs}{s - (x_1 + \cdots + x_n)} = C \sum_{\alpha} \frac{|\alpha|!}{s^{|\alpha|}\alpha!} x^{\alpha}$$

majorizes f for $|x| < s/\sqrt{n}$. □

Remark. We will later need to extend our notation to vector-valued series. So given power series $\{f^k\}_{k=1}^m, \{g^k\}_{k=1}^m$, we set $\mathbf{f} = (f^1, \dots, f^m)$, $\mathbf{g} = (g^1, \dots, g^m)$ and write

$$\mathbf{g} \gg \mathbf{f}$$

to mean

$$g^k \gg f^k \quad (k = 1, \dots, m).$$

4.6.3. Cauchy–Kovalevskaya Theorem.

We turn now to our primary task of building a power series solution for the k^{th} -order quasilinear partial differential equation (1), with analytic Cauchy data (2) specified on an analytic, noncharacteristic hypersurface Γ .

a. Reduction to a first-order system. We intend to construct a solution u as a power series, but must first transform the boundary-value problem (1), (2) into a more convenient form.

First of all, upon flattening out the boundary by an analytic mapping (as in §4.6.1), we can reduce to the situation that $\Gamma \subset \{x_n = 0\}$. Additionally, by subtracting off appropriate analytic functions, we may assume the Cauchy data are identically zero. Consequently we may assume without loss that our problem reads:

$$(14) \quad \left\{ \begin{array}{l} \sum_{|\alpha|=k} a_\alpha(D^{k-1}u, \dots, u, x)D^\alpha u \\ \quad + a_0(D^{k-1}u, \dots, u, x) = 0 \quad \text{for } |x| < r \\ u = \frac{\partial u}{\partial x_n} = \dots = \frac{\partial^{k-1}u}{\partial x_n^{k-1}} = 0 \quad \text{for } |x'| < r, x_n = 0, \end{array} \right.$$

$r > 0$ to be found. Here a_α ($|\alpha| = k$) and a_0 are analytic, and as usual we write $x' = (x_1, \dots, x_{n-1})$.

Finally we transform to a first-order *system*. To do so, we introduce the function

$$\mathbf{u} := (u, \frac{\partial u}{\partial x_1}, \dots, \frac{\partial u}{\partial x_n}, \frac{\partial^2 u}{\partial x_1^2}, \dots, \frac{\partial^{k-1}u}{\partial x_n^{k-1}}),$$

the components of which are all the partial derivatives of u of order less than k . Let m hereafter denote the number of components of \mathbf{u} , so that $\mathbf{u} : \mathbb{R}^n \rightarrow \mathbb{R}^m$, $\mathbf{u} = (u^1, \dots, u^m)$. Observe from the boundary condition in (14) that $\mathbf{u} = 0$ for $|x'| < r, x_n = 0$.

Now for $k \in \{1, \dots, m-1\}$, we can compute $u_{x_n}^k$ in terms of $\{\mathbf{u}_{x_j}\}_{j=1}^{n-1}$. Furthermore in view of the noncharacteristic condition $a_{(0, \dots, 0, k)} \neq 0$ near 0, we can utilize the PDE in (14) also to solve for $u_{x_n}^m$ in terms of \mathbf{u} and $\{\mathbf{u}_{x_j}\}_{j=1}^{n-1}$.

Employing these relations, we can consequently transform (14) into a boundary-value problem for a first-order system for \mathbf{u} , the coefficients of which are analytic functions. This system is of the general form:

$$(15) \quad \begin{cases} \mathbf{u}_{x_n} = \sum_{j=1}^{n-1} \mathbf{B}_j(\mathbf{u}, x') \mathbf{u}_{x_j} + \mathbf{c}(\mathbf{u}, x') & \text{for } |x| < r \\ \mathbf{u} = 0 & \text{for } |x'| < r, x_n = 0, \end{cases}$$

where we are given the analytic functions $\mathbf{B}_j : \mathbb{R}^m \times \mathbb{R}^{n-1} \rightarrow \mathbb{M}^{m \times m}$ ($j = 1, \dots, n-1$) and $\mathbf{c} : \mathbb{R}^m \times \mathbb{R}^{n-1} \rightarrow \mathbb{R}^m$. We will write $\mathbf{B}_j = ((b_j^{kl}))$ and $\mathbf{c} = (c^1, \dots, c^m)$. Carefully note that we have assumed $\{\mathbf{B}_j\}_{j=1}^{n-1}$ and \mathbf{c} do not depend on x_n . We can always reduce to this situation by introducing if necessary a new component u^{m+1} of the unknown \mathbf{u} , with $u^{m+1} \equiv x_n$.

In particular, the components of the system of partial differential equations in (15) read

$$(16) \quad u_{x_n}^k = \sum_{j=1}^{n-1} \sum_{l=1}^m b_j^{kl}(\mathbf{u}, x') u_{x_j}^l + c^k(\mathbf{u}, x') \quad (k = 1, \dots, m).$$

b. Power series for solutions. Having reduced to the special form (15), we can now expand \mathbf{u} into a power series and, more importantly, verify that this series converges near 0.

THEOREM 2 (Cauchy–Kovalevskaya Theorem). *Assume $\{\mathbf{B}_j\}_{j=1}^{n-1}$ and \mathbf{c} are real analytic functions. Then there exist $r > 0$ and a real analytic function*

$$(17) \quad \mathbf{u} = \sum_{\alpha} \mathbf{u}_{\alpha} x^{\alpha}$$

solving the boundary-value problem (15).

Proof. 1. We must compute the coefficients

$$(18) \quad \mathbf{u}_{\alpha} = \frac{D^{\alpha} \mathbf{u}(0)}{\alpha!}$$

in terms of $\{\mathbf{B}_j\}_{j=1}^{n-1}$ and \mathbf{c} and then show that the power series (18) so obtained in fact converges if $|x| < r$ and $r > 0$ is small enough.

2. As the functions $\{\mathbf{B}_j\}_{j=1}^{n-1}$ and \mathbf{c} are analytic, we can write

$$(19) \quad \mathbf{B}_j(z, x') = \sum_{\gamma, \delta} \mathbf{B}_{j, \gamma, \delta} z^{\gamma} x^{\delta} \quad (j = 1, \dots, n-1)$$

and

$$(20) \quad \mathbf{c}(z, x') = \sum_{\gamma, \delta} \mathbf{c}_{\gamma, \delta} z^{\gamma} x^{\delta},$$

these power series convergent if $|z| + |x'| < s$ for some small $s > 0$. Thus

$$(21) \quad \mathbf{B}_{j, \gamma, \delta} = \frac{D_z^{\gamma} D_x^{\delta} \mathbf{B}_j(0, 0)}{(\gamma + \delta)!}, \quad \mathbf{c}_{\gamma, \delta} = \frac{D_z^{\gamma} D_x^{\delta} \mathbf{c}(0, 0)}{(\gamma + \delta)!}$$

for $j = 1, \dots, n - 1$ and all multiindices γ, δ .

3. Since $\mathbf{u} \equiv 0$ on $\{x_n = 0\}$, we have

$$(22) \quad \mathbf{u}_{\alpha} = \frac{D^{\alpha} \mathbf{u}(0)}{\alpha!} = 0 \quad \text{for all multiindices } \alpha \text{ with } \alpha_n = 0.$$

Now fix $i \in \{1, \dots, n - 1\}$ and differentiate (16) with respect to x_i :

$$u_{x_n x_i}^k = \sum_{j=1}^{n-1} \sum_{l=1}^m \left(b_j^{kl} u_{x_i x_j}^l + b_{j, x_i}^{kl} u_{x_j}^l + \sum_{p=1}^m b_{j, z_p}^{kl} u_{x_i}^p u_{x_j}^l \right) + c_{x_i}^k + \sum_{p=1}^m c_{z_p}^k u_{x_i}^p.$$

In view of (22), we conclude $u_{x_n x_i}^k(0) = c_{x_i}^k(0, 0)$.

If α is a multiindex having the form $\alpha = (\alpha_1, \dots, \alpha_{n-1}, 1) = (\alpha', 1)$, we likewise prove by induction that

$$D^{\alpha} u^k(0) = D^{\alpha'} c^k(0, 0).$$

Next suppose $\alpha = (\alpha', 2)$. Then

$$\begin{aligned} D^{\alpha} u^k &= D^{\alpha'} (u_{x_n}^k)_{x_n} \\ &= D^{\alpha'} \left(\sum_{j=1}^{n-1} \sum_{l=1}^m b_j^{kl} u_{x_j}^l + c^k \right)_{x_n} \quad \text{by (16)} \\ &= D^{\alpha'} \left(\sum_{j=1}^{n-1} \sum_{l=1}^m (b_j^{kl} u_{x_j x_n}^l + \sum_{p=1}^m b_{j, z_p}^{kl} u_{x_n}^p u_{x_j}^l) + \sum_{p=1}^m c_{z_p}^k u_{x_n}^p \right). \end{aligned}$$

Thus

$$D^{\alpha} u^k(0) = D^{\alpha'} \left(\sum_{j=1}^{n-1} \sum_{l=1}^m b_j^{kl} u_{x_j x_n}^l + \sum_{p=1}^m c_{z_p}^k u_{x_n}^p \right) \Big|_{\mathbf{x}=\mathbf{u}=0}.$$

The expression on the right-hand side can be worked out to be a polynomial *with nonnegative coefficients* involving various derivatives of $\{\mathbf{B}_j\}_{j=1}^{n-1}$ and \mathbf{c} and the derivatives $D^{\beta} \mathbf{u}$, where $\beta_n \leq 1$.

More generally, for each multiindex α and each $k \in \{1, \dots, m\}$, we compute

$$D^\alpha u^k(0) = p_\alpha^k(\dots, D_z^\gamma D_x^\delta \mathbf{B}_j, \dots, D_z^\gamma D_x^\delta \mathbf{c}, \dots, D^\beta \mathbf{u}, \dots) \Big|_{x=\mathbf{u}=0},$$

where p_α^k denotes some polynomial with nonnegative coefficients.

Recalling (18)–(21), we deduce for each α, k that

$$(23) \quad u_\alpha^k = q_\alpha^k(\dots, \mathbf{B}_{j,\gamma,\delta}, \dots, \mathbf{c}_{\gamma,\delta}, \dots, \mathbf{u}_\beta, \dots),$$

where

$$(24) \quad q_\alpha^k \text{ is a polynomial with nonnegative coefficients}$$

and

$$(25) \quad \beta_n \leq \alpha_n - 1 \text{ for each multiindex } \beta \text{ on the right-hand side of (23).}$$

4. Thus far we have merely demonstrated that if there is a smooth solution of (15), then we can compute all of its derivatives at 0 in terms of known quantities. This of course we already know from the discussion in §4.6.1, since the plane $\{x_n = 0\}$ is noncharacteristic.

We now intend to employ (22)–(25) and the *method of majorants* to show the power series (17) actually converges if $|x| < r$ and r is small. For this, let us first suppose

$$(26) \quad \mathbf{B}_j^* \gg \mathbf{B}_j \quad (j = 1, \dots, n-1)$$

and

$$(27) \quad \mathbf{c}^* \gg \mathbf{c},$$

where

$$\mathbf{B}_j^* := \sum_{\gamma,\delta} \mathbf{B}_{j,\gamma,\delta}^* z^\gamma x^\delta \quad (j = 1, \dots, n-1)$$

and

$$\mathbf{c}^* := \sum_{\gamma,\delta} \mathbf{c}_{\gamma,\delta}^* z^\gamma x^\delta,$$

these power series convergent for $|z| + |x'| < s$. Then for all j, γ, δ ,

$$(28) \quad 0 \leq |\mathbf{B}_{j,\gamma,\delta}| \leq \mathbf{B}_{j,\gamma,\delta}^*, \quad 0 \leq |\mathbf{c}_{\gamma,\delta}| \leq \mathbf{c}_{\gamma,\delta}^*.$$

We consider next the new boundary-value problem

$$(29) \quad \begin{cases} \mathbf{u}_{x_n}^* = \sum_{j=1}^{n-1} \mathbf{B}_j^*(\mathbf{u}^*, x') \mathbf{u}_{x_j}^* + \mathbf{c}^*(\mathbf{u}^*, x') & \text{for } |x| < r \\ \mathbf{u}^* = 0 & \text{for } |x'| < r, x_n = 0, \end{cases}$$

and, as above, look for a solution having the form

$$(30) \quad \mathbf{u}^* = \sum_{\alpha} \mathbf{u}_{\alpha}^* x^{\alpha},$$

where

$$(31) \quad \mathbf{u}_{\alpha}^* = \frac{D^{\alpha} \mathbf{u}^*(0)}{\alpha!}.$$

5. We claim

$$0 \leq |u_{\alpha}^k| \leq u_{\alpha}^{k*} \quad \text{for each multiindex } \alpha.$$

The proof is by induction. The general step follows since

$$\begin{aligned} |u_{\alpha}^k| &= |q_{\alpha}^k(\dots, \mathbf{B}_{j,\gamma,\delta}, \dots, \mathbf{c}_{\gamma,\delta}, \dots, \mathbf{u}_{\beta}, \dots)| \quad \text{by (23)} \\ &\leq q_{\alpha}^k(\dots, |\mathbf{B}_{j,\gamma,\delta}|, \dots, |\mathbf{c}_{\gamma,\delta}|, \dots, |\mathbf{u}_{\beta}|, \dots) \quad \text{by (24)} \\ &\leq q_{\alpha}^k(\dots, \mathbf{B}_{j,\gamma,\delta}^*, \dots, \mathbf{c}_{\gamma,\delta}^*, \dots, \mathbf{u}_{\beta}^*, \dots) \quad \text{by (24), (28) and induction} \\ &= u_{\alpha}^{k*}. \end{aligned}$$

Thus

$$(32) \quad \mathbf{u}^* \gg \mathbf{u},$$

and so it suffices to prove that the power series (30) converges near zero.

6. As demonstrated in the proof of assertion (ii) of the lemma in §4.6.2, if we choose

$$\mathbf{B}_j^* := \frac{Cr}{r - (x_1 + \dots + x_{n-1}) - (z_1 + \dots + z_m)} \begin{pmatrix} 1 & \dots & 1 \\ & \ddots & \\ 1 & \dots & 1 \end{pmatrix}$$

for $j = 1, \dots, n - 1$, and

$$\mathbf{c}^* := \frac{Cr}{r - (x_1 + \dots + x_{n-1}) - (z_1 + \dots + z_m)} (1, \dots, 1),$$

then (26), (27) will hold if C is large enough, $r > 0$ is small enough, and $|x'| + |z| < r$.

Hence the problem (29) reads

$$\begin{cases} \mathbf{u}_{x_n}^* = \frac{Cr}{r - (x_1 + \cdots + x_{n-1}) - (u^{1*} + \cdots + u^{m*})} \left(\sum_{j,l} \mathbf{u}_{x_j}^{l*} + 1 \right) & \text{for } |x| < r \\ \mathbf{u}^* = 0 & \text{for } |x'| < r, x_n = 0. \end{cases}$$

However, this problem has an explicit solution, namely

$$(33) \quad \mathbf{u}^* = v^*(1, \dots, 1),$$

for

$$(34) \quad v^*(x) := \frac{1}{mn} (r - (x_1 + \cdots + x_{n-1}) - [(r - (x_1 + \cdots + x_{n-1}))^2 - 2mnCr x_n]^{1/2}).$$

This expression is analytic for $|x| < r$, provided $r > 0$ is sufficiently small. Thus \mathbf{u}^* defined by (33) necessarily has the form (30), (31), the power series (30) converging for $|x| < r$. As $\mathbf{u}^* \gg \mathbf{u}$, the power series (17) converges as well for $|x| < r$.

This defines the analytic function \mathbf{u} near 0. Since the Taylor expansions of the analytic functions \mathbf{u}_{x_n} and $\sum_{j=1}^{n-1} \mathbf{B}_j(\mathbf{u}, x) \mathbf{u}_{x_j} + \mathbf{c}(\mathbf{u}, x)$ agree at 0, they agree as well throughout the region $|x| < r$. \square

The Cauchy–Kovalevskaya Theorem is valid also for fully nonlinear, analytic PDE: see Folland [F1].

4.7. PROBLEMS

In the following exercises, all given functions are assumed smooth, unless otherwise stated.

1. Use separation of variables to find a nontrivial solution u of the PDE

$$u_{x_1}^2 u_{x_1 x_1} + 2u_{x_1} u_{x_2} u_{x_1 x_2} + u_{x_2}^2 u_{x_2 x_2} = 0 \quad \text{in } \mathbb{R}^2.$$

(G. Aronsson, *Manuscripta Math.* 47 (1984), 133–151)

2. Consider Laplace's equation $\Delta u = 0$ in \mathbb{R}^2 , taken with the Cauchy data

$$u = 0, \quad \frac{\partial u}{\partial x_2} = \frac{1}{n} \sin(nx_1) \quad \text{on } \{x_2 = 0\}.$$

Employ separation of variables to derive the solution

$$u = \frac{1}{n^2} \sin(nx_1) \sinh(nx_2).$$

What happens to u as $n \rightarrow \infty$? Is the Cauchy problem for Laplace's equation well-posed? (This example is due to Hadamard.)

3. Find explicit formulas for v and σ , so that $u(x, t) := v(x - \sigma t)$ is a traveling wave solution of the nonlinear diffusion equation

$$u_t - u_{xx} = f(u),$$

where

$$f(z) = -2z^3 + 3z^2 - z.$$

Assume $\lim_{s \rightarrow \infty} v = 1$, $\lim_{s \rightarrow -\infty} v = 0$, $\lim_{s \rightarrow \pm\infty} v' = 0$.

(Hint: Multiply the equation $v'' + \sigma v' + f(v) = 0$ by v' and integrate, to determine the value of σ .)

4. If we look for a radial solution $u(x) = v(r)$ of the nonlinear elliptic equation

$$-\Delta u = u^p \quad \text{in } \mathbb{R}^n,$$

where $r = |x|$ and $p > 1$, we are led to the nonautonomous ODE

$$(*) \quad v'' + \frac{n-1}{r} v' + v^p = 0.$$

Show that the *Emden-Fowler transformation*

$$t := \log r, \quad x(t) := e^{\frac{2t}{p-1}} v(e^t)$$

converts $(*)$ into an autonomous ODE for the new unknown $x = x(t)$.

5. Find a nonnegative scaling invariant solution having the form

$$u(x, t) = t^{-\alpha} v(xt^{-\beta})$$

for the nonlinear heat equation

$$u_t - \Delta(u^\gamma) = 0,$$

where $\frac{n-2}{n} < \gamma < 1$. Your solution should go to zero algebraically as $|x| \rightarrow \infty$.

6. Find a solution of

$$-\Delta u + u^{\frac{n+2}{n-2}} = 0 \quad \text{in } B(0, 1)$$

having the form $u = \alpha(1 - |x|^2)^{-\beta}$ for positive constants α, β . This example shows that a solution of a nonlinear PDE can be finite within a region and yet approach infinity everywhere on its boundary.

7. Consider the viscous conservation law

$$(*) \quad u_t + F(u)_x - au_{xx} = 0 \quad \text{in } \mathbb{R} \times (0, \infty),$$

where $a > 0$ and F is uniformly convex.

(a) Show u solves $(*)$ if $u(x, t) = v(x - \sigma t)$ and v is defined implicitly by the formula

$$s = \int_c^{v(s)} \frac{a}{F(z) - \sigma z + b} dz \quad (s \in \mathbb{R}),$$

where b and c are constants.

(b) Demonstrate that we can find a traveling wave satisfying

$$\lim_{s \rightarrow -\infty} v(s) = u_l, \quad \lim_{s \rightarrow \infty} v(s) = u_r$$

for $u_l > u_r$, if and only if

$$\sigma = \frac{F(u_l) - F(u_r)}{u_l - u_r}.$$

(c) Let u^ε denote the above traveling wave solution of $(*)$ for $a = \varepsilon$, with $u^\varepsilon(0, 0) = \frac{u_l + u_r}{2}$. Compute $\lim_{\varepsilon \rightarrow 0} u^\varepsilon$ and explain your answer.

8. Prove that if u is the solution of problem (23) for Schrödinger's equation in §4.3 given by formula (20), then

$$\|u(\cdot, t)\|_{L^\infty(\mathbb{R}^n)} \leq \frac{1}{(4\pi|t|)^{n/2}} \|g\|_{L^1(\mathbb{R}^n)}$$

for each $t \neq 0$.

9. Assume that u solves the nonlinear heat equation

$$u_t = \frac{u_{xx}}{u_x^2} \quad \text{in } \mathbb{R} \times (0, \infty)$$

with $u_x > 0$. Let v denote the inverse function to u in the variable x for each time $t > 0$, so that $y = u(x, t)$ if and only if $x = v(y, t)$.

Show that v solves a linear PDE.

10. Find a function $f : \mathbb{R}^3 \rightarrow \mathbb{R}$, $f = f(z, p_1, p_2)$, so that if u is any solution of the rotated wave equation

$$u_{xt} = 0,$$

then $w := f(u, u_x, u_t)$ solves *Liouville's equation*

$$w_{xt} = e^w.$$

(Hint: Show that f must have the form $f(z, p_1, p_2) = a(z) + b(p_1) + c(p_2)$.)

11. (Lax pairs) Assume that $\{L(t)\}_{t \geq 0}$ is a family of symmetric linear operators on some real Hilbert space H , satisfying the evolution equation

$$\dot{L} = [B, L] = BL - LB,$$

for some collection of operators $\{B(t)\}_{t \geq 0}$. Suppose also that we have a corresponding family of eigenvalues $\{\lambda(t)\}_{t \geq 0}$ and eigenvectors $\{w(t)\}_{t \geq 0}$:

$$L(t)w(t) = \lambda(t)w(t).$$

Assume that L, B, λ and w all depend smoothly upon the time parameter t .

Show that

$$\dot{\lambda} \equiv 0.$$

(Hint: Differentiate the identity $Lw = \lambda w$ with respect to t . Calculate $(\dot{\lambda}w, w)$.)

12. (Continuation) Given a function $u = u(x, t)$, define the linear operators $L(t)v := -v_{xx} + uv$ and $B(t)v = -4v_{xxx} + 6uv_x + 3u_xv$.

Show that

$$(\dot{L} - [B, L])v = (u_t + u_{xxx} - 6uu_x)v = 0.$$

Consequently, if u solves this form of the KdV equation:

$$u_t - 6uu_x + u_{xxx} = 0,$$

then the eigenvalues of the linear operators $L(t)$ do not change with time.

13. Show that we can construct an explicit solution of the initial-value problem (17), (19) from Example 1 in §4.5.1, having the form

$$v(z, t) = \frac{1}{\sigma(t)} \frac{1}{(\pi\gamma(t))^{1/2}} e^{-z^2/\gamma(t)} \quad (z \in \mathbb{R}, t > 0),$$

the function $\gamma(t)$ to be found. Substitute into the PDE and determine an ODE that γ should satisfy. What is the initial condition for this ODE?

14. Let u^ϵ and v^ϵ solve the system

$$\begin{cases} u_t^\epsilon + \frac{1}{\epsilon} u_x^\epsilon = \frac{(v^\epsilon)^2 - (u^\epsilon)^2}{\epsilon^2} \\ v_t^\epsilon - \frac{1}{\epsilon} v_x^\epsilon = \frac{(u^\epsilon)^2 - (v^\epsilon)^2}{\epsilon^2}. \end{cases}$$

Suppose we can write

$$u^\epsilon = u_0 + \epsilon u_1 + \epsilon u_2^\epsilon, \quad v^\epsilon = v_0 + \epsilon v_1 + \epsilon v_2^\epsilon,$$

where u_0, v_0, u_1, v_1 are smooth, $u_0, v_0 > 0$, and the functions $u_2^\epsilon, v_2^\epsilon$ are bounded, along with their derivatives, uniformly in ϵ .

Show that $u_0 \equiv v_0 =: w$ and w solves the nonlinear heat equation

$$w_t - \frac{1}{4}(\log w)_{xx} = 0.$$

(T. Kurtz, Trans. AMS 186 (1973), 259–272)

15. Justify in the proof of Lemma 2 in §4.5.3 the transformation of the integral of e^{-z^2} over the line Γ to the integral over the real axis.
16. Utilize Lemma 2 in §4.5.3 to discuss the sense in which u defined by formula (20) in §4.3.1 converges to the initial data g as $t \rightarrow 0^+$.
17. Let $n = 1$ and suppose that u^ϵ solves the problem

$$\begin{cases} -(a(\frac{x}{\epsilon})u_x^\epsilon)_x = f & \text{in } (0, 1) \\ u^\epsilon(0) = u^\epsilon(1) = 0, \end{cases}$$

where a is a smooth, positive function that is 1-periodic. Assume also that $f \in L^2(0, 1)$.

(a) Show that $u^\epsilon \rightharpoonup u$ weakly in $H_0^1(0, 1)$, where u solves

$$\begin{cases} -\bar{a}u_{xx} = f & \text{in } (0, 1) \\ u(0) = u(1) = 0, \end{cases}$$

for $\bar{a} := (\int_0^1 a(y)^{-1} dy)^{-1}$.

(b) Check that this answer agrees with the conclusions (73), (74) in §4.5.4.

(This problem requires knowledge of energy estimates, Sobolev spaces, etc., from Chapters 5, 6.)

18. (Variational principles in homogenization) Let $A(y) = ((a^{ij}(y)))$ be symmetric, positive definite and Q -periodic. Recall from §4.5.4 the expression (74) for the corresponding homogenized coefficients $\bar{A} = ((\bar{a}^{ij}))$.

- (a) Derive for each $\xi \in \mathbb{R}^n$ the variational formula

$$\xi \cdot \bar{A}\xi = \min_w \left\{ \int_Q Dw \cdot A(y)Dw \, dy \mid w = y \cdot \xi + v, v \text{ } Q\text{-periodic} \right\}.$$

(Hint: The minimum is attained by $w = y \cdot \xi - \sum_{i=1}^n \xi_i \chi^i$, for the correctors χ^i introduced in §4.5.4.)

- (b) Derive also the dual variational formula

$$\eta \cdot \bar{A}^{-1}\eta = \min_{\sigma} \left\{ \int_Q \sigma \cdot A(y)^{-1}\sigma \, dy \mid \int_Q \sigma \, dy = \eta, \operatorname{div} \sigma = 0, \sigma \text{ } Q\text{-periodic} \right\}.$$

- (c) Show that therefore

$$\left(\int_Q A(y)^{-1} \, dy \right)^{-1} \leq \bar{A} \leq \int_Q A(y) \, dy.$$

(Remember from §A.1 that for symmetric matrices $R \geq S$ means $R - S$ is nonnegative definite.)

19. Show that the line $\{t = 0\}$ is characteristic for the heat equation $u_t = u_{xx}$. Show there does not exist an analytic solution u of the heat equation in $\mathbb{R} \times \mathbb{R}$, with $u = \frac{1}{1+x^2}$ on $\{t = 0\}$.

(Hint: Assume there is an analytic solution, compute its coefficients, and show that the resulting power series diverges except at $(0, 0)$. This example is due to Kovalevskaya.)

4.8. REFERENCES

- Section 4.1 See for instance Pinsky [P], Strauss [St2], Thoe–Zachmanoglou [T-Z] or Weinberger [We] for more on separation of variables. Grindrod [Gr] and Nishiura [N] discuss Turing instabilities.
- Section 4.2 Theory for nonlinear dispersive PDE may be found in Linares–Ponce [L-P] and Tao [To]. C. Jones provided the discussion of traveling waves for the bistable equation, and J.-L. Vazquez showed me the derivation of Barenblatt’s solution. P. Olver’s book [O] explains much more about symmetry methods for PDE.
- Section 4.3 Stein–Weiss [S-W], Stein [Se], Hörmander [H], Rauch [R] and Treves [T] provide much more information concerning

Fourier transform techniques. Y. Yu improved the proof of Theorem 2, and M. Weinstein helped me with Schrödinger's equation. The asymptotic equipartition of energy formula (28) is based upon Brodsky (Proc. AMS 18 (1967), 207–208). Example 5 is from Pinsky [P]. See Helgason [Hg] for a more careful study of the applicability of the Radon transform: consult also John [J1] and Lax [Lx3]. The solution of the wave equation in §4.3.2 is from Pinsky–Taylor (J. Fourier Analysis 3 (1997), 647–703.)

- Section 4.4 See Courant–Hilbert [C-H] for more on the hodograph and Legendre transforms.
- Section 4.5 J. Neu contributed §4.5.1. Section 4.5.3 is based upon some classroom lectures of J. Ralston, following Hörmander (Acta Math. 127 (1971), 79–183). The discussion of homogenization in §4.5.4 follows Bensoussan–Lions–Papanicolaou [B-L-P].
- Section 4.6 See Folland [F1, Chapter 1], John [J2, Chapter 3], DiBenedetto [DB1, Chapter 1].
- Section 4.7 Problems 9 and 10 are based upon examples in Zwillinger [Zw]. R. Kohn contributed Problem 18. Problem 19 is from Mikhailov [M].

Part II

**THEORY FOR
LINEAR PARTIAL
DIFFERENTIAL
EQUATIONS**

This page is intentionally left blank.

SOBOLEV SPACES

- 5.1 Hölder spaces
- 5.2 Sobolev spaces
- 5.3 Approximation
- 5.4 Extensions
- 5.5 Traces
- 5.6 Sobolev inequalities
- 5.7 Compactness
- 5.8 Additional topics
- 5.9 Other spaces of functions
- 5.10 Problems
- 5.11 References

This chapter mostly develops the theory of *Sobolev spaces*, which turn out often to be the proper settings in which to apply ideas of functional analysis to glean information concerning PDE. The following material is sometimes subtle and seemingly unmotivated, but ultimately will prove extremely useful.

Since we have in mind eventual applications to rather wide classes of partial differential equations, it is worth sketching out here our overall point of view. Our intention, broadly put, will be to take various specific PDE and to recast them abstractly as operators acting on appropriate linear spaces. We can symbolically write

$$A : X \rightarrow Y,$$

where the operator A encodes the structure of the partial differential equations, including possibly boundary conditions, etc., and X, Y are spaces of functions. The great advantage is that once our PDE problem has been suitably interpreted in this form, we can often employ the general and elegant principles of functional analysis (Appendix D) to study the solvability of various equations involving A . We will later see that the really hard work is not so much the invocation of functional analysis, but rather finding the “right” spaces X, Y and the “right” abstract operators A . Sobolev spaces are designed precisely to make all this work out properly, and so these are usually the proper choices for X, Y .

We will utilize Sobolev spaces for studying linear elliptic, parabolic and hyperbolic PDE in Chapters 6–7 and for studying nonlinear elliptic and parabolic equations in Chapters 8–9.

The reader may wish to look over some of the terminology for functional analysis in Appendix D before going further.

5.1. HÖLDER SPACES

Before turning to Sobolev spaces, we first discuss the simpler *Hölder spaces*.

Assume $U \subset \mathbb{R}^n$ is open and $0 < \gamma \leq 1$. We have previously considered the class of Lipschitz continuous functions $u : U \rightarrow \mathbb{R}$, which by definition satisfy the estimate

$$(1) \quad |u(x) - u(y)| \leq C|x - y| \quad (x, y \in U)$$

for some constant C . Now (1) of course implies u is continuous and more importantly provides a uniform modulus of continuity. It turns out to be useful to consider also functions u satisfying a variant of (1), namely

$$(2) \quad |u(x) - u(y)| \leq C|x - y|^\gamma \quad (x, y \in U)$$

for some $0 < \gamma \leq 1$ and a constant C . Such a function is said to be *Hölder continuous with exponent γ* .

DEFINITIONS. (i) If $u : U \rightarrow \mathbb{R}$ is bounded and continuous, we write

$$\|u\|_{C(\bar{U})} := \sup_{x \in U} |u(x)|.$$

(ii) The γ^{th} -Hölder seminorm of $u : U \rightarrow \mathbb{R}$ is

$$[u]_{C^{0,\gamma}(\bar{U})} := \sup_{\substack{x,y \in U \\ x \neq y}} \left\{ \frac{|u(x) - u(y)|}{|x - y|^\gamma} \right\},$$

and the γ^{th} -Hölder norm is

$$\|u\|_{C^{0,\gamma}(\bar{U})} := \|u\|_{C(\bar{U})} + [u]_{C^{0,\gamma}(\bar{U})}.$$

DEFINITION. *The Hölder space*

$$C^{k,\gamma}(\bar{U})$$

consists of all functions $u \in C^k(\bar{U})$ for which the norm

$$(3) \quad \|u\|_{C^{k,\gamma}(\bar{U})} := \sum_{|\alpha| \leq k} \|D^\alpha u\|_{C(\bar{U})} + \sum_{|\alpha|=k} [D^\alpha u]_{C^{0,\gamma}(\bar{U})}$$

is finite.

So the space $C^{k,\gamma}(\bar{U})$ consists of those functions u that are k -times continuously differentiable and whose k^{th} -partial derivatives are bounded and Hölder continuous with exponent γ . Such functions are well-behaved, and furthermore the space $C^{k,\gamma}(\bar{U})$ itself possesses a good mathematical structure:

THEOREM 1 (Hölder spaces as function spaces). *The space of functions $C^{k,\gamma}(\bar{U})$ is a Banach space.*

The proof is left as an exercise (Problem 1), but let us pause here to make clear what is being asserted. Recall from §D.1 that if X denotes a real linear space, then a mapping $\|\cdot\| : X \rightarrow [0, \infty)$ is called a *norm* provided

- (i) $\|u + v\| \leq \|u\| + \|v\|$ for all $u, v \in X$,
- (ii) $\|\lambda u\| = |\lambda| \|u\|$ for all $u \in X, \lambda \in \mathbb{R}$,
- (iii) $\|u\| = 0$ if and only if $u = 0$.

A norm provides us with a notion of convergence: we say a sequence $\{u_k\}_{k=1}^\infty \subset X$ converges to $u \in X$, written $u_k \rightarrow u$, if $\lim_{k \rightarrow \infty} \|u_k - u\| = 0$. A *Banach space* is then a normed linear space which is *complete*, that is, within which each Cauchy sequence converges.

So in Theorem 1 we are stating that if we take on the linear space $C^{k,\gamma}(\bar{U})$ the norm $\|\cdot\| = \|\cdot\|_{C^{k,\gamma}(\bar{U})}$, defined by (3), then $\|\cdot\|$ verifies properties (i)–(iii) above, and in addition each Cauchy sequence converges.

5.2. SOBOLEV SPACES

The Hölder spaces introduced in §5.1 are unfortunately not often suitable settings for elementary PDE theory, as we usually cannot make good enough analytic estimates to demonstrate that the solutions we construct actually belong to such spaces. What are needed rather are some other kinds of spaces, containing less smooth functions. In practice we must strike a balance, by designing spaces comprising functions which have some, but not too great, smoothness properties.

5.2.1. Weak derivatives.

We start off by substantially weakening the notion of partial derivatives.

NOTATION. Let $C_c^\infty(U)$ denote the space of infinitely differentiable functions $\phi : U \rightarrow \mathbb{R}$, with compact support in U . We will sometimes call a function ϕ belonging to $C_c^\infty(U)$ a *test function*. \square

Motivation for definition of weak derivative. Assume we are given a function $u \in C^1(U)$. Then if $\phi \in C_c^\infty(U)$, we see from the integration by parts formula that

$$(1) \quad \int_U u \phi_{x_i} dx = - \int_U u_{x_i} \phi dx \quad (i = 1, \dots, n).$$

There are no boundary terms, since ϕ has compact support in U and thus vanishes near ∂U . More generally now, if k is a positive integer, $u \in C^k(U)$, and $\alpha = (\alpha_1, \dots, \alpha_n)$ is a multiindex of order $|\alpha| = \alpha_1 + \dots + \alpha_n = k$, then

$$(2) \quad \int_U u D^\alpha \phi dx = (-1)^{|\alpha|} \int_U D^\alpha u \phi dx.$$

This equality holds since

$$D^\alpha \phi = \frac{\partial^{\alpha_1}}{\partial x_1^{\alpha_1}} \cdots \frac{\partial^{\alpha_n}}{\partial x_n^{\alpha_n}} \phi$$

and we can apply formula (1) $|\alpha|$ times.

We next examine formula (2), valid for $u \in C^k(U)$, and ask whether some variant of it might still be true even if u is not k times continuously differentiable. Now the left-hand side of (2) makes sense if u is only locally summable: the problem is rather that if u is not C^k , then the expression “ $D^\alpha u$ ” on the right-hand side of (2) has no obvious meaning. We resolve this difficulty by asking if there exists a locally summable function v for which formula (2) is valid, with v replacing $D^\alpha u$:

DEFINITION. Suppose $u, v \in L_{\text{loc}}^1(U)$ and α is a multiindex. We say that v is the α^{th} -weak partial derivative of u , written

$$D^\alpha u = v,$$

provided

$$(3) \quad \int_U u D^\alpha \phi dx = (-1)^{|\alpha|} \int_U v \phi dx$$

for all test functions $\phi \in C_c^\infty(U)$.

In other words, if we are given u and if there happens to exist a function v which verifies (3) for all ϕ , we say that $D^\alpha u = v$ in the weak sense. If there does not exist such a function v , then u does not possess a weak α^{th} -partial derivative.

LEMMA (Uniqueness of weak derivatives). *A weak α^{th} -partial derivative of u , if it exists, is uniquely defined up to a set of measure zero.*

Proof. Assume that $v, \tilde{v} \in L^1_{\text{loc}}(U)$ satisfy

$$\int_U u D^\alpha \phi \, dx = (-1)^{|\alpha|} \int_U v \phi \, dx = (-1)^{|\alpha|} \int_U \tilde{v} \phi \, dx$$

for all $\phi \in C_c^\infty(U)$. Then

$$(4) \quad \int_U (v - \tilde{v}) \phi \, dx = 0$$

for all $\phi \in C_c^\infty(U)$, whence $v - \tilde{v} = 0$ a.e. □

Example 1. Let $n = 1$, $U = (0, 2)$, and

$$u(x) = \begin{cases} x & \text{if } 0 < x \leq 1 \\ 1 & \text{if } 1 \leq x < 2. \end{cases}$$

Define

$$v(x) = \begin{cases} 1 & \text{if } 0 < x \leq 1 \\ 0 & \text{if } 1 < x < 2. \end{cases}$$

Let us show $u' = v$ in the weak sense. To see this, choose any $\phi \in C_c^\infty(U)$. We must demonstrate

$$\int_0^2 u \phi' \, dx = - \int_0^2 v \phi \, dx.$$

But we easily calculate

$$\begin{aligned} \int_0^2 u \phi' \, dx &= \int_0^1 x \phi' \, dx + \int_1^2 \phi' \, dx \\ &= - \int_0^1 \phi \, dx + \phi(1) - \phi(1) = - \int_0^2 v \phi \, dx, \end{aligned}$$

as required. □

Example 2. Let $n = 1$, $U = (0, 2)$, and

$$u(x) = \begin{cases} x & \text{if } 0 < x \leq 1 \\ 2 & \text{if } 1 < x < 2. \end{cases}$$

We assert u' does not exist in the weak sense. To check this, we must show there does not exist any function $v \in L^1_{\text{loc}}(U)$ satisfying

$$(5) \quad \int_0^2 u\phi' dx = - \int_0^2 v\phi dx$$

for all $\phi \in C_c^\infty(U)$. Suppose, to the contrary, (5) were valid for some v and all ϕ . Then

$$(6) \quad \begin{aligned} - \int_0^2 v\phi dx &= \int_0^2 u\phi' dx = \int_0^1 x\phi' dx + 2 \int_1^2 \phi' dx \\ &= - \int_0^1 \phi dx - \phi(1). \end{aligned}$$

Choose a sequence $\{\phi_m\}_{m=1}^\infty$ of smooth functions satisfying

$$0 \leq \phi_m \leq 1, \quad \phi_m(1) = 1, \quad \phi_m(x) \rightarrow 0 \text{ for all } x \neq 1.$$

Replacing ϕ by ϕ_m in (6) and sending $m \rightarrow \infty$, we discover

$$1 = \lim_{m \rightarrow \infty} \phi_m(1) = \lim_{m \rightarrow \infty} \left[\int_0^2 v\phi_m dx - \int_0^1 \phi_m dx \right] = 0,$$

a contradiction. □

More sophisticated examples appear in the next subsection.

5.2.2. Definition of Sobolev spaces.

Fix $1 \leq p \leq \infty$ and let k be a nonnegative integer. We define now certain function spaces, whose members have weak derivatives of various orders lying in various L^p spaces.

DEFINITION. *The Sobolev space*

$$W^{k,p}(U)$$

consists of all locally summable functions $u : U \rightarrow \mathbb{R}$ such that for each multiindex α with $|\alpha| \leq k$, $D^\alpha u$ exists in the weak sense and belongs to $L^p(U)$.

Remarks. (i) If $p = 2$, we usually write

$$H^k(U) = W^{k,2}(U) \quad (k = 0, 1, \dots).$$

The letter H is used, since—as we will see— $H^k(U)$ is a Hilbert space. Note that $H^0(U) = L^2(U)$.

(ii) We henceforth identify functions in $W^{k,p}(U)$ which agree a.e.

DEFINITION. If $u \in W^{k,p}(U)$, we define its norm to be

$$\|u\|_{W^{k,p}(U)} := \begin{cases} \left(\sum_{|\alpha| \leq k} \int_U |D^\alpha u|^p dx \right)^{1/p} & (1 \leq p < \infty) \\ \sum_{|\alpha| \leq k} \text{ess sup}_U |D^\alpha u| & (p = \infty). \end{cases}$$

DEFINITIONS. (i) Let $\{u_m\}_{m=1}^\infty$, $u \in W^{k,p}(U)$. We say u_m converges to u in $W^{k,p}(U)$, written

$$u_m \rightarrow u \quad \text{in } W^{k,p}(U),$$

provided

$$\lim_{m \rightarrow \infty} \|u_m - u\|_{W^{k,p}(U)} = 0.$$

(ii) We write

$$u_m \rightarrow u \quad \text{in } W_{\text{loc}}^{k,p}(U)$$

to mean

$$u_m \rightarrow u \quad \text{in } W^{k,p}(V)$$

for each $V \subset\subset U$.

DEFINITION. We denote by

$$W_0^{k,p}(U)$$

the closure of $C_c^\infty(U)$ in $W^{k,p}(U)$.

Thus $u \in W_0^{k,p}(U)$ if and only if there exist functions $u_m \in C_c^\infty(U)$ such that $u_m \rightarrow u$ in $W^{k,p}(U)$. We interpret $W_0^{k,p}(U)$ as comprising those functions $u \in W^{k,p}(U)$ such that

$$"D^\alpha u = 0 \text{ on } \partial U" \text{ for all } |\alpha| \leq k - 1.$$

This will all be made clearer with the discussion of traces in §5.5.

NOTATION. It is customary to write

$$H_0^k(U) = W_0^{k,2}(U).$$

We will see in the exercises that if $n = 1$ and U is an open interval in \mathbb{R}^1 , then $u \in W^{1,p}(U)$ if and only if u equals a.e. an absolutely continuous function whose ordinary derivative (which exists a.e.) belongs to $L^p(U)$. Such a simple characterization is however only available for $n = 1$. In general a function can belong to a Sobolev space and yet be discontinuous and/or unbounded.

Example 3. Take $U = B^0(0, 1)$, the open unit ball in \mathbb{R}^n , and

$$u(x) = |x|^{-\alpha} \quad (x \in U, x \neq 0).$$

For which values of $\alpha > 0, n, p$ does u belong to $W^{1,p}(U)$? To answer, note first that u is smooth away from 0, with

$$u_{x_i}(x) = \frac{-\alpha x_i}{|x|^{\alpha+2}} \quad (x \neq 0),$$

and so

$$|Du(x)| = \frac{|\alpha|}{|x|^{\alpha+1}} \quad (x \neq 0).$$

Let $\phi \in C_c^\infty(U)$ and fix $\varepsilon > 0$. Then

$$\int_{U-B(0,\varepsilon)} u \phi_{x_i} dx = - \int_{U-B(0,\varepsilon)} u_{x_i} \phi dx + \int_{\partial B(0,\varepsilon)} u \phi \nu^i dS,$$

$\nu = (\nu^1, \dots, \nu^n)$ denoting the inward pointing normal on $\partial B(0, \varepsilon)$. Now if $\alpha + 1 < n$, $|Du(x)| \in L^1(U)$. In this case

$$\left| \int_{\partial B(0,\varepsilon)} u \phi \nu^i dS \right| \leq \|\phi\|_{L^\infty} \int_{\partial B(0,\varepsilon)} \varepsilon^{-\alpha} dS \leq C \varepsilon^{n-1-\alpha} \rightarrow 0.$$

Thus

$$\int_U u \phi_{x_i} dx = - \int_U u_{x_i} \phi dx$$

for all $\phi \in C_c^\infty(U)$, provided $0 \leq \alpha < n - 1$. Furthermore $|Du(x)| = \frac{|\alpha|}{|x|^{\alpha+1}} \in L^p(U)$ if and only if $(\alpha + 1)p < n$. Consequently $u \in W^{1,p}(U)$ if and only if $\alpha < \frac{n-p}{p}$. In particular $u \notin W^{1,p}(U)$ for each $p \geq n$. \square

Example 4. Let $\{r_k\}_{k=1}^\infty$ be a countable, dense subset of $U = B^0(0, 1)$. Write

$$u(x) = \sum_{k=1}^{\infty} \frac{1}{2^k} |x - r_k|^{-\alpha} \quad (x \in U).$$

Then $u \in W^{1,p}(U)$ for $\alpha < \frac{n-p}{p}$. If $0 < \alpha < \frac{n-p}{p}$, we see that u belongs to $W^{1,p}(U)$ and yet is unbounded on each open subset of U . \square

This last example illustrates a fundamental fact of life, that although a function u belonging to a Sobolev space possesses certain smoothness properties, it can still be rather badly behaved in other ways.

5.2.3. Elementary properties.

Next we verify certain properties of weak derivatives. Note very carefully that whereas these various rules are obviously true for smooth functions, functions in Sobolev space are not necessarily smooth: we must always rely solely upon the definition of weak derivatives.

THEOREM 1 (Properties of weak derivatives). *Assume $u, v \in W^{k,p}(U)$, $|\alpha| \leq k$. Then*

- (i) $D^\alpha u \in W^{k-|\alpha|,p}(U)$ and $D^\beta(D^\alpha u) = D^\alpha(D^\beta u) = D^{\alpha+\beta}u$ for all multiindices α, β with $|\alpha| + |\beta| \leq k$.
- (ii) For each $\lambda, \mu \in \mathbb{R}$, $\lambda u + \mu v \in W^{k,p}(U)$ and $D^\alpha(\lambda u + \mu v) = \lambda D^\alpha u + \mu D^\alpha v$, $|\alpha| \leq k$.
- (iii) If V is an open subset of U , then $u \in W^{k,p}(V)$.
- (iv) If $\zeta \in C_c^\infty(U)$, then $\zeta u \in W^{k,p}(U)$ and

$$(7) \quad D^\alpha(\zeta u) = \sum_{\beta \leq \alpha} \binom{\alpha}{\beta} D^\beta \zeta D^{\alpha-\beta} u \quad (\text{Leibniz's formula}),$$

$$\text{where } \binom{\alpha}{\beta} = \frac{\alpha!}{\beta!(\alpha-\beta)!}.$$

Proof. 1. To prove (i), first fix $\phi \in C_c^\infty(U)$. Then $D^\beta \phi \in C_c^\infty(U)$, and so

$$\begin{aligned} \int_U D^\alpha u D^\beta \phi \, dx &= (-1)^{|\alpha|} \int_U u D^{\alpha+\beta} \phi \, dx \\ &= (-1)^{|\alpha|} (-1)^{|\alpha+\beta|} \int_U D^{\alpha+\beta} u \phi \, dx \\ &= (-1)^{|\beta|} \int_U D^{\alpha+\beta} u \phi \, dx. \end{aligned}$$

Thus $D^\beta(D^\alpha u) = D^{\alpha+\beta}u$ in the weak sense.

2. Assertions (ii) and (iii) are easy, and the proofs are omitted.

3. We prove (7) by induction on $|\alpha|$. Suppose first $|\alpha| = 1$. Choose any $\phi \in C_c^\infty(U)$. Then

$$\begin{aligned} \int_U \zeta u D^\alpha \phi \, dx &= \int_U u D^\alpha(\zeta \phi) - u(D^\alpha \zeta) \phi \, dx \\ &= - \int_U (\zeta D^\alpha u + u D^\alpha \zeta) \phi \, dx. \end{aligned}$$

Thus $D^\alpha(\zeta u) = \zeta D^\alpha u + u D^\alpha \zeta$, as required.

Next assume $l < k$ and formula (7) is valid for all $|\alpha| \leq l$ and all functions ζ . Choose a multiindex α with $|\alpha| = l + 1$. Then $\alpha = \beta + \gamma$ for some $|\beta| = l$, $|\gamma| = 1$. Then for ϕ as above,

$$\begin{aligned} \int_U \zeta u D^\alpha \phi \, dx &= \int_U \zeta u D^\beta (D^\gamma \phi) \, dx \\ &= (-1)^{|\beta|} \int_U \sum_{\sigma \leq \beta} \binom{\beta}{\sigma} D^\sigma \zeta D^{\beta-\sigma} u D^\gamma \phi \, dx \end{aligned}$$

(by the induction assumption)

$$= (-1)^{|\beta|+|\gamma|} \int_U \sum_{\sigma \leq \beta} \binom{\beta}{\sigma} D^\gamma (D^\sigma \zeta D^{\beta-\sigma} u) \phi \, dx$$

(by the induction assumption again)

$$= (-1)^{|\alpha|} \int_U \sum_{\sigma \leq \beta} \binom{\beta}{\sigma} [D^\rho \zeta D^{\alpha-\rho} u + D^\sigma \zeta D^{\alpha-\sigma} u] \phi \, dx$$

(where $\rho = \sigma + \gamma$)

$$= (-1)^{|\alpha|} \int_U \left[\sum_{\sigma \leq \alpha} \binom{\alpha}{\sigma} D^\sigma \zeta D^{\alpha-\sigma} u \right] \phi \, dx,$$

since

$$\binom{\beta}{\sigma - \gamma} + \binom{\beta}{\sigma} = \binom{\alpha}{\sigma}. \quad \square$$

Not only do many of the usual rules of calculus apply to weak derivatives, but the Sobolev spaces themselves have a good mathematical structure:

THEOREM 2 (Sobolev spaces as function spaces). *For each $k = 1, \dots$ and $1 \leq p \leq \infty$, the Sobolev space $W^{k,p}(U)$ is a Banach space.*

Proof. 1. Let us first of all check that $\|u\|_{W^{k,p}(U)}$ is a norm. (See the discussion at the end of §5.1, or refer to §D.1, for definitions.) Clearly

$$\|\lambda u\|_{W^{k,p}(U)} = |\lambda| \|u\|_{W^{k,p}(U)},$$

and

$$\|u\|_{W^{k,p}(U)} = 0 \text{ if and only if } u = 0 \text{ a.e.}$$

Next assume $u, v \in W^{k,p}(U)$. Then if $1 \leq p < \infty$, Minkowski's inequality (§B.2) implies

$$\begin{aligned} \|u + v\|_{W^{k,p}(U)} &= \left(\sum_{|\alpha| \leq k} \|D^\alpha u + D^\alpha v\|_{L^p(U)}^p \right)^{1/p} \\ &\leq \left(\sum_{|\alpha| \leq k} (\|D^\alpha u\|_{L^p(U)} + \|D^\alpha v\|_{L^p(U)})^p \right)^{1/p} \\ &\leq \left(\sum_{|\alpha| \leq k} \|D^\alpha u\|_{L^p(U)}^p \right)^{1/p} + \left(\sum_{|\alpha| \leq k} \|D^\alpha v\|_{L^p(U)}^p \right)^{1/p} \\ &= \|u\|_{W^{k,p}(U)} + \|v\|_{W^{k,p}(U)}. \end{aligned}$$

2. It remains to show that $W^{k,p}(U)$ is complete. So assume $\{u_m\}_{m=1}^\infty$ is a Cauchy sequence in $W^{k,p}(U)$. Then for each $|\alpha| \leq k$, $\{D^\alpha u_m\}_{m=1}^\infty$ is a Cauchy sequence in $L^p(U)$. Since $L^p(U)$ is complete, there exist functions $u_\alpha \in L^p(U)$ such that

$$D^\alpha u_m \rightarrow u_\alpha \quad \text{in } L^p(U)$$

for each $|\alpha| \leq k$. In particular,

$$u_m \rightarrow u_{(0,\dots,0)} =: u \quad \text{in } L^p(U).$$

3. We now claim

$$(8) \quad u \in W^{k,p}(U), \quad D^\alpha u = u_\alpha \quad (|\alpha| \leq k).$$

To verify this assertion, fix $\phi \in C_c^\infty(U)$. Then

$$\begin{aligned} \int_U u D^\alpha \phi \, dx &= \lim_{m \rightarrow \infty} \int_U u_m D^\alpha \phi \, dx \\ &= \lim_{m \rightarrow \infty} (-1)^{|\alpha|} \int_U D^\alpha u_m \phi \, dx \\ &= (-1)^{|\alpha|} \int_U u_\alpha \phi \, dx. \end{aligned}$$

Thus (8) is valid. Since therefore $D^\alpha u_m \rightarrow D^\alpha u$ in $L^p(U)$ for all $|\alpha| \leq k$, we see that $u_m \rightarrow u$ in $W^{k,p}(U)$, as required. \square

5.3. APPROXIMATION

5.3.1. Interior approximation by smooth functions.

It is awkward to return continually to the definition of weak derivatives. In order to study the deeper properties of Sobolev spaces, we therefore need to develop some systematic procedures for approximating a function in a Sobolev space by smooth functions. The method of mollifiers, set forth in §C.4, provides the tool.

Fix a positive integer k and $1 \leq p < \infty$. Remember that $U_\varepsilon = \{x \in U \mid \text{dist}(x, \partial U) > \varepsilon\}$.

THEOREM 1 (Local approximation by smooth functions). *Assume $u \in W^{k,p}(U)$ for some $1 \leq p < \infty$, and set*

$$u^\varepsilon = \eta_\varepsilon * u \quad \text{in } U_\varepsilon.$$

Then

$$(i) \quad u^\varepsilon \in C^\infty(U_\varepsilon) \quad \text{for each } \varepsilon > 0,$$

and

$$(ii) \quad u^\varepsilon \rightarrow u \quad \text{in } W_{\text{loc}}^{k,p}(U), \text{ as } \varepsilon \rightarrow 0.$$

Proof. 1. Assertion (i) is proved in §C.4.

2. We next claim that if $|\alpha| \leq k$, then

$$(1) \quad D^\alpha u^\varepsilon = \eta_\varepsilon * D^\alpha u \quad \text{in } U_\varepsilon;$$

that is, the ordinary α^{th} -partial derivative of the smooth function u^ε is the ε -mollification of the α^{th} -weak partial derivative of u . To confirm this, we compute for $x \in U_\varepsilon$

$$\begin{aligned} D^\alpha u^\varepsilon(x) &= D^\alpha \int_U \eta_\varepsilon(x-y)u(y) dy \\ &= \int_U D_x^\alpha \eta_\varepsilon(x-y)u(y) dy \\ &= (-1)^{|\alpha|} \int_U D_y^\alpha \eta_\varepsilon(x-y)u(y) dy. \end{aligned}$$

Now for fixed $x \in U_\varepsilon$ the function $\phi(y) := \eta_\varepsilon(x-y)$ belongs to $C_c^\infty(U)$. Consequently the definition of the α^{th} -weak partial derivative implies:

$$\int_U D_y^\alpha \eta_\varepsilon(x-y)u(y) dy = (-1)^{|\alpha|} \int_U \eta_\varepsilon(x-y)D^\alpha u(y) dy.$$

Thus

$$\begin{aligned} D^\alpha u^\varepsilon(x) &= (-1)^{|\alpha|+|\alpha|} \int_U \eta_\varepsilon(x-y) D^\alpha u(y) dy \\ &= [\eta_\varepsilon * D^\alpha u](x). \end{aligned}$$

This establishes (1).

3. Now choose an open set $V \subset\subset U$. In view of (1) and §C.4, $D^\alpha u^\varepsilon \rightarrow D^\alpha u$ in $L^p(V)$ as $\varepsilon \rightarrow 0$, for each $|\alpha| \leq k$. Consequently

$$\|u^\varepsilon - u\|_{W^{k,p}(V)}^p = \sum_{|\alpha| \leq k} \|D^\alpha u^\varepsilon - D^\alpha u\|_{L^p(V)}^p \rightarrow 0$$

as $\varepsilon \rightarrow 0$. This proves assertion (ii). \square

5.3.2. Approximation by smooth functions.

Next we show that we can find smooth functions which approximate in $W^{k,p}(U)$ and not just in $W_{\text{loc}}^{k,p}(U)$. Notice in the following that we make no assumptions about the smoothness of ∂U .

THEOREM 2 (Global approximation by smooth functions). *Assume U is bounded, and suppose as well that $u \in W^{k,p}(U)$ for some $1 \leq p < \infty$. Then there exist functions $u_m \in C^\infty(U) \cap W^{k,p}(U)$ such that*

$$u_m \rightarrow u \quad \text{in } W^{k,p}(U).$$

Note carefully that we do not assert $u_m \in C^\infty(\bar{U})$ (but see Theorem 3 below).

Proof. 1. We have $U = \bigcup_{i=1}^\infty U_i$, where

$$U_i := \{x \in U \mid \text{dist}(x, \partial U) > 1/i\} \quad (i = 1, 2, \dots).$$

Write $V_i := U_{i+3} - \bar{U}_{i+1}$.

Choose also any open set $V_0 \subset\subset U$ so that $U = \bigcup_{i=0}^\infty V_i$. Now let $\{\zeta_i\}_{i=0}^\infty$ be a smooth partition of unity subordinate to the open sets $\{V_i\}_{i=0}^\infty$; that is, suppose

$$(2) \quad \begin{cases} 0 \leq \zeta_i \leq 1, & \zeta_i \in C_c^\infty(V_i) \\ \sum_{i=0}^\infty \zeta_i = 1 & \text{on } U. \end{cases}$$

Next, choose any function $u \in W^{k,p}(U)$. According to Theorem 1(iv) in §5.2, $\zeta_i u \in W^{k,p}(U)$ and $\text{spt}(\zeta_i u) \subset V_i$.

2. Fix $\delta > 0$. Choose then $\varepsilon_i > 0$ so small that $u^i := \eta_{\varepsilon_i} * (\zeta_i u)$ satisfies

$$(3) \quad \begin{cases} \|u^i - \zeta_i u\|_{W^{k,p}(U)} \leq \frac{\delta}{2^{i+1}} & (i = 0, 1, \dots) \\ \text{spt } u^i \subset W_i & (i = 1, \dots), \end{cases}$$

for $W_i := U_{i+4} - \bar{U}_i \supset V_i$ ($i = 1, \dots$).

3. Write $v := \sum_{i=0}^{\infty} u^i$. This function belongs to $C^\infty(U)$, since for each open set $V \subset\subset U$ there are at most finitely many nonzero terms in the sum. Since $u = \sum_{i=0}^{\infty} \zeta_i u$, we have for each $V \subset\subset U$

$$\begin{aligned} \|v - u\|_{W^{k,p}(V)} &\leq \sum_{i=0}^{\infty} \|u^i - \zeta_i u\|_{W^{k,p}(U)} \\ &\leq \delta \sum_{i=0}^{\infty} \frac{1}{2^{i+1}} \quad \text{by (3)} \\ &= \delta. \end{aligned}$$

Take the supremum over sets $V \subset\subset U$, to conclude $\|v - u\|_{W^{k,p}(U)} \leq \delta$. \square

5.3.3. Global approximation by smooth functions.

We now ask when it is possible to approximate a function $u \in W^{k,p}(U)$ by functions belonging to $C^\infty(\bar{U})$, rather than only $C^\infty(U)$. Such an approximation requires some condition to exclude ∂U being wild geometrically.

THEOREM 3 (Global approximation by functions smooth up to the boundary). *Assume U is bounded and ∂U is C^1 . Suppose $u \in W^{k,p}(U)$ for some $1 \leq p < \infty$. Then there exist functions $u_m \in C^\infty(\bar{U})$ such that*

$$u_m \rightarrow u \quad \text{in } W^{k,p}(U).$$


Proof. 1. Fix any point $x^0 \in \partial U$. As ∂U is C^1 , there exist, according to §C.1, a radius $r > 0$ and a C^1 function $\gamma : \mathbb{R}^{n-1} \rightarrow \mathbb{R}$ such that—upon relabeling the coordinate axes if necessary—we have

$$U \cap B(x^0, r) = \{x \in B(x^0, r) \mid x_n > \gamma(x_1, \dots, x_{n-1})\}.$$

Set $V := U \cap B(x^0, r/2)$.

2. Define the shifted point

$$x^\varepsilon := x + \lambda \varepsilon e_n \quad (x \in V, \varepsilon > 0),$$



and observe that for some fixed, sufficiently large number $\lambda > 0$ the ball $B(x^\epsilon, \epsilon)$ lies in $U \cap B(x^0, r)$ for all $x \in V$ and all small $\epsilon > 0$.

Now define $u_\epsilon(x) := u(x^\epsilon)$ ($x \in V$); this is the function u translated a distance $\lambda\epsilon$ in the e_n direction. Next write $v^\epsilon = \eta_\epsilon * u_\epsilon$. The idea is that we have moved up enough so that “there is room to mollify within U ”. Clearly $v^\epsilon \in C^\infty(\bar{V})$.

3. We now claim

$$(4) \quad v^\epsilon \rightarrow u \quad \text{in } W^{k,p}(V).$$

To confirm this, take α to be any multiindex with $|\alpha| \leq k$. Then

$$\|D^\alpha v^\epsilon - D^\alpha u\|_{L^p(V)} \leq \|D^\alpha v^\epsilon - D^\alpha u_\epsilon\|_{L^p(V)} + \|D^\alpha u_\epsilon - D^\alpha u\|_{L^p(V)}.$$

The second term on the right-hand side goes to zero with ϵ , since translation is continuous in the L^p -norm; and the first term also vanishes in the limit, by reasoning similar to that in the proof of Theorem 1.

4. Select $\delta > 0$. Since ∂U is compact, we can find finitely many points $x_i^0 \in \partial U$, radii $r_i > 0$, corresponding sets $V_i = U \cap B^0(x_i^0, \frac{r_i}{2})$, and functions $v_i \in C^\infty(\bar{V}_i)$ ($i = 1, \dots, N$) such that $\partial U \subset \bigcup_{i=1}^N B^0(x_i^0, \frac{r_i}{2})$ and

$$(5) \quad \|v_i - u\|_{W^{k,p}(V_i)} \leq \delta.$$

Take an open set $V_0 \subset\subset U$ such that $U \subset \bigcup_{i=0}^N V_i$ and select, using Theorem 1, a function $v_0 \in C^\infty(\bar{V}_0)$ satisfying

$$(6) \quad \|v_0 - u\|_{W^{k,p}(V_0)} \leq \delta.$$

5. Now let $\{\zeta_i\}_{i=0}^N$ be a smooth partition of unity on \bar{U} , subordinate to the open sets $\{V_0, B^0(x_1^0, \frac{r_1}{2}), \dots, B^0(x_N^0, \frac{r_N}{2})\}$. Define $v := \sum_{i=0}^N \zeta_i v_i$.

Then clearly $v \in C^\infty(\bar{U})$. In addition, since $u = \sum_{i=0}^N \zeta_i u$, we see using Theorem 1 in §5.2.3 that for each $|\alpha| \leq k$

$$\begin{aligned} \|D^\alpha v - D^\alpha u\|_{L^p(U)} &\leq \sum_{i=0}^N \|D^\alpha(\zeta_i v_i) - D^\alpha(\zeta_i u)\|_{L^p(V_i)} \\ &\leq C \sum_{i=0}^N \|v_i - u\|_{W^{k,p}(V_i)} = C(N+1)\delta, \end{aligned}$$

according to (5) and (6). □

5.4. EXTENSIONS

Our goal next is to extend functions in the Sobolev space $W^{1,p}(U)$ to become functions in the Sobolev space $W^{1,p}(\mathbb{R}^n)$. This can be subtle. Observe for instance that our extending $u \in W^{1,p}(U)$ to be zero in $\mathbb{R}^n - U$ will not in general work, as we may thereby create such a bad discontinuity along ∂U that the extended function no longer has a weak first partial derivative. We must instead invent a way to extend u which “preserves the weak derivatives across ∂U ”.

Suppose $1 \leq p \leq \infty$.

THEOREM 1 (Extension Theorem). *Assume U is bounded and ∂U is C^1 . Select a bounded open set V such that $U \subset\subset V$. Then there exists a bounded linear operator*

$$(1) \quad E : W^{1,p}(U) \rightarrow W^{1,p}(\mathbb{R}^n)$$

such that for each $u \in W^{1,p}(U)$:

- (i) $Eu = u$ a.e. in U ,
- (ii) Eu has support within V ,

and

- (iii)


$$\|Eu\|_{W^{1,p}(\mathbb{R}^n)} \leq C\|u\|_{W^{1,p}(U)},$$

the constant C depending only on p , U , and V .

DEFINITION. We call Eu an extension of u to \mathbb{R}^n .

Proof. 1. Fix $x^0 \in \partial U$ and suppose first

$$(2) \quad \partial U \text{ is flat near } x^0, \text{ lying in the plane } \{x_n = 0\}.$$



A half-ball at the boundary

Then we may assume there exists an open ball B , with center x^0 and radius r , such that

$$\begin{cases} B^+ := B \cap \{x_n \geq 0\} \subset \bar{U} \\ B^- := B \cap \{x_n \leq 0\} \subset \mathbb{R}^n - U. \end{cases}$$

2. Temporarily suppose also $u \in C^1(\bar{U})$. We define then

$$(3) \quad \bar{u}(x) := \begin{cases} u(x) & \text{if } x \in B^+ \\ -3u(x_1, \dots, x_{n-1}, -x_n) + 4u(x_1, \dots, x_{n-1}, -\frac{x_n}{2}) & \text{if } x \in B^-. \end{cases}$$

This is called a *higher-order reflection* of u from B^+ to B^- .

3. We claim

$$(4) \quad \bar{u} \in C^1(B).$$

To check this, let us write $u^- := \bar{u}|_{B^-}$, $u^+ := \bar{u}|_{B^+}$. We demonstrate first

$$(5) \quad u_{x_n}^- = u_{x_n}^+ \quad \text{on } \{x_n = 0\}.$$

Indeed according to (3),

$$u_{x_n}^-(x) = 3u_{x_n}(x_1, \dots, x_{n-1}, -x_n) - 2u_{x_n}(x_1, \dots, x_{n-1}, -\frac{x_n}{2})$$

and so


$$u_{x_n}^-|_{\{x_n=0\}} = u_{x_n}^+|_{\{x_n=0\}}.$$

This confirms (5). Now since $u^+ = u^-$ on $\{x_n = 0\}$, we see as well that

$$(6) \quad u_{x_i}^-|_{\{x_n=0\}} = u_{x_i}^+|_{\{x_n=0\}}$$

for $i = 1, \dots, n-1$. But then (5) and (6) together imply

$$D^\alpha u^-|_{\{x_n=0\}} = D^\alpha u^+|_{\{x_n=0\}}$$



Straightening out the boundary

for each $|\alpha| \leq 1$, and so (4) follows.

4. Using these calculations, we readily check as well

$$(7) \quad \|\bar{u}\|_{W^{1,p}(B)} \leq C\|u\|_{W^{1,p}(B^+)}$$

for some constant C which does not depend on u .

5. Let us next consider the situation that ∂U is not necessarily flat near x^0 . Then utilizing the notation and terminology from §C.1, we can find a C^1 mapping Φ , with inverse Ψ , such that Φ “straightens out ∂U near x^0 ”.

We write $y = \Phi(x)$, $x = \Psi(y)$, $u'(y) := u(\Psi(y))$. Choose a small ball B as drawn before. Then utilizing steps 1–3 above, we extend u' from B^+ to a function \bar{u}' defined on all of B , such that \bar{u}' is C^1 and we have the estimate

$$\|\bar{u}'\|_{W^{1,p}(B)} \leq C\|u'\|_{W^{1,p}(B^+)}.$$

Let $W := \Psi(B)$. Then converting back to the x -variables, we obtain an extension \bar{u} of u to W , with

$$(8) \quad \|\bar{u}\|_{W^{1,p}(W)} \leq C\|u\|_{W^{1,p}(U)}.$$

6. Since ∂U is compact, there exist finitely many points $x_i^0 \in \partial U$, open sets W_i , and extensions \bar{u}_i of u to W_i ($i = 1, \dots, N$), as above, such that $\partial U \subset \bigcup_{i=1}^N W_i$. Take $W_0 \subset\subset U$ so that $U \subset \bigcup_{i=0}^N W_i$, and let $\{\zeta_i\}_{i=0}^N$ be an associated partition of unity. Write $\bar{u} := \sum_{i=0}^N \zeta_i \bar{u}_i$, where $\bar{u}_0 = u$. Then utilizing estimate (8) (with u_i in place of u , \bar{u}_i in place of \bar{u}), we obtain the bound

$$(9) \quad \|\bar{u}\|_{W^{1,p}(\mathbb{R}^n)} \leq C\|u\|_{W^{1,p}(U)}$$

for some constant C , depending on U, p, n , etc., but not on u . Furthermore we can arrange for the support of \bar{u} to lie within $V \supset\supset U$.

7. We henceforth write $Eu := \bar{u}$ and observe that the mapping $u \mapsto Eu$ is linear.

Recall that the construction so far assumed $u \in C^\infty(\bar{U})$. Suppose now $1 \leq p < \infty$, $u \in W^{1,p}(U)$, and choose $u_m \in C^\infty(\bar{U})$ converging to u in $W^{1,p}(U)$. Estimate (9) and the linearity of E imply

$$\|Eu_m - Eu_l\|_{W^{1,p}(\mathbb{R}^n)} \leq C\|u_m - u_l\|_{W^{1,p}(U)}.$$

Thus $\{Eu_m\}_{m=1}^\infty$ is a Cauchy sequence and so converges to $\bar{u} =: Eu$. This extension, which does not depend on the particular choice of the approximating sequence $\{u_m\}_{m=1}^\infty$, satisfies the conclusions of the theorem.

The case $p = \infty$ is left as an exercise. \square

Remarks. (i) Assume now that ∂U is C^2 . Then the extension operator E constructed above is also a bounded linear operator from $W^{2,p}(U)$ to $W^{2,p}(\mathbb{R}^n)$. To see this, note first in steps 3, 4 of the proof that although \bar{u} is not in general C^2 , it does belong to $W^{2,p}(B)$. We also have the bound

$$\|\bar{u}\|_{W^{2,p}(B)} \leq C\|u\|_{W^{2,p}(B^+)},$$

which follows from the definition (3). As before, we consequently derive the estimate

$$(10) \quad \|Eu\|_{W^{2,p}(\mathbb{R}^n)} \leq C\|u\|_{W^{2,p}(U)},$$

provided ∂U is C^2 , the constants C depending only on U, V, n and p .

We will need these observations later.

(ii) The above construction does *not* provide us with an extension for the Sobolev spaces $W^{k,p}(U)$, if $k > 2$. This requires a more complicated higher-order reflection technique.

5.5. TRACES

Next we discuss the possibility of assigning “boundary values” along ∂U to a function $u \in W^{1,p}(U)$, assuming that ∂U is C^1 . Now if $u \in C(\bar{U})$, then clearly u has values on ∂U in the usual sense. The problem is that a typical function $u \in W^{1,p}(U)$ is not in general continuous and, even worse, is only defined a.e. in U . Since ∂U has n -dimensional Lebesgue measure zero, there is no direct meaning we can give to the expression “ u restricted to ∂U ”. The notion of a *trace operator* resolves this problem.

For this section we take $1 \leq p < \infty$.

THEOREM 1 (Trace Theorem). *Assume U is bounded and ∂U is C^1 . Then there exists a bounded linear operator*

$$T : W^{1,p}(U) \rightarrow L^p(\partial U)$$

such that

$$(i) \quad Tu = u|_{\partial U} \quad \text{if } u \in W^{1,p}(U) \cap C(\bar{U})$$

and

(ii)

$$\|Tu\|_{L^p(\partial U)} \leq C\|u\|_{W^{1,p}(U)},$$

for each $u \in W^{1,p}(U)$, with the constant C depending only on p and U .

DEFINITION. *We call Tu the trace of u on ∂U .*

Proof. 1. Assume first $u \in C^1(\bar{U})$. As in the first part of the proof of Theorem 1 in §5.4 let us also initially suppose $x^0 \in \partial U$ and ∂U is flat near x^0 , lying in the plane $\{x_n = 0\}$. Choose an open ball B as in the previous proof and let \hat{B} denote the concentric ball with radius $r/2$.

Select $\zeta \in C_c^\infty(B)$, with $\zeta \geq 0$ in B , $\zeta \equiv 1$ on \hat{B} . Denote by Γ that portion of ∂U within \hat{B} . Set $x' = (x_1, \dots, x_{n-1}) \in \mathbb{R}^{n-1} = \{x_n = 0\}$.

Then

$$\begin{aligned} \int_{\Gamma} |u|^p dx' &\leq \int_{\{x_n=0\}} \zeta |u|^p dx' = - \int_{B^+} (\zeta |u|^p)_{x_n} dx \\ (1) \quad &= - \int_{B^+} |u|^p \zeta_{x_n} + p |u|^{p-1} (\text{sgn } u) u_{x_n} \zeta dx \\ &\leq C \int_{B^+} |u|^p + |Du|^p dx, \end{aligned}$$

where we employed Young's inequality, from §B.2.

2. If $x^0 \in \partial U$, but ∂U is not flat near x^0 , we as usual straighten out the boundary near x^0 to obtain the setting above. Applying estimate (1) and changing variables, we obtain the bound

$$\int_{\Gamma} |u|^p dS \leq C \int_U |u|^p + |Du|^p dx,$$

where Γ is some open subset of ∂U containing x^0 .

3. Since ∂U is compact, there exist finitely many points $x_i^0 \in \partial U$ and open subsets $\Gamma_i \subset \partial U$ ($i = 1, \dots, N$) such that $\partial U = \bigcup_{i=1}^N \Gamma_i$ and

$$\|u\|_{L^p(\Gamma_i)} \leq C\|u\|_{W^{1,p}(U)} \quad (i = 1, \dots, N).$$

Consequently, if we write

$$Tu := u|_{\partial U},$$

then

$$(2) \quad \|Tu\|_{L^p(\partial U)} \leq C\|u\|_{W^{1,p}(U)}$$

for some appropriate constant C , which does not depend on u .

4. Inequality (2) holds for $u \in C^1(\bar{U})$. Assume now $u \in W^{1,p}(U)$. Then there exist functions $u_m \in C^\infty(\bar{U})$ converging to u in $W^{1,p}(U)$. According to (2) we have

$$(3) \quad \|Tu_m - Tu_l\|_{L^p(\partial U)} \leq C\|u_m - u_l\|_{W^{1,p}(U)};$$

so that $\{Tu_m\}_{m=1}^\infty$ is a Cauchy sequence in $L^p(\partial U)$. We define

$$Tu := \lim_{m \rightarrow \infty} Tu_m,$$

the limit taken in $L^p(\partial U)$. According to (3) this definition does not depend on the particular choice of smooth functions approximating u .

Finally if $u \in W^{1,p}(U) \cap C(\bar{U})$, we note that the functions $u_m \in C^\infty(\bar{U})$ constructed in the proof of Theorem 3 in §5.3.3 converge uniformly to u on \bar{U} . Hence $Tu = u|_{\partial U}$. \square

We next examine more closely what it means for a function to have zero trace.

THEOREM 2 (Trace-zero functions in $W^{1,p}$). *Assume U is bounded and ∂U is C^1 . Suppose furthermore that $u \in W^{1,p}(U)$. Then*

$$(4) \quad u \in W_0^{1,p}(U) \quad \text{if and only if} \quad Tu = 0 \quad \text{on} \quad \partial U.$$

Proof*. 1. Suppose first $u \in W_0^{1,p}(U)$. Then by definition there exist functions $u_m \in C_c^\infty(U)$ such that

$$u_m \rightarrow u \quad \text{in} \quad W^{1,p}(U).$$

As $Tu_m = 0$ on ∂U ($m = 1, \dots$) and $T : W^{1,p}(U) \rightarrow L^p(\partial U)$ is a bounded linear operator, we deduce $Tu = 0$ on ∂U .

2. The converse statement is more difficult. Let us assume that

$$(5) \quad Tu = 0 \quad \text{on} \quad \partial U.$$

*Omit on first reading.

Using partitions of unity and flattening out ∂U as usual, we may as well assume

$$(6) \quad \begin{cases} u \in W^{1,p}(\mathbb{R}_+^n), & u \text{ has compact support in } \bar{\mathbb{R}}_+^n, \\ Tu = 0 \text{ on } \partial\mathbb{R}_+^n = \mathbb{R}^{n-1}. \end{cases}$$

Then since $Tu = 0$ on \mathbb{R}^{n-1} , there exist functions $u_m \in C^1(\bar{\mathbb{R}}_+^n)$ such that

$$(7) \quad u_m \rightarrow u \quad \text{in } W^{1,p}(\mathbb{R}_+^n)$$

and

$$(8) \quad Tu_m = u_m|_{\mathbb{R}^{n-1}} \rightarrow 0 \quad \text{in } L^p(\mathbb{R}^{n-1}).$$

Now if $x' \in \mathbb{R}^{n-1}$, $x_n \geq 0$, we have

$$|u_m(x', x_n)| \leq |u_m(x', 0)| + \int_0^{x_n} |u_{m,x_n}(x', t)| dt.$$

Thus

$$\begin{aligned} \int_{\mathbb{R}^{n-1}} |u_m(x', x_n)|^p dx' &\leq C \left(\int_{\mathbb{R}^{n-1}} |u_m(x', 0)|^p dx' \right. \\ &\quad \left. + x_n^{p-1} \int_0^{x_n} \int_{\mathbb{R}^{n-1}} |Du_m(x', t)|^p dx' dt \right). \end{aligned}$$

Letting $m \rightarrow \infty$ and recalling (7), (8), we deduce

$$(9) \quad \int_{\mathbb{R}^{n-1}} |u(x', x_n)|^p dx' \leq C x_n^{p-1} \int_0^{x_n} \int_{\mathbb{R}^{n-1}} |Du|^p dx' dt$$

for a.e. $x_n > 0$.

3. Next let $\zeta \in C^\infty(\mathbb{R}_+)$ satisfy

$$\zeta \equiv 1 \text{ on } [0, 1], \quad \zeta \equiv 0 \text{ on } \mathbb{R}_+ - [0, 2], \quad 0 \leq \zeta \leq 1,$$

and write

$$\begin{cases} \zeta_m(x) := \zeta(mx_n) & (x \in \mathbb{R}_+^n) \\ w_m := u(x)(1 - \zeta_m). \end{cases}$$

Then

$$\begin{cases} w_{m,x_n} = u_{x_n}(1 - \zeta_m) - mu\zeta' \\ D_{x'} w_m = D_{x'} u(1 - \zeta_m). \end{cases}$$

Consequently

$$\begin{aligned}
 \int_{\mathbb{R}_+^n} |Dw_m - Du|^p dx &\leq C \int_{\mathbb{R}_+^n} |\zeta_m|^p |Du|^p dx \\
 (10) \qquad \qquad \qquad &+ Cm^p \int_0^{2/m} \int_{\mathbb{R}^{n-1}} |u|^p dx' dt \\
 &=: A + B.
 \end{aligned}$$

Now

$$(11) \qquad \qquad \qquad A \rightarrow 0 \quad \text{as } m \rightarrow \infty,$$

since $\zeta_m \neq 0$ only if $0 \leq x_n \leq 2/m$. To estimate the term B , we utilize inequality (9):

$$\begin{aligned}
 (12) \qquad B &\leq Cm^p \left(\int_0^{2/m} t^{p-1} dt \right) \left(\int_0^{2/m} \int_{\mathbb{R}^{n-1}} |Du|^p dx' dx_n \right) \\
 &\leq C \int_0^{2/m} \int_{\mathbb{R}^{n-1}} |Du|^p dx' dx_n \rightarrow 0 \quad \text{as } m \rightarrow \infty.
 \end{aligned}$$

Employing (10)–(12), we deduce $Dw_m \rightarrow Du$ in $L^p(\mathbb{R}_+^n)$. Since clearly $w_m \rightarrow u$ in $L^p(\mathbb{R}_+^n)$, we conclude

$$w_m \rightarrow u \quad \text{in } W^{1,p}(\mathbb{R}_+^n).$$

But $w_m = 0$ if $0 < x_n < 1/m$. We can therefore mollify the w_m to produce functions $u_m \in C_c^\infty(\mathbb{R}_+^n)$ such that $u_m \rightarrow u$ in $W^{1,p}(\mathbb{R}_+^n)$. Hence $u \in W_0^{1,p}(\mathbb{R}_+^n)$. \square

5.6. SOBOLEV INEQUALITIES

Our goal in this section is to discover embeddings of various Sobolev spaces into others. The crucial analytic tools here will be certain so-called ‘‘Sobolev-type inequalities’’, which we will prove below for smooth functions. These will then establish the estimates for arbitrary functions in the various relevant Sobolev spaces, since—as we saw in §5.3—smooth functions are dense.

To clarify the presentation we will consider first only the Sobolev space $W^{1,p}(U)$ and ask the following basic question: *if a function u belongs to $W^{1,p}(U)$, does u automatically belong to certain other spaces?* The answer will be ‘‘yes’’, but which other spaces depends upon whether

- (1) $1 \leq p < n,$
- (2) $p = n,$
- (3) $n < p \leq \infty.$

We study case (1) in §5.6.1, case (3) in §5.6.2, and the borderline case (2) only later in §5.8.1.

5.6.1. Gagliardo–Nirenberg–Sobolev inequality.

For this section let us assume

$$(4) \quad 1 \leq p < n$$

and first ask whether we can establish an estimate of the form

$$(5) \quad \|u\|_{L^q(\mathbb{R}^n)} \leq C \|Du\|_{L^p(\mathbb{R}^n)},$$

for certain constants $C > 0$, $1 \leq q < \infty$ and all functions $u \in C_c^\infty(\mathbb{R}^n)$. The point is that the constants C and q should not depend on u .

Motivation. Let us first demonstrate that *if* any inequality of the form (5) holds, then the number q cannot be arbitrary but must in fact have a very specific form. For this, choose a function $u \in C_c^\infty(\mathbb{R}^n)$, $u \not\equiv 0$, and define for $\lambda > 0$ the rescaled function

$$u_\lambda(x) := u(\lambda x) \quad (x \in \mathbb{R}^n).$$

Applying (5) to u_λ , we find

$$(6) \quad \|u_\lambda\|_{L^q(\mathbb{R}^n)} \leq C \|Du_\lambda\|_{L^p(\mathbb{R}^n)}.$$

Now

$$\int_{\mathbb{R}^n} |u_\lambda|^q dx = \int_{\mathbb{R}^n} |u(\lambda x)|^q dx = \frac{1}{\lambda^n} \int_{\mathbb{R}^n} |u(y)|^q dy,$$

and

$$\int_{\mathbb{R}^n} |Du_\lambda|^p dx = \lambda^p \int_{\mathbb{R}^n} |Du(\lambda x)|^p dx = \frac{\lambda^p}{\lambda^n} \int_{\mathbb{R}^n} |Du(y)|^p dy.$$

Inserting these equalities into (6), we discover

$$\frac{1}{\lambda^{n/q}} \|u\|_{L^q(\mathbb{R}^n)} \leq C \frac{\lambda}{\lambda^{n/p}} \|Du\|_{L^p(\mathbb{R}^n)},$$

and so

$$(7) \quad \|u\|_{L^q(\mathbb{R}^n)} \leq C \lambda^{1 - \frac{n}{p} + \frac{n}{q}} \|Du\|_{L^p(\mathbb{R}^n)}.$$

But then if $1 - \frac{n}{p} + \frac{n}{q} \neq 0$, we can upon sending λ to either 0 or ∞ in (7) obtain a contradiction. Thus *if* in fact the desired inequality (5) holds, we must necessarily have $1 - \frac{n}{p} + \frac{n}{q} = 0$; so that $\frac{1}{q} = \frac{1}{p} - \frac{1}{n}$, $q = \frac{np}{n-p}$.

This observation motivates the following

DEFINITION. If $1 \leq p < n$, the Sobolev conjugate of p is

$$(8) \quad p^* := \frac{np}{n-p}.$$

Note that

$$(9) \quad \frac{1}{p^*} = \frac{1}{p} - \frac{1}{n}, \quad p^* > p.$$

The foregoing scaling analysis shows that the estimate (5) can only possibly be true for $q = p^*$. Next we prove this inequality is in fact valid.

THEOREM 1 (Gagliardo–Nirenberg–Sobolev inequality). Assume $1 \leq p < n$. There exists a constant C , depending only on p and n , such that

$$(10) \quad \|u\|_{L^{p^*}(\mathbb{R}^n)} \leq C \|Du\|_{L^p(\mathbb{R}^n)},$$

for all $u \in C_c^1(\mathbb{R}^n)$.

Now we really do need u to have compact support for (10) to hold, as the example $u \equiv 1$ shows. But remarkably the constant here does not depend at all upon the size of the support of u .

Proof. 1. First assume $p = 1$.

Since u has compact support, for each $i = 1, \dots, n$ and $x \in \mathbb{R}^n$ we have

$$u(x) = \int_{-\infty}^{x_i} u_{x_i}(x_1, \dots, x_{i-1}, y_i, x_{i+1}, \dots, x_n) dy_i;$$

and so

$$|u(x)| \leq \int_{-\infty}^{\infty} |Du(x_1, \dots, y_i, \dots, x_n)| dy_i \quad (i = 1, \dots, n).$$

Consequently

$$(11) \quad |u(x)|^{\frac{n}{n-1}} \leq \prod_{i=1}^n \left(\int_{-\infty}^{\infty} |Du(x_1, \dots, y_i, \dots, x_n)| dy_i \right)^{\frac{1}{n-1}}.$$

Integrate this inequality with respect to x_1 :

$$(12) \quad \begin{aligned} \int_{-\infty}^{\infty} |u|^{\frac{n}{n-1}} dx_1 &\leq \int_{-\infty}^{\infty} \prod_{i=1}^n \left(\int_{-\infty}^{\infty} |Du| dy_i \right)^{\frac{1}{n-1}} dx_1 \\ &= \left(\int_{-\infty}^{\infty} |Du| dy_1 \right)^{\frac{1}{n-1}} \int_{-\infty}^{\infty} \prod_{i=2}^n \left(\int_{-\infty}^{\infty} |Du| dy_i \right)^{\frac{1}{n-1}} dx_1 \\ &\leq \left(\int_{-\infty}^{\infty} |Du| dy_1 \right)^{\frac{1}{n-1}} \left(\prod_{i=2}^n \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} |Du| dx_1 dy_i \right)^{\frac{1}{n-1}}, \end{aligned}$$

the last inequality resulting from the general Hölder inequality (§B.2).

Now integrate (12) with respect to x_2 :

$$\begin{aligned} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} |u|^{\frac{n}{n-1}} dx_1 dx_2 \\ \leq \left(\int_{-\infty}^{\infty} \int_{-\infty}^{\infty} |Du| dx_1 dy_2 \right)^{\frac{1}{n-1}} \int_{-\infty}^{\infty} \prod_{\substack{i=1 \\ i \neq 2}}^n I_i^{\frac{1}{n-1}} dx_2, \end{aligned}$$

for

$$I_1 := \int_{-\infty}^{\infty} |Du| dy_1, \quad I_i := \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} |Du| dx_1 dy_i \quad (i = 3, \dots, n).$$

Applying once more the extended Hölder inequality, we find

$$\begin{aligned} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} |u|^{\frac{n}{n-1}} dx_1 dx_2 \\ \leq \left(\int_{-\infty}^{\infty} \int_{-\infty}^{\infty} |Du| dx_1 dy_2 \right)^{\frac{1}{n-1}} \left(\int_{-\infty}^{\infty} \int_{-\infty}^{\infty} |Du| dy_1 dx_2 \right)^{\frac{1}{n-1}} \\ \prod_{i=3}^n \left(\int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} |Du| dx_1 dx_2 dy_i \right)^{\frac{1}{n-1}}. \end{aligned}$$

We continue by integrating with respect to x_3, \dots, x_n , eventually to find

$$\begin{aligned} (13) \quad \int_{\mathbb{R}^n} |u|^{\frac{n}{n-1}} dx &\leq \prod_{i=1}^n \left(\int_{-\infty}^{\infty} \cdots \int_{-\infty}^{\infty} |Du| dx_1 \dots dy_i \dots dx_n \right)^{\frac{1}{n-1}} \\ &= \left(\int_{\mathbb{R}^n} |Du| dx \right)^{\frac{n}{n-1}}. \end{aligned}$$

This is estimate (10) for $p = 1$.

2. Consider now the case that $1 < p < n$. We apply estimate (13) to $v := |u|^\gamma$, where $\gamma > 1$ is to be selected. Then

$$\begin{aligned} (14) \quad \left(\int_{\mathbb{R}^n} |u|^{\frac{\gamma n}{n-1}} dx \right)^{\frac{n-1}{n}} &\leq \int_{\mathbb{R}^n} |D|u|^\gamma| dx = \gamma \int_{\mathbb{R}^n} |u|^{\gamma-1} |Du| dx \\ &\leq \gamma \left(\int_{\mathbb{R}^n} |u|^{(\gamma-1)\frac{p}{p-1}} dx \right)^{\frac{p-1}{p}} \left(\int_{\mathbb{R}^n} |Du|^p dx \right)^{\frac{1}{p}}. \end{aligned}$$

We choose γ so that $\frac{\gamma n}{n-1} = (\gamma-1)\frac{p}{p-1}$. That is, we set

$$\gamma := \frac{p(n-1)}{n-p} > 1,$$

in which case $\frac{\gamma n}{n-1} = (\gamma - 1) \frac{p}{p-1} = \frac{np}{n-p} = p^*$. Thus, in view of (5), estimate (14) becomes

$$\left(\int_{\mathbb{R}^n} |u|^{p^*} dx \right)^{\frac{1}{p^*}} \leq C \left(\int_{\mathbb{R}^n} |Du|^p dx \right)^{\frac{1}{p}}. \quad \square$$

THEOREM 2 (Estimates for $W^{1,p}$, $1 \leq p < n$). *Let U be a bounded, open subset of \mathbb{R}^n , and suppose ∂U is C^1 . Assume $1 \leq p < n$, and $u \in W^{1,p}(U)$. Then $u \in L^{p^*}(U)$, with the estimate*

$$(15) \quad \|u\|_{L^{p^*}(U)} \leq C \|u\|_{W^{1,p}(U)},$$

the constant C depending only on p, n , and U .

Proof. Since ∂U is C^1 , there exists according to Theorem 1 in §5.4 an extension $Eu = \bar{u} \in W^{1,p}(\mathbb{R}^n)$, such that

$$(16) \quad \begin{cases} \bar{u} = u \text{ in } U, \bar{u} \text{ has compact support, and} \\ \|\bar{u}\|_{W^{1,p}(\mathbb{R}^n)} \leq C \|u\|_{W^{1,p}(U)}. \end{cases}$$

Because \bar{u} has compact support, we know from Theorem 1 in §5.3 that there exist functions $u_m \in C_c^\infty(\mathbb{R}^n)$ ($m = 1, 2, \dots$) such that

$$(17) \quad u_m \rightarrow \bar{u} \quad \text{in } W^{1,p}(\mathbb{R}^n).$$

Now according to Theorem 1, $\|u_m - u_l\|_{L^{p^*}(\mathbb{R}^n)} \leq C \|Du_m - Du_l\|_{L^p(\mathbb{R}^n)}$ for all $l, m \geq 1$. Thus

$$(18) \quad u_m \rightarrow \bar{u} \quad \text{in } L^{p^*}(\mathbb{R}^n)$$

as well. Since Theorem 1 also implies $\|u_m\|_{L^{p^*}(\mathbb{R}^n)} \leq C \|Du_m\|_{L^p(\mathbb{R}^n)}$, assertions (17) and (18) yield the bound

$$\|\bar{u}\|_{L^{p^*}(\mathbb{R}^n)} \leq C \|D\bar{u}\|_{L^p(\mathbb{R}^n)}.$$

This inequality and (16) complete the proof. □

THEOREM 3 (Estimates for $W_0^{1,p}$, $1 \leq p < n$). *Assume U is a bounded, open subset of \mathbb{R}^n . Suppose $u \in W_0^{1,p}(U)$ for some $1 \leq p < n$. Then we have the estimate*

$$\|u\|_{L^q(U)} \leq C \|Du\|_{L^p(U)}$$

for each $q \in [1, p^*]$, the constant C depending only on p, q, n and U .

In particular, for all $1 \leq p \leq \infty$,

$$\|u\|_{L^p(U)} \leq C \|Du\|_{L^p(U)}.$$

This estimate is sometimes called *Poincaré's inequality*. The difference with Theorem 2 is that only the gradient of u appears on the right-hand side of the inequality. (Other Poincaré-type inequalities will be established later, in §5.8.1.)

In view of this estimate, on $W_0^{1,p}(U)$ the norm $\|Du\|_{L^p(U)}$ is equivalent to $\|u\|_{W^{1,p}(U)}$, if U is bounded.

Proof. Since $u \in W_0^{1,p}(U)$, there exist functions $u_m \in C_c^\infty(U)$ ($m = 1, 2, \dots$) converging to u in $W^{1,p}(U)$. We extend each function u_m to be 0 on $\mathbb{R}^n - \bar{U}$ and apply Theorem 1 to discover $\|u\|_{L^{p^*}(U)} \leq C \|Du\|_{L^p(U)}$. As $|U| < \infty$, we furthermore have $\|u\|_{L^q(U)} \leq C \|u\|_{L^{p^*}(U)}$ if $1 \leq q \leq p^*$. \square

The borderline case $p=n$. We assume next that

$$p = n.$$

Owing to Theorem 2 and the fact that $p^* = \frac{np}{n-p} \rightarrow +\infty$ as $p \rightarrow n$, we might expect $u \in L^\infty(U)$, provided $u \in W^{1,n}(U)$. *This is however false if $n > 1$:* for example, if $U = B^0(0, 1)$, the function $u = \log \log \left(1 + \frac{1}{|x|}\right)$ belongs to $W^{1,n}(U)$ but not to $L^\infty(U)$. We will return to this borderline situation in §5.8.1 below.

5.6.2. Morrey's inequality.

Now let us suppose

$$(19) \quad n < p < \infty.$$

We will show that if $u \in W^{1,p}(U)$, then u is in fact Hölder continuous, after possibly being redefined on a set of measure zero.

THEOREM 4 (Morrey's inequality). *Assume $n < p \leq \infty$. Then there exists a constant C , depending only on p and n , such that*

$$(20) \quad \|u\|_{C^{0,\gamma}(\mathbb{R}^n)} \leq C \|u\|_{W^{1,p}(\mathbb{R}^n)}$$

for all $u \in C^1(\mathbb{R}^n)$, where

$$\gamma := 1 - n/p.$$

Proof. 1. We claim there exists a constant C , depending only on n , such that

$$(21) \quad \int_{B(x,r)} |u(y) - u(x)| dy \leq C \int_{B(x,r)} \frac{|Du(y)|}{|y-x|^{n-1}} dy$$

for each ball $B(x, r) \subset \mathbb{R}^n$.

To prove this, fix any point $w \in \partial B(0, 1)$. Then if $0 < s < r$,

$$\begin{aligned} |u(x + sw) - u(x)| &= \left| \int_0^s \frac{d}{dt} u(x + tw) dt \right| \\ &= \left| \int_0^s Du(x + tw) \cdot w dt \right| \\ &\leq \int_0^s |Du(x + tw)| dt. \end{aligned}$$

Hence

$$(22) \quad \int_{\partial B(0,1)} |u(x + sw) - u(x)| dS(w) \leq \int_0^s \int_{\partial B(0,1)} |Du(x + tw)| dS(w).$$

Now

$$\begin{aligned} \int_0^s \int_{\partial B(0,1)} |Du(x + tw)| dS(w) dt &= \int_0^s \int_{\partial B(x,t)} \frac{|Du(y)|}{t^{n-1}} dS(y) dt \\ &= \int_{B(x,s)} \frac{|Du(y)|}{|x-y|^{n-1}} dy \\ &\leq \int_{B(x,r)} \frac{|Du(y)|}{|x-y|^{n-1}} dy, \end{aligned}$$

where we put $y = x + tw$, $t = |x - y|$. Furthermore

$$\int_{\partial B(0,1)} |u(x + sw) - u(x)| dS(w) = \frac{1}{s^{n-1}} \int_{\partial B(x,s)} |u(z) - u(x)| dS(z),$$

for $z = x + sw$. Using the preceding two calculations in (22), we obtain the estimate

$$\int_{\partial B(x,s)} |u(z) - u(x)| dS(z) \leq s^{n-1} \int_{B(x,r)} \frac{|Du(y)|}{|x-y|^{n-1}} dy.$$

Now integrate with respect to s from 0 to r :

$$\int_{B(x,r)} |u(y) - u(x)| dy \leq \frac{r^n}{n} \int_{B(x,r)} \frac{|Du(y)|}{|x-y|^{n-1}} dy.$$

This implies (21).

2. Now fix $x \in \mathbb{R}^n$. We apply inequality (21) as follows:

$$\begin{aligned}
 |u(x)| &\leq \int_{B(x,1)} |u(x) - u(y)| dy + \int_{B(x,1)} |u(y)| dy \\
 &\leq C \int_{B(x,1)} \frac{|Du(y)|}{|x-y|^{n-1}} dy + C \|u\|_{L^p(B(x,1))} \\
 &\leq C \left(\int_{\mathbb{R}^n} |Du|^p dy \right)^{1/p} \left(\int_{B(x,1)} \frac{1}{|x-y|^{(n-1)\frac{p}{p-1}}} dy \right)^{\frac{p-1}{p}} \\
 &\qquad\qquad\qquad + C \|u\|_{L^p(\mathbb{R}^n)} \\
 &\leq C \|u\|_{W^{1,p}(\mathbb{R}^n)}.
 \end{aligned}$$

The last estimate holds since $p > n$ implies $(n-1)\frac{p}{p-1} < n$, so that

$$\int_{B(x,1)} \frac{1}{|x-y|^{(n-1)\frac{p}{p-1}}} dy < \infty.$$

As $x \in \mathbb{R}^n$ is arbitrary, it follows that

$$(23) \quad \sup_{\mathbb{R}^n} |u| \leq C \|u\|_{W^{1,p}(\mathbb{R}^n)}.$$

3. Next, choose any two points $x, y \in \mathbb{R}^n$ and write $r := |x - y|$. Let $W := B(x, r) \cap B(y, r)$. Then

$$(24) \quad |u(x) - u(y)| \leq \int_W |u(x) - u(z)| dz + \int_W |u(y) - u(z)| dz.$$

But inequality (21) allows us to estimate

$$\begin{aligned}
 \int_W |u(x) - u(z)| dz &\leq C \int_{B(x,r)} |u(x) - u(z)| dz \\
 (25) \quad &\leq C \left(\int_{B(x,r)} |Du|^p dz \right)^{1/p} \left(\int_{B(x,r)} \frac{dz}{|x-z|^{(n-1)\frac{p}{p-1}}} \right)^{\frac{p-1}{p}} \\
 &\leq C \left(r^{n-(n-1)\frac{p}{p-1}} \right)^{\frac{p-1}{p}} \|Du\|_{L^p(\mathbb{R}^n)} \\
 &= Cr^{1-\frac{n}{p}} \|Du\|_{L^p(\mathbb{R}^n)}.
 \end{aligned}$$

Likewise,

$$\int_W |u(y) - u(z)| dz \leq Cr^{1-\frac{n}{p}} \|Du\|_{L^p(\mathbb{R}^n)}.$$

Our substituting this estimate and (25) into (24) yields

$$|u(x) - u(y)| \leq Cr^{1-\frac{n}{p}} \|Du\|_{L^p(\mathbb{R}^n)} = C|x - y|^{1-\frac{n}{p}} \|Du\|_{L^p(\mathbb{R}^n)}.$$

Thus

$$[u]_{C^{0,1-n/p}(\mathbb{R}^n)} = \sup_{x \neq y} \left\{ \frac{|u(x) - u(y)|}{|x - y|^{1-n/p}} \right\} \leq C \|Du\|_{L^p(\mathbb{R}^n)}.$$

This inequality and (23) complete the proof of (20). \square

Remark. A slight variant of the proof above provides the estimate

$$|u(y) - u(x)| \leq Cr^{1-\frac{n}{p}} \left(\int_{B(x,2r)} |Du(z)|^p dz \right)^{1/p}$$

for all $u \in C^1(B(x, 2r))$, $y \in B(x, r)$, $n < p < \infty$. By an approximation the same bound is valid for $u \in W^{1,p}(B(x, 2r))$, $n < p < \infty$. We will use this inequality later in §5.8.2. (This estimate is in fact valid if on the right-hand side we integrate over $B(x, r)$, instead of $B(x, 2r)$, but the proof is a bit trickier.)

DEFINITION. We say u^* is a version of a given function u provided

$$u = u^* \text{ a.e.}$$

THEOREM 5 (Estimates for $W^{1,p}$, $n < p \leq \infty$). *Let U be a bounded, open subset of \mathbb{R}^n , and suppose ∂U is C^1 . Assume $n < p \leq \infty$ and $u \in W^{1,p}(U)$. Then u has a version $u^* \in C^{0,\gamma}(\bar{U})$, for $\gamma = 1 - \frac{n}{p}$, with the estimate*

$$\|u^*\|_{C^{0,\gamma}(\bar{U})} \leq C \|u\|_{W^{1,p}(U)}.$$

The constant C depends only on p, n and U .

In view of Theorem 5, we will henceforth always identify a function $u \in W^{1,p}(U)$ ($p > n$) with its continuous version.

Proof. Since ∂U is C^1 , there exists according to Theorem 1 in §5.4 an extension $Eu = \bar{u} \in W^{1,p}(\mathbb{R}^n)$ such that

$$(26) \quad \begin{cases} \bar{u} = u \text{ in } U, \\ \bar{u} \text{ has compact support, and} \\ \|\bar{u}\|_{W^{1,p}(\mathbb{R}^n)} \leq C \|u\|_{W^{1,p}(U)}. \end{cases}$$

Assume first $n < p < \infty$. Since \bar{u} has compact support, we obtain from Theorem 1 in §5.3 the existence of functions $u_m \in C_c^\infty(\mathbb{R}^n)$ such that

$$(27) \quad u_m \rightarrow \bar{u} \quad \text{in } W^{1,p}(\mathbb{R}^n).$$

Now according to Theorem 4, $\|u_m - u_l\|_{C^{0,1-n/p}(\mathbb{R}^n)} \leq C\|u_m - u_l\|_{W^{1,p}(\mathbb{R}^n)}$ for all $l, m \geq 1$, whence there exists a function $u^* \in C^{0,1-n/p}(\mathbb{R}^n)$ such that

$$(28) \quad u_m \rightarrow u^* \quad \text{in } C^{0,1-n/p}(\mathbb{R}^n).$$

Owing to (27) and (28), we see that $u^* = u$ a.e. on U , so that u^* is a version of u . Since Theorem 4 also implies $\|u_m\|_{C^{0,1-n/p}(\mathbb{R}^n)} \leq C\|u_m\|_{W^{1,p}(\mathbb{R}^n)}$, assertions (27) and (28) yield

$$\|u^*\|_{C^{0,1-n/p}(\mathbb{R}^n)} \leq C\|\bar{u}\|_{W^{1,p}(\mathbb{R}^n)}.$$

This inequality and (26) complete the proof if $n < p < \infty$. The case $p = \infty$ is easy to prove directly. \square

5.6.3. General Sobolev inequalities.

We can now concatenate the estimates established in §§5.6.1 and 5.6.2 to obtain more complicated (and hard-to-remember) inequalities.

THEOREM 6 (General Sobolev inequalities). *Let U be a bounded open subset of \mathbb{R}^n , with a C^1 boundary. Assume $u \in W^{k,p}(U)$.*

(i) *If*

$$(29) \quad k < \frac{n}{p},$$

then $u \in L^q(U)$, where

$$\frac{1}{q} = \frac{1}{p} - \frac{k}{n}.$$

We have in addition the estimate

$$(30) \quad \|u\|_{L^q(U)} \leq C\|u\|_{W^{k,p}(U)},$$

the constant C depending only on k, p, n and U .

(ii) *If*

$$(31) \quad k > \frac{n}{p},$$

then $u \in C^{k - [\frac{n}{p}] - 1, \gamma}(\bar{U})$, where

$$\gamma = \begin{cases} [\frac{n}{p}] + 1 - \frac{n}{p}, & \text{if } \frac{n}{p} \text{ is not an integer} \\ \text{any positive number} < 1, & \text{if } \frac{n}{p} \text{ is an integer.} \end{cases}$$

We have in addition the estimate

$$(32) \quad \|u\|_{C^{k - [\frac{n}{p}] - 1, \gamma}(\bar{U})} \leq C \|u\|_{W^{k, p}(U)},$$

the constant C depending only on k, p, n, γ and U .

Proof. 1. Assume (29). Then since $D^\alpha u \in L^p(U)$ for all $|\alpha| \leq k$, the Gagliardo–Nirenberg–Sobolev inequality implies

$$\|D^\beta u\|_{L^{p^*}(U)} \leq C \|u\|_{W^{k, p}(U)} \quad \text{if } |\beta| \leq k - 1,$$

and so $u \in W^{k-1, p^*}(U)$. Similarly, we find $u \in W^{k-2, p^{**}}(U)$, where $\frac{1}{p^{**}} = \frac{1}{p^*} - \frac{1}{n} = \frac{1}{p} - \frac{2}{n}$. Continuing, we eventually discover after k steps that $u \in W^{0, q}(U) = L^q(U)$, for $\frac{1}{q} = \frac{1}{p} - \frac{k}{n}$. The stated estimate (30) follows from multiplying the relevant estimates at each stage of the above argument.

2. Assume now condition (31) holds and $\frac{n}{p}$ is not an integer. Then as above we see

$$(33) \quad u \in W^{k-l, r}(U),$$

for

$$(34) \quad \frac{1}{r} = \frac{1}{p} - \frac{l}{n},$$

provided $lp < n$. We choose the integer l so that

$$(35) \quad l < \frac{n}{p} < l + 1;$$

that is, we set $l = [\frac{n}{p}]$. Consequently (34) and (35) imply $r = \frac{pn}{n-pl} > n$. Hence (33) and Morrey's inequality imply that $D^\alpha u \in C^{0, 1 - \frac{n}{r}}(\bar{U})$ for all $|\alpha| \leq k - l - 1$. Observe also that $1 - \frac{n}{r} = 1 - \frac{n}{p} + l = [\frac{n}{p}] + 1 - \frac{n}{p}$. Thus $u \in C^{k - [\frac{n}{p}] - 1, [\frac{n}{p}] + 1 - \frac{n}{p}}(\bar{U})$, and the stated estimate follows easily.

3. Finally, suppose (31) holds, with $\frac{n}{p}$ an integer. Set $l = [\frac{n}{p}] - 1 = \frac{n}{p} - 1$. Consequently, we have as above $u \in W^{k-l, r}(U)$ for $r = \frac{pn}{n-pl} = n$. Hence

the Gagliardo–Nirenberg–Sobolev inequality shows $D^\alpha u \in L^q(U)$ for all $n \leq q < \infty$ and all $|\alpha| \leq k - l - 1 = k - \left\lfloor \frac{n}{p} \right\rfloor$. Therefore Morrey's inequality further implies $D^\alpha u \in C^{0,1-\frac{n}{q}}(\bar{U})$ for all $n < q < \infty$ and all $|\alpha| \leq k - \left\lfloor \frac{n}{p} \right\rfloor - 1$. Consequently $u \in C^{k-\left\lfloor \frac{n}{p} \right\rfloor-1,\gamma}(\bar{U})$ for each $0 < \gamma < 1$. As before, the stated estimate follows as well. \square

Various general Sobolev-type inequalities can also be proved using the Fourier transform: see Problem 20.

5.7. COMPACTNESS

We have seen in §5.6 that the Gagliardo–Nirenberg–Sobolev inequality implies the embedding of $W^{1,p}(U)$ into $L^{p^*}(U)$ for $1 \leq p < n$, $p^* = \frac{pn}{n-p}$. We will now demonstrate that $W^{1,p}(U)$ is in fact *compactly* embedded in $L^q(U)$ for $1 \leq q < p^*$. This compactness will be fundamental for our applications of linear and nonlinear functional analysis to PDE in Chapters 6–9.

DEFINITION. *Let X and Y be Banach spaces, $X \subset Y$. We say that X is compactly embedded in Y , written*

$$X \subset\subset Y,$$

provided

(i) $\|u\|_Y \leq C\|u\|_X$ ($u \in X$) for some constant C

and

(ii) *each bounded sequence in X is precompact in Y .*

More precisely, condition (ii) means that if $\{u_k\}_{k=1}^\infty$ is a sequence in X with $\sup_k \|u_k\|_X < \infty$, then some subsequence $\{u_{k_j}\}_{j=1}^\infty \subseteq \{u_k\}_{k=1}^\infty$ converges in Y to some limit u :

$$\lim_{j \rightarrow \infty} \|u_{k_j} - u\|_Y = 0.$$

THEOREM 1 (Rellich–Kondrachov Compactness Theorem). *Assume U is a bounded open subset of \mathbb{R}^n and ∂U is C^1 . Suppose $1 \leq p < n$. Then*

$$W^{1,p}(U) \subset\subset L^q(U)$$

for each $1 \leq q < p^$.*

Proof. 1. Fix $1 \leq q < p^*$ and note that since U is bounded, Theorem 2 in §5.6.1 implies

$$W^{1,p}(U) \subset L^q(U), \quad \|u\|_{L^q(U)} \leq C\|u\|_{W^{1,p}(U)}.$$

It remains therefore to show that if $\{u_m\}_{m=1}^\infty$ is a bounded sequence in $W^{1,p}(U)$, there exists a subsequence $\{u_{m_j}\}_{j=1}^\infty$ which converges in $L^q(U)$.

2. In view of the Extension Theorem from §5.4 we may with no loss of generality assume that $U = \mathbb{R}^n$ and the functions $\{u_m\}_{m=1}^\infty$ all have compact support in some bounded open set $V \subset \mathbb{R}^n$. We also may assume

$$(1) \quad \sup_m \|u_m\|_{W^{1,p}(V)} < \infty.$$

3. Let us first study the smoothed functions

$$u_m^\varepsilon := \eta_\varepsilon * u_m \quad (\varepsilon > 0, m = 1, 2, \dots),$$

η_ε denoting the usual mollifier. We may suppose the functions $\{u_m^\varepsilon\}_{m=1}^\infty$ all have support in V as well.

4. We first claim

$$(2) \quad u_m^\varepsilon \rightarrow u_m \text{ in } L^q(V) \text{ as } \varepsilon \rightarrow 0, \text{ uniformly in } m.$$

To prove this, we first note that if u_m is smooth, then

$$\begin{aligned} u_m^\varepsilon(x) - u_m(x) &= \frac{1}{\varepsilon^n} \int_{B(x,\varepsilon)} \eta\left(\frac{x-z}{\varepsilon}\right) (u_m(z) - u_m(x)) dz \\ &= \int_{B(0,1)} \eta(y) (u_m(x - \varepsilon y) - u_m(x)) dy \\ &= \int_{B(0,1)} \eta(y) \int_0^1 \frac{d}{dt} (u_m(x - \varepsilon ty)) dt dy \\ &= -\varepsilon \int_{B(0,1)} \eta(y) \int_0^1 Du_m(x - \varepsilon ty) \cdot y dt dy. \end{aligned}$$

Thus

$$\begin{aligned} \int_V |u_m^\varepsilon(x) - u_m(x)| dx &\leq \varepsilon \int_{B(0,1)} \eta(y) \int_0^1 \int_V |Du_m(x - \varepsilon ty)| dx dt dy \\ &\leq \varepsilon \int_V |Du_m(z)| dz. \end{aligned}$$

By approximation this estimate holds if $u_m \in W^{1,p}(V)$. Hence

$$\|u_m^\varepsilon - u_m\|_{L^1(V)} \leq \varepsilon \|Du_m\|_{L^1(V)} \leq \varepsilon C \|Du_m\|_{L^p(V)},$$

the latter inequality holding since V is bounded. Owing to (1), we thereby discover

$$(3) \quad u_m^\varepsilon \rightarrow u_m \quad \text{in } L^1(V), \text{ uniformly in } m.$$

But then since $1 \leq q < p^*$, we see using the interpolation inequality for L^p -norms (§B.2) that

$$\|u_m^\varepsilon - u_m\|_{L^q(V)} \leq \|u_m^\varepsilon - u_m\|_{L^1(V)}^\theta \|u_m^\varepsilon - u_m\|_{L^{p^*}(V)}^{1-\theta},$$

where $\frac{1}{q} = \theta + \frac{(1-\theta)}{p^*}$, $0 < \theta < 1$. Consequently (1) and the Gagliardo–Nirenberg–Sobolev inequality imply

$$\|u_m^\varepsilon - u_m\|_{L^q(V)} \leq C \|u_m^\varepsilon - u_m\|_{L^1(V)}^\theta,$$

whence assertion (2) follows from (3).

5. Next we claim

$$(4) \quad \begin{cases} \text{for each fixed } \varepsilon > 0, \text{ the sequence } \{u_m^\varepsilon\}_{m=1}^\infty \\ \text{is uniformly bounded and equicontinuous.} \end{cases}$$

Indeed, if $x \in \mathbb{R}^n$, then

$$\begin{aligned} |u_m^\varepsilon(x)| &\leq \int_{B(x,\varepsilon)} \eta_\varepsilon(x-y) |u_m(y)| dy \\ &\leq \|\eta_\varepsilon\|_{L^\infty(\mathbb{R}^n)} \|u_m\|_{L^1(V)} \leq \frac{C}{\varepsilon^n} < \infty \end{aligned}$$

for $m = 1, 2, \dots$. Similarly

$$\begin{aligned} |Du_m^\varepsilon(x)| &\leq \int_{B(x,\varepsilon)} |D\eta_\varepsilon(x-y)| |u_m(y)| dy \\ &\leq \|D\eta_\varepsilon\|_{L^\infty(\mathbb{R}^n)} \|u_m\|_{L^1(V)} \leq \frac{C}{\varepsilon^{n+1}} < \infty, \end{aligned}$$

for $m = 1, \dots$. Assertion (4) follows from these two estimates.

6. Now fix $\delta > 0$. We will show there exists a subsequence $\{u_{m_j}\}_{j=1}^\infty \subset \{u_m\}_{m=1}^\infty$ such that

$$(5) \quad \limsup_{j,k \rightarrow \infty} \|u_{m_j} - u_{m_k}\|_{L^q(V)} \leq \delta.$$

To see this, let us first employ assertion (2) to select $\varepsilon > 0$ so small that

$$(6) \quad \|u_m^\varepsilon - u_m\|_{L^q(V)} \leq \delta/2$$

for $m = 1, 2, \dots$

We now observe that since the functions $\{u_m\}_{m=1}^\infty$, and thus the functions $\{u_m^\varepsilon\}_{m=1}^\infty$, have support in some fixed bounded set $V \subset \mathbb{R}^n$, we may utilize (4) and the Arzela–Ascoli compactness criterion, §C.7, to obtain a subsequence $\{u_{m_j}^\varepsilon\}_{j=1}^\infty \subset \{u_m^\varepsilon\}_{m=1}^\infty$ which converges *uniformly* on V . In particular therefore

$$(7) \quad \limsup_{j,k \rightarrow \infty} \|u_{m_j}^\varepsilon - u_{m_k}^\varepsilon\|_{L^q(V)} = 0.$$

But then (6) and (7) imply

$$\limsup_{j,k \rightarrow \infty} \|u_{m_j} - u_{m_k}\|_{L^q(V)} \leq \delta,$$

and so (5) is proved.

7. We next employ assertion (5) with $\delta = 1, \frac{1}{2}, \frac{1}{3}, \dots$ and use a standard diagonal argument to extract a subsequence $\{u_{m_l}\}_{l=1}^\infty \subset \{u_m\}_{m=1}^\infty$ satisfying

$$\limsup_{l,k \rightarrow \infty} \|u_{m_l} - u_{m_k}\|_{L^q(V)} = 0. \quad \square$$

Remark. Observe that since $p^* > p$ and $p^* \rightarrow \infty$ as $p \rightarrow n$, we have in particular

$$W^{1,p}(U) \subset\subset L^p(U)$$

for all $1 \leq p \leq \infty$. Observe that if $n < p \leq \infty$, this follows from Morrey's inequality and the Arzela–Ascoli compactness criterion (§C.8). Note also that

$$W_0^{1,p}(U) \subset\subset L^p(U),$$

even if we do not assume ∂U to be C^1 .

5.8. ADDITIONAL TOPICS

5.8.1. Poincaré's inequalities.

We now illustrate how the compactness assertion in §5.7 can be used to generate new inequalities.

NOTATION. $(u)_U = \int_U u \, dy =$ average of u over U .

THEOREM 1 (Poincaré's inequality). *Let U be a bounded, connected, open subset of \mathbb{R}^n , with a C^1 boundary ∂U . Assume $1 \leq p \leq \infty$. Then there exists a constant C , depending only on n, p and U , such that*

$$(1) \quad \|u - (u)_U\|_{L^p(U)} \leq C \|Du\|_{L^p(U)}$$

for each function $u \in W^{1,p}(U)$.

The significance of (1) is that only the gradient of u appears on the right-hand side.

Proof. We argue by contradiction. Were the stated estimate false, there would exist for each integer $k = 1, \dots$ a function $u_k \in W^{1,p}(U)$ satisfying

$$(2) \quad \|u_k - (u_k)_U\|_{L^p(U)} > k \|Du_k\|_{L^p(U)}.$$

We renormalize by defining

$$(3) \quad v_k := \frac{u_k - (u_k)_U}{\|u_k - (u_k)_U\|_{L^p(U)}} \quad (k = 1, \dots).$$

Then

$$(v_k)_U = 0, \quad \|v_k\|_{L^p(U)} = 1;$$

and (2) implies

$$(4) \quad \|Dv_k\|_{L^p(U)} < \frac{1}{k} \quad (k = 1, 2, \dots).$$

In particular the functions $\{v_k\}_{k=1}^\infty$ are bounded in $W^{1,p}(U)$.

In view of the remark after the proof of the Rellich–Kondrachov Theorem in §5.7, there exist a subsequence $\{v_{k_j}\}_{j=1}^\infty \subset \{v_k\}_{k=1}^\infty$ and a function $v \in L^p(U)$ such that

$$(5) \quad v_{k_j} \rightarrow v \quad \text{in } L^p(U).$$

From (3) it follows that

$$(6) \quad (v)_U = 0, \quad \|v\|_{L^p(U)} = 1.$$

On the other hand, (4) implies for each $i = 1, \dots, n$ and $\phi \in C_c^\infty(U)$ that

$$\int_U v \phi_{x_i} \, dx = \lim_{k_j \rightarrow \infty} \int_U v_{k_j} \phi_{x_i} \, dx = - \lim_{k_j \rightarrow \infty} \int_U v_{k_j, x_i} \phi \, dx = 0.$$

Consequently $v \in W^{1,p}(U)$, with $Dv = 0$ a.e. Thus v is constant, since U is connected (see Problem 11). However this conclusion is at variance with (6): since v is constant and $(v)_U = 0$, we must have $v \equiv 0$, in which case $\|v\|_{L^p(U)} = 0$. This contradiction establishes estimate (1). \square

A particularly important special case follows.

NOTATION. $(u)_{x,r} = \int_{B(x,r)} u \, dy =$ average of u over the ball $B(x,r)$.

THEOREM 2 (Poincaré's inequality for a ball). *Assume $1 \leq p \leq \infty$. Then there exists a constant C , depending only on n and p , such that*

$$(7) \quad \|u - (u)_{x,r}\|_{L^p(B(x,r))} \leq Cr \|Du\|_{L^p(B(x,r))}$$

for each ball $B(x,r) \subset \mathbb{R}^n$ and each function $u \in W^{1,p}(B^0(x,r))$.

Proof. The case $U = B^0(0,1)$ follows from Theorem 1. In general, if $u \in W^{1,p}(B^0(x,r))$, write

$$v(y) := u(x + ry) \quad (y \in B(0,1)).$$

Then $v \in W^{1,p}(B^0(0,1))$, and we have

$$\|v - (v)_{0,1}\|_{L^p(B(0,1))} \leq C \|Dv\|_{L^p(B(0,1))}.$$

Changing variables, we recover estimate (7). □

BMO and $W^{1,n}$. Assume $u \in W^{1,n}(\mathbb{R}^n) \cap L^1(\mathbb{R}^n)$, and let $B(x,r)$ be any ball. Then Theorem 2 with $p = 1$ implies

$$\begin{aligned} \int_{B(x,r)} |u - (u)_{x,r}| \, dy &\leq Cr \int_{B(x,r)} |Du| \, dy \\ &\leq Cr \left(\int_{B(x,r)} |Du|^n \, dy \right)^{1/n} \leq C \left(\int_{\mathbb{R}^n} |Du|^n \, dy \right)^{1/n}. \end{aligned}$$

Thus $u \in BMO(\mathbb{R}^n)$, the space of functions of *bounded mean oscillation* in \mathbb{R}^n , with the seminorm

$$[u]_{BMO(\mathbb{R}^n)} := \sup_{B(x,r) \subset \mathbb{R}^n} \left\{ \int_{B(x,r)} |u - (u)_{x,r}| \, dy \right\}.$$

See Stein [Se, Chapter IV] for the theory of the space BMO .

5.8.2. Difference quotients.

When we later apply Sobolev space theory to PDE, we will be forced to study difference quotient approximations to weak derivatives. Following is the relevant theory, which the reader may wish to postpone studying until the need arises, in §6.3.

a. Difference quotients and $W^{1,p}$.

Assume $u : U \rightarrow \mathbb{R}$ is a locally summable function and $V \subset\subset U$.

DEFINITIONS.

(i) The i^{th} -difference quotient of size h is

$$D_i^h u(x) = \frac{u(x + he_i) - u(x)}{h} \quad (i = 1, \dots, n)$$

for $x \in V$ and $h \in \mathbb{R}$, $0 < |h| < \text{dist}(V, \partial U)$.

(ii) $D^h u := (D_1^h u, \dots, D_n^h u)$.

THEOREM 3 (Difference quotients and weak derivatives).

(i) Suppose $1 \leq p < \infty$ and $u \in W^{1,p}(U)$. Then for each $V \subset\subset U$

$$(8) \quad \|D^h u\|_{L^p(V)} \leq C \|Du\|_{L^p(U)}$$

for some constant C and all $0 < |h| < \frac{1}{2} \text{dist}(V, \partial U)$.

(ii) Assume $1 < p < \infty$, $u \in L^p(V)$, and there exists a constant C such that

$$(9) \quad \|D^h u\|_{L^p(V)} \leq C$$

for all $0 < |h| < \frac{1}{2} \text{dist}(V, \partial U)$. Then

$$u \in W^{1,p}(V), \quad \text{with } \|Du\|_{L^p(V)} \leq C.$$

Assertion (ii) is false for $p = 1$ (Problem 12).

Proof. 1. Assume $1 \leq p < \infty$, and temporarily suppose u is smooth. Then for each $x \in V$, $i = 1, \dots, n$, and $0 < |h| < \frac{1}{2} \text{dist}(V, \partial U)$, we have

$$u(x + he_i) - u(x) = h \int_0^1 u_{x_i}(x + the_i) dt,$$

so that

$$|u(x + he_i) - u(x)| \leq |h| \int_0^1 |Du(x + the_i)| dt.$$

Consequently

$$\begin{aligned} \int_V |D^h u|^p dx &\leq C \sum_{i=1}^n \int_V \int_0^1 |Du(x + the_i)|^p dt dx \\ &= C \sum_{i=1}^n \int_0^1 \int_V |Du(x + the_i)|^p dx dt. \end{aligned}$$

Thus

$$\int_V |D^h u|^p dx \leq C \int_U |Du|^p dx.$$

This estimate holds should u be smooth, and thus is valid by approximation for arbitrary $u \in W^{1,p}(U)$.

2. Now suppose estimate (9) holds for all $0 < |h| < \frac{1}{2} \text{dist}(V, \partial U)$ and some constant C . Choose $i = 1, \dots, n$, $\phi \in C_c^\infty(V)$, and note for small enough h that

$$\int_V u(x) \left[\frac{\phi(x + he_i) - \phi(x)}{h} \right] dx = - \int_V \left[\frac{u(x) - u(x - he_i)}{h} \right] \phi(x) dx;$$

that is,

$$(10) \quad \int_V u(D_i^h \phi) dx = - \int_V (D_i^{-h} u) \phi dx.$$

This is the “integration-by-parts” formula for difference quotients. Estimate (9) implies

$$\sup_h \|D_i^{-h} u\|_{L^p(V)} < \infty;$$

and therefore, since $1 < p < \infty$, there exists a function $v_i \in L^p(V)$ and a subsequence $h_k \rightarrow 0$ such that

$$(11) \quad D_i^{-h_k} u \rightharpoonup v_i \quad \text{weakly in } L^p(V).$$

(See §D.4 for weak convergence.) But then

$$\begin{aligned} \int_V u \phi_{x_i} dx &= \int_U u \phi_{x_i} dx = \lim_{h_k \rightarrow 0} \int_U u D_i^{h_k} \phi dx \\ &= - \lim_{h_k \rightarrow 0} \int_V D_i^{-h_k} u \phi dx \\ &= - \int_V v_i \phi dx = - \int_U v_i \phi dx. \end{aligned}$$

Thus $v_i = u_{x_i}$ in the weak sense ($i = 1, \dots, n$), and so $Du \in L^p(V)$. As $u \in L^p(V)$, we deduce therefore that $u \in W^{1,p}(V)$. \square

Difference quotients near the boundary. Variants of Theorem 3 can be valid even if it is not true that $V \subset\subset U$. For example if U is the open half-ball $B^0(0, 1) \cap \{x_n > 0\}$, $V = B^0(0, 1/2) \cap \{x_n > 0\}$, we have the bound $\int_V |D_i^h u|^p dx \leq \int_U |u_{x_i}|^p dx$ for $i = 1, \dots, n-1$. The proof is similar to that just given.

We will need this comment in Chapter 6, §6.3.2.

b. Lipschitz functions and $W^{1,\infty}$.

THEOREM 4 (Characterization of $W^{1,\infty}$). *Let U be open and bounded, with ∂U of class C^1 . Then $u : U \rightarrow \mathbb{R}$ is Lipschitz continuous if and only if $u \in W^{1,\infty}(U)$.*

Proof. 1. First assume $U = \mathbb{R}^n$ and u has compact support.

Suppose $u \in W^{1,\infty}(\mathbb{R}^n)$. Then $u^\varepsilon := \eta_\varepsilon * u$, where η_ε is the usual mollifier, is smooth and satisfies

$$\begin{cases} u^\varepsilon \rightarrow u \text{ uniformly as } \varepsilon \rightarrow 0, \\ \|Du^\varepsilon\|_{L^\infty(\mathbb{R}^n)} \leq \|Du\|_{L^\infty(\mathbb{R}^n)}. \end{cases}$$

Choose any two points $x, y \in \mathbb{R}^n$, $x \neq y$. We have

$$\begin{aligned} u^\varepsilon(x) - u^\varepsilon(y) &= \int_0^1 \frac{d}{dt} u^\varepsilon(tx + (1-t)y) dt \\ &= \int_0^1 Du^\varepsilon(tx + (1-t)y) dt \cdot (x - y), \end{aligned}$$

and so

$$|u^\varepsilon(x) - u^\varepsilon(y)| \leq \|Du^\varepsilon\|_{L^\infty(\mathbb{R}^n)} |x - y| \leq \|Du\|_{L^\infty(\mathbb{R}^n)} |x - y|.$$

We let $\varepsilon \rightarrow 0$ to discover

$$|u(x) - u(y)| \leq \|Du\|_{L^\infty(\mathbb{R}^n)} |x - y|.$$

Hence u is Lipschitz continuous.

2. On the other hand assume now u is Lipschitz continuous; we must prove that u has essentially bounded weak first derivatives. Since u is Lipschitz, we see

$$\|D_i^{-h} u\|_{L^\infty(\mathbb{R}^n)} \leq \text{Lip}(u),$$

and thus there exists a function $v_i \in L^\infty(\mathbb{R}^n)$ and a subsequence $h_k \rightarrow 0$ such that

$$(12) \quad D_i^{-h_k} u \rightharpoonup v_i \quad \text{weakly in } L_{\text{loc}}^2(\mathbb{R}^n).$$

Consequently

$$\begin{aligned} \int_{\mathbb{R}^n} u \phi_{x_i} dx &= \lim_{h_k \rightarrow 0} \int_{\mathbb{R}^n} u D_i^{h_k} \phi dx \\ &= - \lim_{h_k \rightarrow 0} \int_{\mathbb{R}^n} D_i^{-h_k} u \phi dx = - \int_{\mathbb{R}^n} v_i \phi dx \end{aligned}$$

by (12). The above equality holds for all $\phi \in C_c^\infty(\mathbb{R}^n)$, and so $v_i = u_{x_i}$ in the weak sense ($i = 1, \dots, n$). Consequently $u \in W^{1,\infty}(\mathbb{R}^n)$.

3. In the general case that U is bounded, with ∂U of class C^1 , we as usual extend u to $Eu = \bar{u}$ and apply the above argument. \square

Remark. The argument above adapts easily to prove that for any open set U , $u \in W_{\text{loc}}^{1,\infty}(U)$ if and only if u is locally Lipschitz continuous in U . There is no corresponding characterization of the spaces $W^{1,p}$ for $1 \leq p < \infty$. If $n < p < \infty$, then each function $u \in W^{1,p}$ belongs to $C^{0,1-n/p}$; but a function Hölder continuous with exponent less than one need not belong to any Sobolev space $W^{1,p}$.

5.8.3. Differentiability a.e.

Next we examine more closely the connections between weak partial derivatives and partial derivatives in the usual calculus sense.

DEFINITION. A function $u : U \rightarrow \mathbb{R}$ is differentiable at $x \in U$ if there exists $a \in \mathbb{R}^n$ such that

$$(13) \quad u(y) = u(x) + a \cdot (y - x) + o(|y - x|) \quad \text{as } y \rightarrow x.$$

In other words,

$$\lim_{y \rightarrow x} \frac{|u(y) - u(x) - a \cdot (y - x)|}{|y - x|} = 0.$$

It is easy to check that a , if it exists, is unique. We henceforth write

$$Du(x)$$

for a and call Du the gradient of u .

To be sure that this notation is consistent, we need to study the relationships between the various notions of derivatives:

THEOREM 5 (Differentiability almost everywhere). Assume $u \in W_{\text{loc}}^{1,p}(U)$ for some $n < p \leq \infty$. Then u is differentiable a.e. in U , and its gradient equals its weak gradient a.e.

Recall that we always identify u with its continuous version.

Proof. 1. Assume first $n < p < \infty$. From the remark after the proof of Theorem 4 in §5.6.2, we recall Morrey's estimate

$$(14) \quad |v(y) - v(x)| \leq Cr^{1-\frac{n}{p}} \left(\int_{B(x,2r)} |Dv(z)|^p dz \right)^{1/p} \quad (y \in B(x,r)),$$

valid for any C^1 function v and thus, by approximation, for any $v \in W^{1,p}$.

2. Choose $u \in W_{\text{loc}}^{1,p}(U)$. Now for a.e. $x \in U$, a version of Lebesgue's Differentiation Theorem (§E.4) implies

$$(15) \quad \int_{B(x,r)} |Du(x) - Du(z)|^p dz \rightarrow 0$$

as $r \rightarrow 0$, Du denoting as usual the weak derivative of u . Fix any such point x and set

$$v(y) := u(y) - u(x) - Du(x) \cdot (y - x)$$

in estimate (14), where

$$(16) \quad r = |x - y|.$$

We find

$$\begin{aligned} & |u(y) - u(x) - Du(x) \cdot (y - x)| \\ & \leq Cr^{1-n/p} \left(\int_{B(x,2r)} |Du(x) - Du(z)|^p dz \right)^{1/p} \\ & \leq Cr \left(\int_{B(x,2r)} |Du(x) - Du(z)|^p dz \right)^{1/p} \\ & = o(r) \quad \text{by (15)} \\ & = o(|x - y|) \quad \text{by (16)}. \end{aligned}$$

Thus u is differentiable at x , and its gradient equals its weak gradient at x .

3. In case $p = \infty$, we note $W_{\text{loc}}^{1,\infty}(U) \subset W_{\text{loc}}^{1,p}(U)$ for all $1 \leq p < \infty$ and apply the reasoning above. \square

Finally, in view of Theorem 5, we obtain

THEOREM 6 (Rademacher's Theorem). *Let u be locally Lipschitz continuous in U . Then u is differentiable almost everywhere in U .*

5.8.4. Hardy's inequality.

In Chapter 12 we will need

THEOREM 7 (Hardy's inequality). *Assume $n \geq 3$ and $r > 0$. Suppose that $u \in H^1(B(0,r))$.*

Then $\frac{u}{|x|} \in L^2(B(0,r))$, with the estimate

$$(17) \quad \int_{B(0,r)} \frac{u^2}{|x|^2} dx \leq C \int_{B(0,r)} |Du|^2 + \frac{u^2}{r^2} dx.$$

Observe that this is not a consequence of the Gagliardo–Nirenberg–Sobolev inequality.

Proof. We may assume $u \in C^\infty(B(0, r))$. Note that $D\left(\frac{1}{|x|}\right) = -\frac{x}{|x|^3}$. Thus

$$\begin{aligned} \int_{B(0,r)} \frac{u^2}{|x|^2} dx &= - \int_{B(0,r)} u^2 D\left(\frac{1}{|x|}\right) \cdot \frac{x}{|x|} dx \\ &= \int_{B(0,r)} 2uD u \cdot \frac{x}{|x|^2} + (n-1) \frac{u^2}{|x|^2} dx - \int_{\partial B(0,r)} u^2 \nu \cdot \frac{x}{|x|^2} dS. \end{aligned}$$

Therefore

$$(2-n) \int_{B(0,r)} \frac{u^2}{|x|^2} dx = 2 \int_{B(0,r)} u D u \cdot \frac{x}{|x|^2} dx - \frac{1}{r} \int_{\partial B(0,r)} u^2 dS,$$

and consequently

$$(18) \quad \int_{B(0,r)} \frac{u^2}{|x|^2} dx \leq C \int_{B(0,r)} |D u|^2 dx + \frac{C}{r} \int_{\partial B(0,r)} u^2 dS.$$

Observe next that

$$\begin{aligned} r \int_{\partial B(0,r)} u^2 dS &= \int_{B(0,r)} \operatorname{div}(x u^2) dx = \int_{B(0,r)} n u^2 + 2u D u \cdot x dx \\ &\leq C \int_{B(0,r)} u^2 + r^2 |D u|^2 dx. \end{aligned}$$

Dividing by r^2 , we obtain the trace inequality

$$\frac{1}{r} \int_{\partial B(0,r)} u^2 dS \leq C \int_{B(0,r)} |D u|^2 + \frac{u^2}{r^2} dx.$$

Employing this inequality in (18) finishes the proof of (17). \square

5.8.5. Fourier transform methods.

Next we employ the Fourier transform (§4.3) to give an alternate characterization of the spaces $H^k(\mathbb{R}^n)$. For this subsection all functions are complex-valued.

THEOREM 8 (Characterization of H^k by Fourier transform).

Let k be a nonnegative integer.

(i) A function $u \in L^2(\mathbb{R}^n)$ belongs to $H^k(\mathbb{R}^n)$ if and only if

$$(19) \quad (1 + |y|^k) \hat{u} \in L^2(\mathbb{R}^n).$$

(ii) In addition, there exists a positive constant C such that

$$(20) \quad \frac{1}{C} \|u\|_{H^k(\mathbb{R}^n)} \leq \|(1 + |y|^k) \hat{u}\|_{L^2(\mathbb{R}^n)} \leq C \|u\|_{H^k(\mathbb{R}^n)}$$

for each $u \in H^k(\mathbb{R}^n)$.

Proof. 1. Assume first $u \in H^k(\mathbb{R}^n)$. Then for each multiindex $|\alpha| \leq k$, we have $D^\alpha u \in L^2(\mathbb{R}^n)$. Now if $u \in C^k$ has compact support, we have

$$(21) \quad \widehat{D^\alpha u} = (iy)^\alpha \hat{u}$$

according to Theorem 2 in §4.3.1. Approximating by smooth functions, we deduce formula (21) provided $u \in H^k(\mathbb{R}^n)$. Thus $(iy)^\alpha \hat{u} \in L^2(\mathbb{R}^n)$ for each $|\alpha| \leq k$. In particular choosing $\alpha = (k, 0, \dots, 0), (0, k, \dots, 0), \dots, (0, \dots, k)$, we deduce

$$\int_{\mathbb{R}^n} |y|^{2k} |\hat{u}|^2 dy \leq C \int_{\mathbb{R}^n} |D^k u|^2 dx < \infty.$$

Thus

$$\int_{\mathbb{R}^n} (1 + |y|^k)^2 |\hat{u}|^2 dy \leq C \|u\|_{H^k(\mathbb{R}^n)}^2,$$

and so $(1 + |y|^k) \hat{u} \in L^2(\mathbb{R}^n)$.

2. Suppose conversely $(1 + |y|^k) \hat{u} \in L^2(\mathbb{R}^n)$ and $|\alpha| \leq k$. Then

$$(22) \quad \|(iy)^\alpha \hat{u}\|_{L^2(\mathbb{R}^n)}^2 \leq \int_{\mathbb{R}^n} |y|^{2|\alpha|} |\hat{u}|^2 dy \leq C \|(1 + |y|^k) \hat{u}\|_{L^2(\mathbb{R}^n)}^2.$$

Set

$$u_\alpha := ((iy)^\alpha \hat{u})^\vee.$$

Then for each $\phi \in C_c^\infty(\mathbb{R}^n)$

$$\begin{aligned} \int_{\mathbb{R}^n} (D^\alpha \phi) \bar{u} dx &= \int_{\mathbb{R}^n} (\widehat{D^\alpha \phi}) \bar{\hat{u}} dy = \int_{\mathbb{R}^n} (iy)^\alpha \hat{\phi} \bar{\hat{u}} dy \\ &= (-1)^{|\alpha|} \int_{\mathbb{R}^n} \phi \bar{u}_\alpha dx. \end{aligned}$$

Thus $u_\alpha = D^\alpha u$ in the weak sense and, by (22), $D^\alpha u \in L^2(\mathbb{R}^n)$. Hence $u \in H^k(U)$, as required. \square

It is sometimes useful to define also *fractional* Sobolev spaces.

DEFINITION. Assume $0 < s < \infty$ and $u \in L^2(\mathbb{R}^n)$. Then $u \in H^s(\mathbb{R}^n)$ if $(1 + |y|^s) \hat{u} \in L^2(\mathbb{R}^n)$. For noninteger s , we set

$$\|u\|_{H^s(\mathbb{R}^n)} := \|(1 + |y|^s) \hat{u}\|_{L^2(\mathbb{R}^n)}.$$

5.9. OTHER SPACES OF FUNCTIONS

5.9.1. The space H^{-1} .

As we will see later in our systematic study in Chapters 6 and 7 of linear elliptic, parabolic and hyperbolic PDE, it is important to have an explicit characterization of the dual space of H_0^1 . (See Appendix D for definitions.)

DEFINITION. We denote by $H^{-1}(U)$ the dual space to $H_0^1(U)$.

In other words f belongs to $H^{-1}(U)$ provided f is a bounded linear functional on $H_0^1(U)$. Note very carefully that we do not identify the space H_0^1 with its dual. Instead, as we will see in a moment, we have

$$H_0^1(U) \subset L^2(U) \subset H^{-1}(U).$$

NOTATION. We will write $\langle \cdot, \cdot \rangle$ to denote the pairing between $H^{-1}(U)$ and $H_0^1(U)$.

DEFINITION. If $f \in H^{-1}(U)$, we define the norm

$$\|f\|_{H^{-1}(U)} := \sup \left\{ \langle f, u \rangle \mid u \in H_0^1(U), \|u\|_{H_0^1(U)} \leq 1 \right\}.$$

THEOREM 1 (Characterization of H^{-1}).

(i) Assume $f \in H^{-1}(U)$. Then there exist functions f^0, f^1, \dots, f^n in $L^2(U)$ such that

$$(1) \quad \langle f, v \rangle = \int_U f^0 v + \sum_{i=1}^n f^i v_{x_i} dx \quad (v \in H_0^1(U)).$$

(ii) Furthermore,

$$\|f\|_{H^{-1}(U)} = \inf \left\{ \left(\int_U \sum_{i=0}^n |f^i|^2 dx \right)^{1/2} \mid f \text{ satisfies (1) for } f^0, \dots, f^n \in L^2(U) \right\}.$$

(iii) In particular, we have

$$(2) \quad (v, u)_{L^2(U)} = \langle v, u \rangle$$

for all $u \in H_0^1(U)$, $v \in L^2(U) \subset H^{-1}(U)$.

NOTATION. We write " $f = f^0 - \sum_{i=1}^n f_{x_i}^i$ " whenever (1) holds.

Proof. 1. Given $u, v \in H_0^1(U)$, we define the inner product $(u, v) := \int_U Du \cdot Dv + uv \, dx$. Let $f \in H^{-1}(U)$. We apply the Riesz Representation Theorem (§D.3) to deduce the existence of a unique function $u \in H_0^1(U)$ satisfying $(u, v) = \langle f, v \rangle$ for all $v \in H_0^1(U)$; that is,

$$(3) \quad \int_U Du \cdot Dv + uv \, dx = \langle f, v \rangle$$

for each $v \in H_0^1(U)$. This establishes (1) for

$$(4) \quad \begin{cases} f^0 = u \\ f^i = u_{x_i} \quad (i = 1, \dots, n). \end{cases}$$

2. Assume now $f \in H^{-1}(U)$,

$$(5) \quad \langle f, v \rangle = \int_U g^0 v + \sum_{i=1}^n g^i v_{x_i} \, dx$$

for $g^0, g^1, \dots, g^n \in L^2(U)$. Setting $v = u$ in (3) and using (5), we deduce

$$\int_U |Du|^2 + |u|^2 \, dx \leq \int_U \sum_{i=0}^n |g^i|^2 \, dx.$$

Thus (4) implies

$$(6) \quad \int_U \sum_{i=0}^n |f^i|^2 \, dx \leq \int_U \sum_{i=0}^n |g^i|^2 \, dx.$$

3. From (1) it follows that

$$|\langle f, v \rangle| \leq \left(\int_U \sum_{i=0}^n |f^i|^2 \, dx \right)^{1/2}$$

if $\|v\|_{H_0^1(U)} \leq 1$. Consequently

$$\|f\|_{H^{-1}(U)} \leq \left(\int_U \sum_{i=0}^n |f^i|^2 \, dx \right)^{1/2}.$$

Setting $v = \frac{u}{\|u\|_{H_0^1(U)}}$ in (3), we deduce that in fact

$$\|f\|_{H^{-1}(U)} = \left(\int_U \sum_{i=0}^n |f^i|^2 \, dx \right)^{1/2}.$$

Assertion (ii) follows now from (4)–(6).

4. The identity (2) follows from assertion (i). □

5.9.2. Spaces involving time.

We study next some other sorts of Sobolev spaces, these comprising functions mapping time into Banach spaces. These will prove essential in our constructions of weak solutions to linear parabolic and hyperbolic PDE in Chapter 7 and to nonlinear parabolic PDE in Chapter 9.

Let X denote a real Banach space, with norm $\| \cdot \|$. The reader should first of all read §E.5 about measure and integration theory for mappings taking values in X .

DEFINITION. *The space*

$$L^p(0, T; X)$$

consists of all strongly measurable functions $\mathbf{u} : [0, T] \rightarrow X$ with

$$(i) \quad \|\mathbf{u}\|_{L^p(0, T; X)} := \left(\int_0^T \|\mathbf{u}(t)\|^p dt \right)^{1/p} < \infty$$

for $1 \leq p < \infty$ and

$$(ii) \quad \|\mathbf{u}\|_{L^\infty(0, T; X)} := \operatorname{ess\,sup}_{0 \leq t \leq T} \|\mathbf{u}(t)\| < \infty.$$

DEFINITION. *The space*

$$C([0, T]; X)$$

comprises all continuous functions $\mathbf{u} : [0, T] \rightarrow X$ with

$$\|\mathbf{u}\|_{C([0, T]; X)} := \max_{0 \leq t \leq T} \|\mathbf{u}(t)\| < \infty.$$

DEFINITION. *Let $\mathbf{u} \in L^1(0, T; X)$. We say $\mathbf{v} \in L^1(0, T; X)$ is the weak derivative of \mathbf{u} , written*

$$\mathbf{u}' = \mathbf{v},$$

provided

$$\int_0^T \phi'(t) \mathbf{u}(t) dt = - \int_0^T \phi(t) \mathbf{v}(t) dt$$

for all scalar test functions $\phi \in C_c^\infty(0, T)$.

DEFINITIONS. (i) *The Sobolev space*

$$W^{1,p}(0, T; X)$$

consists of all functions $\mathbf{u} \in L^p(0, T; X)$ such that \mathbf{u}' exists in the weak sense and belongs to $L^p(0, T; X)$. Furthermore,

$$\|\mathbf{u}\|_{W^{1,p}(0,T;X)} := \begin{cases} \left(\int_0^T \|\mathbf{u}(t)\|^p + \|\mathbf{u}'(t)\|^p dt \right)^{1/p} & (1 \leq p < \infty) \\ \text{ess sup}_{0 \leq t \leq T} (\|\mathbf{u}(t)\| + \|\mathbf{u}'(t)\|) & (p = \infty). \end{cases}$$

(ii) *We write $H^1(0, T; X) = W^{1,2}(0, T; X)$.*

THEOREM 2 (Calculus in an abstract space). *Let $\mathbf{u} \in W^{1,p}(0, T; X)$ for some $1 \leq p \leq \infty$. Then*

(i) $\mathbf{u} \in C([0, T]; X)$ (after possibly being redefined on a set of measure zero).

(ii) $\mathbf{u}(t) = \mathbf{u}(s) + \int_s^t \mathbf{u}'(\tau) d\tau$ for all $0 \leq s \leq t \leq T$.

(iii) *Furthermore, we have the estimate*

$$(7) \quad \max_{0 \leq t \leq T} \|\mathbf{u}(t)\| \leq C \|\mathbf{u}\|_{W^{1,p}(0,T;X)},$$

the constant C depending only on T .

Proof. 1. Extend \mathbf{u} to be $\mathbf{0}$ on $(-\infty, 0)$ and (T, ∞) , and then set $\mathbf{u}^\varepsilon = \eta_\varepsilon * \mathbf{u}$, η_ε denoting the usual mollifier on \mathbb{R}^1 . We check as in the proof of Theorem 1 in §5.3.1 that $(\mathbf{u}^\varepsilon)' = \eta_\varepsilon * \mathbf{u}'$ on $(\varepsilon, T - \varepsilon)$.

Then as $\varepsilon \rightarrow 0$,

$$(8) \quad \begin{cases} \mathbf{u}^\varepsilon \rightarrow \mathbf{u} & \text{in } L^p(0, T; X) \\ (\mathbf{u}^\varepsilon)' \rightarrow \mathbf{u}' & \text{in } L^p_{loc}(0, T; X). \end{cases}$$

Fixing $0 < s < t < T$, we compute

$$\mathbf{u}^\varepsilon(t) = \mathbf{u}^\varepsilon(s) + \int_s^t \mathbf{u}^{\varepsilon'}(\tau) d\tau.$$

Thus

$$(9) \quad \mathbf{u}(t) = \mathbf{u}(s) + \int_s^t \mathbf{u}'(\tau) d\tau$$

for a.e. $0 < s < t < T$, according to (8). As the mapping $t \mapsto \int_0^t \mathbf{u}'(\tau) d\tau$ is continuous, assertions (i), (ii) follow.

2. Estimate (7) follows easily from (9). □

The next two propositions concern what happens when \mathbf{u} and \mathbf{u}' lie in different spaces.

THEOREM 3 (More calculus). *Suppose $\mathbf{u} \in L^2(0, T; H_0^1(U))$, with $\mathbf{u}' \in L^2(0, T; H^{-1}(U))$.*

(i) *Then*

$$\mathbf{u} \in C([0, T]; L^2(U))$$

(after possibly being redefined on a set of measure zero).

(ii) *The mapping*

$$t \mapsto \|\mathbf{u}(t)\|_{L^2(U)}^2$$

is absolutely continuous, with

$$\frac{d}{dt} \|\mathbf{u}(t)\|_{L^2(U)}^2 = 2\langle \mathbf{u}'(t), \mathbf{u}(t) \rangle$$

for a.e. $0 \leq t \leq T$.

(iii) *Furthermore, we have the estimate*

$$(10) \quad \max_{0 \leq t \leq T} \|\mathbf{u}(t)\|_{L^2(U)} \leq C(\|\mathbf{u}\|_{L^2(0, T; H_0^1(U))} + \|\mathbf{u}'\|_{L^2(0, T; H^{-1}(U))}),$$

the constant C depending only on T .

Proof. 1. Extend \mathbf{u} to the larger interval $[-\sigma, T + \sigma]$ for $\sigma > 0$, and define the regularizations $\mathbf{u}^\varepsilon = \eta_\varepsilon * \mathbf{u}$, as in the earlier proof. Then for $\varepsilon, \delta > 0$,

$$\frac{d}{dt} \|\mathbf{u}^\varepsilon(t) - \mathbf{u}^\delta(t)\|_{L^2(U)}^2 = 2\langle \mathbf{u}^{\varepsilon'}(t) - \mathbf{u}^{\delta'}(t), \mathbf{u}^\varepsilon(t) - \mathbf{u}^\delta(t) \rangle.$$

Thus

$$(11) \quad \begin{aligned} \|\mathbf{u}^\varepsilon(t) - \mathbf{u}^\delta(t)\|_{L^2(U)}^2 &= \|\mathbf{u}^\varepsilon(s) - \mathbf{u}^\delta(s)\|_{L^2(U)}^2 \\ &+ 2 \int_s^t \langle \mathbf{u}^{\varepsilon'}(\tau) - \mathbf{u}^{\delta'}(\tau), \mathbf{u}^\varepsilon(\tau) - \mathbf{u}^\delta(\tau) \rangle d\tau \end{aligned}$$

for all $0 \leq s, t \leq T$. Fix any point $s \in (0, T)$ for which

$$\mathbf{u}^\varepsilon(s) \rightarrow \mathbf{u}(s) \quad \text{in } L^2(U).$$

Consequently (11) implies

$$\begin{aligned} \limsup_{\varepsilon, \delta \rightarrow 0} \sup_{0 \leq t \leq T} \|\mathbf{u}^\varepsilon(t) - \mathbf{u}^\delta(t)\|_{L^2(U)}^2 &\leq \lim_{\varepsilon, \delta \rightarrow 0} \int_0^T \|\mathbf{u}^{\varepsilon'}(\tau) - \mathbf{u}^{\delta'}(\tau)\|_{H^{-1}(U)}^2 \\ &\quad + \|\mathbf{u}^\varepsilon(\tau) - \mathbf{u}^\delta(\tau)\|_{H_0^1(U)}^2 d\tau \\ &= 0. \end{aligned}$$

Thus the smoothed functions $\{\mathbf{u}^\varepsilon\}_{0 < \varepsilon \leq 1}$ converge in $C([0, T]; L^2(U))$ to a limit $\mathbf{v} \in C([0, T]; L^2(U))$. Since we also know $\mathbf{u}^\varepsilon(t) \rightarrow \mathbf{u}(t)$ for a.e. t , we deduce $\mathbf{u} = \mathbf{v}$ a.e.

2. We similarly have

$$\|\mathbf{u}^\varepsilon(t)\|_{L^2(U)}^2 = \|\mathbf{u}^\varepsilon(s)\|_{L^2(U)}^2 + 2 \int_s^t \langle \mathbf{u}^{\varepsilon'}(\tau), \mathbf{u}^\varepsilon(\tau) \rangle d\tau,$$

and so, identifying \mathbf{u} with \mathbf{v} above,

$$(12) \quad \|\mathbf{u}(t)\|_{L^2(U)}^2 = \|\mathbf{u}(s)\|_{L^2(U)}^2 + 2 \int_s^t \langle \mathbf{u}'(\tau), \mathbf{u}(\tau) \rangle d\tau$$

for all $0 \leq s, t \leq T$.

3. To obtain (10), we integrate (12) with respect to s , recall the inequality $|\langle \mathbf{u}', \mathbf{u} \rangle| \leq \|\mathbf{u}'\|_{H^{-1}(U)} \|\mathbf{u}\|_{H_0^1(U)}$, and make some simple estimates. \square

For use later in the regularity theory for second-order parabolic and hyperbolic equations in Chapter 7, we will also need this extension of Theorem 3.

THEOREM 4 (Mappings into better spaces). *Assume that U is open, bounded, and ∂U is smooth. Take m to be a nonnegative integer.*

Suppose $\mathbf{u} \in L^2(0, T; H^{m+2}(U))$, with $\mathbf{u}' \in L^2(0, T; H^m(U))$.

(i) *Then*

$$\mathbf{u} \in C([0, T]; H^{m+1}(U))$$

(after possibly being redefined on a set of measure zero).

(ii) *Furthermore, we have the estimate*

$$(13) \quad \max_{0 \leq t \leq T} \|\mathbf{u}(t)\|_{H^{m+1}(U)} \leq C(\|\mathbf{u}\|_{L^2(0, T; H^{m+2}(U))} + \|\mathbf{u}'\|_{L^2(0, T; H^m(U))}),$$

the constant C depending only on T , U , and m .

Proof. 1. Suppose first that $m = 0$, in which case

$$\mathbf{u} \in L^2(0, T; H^2(U)), \quad \mathbf{u}' \in L^2(0, T; L^2(U)).$$

We select a bounded open set $V \supset \supset U$ and then construct a corresponding extension $\bar{\mathbf{u}} = E\mathbf{u}$, as in §5.4. In view of estimate (10) from that section, we see

$$\bar{\mathbf{u}} \in L^2(0, T; H^2(V))$$

and

$$(14) \quad \|\bar{\mathbf{u}}\|_{L^2(0,T;H^2(V))} \leq C\|\mathbf{u}\|_{L^2(0,T;H^2(U))},$$

for an appropriate constant C . In addition, $\bar{\mathbf{u}}' \in L^2(0,T;L^2(V))$, with the estimate

$$(15) \quad \|\bar{\mathbf{u}}'\|_{L^2(0,T;L^2(V))} \leq C\|\mathbf{u}'\|_{L^2(0,T;L^2(U))}.$$

This follows if we consider difference quotients in the t -variable, remember the methods in §5.8.2, and observe also that E is a bounded linear operator from $L^2(U)$ into $L^2(V)$.

2. Assume for the moment that $\bar{\mathbf{u}}$ is smooth. We then compute

$$\begin{aligned} \left| \frac{d}{dt} \left(\int_V |D\bar{\mathbf{u}}|^2 dx \right) \right| &= 2 \left| \int_V D\bar{\mathbf{u}} \cdot D\bar{\mathbf{u}}' dx \right| = 2 \left| \int_V \Delta \bar{\mathbf{u}} \bar{\mathbf{u}}' dx \right| \\ &\leq C(\|\bar{\mathbf{u}}\|_{H^2(V)}^2 + \|\bar{\mathbf{u}}'\|_{L^2(V)}^2). \end{aligned}$$

There is no boundary term when we integrate by parts, since the extension $\bar{\mathbf{u}} = E\mathbf{u}$ has compact support within V . Integrating and recalling (14), (15), it follows that

$$(16) \quad \max_{0 \leq t \leq T} \|\mathbf{u}(t)\|_{H^1(U)} \leq C(\|\mathbf{u}\|_{L^2(0,T;H^2(U))} + \|\mathbf{u}'\|_{L^2(0,T;L^2(U))}).$$

We obtain the same estimate if \mathbf{u} is not smooth, upon approximating by $\mathbf{u}^\varepsilon := \eta_\varepsilon * \mathbf{u}$, as before. As in the previous proofs, it also follows that $\mathbf{u} \in C([0, T]; H^1(U))$.

3. In the general case that $m \geq 1$, we let α be a multiindex of order $|\alpha| \leq m$ and set $\mathbf{v} := D^\alpha \mathbf{u}$. Then

$$\mathbf{v} \in L^2(0, T; H^2(U)), \quad \mathbf{v}' \in L^2(0, T; L^2(U)).$$

We apply estimate (16), with \mathbf{v} replacing \mathbf{u} , and sum over all indices $|\alpha| \leq m$, to derive (13). \square

5.10. PROBLEMS

In these exercises U always denotes an open subset of \mathbb{R}^n , with a smooth boundary ∂U . As usual, all given functions are assumed smooth, unless otherwise stated.

1. Suppose $k \in \{0, 1, \dots\}$, $0 < \gamma \leq 1$. Prove $C^{k,\gamma}(\bar{U})$ is a Banach space.

2. Assume $0 < \beta < \gamma \leq 1$. Prove the interpolation inequality

$$\|u\|_{C^{0,\gamma}(U)} \leq \|u\|_{C^{0,\beta}(U)}^{\frac{1-\gamma}{1-\beta}} \|u\|_{C^{0,1}(U)}^{\frac{\gamma-\beta}{1-\beta}}.$$

3. Denote by U the open square $\{x \in \mathbb{R}^2 \mid |x_1| < 1, |x_2| < 1\}$. Define

$$u(x) = \begin{cases} 1 - x_1 & \text{if } x_1 > 0, \quad |x_2| < x_1 \\ 1 + x_1 & \text{if } x_1 < 0, \quad |x_2| < -x_1 \\ 1 - x_2 & \text{if } x_2 > 0, \quad |x_1| < x_2 \\ 1 + x_2 & \text{if } x_2 < 0, \quad |x_1| < -x_2. \end{cases}$$

For which $1 \leq p \leq \infty$ does u belong to $W^{1,p}(U)$?

4. Assume $n = 1$ and $u \in W^{1,p}(0, 1)$ for some $1 \leq p < \infty$.
- (a) Show that u is equal a.e. to an absolutely continuous function and u' (which exists a.e.) belongs to $L^p(0, 1)$.
- (b) Prove that if $1 < p < \infty$, then

$$|u(x) - u(y)| \leq |x - y|^{1-\frac{1}{p}} \left(\int_0^1 |u'|^p dt \right)^{1/p}$$

for a.e. $x, y \in [0, 1]$.

5. Let U, V be open sets, with $V \subset\subset U$. Show there exists a smooth function ζ such that $\zeta \equiv 1$ on V , $\zeta = 0$ near ∂U . (Hint: Take $V \subset\subset W \subset\subset U$ and mollify χ_W .)
6. Assume U is bounded and $U \subset\subset \bigcup_{i=1}^N V_i$. Show there exist C^∞ functions ζ_i ($i = 1, \dots, N$) such that

$$\begin{cases} 0 \leq \zeta_i \leq 1, \text{ spt } \zeta_i \subset V_i & (i = 1, \dots, N) \\ \sum_{i=1}^N \zeta_i = 1 & \text{on } U. \end{cases}$$

The functions $\{\zeta_i\}_{i=1}^N$ form a *partition of unity*.

7. Assume that U is bounded and there exists a smooth vector field α such that $\alpha \cdot \nu \geq 1$ along ∂U , where ν as usual denotes the outward unit normal. Assume $1 \leq p < \infty$.

Apply the Gauss–Green Theorem to $\int_{\partial U} |u|^p \alpha \cdot \nu dS$, to derive a new proof of the trace inequality

$$\int_{\partial U} |u|^p dS \leq C \int_U |Du|^p + |u|^p dx$$

for all $u \in C^1(\bar{U})$.

8. Let U be bounded, with a C^1 boundary. Show that a “typical” function $u \in L^p(U)$ ($1 \leq p < \infty$) does not have a trace on ∂U . More precisely, prove there does not exist a bounded linear operator

$$T : L^p(U) \rightarrow L^p(\partial U)$$

such that $Tu = u|_{\partial U}$ whenever $u \in C(\bar{U}) \cap L^p(U)$.

9. Integrate by parts to prove the interpolation inequality:

$$\|Du\|_{L^2} \leq C \|u\|_{L^2}^{1/2} \|D^2u\|_{L^2}^{1/2}$$

for all $u \in C_c^\infty(U)$. Assume U is bounded, ∂U is smooth, and prove this inequality if $u \in H^2(U) \cap H_0^1(U)$.

(Hint: Take sequences $\{v_k\}_{k=1}^\infty \subset C_c^\infty(U)$ converging to u in $H_0^1(U)$ and $\{w_k\}_{k=1}^\infty \subset C^\infty(\bar{U})$ converging to u in $H^2(U)$.)

10. (a) Integrate by parts to prove

$$\|Du\|_{L^p} \leq C \|u\|_{L^p}^{1/2} \|D^2u\|_{L^p}^{1/2}$$

for $2 \leq p < \infty$ and all $u \in C_c^\infty(U)$.

(Hint: $\int_U |Du|^p dx = \sum_{i=1}^n \int_U u_{x_i} u_{x_i} |Du|^{p-2} dx$.)

- (b) Prove

$$\|Du\|_{L^{2p}} \leq C \|u\|_{L^\infty}^{1/2} \|D^2u\|_{L^p}^{1/2}$$

for $1 \leq p < \infty$ and all $u \in C_c^\infty(U)$.

11. Suppose U is connected and $u \in W^{1,p}(U)$ satisfies

$$Du = 0 \quad \text{a.e. in } U.$$

Prove u is constant a.e. in U .

12. Show by example that if we have $\|D^h u\|_{L^1(V)} \leq C$ for all $0 < |h| < \frac{1}{2} \text{dist}(V, \partial U)$, it does not necessarily follow that $u \in W^{1,1}(V)$.
13. Give an example of an open set $U \subset \mathbb{R}^n$ and a function $u \in W^{1,\infty}(U)$, such that u is *not* Lipschitz continuous on U . (Hint: Take U to be the open unit disk in \mathbb{R}^2 , with a slit removed.)
14. Verify that if $n > 1$, the unbounded function $u = \log \log \left(1 + \frac{1}{|x|}\right)$ belongs to $W^{1,n}(U)$, for $U = B^0(0, 1)$.
15. Fix $\alpha > 0$ and let $U = B^0(0, 1)$. Show there exists a constant C , depending only on n and α , such that

$$\int_U u^2 dx \leq C \int_U |Du|^2 dx,$$

provided

$$|\{x \in U \mid u(x) = 0\}| \geq \alpha, \quad u \in H^1(U).$$

16. (Variant of Hardy's inequality) Show that for each $n \geq 3$ there exists a constant C so that

$$\int_{\mathbb{R}^n} \frac{u^2}{|x|^2} dx \leq C \int_{\mathbb{R}^n} |Du|^2 dx$$

for all $u \in H^1(\mathbb{R}^n)$.

(Hint: $|Du + \lambda \frac{x}{|x|^2} u|^2 \geq 0$ for each $\lambda \in \mathbb{R}$.)

17. (Chain rule) Assume $F : \mathbb{R} \rightarrow \mathbb{R}$ is C^1 , with F' bounded. Suppose U is bounded and $u \in W^{1,p}(U)$ for some $1 \leq p \leq \infty$. Show

$$v := F(u) \in W^{1,p}(U) \quad \text{and} \quad v_{x_i} = F'(u)u_{x_i} \quad (i = 1, \dots, n).$$

18. Assume $1 \leq p \leq \infty$ and U is bounded.

- (a) Prove that if $u \in W^{1,p}(U)$, then $|u| \in W^{1,p}(U)$.
 (b) Prove $u \in W^{1,p}(U)$ implies $u^+, u^- \in W^{1,p}(U)$, and

$$Du^+ = \begin{cases} Du & \text{a.e. on } \{u > 0\} \\ 0 & \text{a.e. on } \{u \leq 0\}, \end{cases}$$

$$Du^- = \begin{cases} 0 & \text{a.e. on } \{u \geq 0\} \\ -Du & \text{a.e. on } \{u < 0\}. \end{cases}$$

(Hint: $u^+ = \lim_{\epsilon \rightarrow 0} F_\epsilon(u)$, for

$$F_\epsilon(z) := \begin{cases} (z^2 + \epsilon^2)^{1/2} - \epsilon & \text{if } z \geq 0 \\ 0 & \text{if } z < 0. \end{cases}$$

- (c) Prove that if $u \in W^{1,p}(U)$, then

$$Du = 0 \text{ a.e. on the set } \{u = 0\}.$$

19. Provide details for the following alternative proof that if $u \in H^1(U)$, then

$$Du = 0 \text{ a.e. on the set } \{u = 0\}.$$

Let ϕ be a smooth, bounded and nondecreasing function, such that ϕ' is bounded and $\phi(z) = z$ if $|z| \leq 1$. Set

$$u^\epsilon(x) := \epsilon \phi(u/\epsilon).$$

Show that $u^\epsilon \rightarrow 0$ weakly in $H^1(U)$ and therefore

$$\int_U Du^\epsilon \cdot Du \, dx = \int_U \phi'(u/\epsilon) |Du|^2 \, dx \rightarrow 0.$$

Employ this observation to finish the proof.

20. Use the Fourier transform to prove that if $u \in H^s(\mathbb{R}^n)$ for $s > n/2$, then $u \in L^\infty(\mathbb{R}^n)$, with the bound

$$\|u\|_{L^\infty(\mathbb{R}^n)} \leq C \|u\|_{H^s(\mathbb{R}^n)}$$

for a constant C depending only on s and n .

21. Show that if $u, v \in H^s(\mathbb{R}^n)$ for $s > n/2$, then $uv \in H^s(\mathbb{R}^n)$ and

$$\|uv\|_{H^s(\mathbb{R}^n)} \leq C \|u\|_{H^s(\mathbb{R}^n)} \|v\|_{H^s(\mathbb{R}^n)},$$

the constant C depending only on s and n .

5.11. REFERENCES

- Sections 5.2–8 See Gilbarg–Trudinger [G-T, Chapter 7], Lieb–Loss [L-L], Ziemer [Z] and [E-G] for more on Sobolev spaces.
- Section 5.5 W. Schlag showed me the proof of Theorem 2.
- Section 5.6 J. Ralston suggested an improvement in the proof of Theorem 4.
- Section 5.9 See Temam [Te, pp. 248–273].
- Section 5.10 Problem 16: see Tartar [Tr, Chapter 17]. H. Brezis taught me the trick in Problem 19.

This page is intentionally left blank.

SECOND-ORDER ELLIPTIC EQUATIONS

- 6.1 Definitions
- 6.2 Existence of weak solutions
- 6.3 Regularity
- 6.4 Maximum principles
- 6.5 Eigenvalues and eigenfunctions
- 6.6 Problems
- 6.7 References

This chapter investigates the solvability of uniformly elliptic, second-order partial differential equations, subject to prescribed boundary conditions. We will exploit two essentially distinct techniques, energy methods within Sobolev spaces (§§6.1–6.3) and maximum principle methods (§6.4).

6.1. DEFINITIONS

6.1.1. Elliptic equations.

We will in this chapter mostly study the boundary-value problem

$$(1) \quad \begin{cases} Lu = f & \text{in } U \\ u = 0 & \text{on } \partial U, \end{cases}$$

where U is an open, bounded subset of \mathbb{R}^n and $u : \bar{U} \rightarrow \mathbb{R}$ is the unknown, $u = u(x)$. Here $f : U \rightarrow \mathbb{R}$ is given, and L denotes a second-order partial

differential operator having either the form

$$(2) \quad Lu = - \sum_{i,j=1}^n (a^{ij}(x)u_{x_i})_{x_j} + \sum_{i=1}^n b^i(x)u_{x_i} + c(x)u$$

or else

$$(3) \quad Lu = - \sum_{i,j=1}^n a^{ij}(x)u_{x_i x_j} + \sum_{i=1}^n b^i(x)u_{x_i} + c(x)u,$$

for given coefficient functions a^{ij}, b^i, c ($i, j = 1, \dots, n$).

We say that the PDE $Lu = f$ is in *divergence form* if L is given by (2) and is in *nondivergence form* provided L is given by (3). The requirement that $u = 0$ on ∂U in (1) is sometimes called *Dirichlet's boundary condition*.

Remark. If the highest-order coefficients a^{ij} ($i, j = 1, \dots, n$) are C^1 functions, then an operator given in divergence form can be rewritten into non-divergence structure, and vice versa. Indeed the divergence form equation (2) becomes

$$(2') \quad Lu = - \sum_{i,j=1}^n a^{ij}(x)u_{x_i x_j} + \sum_{i=1}^n \tilde{b}^i(x)u_{x_i} + c(x)u$$

for $\tilde{b}^i := b^i - \sum_{j=1}^n a_{x_j}^{ij}$ ($i = 1, \dots, n$), and (2') is obviously in nondivergence form. We will see, however, that there are definite advantages to considering the two different representations of L separately. The divergence form is most natural for energy methods, based upon integration by parts (§§6.1–6.3), and the nondivergence form is most appropriate for maximum principle techniques (§6.4).

We henceforth assume as well the symmetry condition

$$a^{ij} = a^{ji} \quad (i, j = 1, \dots, n).$$

DEFINITION. We say the partial differential operator L is (uniformly) elliptic if there exists a constant $\theta > 0$ such that

$$(4) \quad \sum_{i,j=1}^n a^{ij}(x)\xi_i\xi_j \geq \theta|\xi|^2$$

for a.e. $x \in U$ and all $\xi \in \mathbb{R}^n$.

Ellipticity thus means that for each point $x \in U$, the symmetric $n \times n$ matrix $\mathbf{A}(x) = ((a^{ij}(x)))$ is positive definite, with smallest eigenvalue greater than or equal to θ .

An obvious example is $a^{ij} \equiv \delta_{ij}$, $b^i \equiv 0$, $c \equiv 0$, in which case the operator L is $-\Delta$. Indeed we will see that solutions of the general second-order elliptic PDE $Lu = 0$ are similar in many ways to harmonic functions. However, for these partial differential equations we do not have available the various explicit formulas developed for harmonic functions in Chapter 2: we must instead work directly with the PDE. Readers should continually be alert in the following calculations for uses of the structural condition of ellipticity (4).

Physical interpretation. As just noted, second-order elliptic PDE generalize Laplace's and Poisson's equations. As in the derivation of Laplace's equation set forth in §2.2, u in applications typically represents the density of some quantity, say a chemical concentration, at equilibrium within a region U . The second-order term $\mathbf{A} : D^2u = \sum_{i,j=1}^n a^{ij}u_{x_i x_j}$ represents the *diffusion* of u within U , the coefficients $((a^{ij}))$ describing the anisotropic, heterogeneous nature of the medium. In particular, $\mathbf{F} := -\mathbf{A}Du$ is the diffusive flux density, and the ellipticity condition implies

$$\mathbf{F} \cdot Du \leq 0;$$

that is, the flow is from regions of higher to lower concentration. The first-order term $\mathbf{b} \cdot Du = \sum_{i=1}^n b^i u_{x_i}$ represents *transport* within U , and the zeroth-order term cu describes the local *increase* or *depletion* of the chemical (owing, say, to reactions). A careful analysis of these interpretations requires the probabilistic study of diffusion processes.

Nonlinear second-order elliptic PDE also arise naturally in the calculus of variations (as the Euler–Lagrange equations of convex energy integrands) and in differential geometry (as expressions involving curvatures). We will encounter some such nonlinear equations later, in Chapters 8 and 9.

6.1.2. Weak solutions.

Let us consider first the boundary-value problem (1) when L has the divergence form (2). Our overall plan is first to define and then construct an appropriate weak solution u of (1) and only later to investigate the smoothness and other properties of u .

We will assume in the following exposition that

$$(5) \quad a^{ij}, b^i, c \in L^\infty(U) \quad (i, j = 1, \dots, n)$$

and

$$(6) \quad f \in L^2(U).$$

Motivation for definition of weak solution. How should we define a weak or generalized solution? Assuming for the moment u is really a smooth solution, let us multiply the PDE $Lu = f$ by a smooth test function $v \in C_c^\infty(U)$ and integrate over U , to find

$$(7) \quad \int_U \sum_{i,j=1}^n a^{ij} u_{x_i} v_{x_j} + \sum_{i=1}^n b^i u_{x_i} v + cuv \, dx = \int_U f v \, dx,$$

where we have integrated by parts in the first term on the left-hand side. There are no boundary terms since $v = 0$ on ∂U . By approximation we find the same identity holds with the smooth function v replaced by any $v \in H_0^1(U)$, and the resulting identity makes sense if only $u \in H_0^1(U)$. (We choose the space $H_0^1(U)$ to incorporate the boundary condition from (1) that “ $u = 0$ on ∂U ”.)

DEFINITIONS. (i) *The bilinear form $B[\ , \]$ associated with the divergence form elliptic operator L defined by (2) is*

$$(8) \quad B[u, v] := \int_U \sum_{i,j=1}^n a^{ij} u_{x_i} v_{x_j} + \sum_{i=1}^n b^i u_{x_i} v + cuv \, dx$$

for $u, v \in H_0^1(U)$.

(ii) *We say that $u \in H_0^1(U)$ is a weak solution of the boundary-value problem (1) if*

$$(9) \quad B[u, v] = (f, v)$$

for all $v \in H_0^1(U)$, where $(\ , \)$ denotes the inner product in $L^2(U)$.

The identity (9) is sometimes called the *variational formulation* of (1). This terminology will be explained later, in Example 2 of §8.1.2.

More generally, let us consider the boundary-value problem

$$(10) \quad \begin{cases} Lu = f^0 - \sum_{i=1}^n f_{x_i}^i & \text{in } U \\ u = 0 & \text{on } \partial U, \end{cases}$$

where L is defined by (2) and $f^i \in L^2(U)$ ($i = 0, \dots, n$). In view of the theory set forth in §5.9.1 we see that the right-hand term $f = f^0 - \sum_{i=1}^n f_{x_i}^i$ belongs to $H^{-1}(U)$, the dual space of $H_0^1(U)$.

DEFINITION. We say $u \in H_0^1(U)$ is a weak solution of problem (10) provided

$$B[u, v] = \langle f, v \rangle$$

for all $v \in H_0^1(U)$, where $\langle f, v \rangle = \int_U f^0 v + \sum_{i=1}^n f^i v_{x_i} dx$ and $\langle \cdot, \cdot \rangle$ is the pairing of $H^{-1}(U)$ and $H_0^1(U)$.

Other boundary conditions. We will hereafter, as above, focus our attention exclusively on the case of zero boundary conditions, but in fact a problem with prescribed, nonzero boundary values can easily be transformed into this setting. We spell this out by supposing now that ∂U is C^1 and $u \in H^1(U)$ is a weak solution of

$$\begin{cases} Lu = f & \text{in } U \\ u = g & \text{on } \partial U. \end{cases}$$

This means that $u = g$ on ∂U in the trace sense and furthermore that the bilinear form identity (9) holds for all $v \in H_0^1(U)$. For this to be possible, it is necessary for g to be the trace of some H^1 function, say w . But then $\tilde{u} := u - w$ belongs to $H_0^1(U)$ and is a weak solution of the boundary-value problem

$$\begin{cases} L\tilde{u} = \tilde{f} & \text{in } U \\ \tilde{u} = 0 & \text{on } \partial U, \end{cases}$$

where $\tilde{f} := f - Lw \in H^{-1}(U)$.

See Problems 3-6 to learn how to cast some other sorts of PDE and boundary conditions into weak formulations.

6.2. EXISTENCE OF WEAK SOLUTIONS

6.2.1. Lax–Milgram Theorem.

We now introduce a fairly simple abstract principle from linear functional analysis, which will later in §6.2.2 provide in certain circumstances the existence and uniqueness of a weak solution to our boundary-value problem.

We assume for this section H is a real Hilbert space, with norm $\| \cdot \|$ and inner product (\cdot, \cdot) . We let $\langle \cdot, \cdot \rangle$ denote the pairing of H with its dual space. Readers should review as necessary the basic Hilbert space theory described in §D.2–D.3.

THEOREM 1 (Lax–Milgram Theorem). *Assume that*

$$B : H \times H \rightarrow \mathbb{R}$$

is a bilinear mapping, for which there exist constants $\alpha, \beta > 0$ such that

$$(i) \quad |B[u, v]| \leq \alpha \|u\| \|v\| \quad (u, v \in H)$$

and

$$(ii) \quad \beta \|u\|^2 \leq B[u, u] \quad (u \in H).$$

Finally, let $f : H \rightarrow \mathbb{R}$ be a bounded linear functional on H .

Then there exists a unique element $u \in H$ such that

$$(1) \quad B[u, v] = \langle f, v \rangle$$

for all $v \in H$.

Proof. 1. For each fixed element $u \in H$, the mapping $v \mapsto B[u, v]$ is a bounded linear functional on H , whence the Riesz Representation Theorem (§D.3) asserts the existence of a unique element $w \in H$ satisfying

$$(2) \quad B[u, v] = (w, v) \quad (v \in H).$$

Let us write $Au = w$ whenever (2) holds, so that

$$(3) \quad B[u, v] = (Au, v) \quad (u, v \in H).$$

2. We first claim $A : H \rightarrow H$ is a bounded linear operator. Indeed if $\lambda_1, \lambda_2 \in \mathbb{R}$ and $u_1, u_2 \in H$, we see for each $v \in H$ that

$$\begin{aligned} (A(\lambda_1 u_1 + \lambda_2 u_2), v) &= B[\lambda_1 u_1 + \lambda_2 u_2, v] \quad \text{by (3)} \\ &= \lambda_1 B[u_1, v] + \lambda_2 B[u_2, v] \\ &= \lambda_1 (Au_1, v) + \lambda_2 (Au_2, v) \quad \text{by (3) again} \\ &= (\lambda_1 Au_1 + \lambda_2 Au_2, v). \end{aligned}$$

This equality obtains for each $v \in H$, and so A is linear. Furthermore

$$\|Au\|^2 = (Au, Au) = B[u, Au] \leq \alpha \|u\| \|Au\|.$$

Consequently $\|Au\| \leq \alpha \|u\|$ for all $u \in H$, and so A is bounded.

3. Next we assert

$$(4) \quad \begin{cases} A \text{ is one-to-one and} \\ R(A), \text{ the range of } A, \text{ is closed in } H. \end{cases}$$

To prove this, let us compute

$$\beta\|u\|^2 \leq B[u, u] = (Au, u) \leq \|Au\| \|u\|.$$

Hence $\beta\|u\| \leq \|Au\|$. This inequality easily implies (4).

4. We demonstrate now

$$(5) \quad R(A) = H.$$

For if not, then, since $R(A)$ is closed, there would exist a nonzero element $w \in H$ with $w \in R(A)^\perp$. But this fact in turn implies the contradiction $\beta\|w\|^2 \leq B[w, w] = (Aw, w) = 0$.

5. Next, we observe once more from the Riesz Representation Theorem that

$$\langle f, v \rangle = (w, v) \quad \text{for all } v \in H$$

for some element $w \in H$. We then utilize (4) and (5) to find $u \in H$ satisfying $Au = w$. Then

$$B[u, v] = (Au, v) = (w, v) = \langle f, v \rangle \quad (v \in H),$$

and this is (1).

6. Finally, we show there is at most one element $u \in H$ verifying (1). For if both $B[u, v] = \langle f, v \rangle$ and $B[\tilde{u}, v] = \langle f, v \rangle$, then $B[u - \tilde{u}, v] = 0$ ($v \in H$). We set $v = u - \tilde{u}$ to find $\beta\|u - \tilde{u}\|^2 \leq B[u - \tilde{u}, u - \tilde{u}] = 0$. \square

Remark. If the bilinear form $B[\ , \]$ is symmetric, that is, if

$$B[u, v] = B[v, u] \quad (u, v \in H),$$

we can fashion a much simpler proof by noting $((u, v)) := B[u, v]$ is a new inner product on H , to which the Riesz Representation Theorem directly applies. Consequently, *the Lax–Milgram Theorem is primarily significant in that it does not require symmetry of $B[\ , \]$.*

6.2.2. Energy estimates.

We return now to the specific bilinear form $B[\ , \]$, defined in §6.1.2 by the formula

$$B[u, v] = \int_U \sum_{i,j=1}^n a^{ij} u_{x_i} v_{x_j} + \sum_{i=1}^n b^i u_{x_i} v + cuv \, dx$$

for $u, v \in H_0^1(U)$, and try to verify the hypothesis of the Lax–Milgram Theorem.

THEOREM 2 (Energy estimates). *There exist constants $\alpha, \beta > 0$ and $\gamma \geq 0$ such that*

$$(i) \quad |B[u, v]| \leq \alpha \|u\|_{H_0^1(U)} \|v\|_{H_0^1(U)}$$

and

$$(ii) \quad \beta \|u\|_{H_0^1(U)}^2 \leq B[u, u] + \gamma \|u\|_{L^2(U)}^2$$

for all $u, v \in H_0^1(U)$.

Proof. 1. We readily check

$$\begin{aligned} |B[u, v]| &\leq \sum_{i,j=1}^n \|a^{ij}\|_{L^\infty} \int_U |Du| |Dv| dx \\ &\quad + \sum_{i=1}^n \|b^i\|_{L^\infty} \int_U |Du| |v| dx + \|c\|_{L^\infty} \int_U |u| |v| dx \\ &\leq \alpha \|u\|_{H_0^1(U)} \|v\|_{H_0^1(U)}, \end{aligned}$$

for some appropriate constant α .

2. Furthermore, in view of the ellipticity condition (4) from §6.1 we have

$$\begin{aligned} \theta \int_U |Du|^2 dx &\leq \int_U \sum_{i,j=1}^n a^{ij} u_{x_i} u_{x_j} dx \\ (6) \quad &= B[u, u] - \int_U \sum_{i=1}^n b^i u_{x_i} u + cu^2 dx \\ &\leq B[u, u] + \sum_{i=1}^n \|b^i\|_{L^\infty} \int_U |Du| |u| dx + \|c\|_{L^\infty} \int_U u^2 dx. \end{aligned}$$

Now from Cauchy's inequality with ε (§B.2), we observe

$$\int_U |Du| |u| dx \leq \varepsilon \int_U |Du|^2 dx + \frac{1}{4\varepsilon} \int_U u^2 dx \quad (\varepsilon > 0).$$

We insert this estimate into (6) and then choose $\varepsilon > 0$ so small that

$$\varepsilon \sum_{i=1}^n \|b^i\|_{L^\infty} < \frac{\theta}{2}.$$

Thus

$$\frac{\theta}{2} \int_U |Du|^2 dx \leq B[u, u] + C \int_U u^2 dx$$

for some appropriate constant C . In addition we recall from Poincaré's inequality in §5.6.1 that

$$\|u\|_{L^2(U)} \leq C\|Du\|_{L^2(U)}.$$

It easily follows that

$$\beta\|u\|_{H_0^1(U)}^2 \leq B[u, u] + \gamma\|u\|_{L^2(U)}^2$$

for appropriate constants $\beta > 0$, $\gamma \geq 0$. □

Observe now that if $\gamma > 0$ in these energy estimates, then $B[\cdot, \cdot]$ does not precisely satisfy the hypotheses of the Lax–Milgram Theorem. The following existence assertion for weak solutions must confront this possibility:

THEOREM 3 (First Existence Theorem for weak solutions). *There is a number $\gamma \geq 0$ such that for each*

$$(7) \quad \mu \geq \gamma$$

and each function

$$f \in L^2(U),$$

there exists a unique weak solution $u \in H_0^1(U)$ of the boundary-value problem

$$(8) \quad \begin{cases} Lu + \mu u = f & \text{in } U \\ u = 0 & \text{on } \partial U. \end{cases}$$

Proof. 1. Take γ from Theorem 2, let $\mu \geq \gamma$, and define then the bilinear form

$$B_\mu[u, v] := B[u, v] + \mu(u, v) \quad (u, v \in H_0^1(U)),$$

which corresponds as in §6.1 to the operator $L_\mu u := Lu + \mu u$. As before (\cdot, \cdot) means the inner product in $L^2(U)$. Then $B_\mu[\cdot, \cdot]$ satisfies the hypotheses of the Lax–Milgram Theorem.

2. Now fix $f \in L^2(U)$ and set $\langle f, v \rangle := (f, v)_{L^2(U)}$. This is a bounded linear functional on $L^2(U)$ and thus on $H_0^1(U)$.

We apply the Lax–Milgram Theorem to find a unique function $u \in H_0^1(U)$ satisfying

$$B_\mu[u, v] = \langle f, v \rangle$$

for all $v \in H_0^1(U)$; u is consequently the unique weak solution of (8). □

Mapping H_0^1 to H^{-1} . We can similarly show that for all

$$f^i \in L^2(U) \quad (i = 0, \dots, n),$$

there exists a unique weak solution u of the PDE

$$(9) \quad \begin{cases} Lu + \mu u = f^0 - \sum_{i=1}^n f_{x_i}^i & \text{in } U \\ u = 0 & \text{on } \partial U. \end{cases}$$

Indeed, it is enough to note $\langle f, v \rangle = \int_U f^0 v + \sum_{i=1}^n f^i v_{x_i} dx$ is a bounded linear functional on $H_0^1(U)$, as previously discussed in §5.9.1.

In particular, we deduce that the mapping

$$L_\mu := L + \mu I : H_0^1(U) \rightarrow H^{-1}(U) \quad (\mu \geq \gamma)$$

is an isomorphism.

Examples. In the case $Lu = -\Delta u$, so that $B[u, v] = \int_U Du \cdot Dv dx$, we easily check using Poincaré's inequality that Theorem 2 holds with $\gamma = 0$. A similar assertion holds for the general operator $Lu = -\sum_{i,j=1}^n (a^{ij} u_{x_i})_{x_j} + cu$, provided $c \geq 0$ in U . \square

6.2.3. Fredholm alternative.

We next employ the Fredholm theory for compact operators (discussed in §D.5) to glean more detailed information regarding the solvability of second-order elliptic PDE.

DEFINITIONS. (i) *The operator L^* , the formal adjoint of L , is*

$$L^*v := - \sum_{i,j=1}^n (a^{ij} v_{x_j})_{x_i} - \sum_{i=1}^n b^i v_{x_i} + (c - \sum_{i=1}^n b_{x_i}^i)v,$$

provided $b^i \in C^1(\bar{U})$ ($i = 1, \dots, n$).

(ii) *The adjoint bilinear form*

$$B^* : H_0^1(U) \times H_0^1(U) \rightarrow \mathbb{R}$$

is defined by

$$B^*[v, u] := B[u, v]$$

for all $u, v \in H_0^1(U)$.

(iii) *We say that $v \in H_0^1(U)$ is a weak solution of the adjoint problem*

$$\begin{cases} L^*v = f & \text{in } U \\ v = 0 & \text{on } \partial U, \end{cases}$$

provided

$$B^*[v, u] = (f, u)$$

for all $u \in H_0^1(U)$.

THEOREM 4 (Second Existence Theorem for weak solutions).

(i) *Precisely one of the following statements holds:
either*

$$(\alpha) \quad \left\{ \begin{array}{l} \text{for each } f \in L^2(U) \text{ there exists a unique} \\ \text{weak solution } u \text{ of the boundary-value problem} \\ (10) \quad \begin{cases} Lu = f & \text{in } U \\ u = 0 & \text{on } \partial U \end{cases} \end{array} \right.$$

or else

$$(\beta) \quad \left\{ \begin{array}{l} \text{there exists a weak solution } u \neq 0 \text{ of} \\ \text{the homogeneous problem} \\ (11) \quad \begin{cases} Lu = 0 & \text{in } U \\ u = 0 & \text{on } \partial U. \end{cases} \end{array} \right.$$

(ii) *Furthermore, should assertion (β) hold, the dimension of the subspace $N \subset H_0^1(U)$ of weak solutions of (11) is finite and equals the dimension of the subspace $N^* \subset H_0^1(U)$ of weak solutions of*

$$(12) \quad \begin{cases} L^*v = 0 & \text{in } U \\ v = 0 & \text{on } \partial U. \end{cases}$$

(iii) *Finally, the boundary-value problem (10) has a weak solution if and only if*

$$(f, v) = 0 \quad \text{for all } v \in N^*.$$

The dichotomy $(\alpha), (\beta)$ is the *Fredholm alternative*.

Proof. 1. Choose $\mu = \gamma$ as in Theorem 3 and define the bilinear form

$$B_\gamma[u, v] := B[u, v] + \gamma(u, v),$$

corresponding to the operator $L_\gamma u := Lu + \gamma u$. Then for each $g \in L^2(U)$ there exists a unique function $u \in H_0^1(U)$ solving

$$(13) \quad B_\gamma[u, v] = (g, v) \quad \text{for all } v \in H_0^1(U).$$

Let us write

$$(14) \quad u = L_\gamma^{-1}g$$

whenever (13) holds.

2. Observe next $u \in H_0^1(U)$ is a weak solution of (10) if and only if

$$(15) \quad B_\gamma[u, v] = (\gamma u + f, v) \quad \text{for all } v \in H_0^1(U),$$

that is, if and only if

$$(16) \quad u = L_\gamma^{-1}(\gamma u + f).$$

We rewrite this equality to read

$$(17) \quad u - Ku = h,$$

for

$$(18) \quad Ku := \gamma L_\gamma^{-1}u$$

and

$$(19) \quad h := L_\gamma^{-1}f.$$

3. We now claim $K : L^2(U) \rightarrow L^2(U)$ is a bounded, linear, compact operator. Indeed, from our choice of γ and the energy estimates from §6.2.2 we note that if (13) holds, then

$$\beta \|u\|_{H_0^1(U)}^2 \leq B_\gamma[u, u] = (g, u) \leq \|g\|_{L^2(U)} \|u\|_{L^2(U)} \leq \|g\|_{L^2(U)} \|u\|_{H_0^1(U)},$$

so that (18) implies

$$\|Kg\|_{H_0^1(U)} \leq C \|g\|_{L^2(U)} \quad (g \in L^2(U))$$

for some appropriate constant C . But since $H_0^1(U) \subset\subset L^2(U)$ according to the Rellich–Kondrachov compactness theorem (§5.7), we deduce that K is a compact operator.

4. We may consequently apply the Fredholm alternative from §D.5: either

$$(20) \quad (\alpha) \quad \begin{cases} \text{for each } h \in L^2(U) \text{ the equation} \\ u - Ku = h \\ \text{has a unique solution } u \in L^2(U) \end{cases}$$

or else

$$(21) \quad (\beta) \quad \begin{cases} \text{the equation} \\ u - Ku = 0 \\ \text{has nonzero solutions in } L^2(U). \end{cases}$$

Should assertion (α) hold, then according to (15)–(19) there exists a unique weak solution of problem (10). On the other hand, should assertion

(β) be valid, then necessarily $\gamma \neq 0$ and we recall further from §D.5 that the dimension of the space N of the solutions of (21) is finite and equals the dimension of the space N^* of solutions of

$$(22) \quad v - K^*v = 0.$$

We readily check however that (21) holds if and only if u is a weak solution of (11) and that (22) holds if and only if v is a weak solution of (12).

5. Finally, we recall (20) has a solution if and only if

$$(23) \quad (h, v) = 0$$

for all v solving (22). But from (18), (19) and (22) we compute

$$(h, v) = \frac{1}{\gamma}(Kf, v) = \frac{1}{\gamma}(f, K^*v) = \frac{1}{\gamma}(f, v).$$

Consequently the boundary-value problem (10) has a solution if and only if $(f, v) = 0$ for all weak solutions v of (12). \square

THEOREM 5 (Third Existence Theorem for weak solutions).

(i) *There exists an at most countable set $\Sigma \subset \mathbb{R}$ such that the boundary-value problem*

$$(24) \quad \begin{cases} Lu = \lambda u + f & \text{in } U \\ u = 0 & \text{on } \partial U \end{cases}$$

has a unique weak solution for each $f \in L^2(U)$ if and only if $\lambda \notin \Sigma$.

(ii) *If Σ is infinite, then $\Sigma = \{\lambda_k\}_{k=1}^{\infty}$, the values of a nondecreasing sequence with*

$$\lambda_k \rightarrow +\infty.$$

DEFINITION. *We call Σ the (real) spectrum of the operator L .*

Note in particular that the boundary-value problem

$$\begin{cases} Lu = \lambda u & \text{in } U \\ u = 0 & \text{on } \partial U \end{cases}$$

has a nontrivial solution $w \neq 0$ if and only if $\lambda \in \Sigma$, in which case λ is called an *eigenvalue* of L , w a corresponding *eigenfunction*. The partial differential equation $Lu = \lambda u$ for $L = -\Delta$ is sometimes called *Helmholtz's equation*.

Proof. 1. Let γ be the constant from Theorem 2 and assume

$$(25) \quad \lambda > -\gamma.$$

Assume also with no loss of generality that $\gamma > 0$.

2. According to the Fredholm alternative, the boundary-value problem (24) has a unique weak solution for each $f \in L^2(U)$ if and only if $u \equiv 0$ is the only weak solution of the homogeneous problem

$$\begin{cases} Lu = \lambda u & \text{in } U \\ u = 0 & \text{on } \partial U. \end{cases}$$

This is in turn true if and only if $u \equiv 0$ is the only weak solution of

$$(26) \quad \begin{cases} Lu + \gamma u = (\gamma + \lambda)u & \text{in } U \\ u = 0 & \text{on } \partial U. \end{cases}$$

Now (26) holds exactly when

$$(27) \quad u = L_\gamma^{-1}(\gamma + \lambda)u = \frac{\gamma + \lambda}{\gamma}Ku,$$

where, as in the proof of Theorem 4, we have set $Ku = \gamma L_\gamma^{-1}u$. Recall also from that proof that $K : L^2(U) \rightarrow L^2(U)$ is a bounded, linear, compact operator.

Now if $u \equiv 0$ is the only solution of (27), we see

$$(28) \quad \frac{\gamma}{\gamma + \lambda} \text{ is not an eigenvalue of } K.$$

Consequently we see the PDE (24) has a unique weak solution for each $f \in L^2(U)$ if and only if (28) holds.

3. According to Theorem 6 in §D.5 the collection of all eigenvalues of K comprises either a finite set or else the values of a sequence converging to zero. In the second case we see, according to (25) and (27), that the PDE (24) has a unique weak solution for all $f \in L^2(U)$, except for a sequence $\lambda_k \rightarrow +\infty$. \square

Finally, we explicitly note:

THEOREM 6 (Boundedness of the inverse). *If $\lambda \notin \Sigma$, there exists a constant C such that*

$$(29) \quad \|u\|_{L^2(U)} \leq C\|f\|_{L^2(U)},$$

whenever $f \in L^2(U)$ and $u \in H_0^1(U)$ is the unique weak solution of

$$\begin{cases} Lu = \lambda u + f & \text{in } U \\ u = 0 & \text{on } \partial U. \end{cases}$$

The constant C depends only on λ , U and the coefficients of L .

This constant will blow up if λ approaches an eigenvalue.

Proof. If not, there would exist sequences $\{f_k\}_{k=1}^\infty \subset L^2(U)$ and $\{u_k\}_{k=1}^\infty \subset H_0^1(U)$ such that

$$\begin{cases} Lu_k = \lambda u_k + f_k & \text{in } U \\ u_k = 0 & \text{on } \partial U \end{cases}$$

in the weak sense, but

$$\|u_k\|_{L^2(U)} > k \|f_k\|_{L^2(U)} \quad (k = 1, \dots).$$

As we may with no loss suppose $\|u_k\|_{L^2(U)} = 1$, we see $f_k \rightarrow 0$ in $L^2(U)$. According to the usual energy estimates the sequence $\{u_k\}_{k=1}^\infty$ is bounded in $H_0^1(U)$. Thus there exists a subsequence $\{u_{k_j}\}_{j=1}^\infty \subset \{u_k\}_{k=1}^\infty$ such that

$$(30) \quad \begin{cases} u_{k_j} \rightharpoonup u & \text{weakly in } H_0^1(U), \\ u_{k_j} \rightarrow u & \text{in } L^2(U). \end{cases}$$

(See §D.4 for weak convergence.) Then u is a weak solution of

$$\begin{cases} Lu = \lambda u & \text{in } U \\ u = 0 & \text{on } \partial U. \end{cases}$$

Since $\lambda \notin \Sigma$, $u \equiv 0$. However (30) implies as well that $\|u\|_{L^2(U)} = 1$, a contradiction. \square

Complex solutions. The foregoing theory extends easily to include complex-valued solutions. Given complex-valued $u, v \in H^1(U)$, write

$$(u, v)_{L^2(U)} := \int_U u \bar{v} \, dx, \quad (u, v)_{H^1(U)} := \int_U Du \cdot D\bar{v} + u \bar{v} \, dx,$$

and set

$$B[u, v] := \int_U \sum_{i,j=1}^n a^{ij} u_{x_i} \bar{v}_{x_j} + \sum_{i=1}^n b^i u_{x_i} \bar{v} + cu \bar{v} \, dx,$$

where $\bar{}$ denotes complex conjugate. We check

$$|B[u, v]| \leq \alpha \|u\|_{H_0^1(U)} \|v\|_{H_0^1(U)},$$

$$\beta \|u\|_{H_0^1(U)}^2 \leq \operatorname{Re} B[u, u] + \gamma \|u\|_{L^2(U)}^2 \quad (u, v \in H_0^1(U))$$

for appropriate constants $\alpha, \beta > 0$, $\gamma \geq 0$. Complex variants of the Lax–Milgram Theorem and Fredholm alternative lead to analogues of Theorems 3–6 above. \square

6.3. REGULARITY

We now address the question as to whether a weak solution u of the PDE

$$(1) \quad Lu = f \quad \text{in } U$$

is in fact smooth: this is the *regularity* problem for weak solutions.

Motivation: formal derivation of estimates. To see that there is some hope that a weak solution may be better than a typical function in $H_0^1(U)$, let us consider the model problem

$$(2) \quad -\Delta u = f \quad \text{in } \mathbb{R}^n.$$

We assume for heuristic purposes that u is smooth and vanishes sufficiently rapidly as $|x| \rightarrow \infty$ to justify the following calculations. We then compute

$$(3) \quad \begin{aligned} \int_{\mathbb{R}^n} f^2 dx &= \int_{\mathbb{R}^n} (\Delta u)^2 dx = \sum_{i,j=1}^n \int_{\mathbb{R}^n} u_{x_i x_i} u_{x_j x_j} dx \\ &= - \sum_{i,j=1}^n \int_{\mathbb{R}^n} u_{x_i x_i x_j} u_{x_j} dx \\ &= \sum_{i,j=1}^n \int_{\mathbb{R}^n} u_{x_i x_j} u_{x_i x_j} dx = \int_{\mathbb{R}^n} |D^2 u|^2 dx. \end{aligned}$$

Thus we see the L^2 -norm of the *second* derivatives of u can be estimated by (and in fact equals) the L^2 -norm of f . Similarly, we can differentiate the PDE (2), to find

$$-\Delta \tilde{u} = \tilde{f},$$

for $\tilde{u} := u_{x_k}$ and $\tilde{f} := f_{x_k}$ ($k = 1, \dots, n$). Applying the same method, we discover that the L^2 -norm of the third derivatives of u can be estimated by the first derivatives of f . Continuing, we see the L^2 -norm of the $(m+2)$ nd derivatives of u can be controlled by the L^2 -norm of the m th derivatives of f , for $m = 0, 1, \dots$ □

These computations suggest that for Poisson's equation (2), we can expect a weak solution $u \in H_0^1$ to belong to H^{m+2} whenever the inhomogeneous term f belongs to H^m ($m = 1, \dots$). Informally we say that u has "two more derivatives in L^2 than f has". This will be particularly interesting for $m = \infty$, in which case u belongs to H^m for all $m = 1, \dots$ and thus belongs to C^∞ .

Observe, however, the calculations above do not really constitute a proof. We assumed u was smooth, or at least say C^3 , in order to carry out the

calculation (3), whereas if we start with merely a weak solution in H_0^1 , we cannot immediately justify these computations. We will instead have to rely upon an analysis of certain difference quotients.

The following calculations are often technically difficult but eventually yield extremely powerful and useful assertions concerning the smoothness of weak solutions. As always, the heart of each computation is the invocation of ellipticity: *the point is to derive analytic estimates from the structural, algebraic assumption of ellipticity.*

6.3.1. Interior regularity.

We as always assume that $U \subset \mathbb{R}^n$ is a bounded, open set. Suppose also $u \in H_0^1(U)$ is a weak solution of the PDE (1), where L has the divergence form

$$(4) \quad Lu = - \sum_{i,j=1}^n (a^{ij}(x)u_{x_i})_{x_j} + \sum_{i=1}^n b^i(x)u_{x_i} + c(x)u.$$

We continue to require the uniform ellipticity condition from §6.1.1 and will as necessary make various additional assumptions about the smoothness of the coefficients a^{ij}, b^i, c .

THEOREM 1 (Interior H^2 -regularity). *Assume*

$$(5) \quad a^{ij} \in C^1(U), \quad b^i, c \in L^\infty(U) \quad (i, j = 1, \dots, n)$$

and

$$(6) \quad f \in L^2(U).$$

Suppose furthermore that $u \in H^1(U)$ is a weak solution of the elliptic PDE

$$Lu = f \quad \text{in } U.$$

Then

$$(7) \quad u \in H_{\text{loc}}^2(U) ;$$

and for each open subset $V \subset\subset U$ we have the estimate

$$(8) \quad \|u\|_{H^2(V)} \leq C (\|f\|_{L^2(U)} + \|u\|_{L^2(U)}),$$

the constant C depending only on V, U , and the coefficients of L .

Remarks. (i) Note carefully that we do not require $u \in H_0^1(U)$; that is, we are not necessarily assuming the boundary condition $u = 0$ on ∂U in the trace sense.

(ii) Observe additionally that since $u \in H_{\text{loc}}^2(U)$, we have

$$Lu = f \quad \text{a.e. in } U.$$

Thus u actually solves the PDE, at least for a.e. point within U . (To see this, note that for each $v \in C_c^\infty(U)$, we have

$$B[u, v] = (f, v).$$

Since $u \in H_{\text{loc}}^2(U)$, we can integrate by parts:

$$B[u, v] = (Lu, v).$$

Thus $(Lu - f, v) = 0$ for all $v \in C_c^\infty(U)$, and so $Lu = f$ a.e.)

Proof. 1. Fix any open set $V \subset\subset U$, and choose an open set W such that $V \subset\subset W \subset\subset U$. Then select a smooth function ζ satisfying

$$\begin{cases} \zeta \equiv 1 \text{ on } V, \zeta \equiv 0 \text{ on } \mathbb{R}^n - W, \\ 0 \leq \zeta \leq 1. \end{cases}$$

We call ζ a *cutoff* function. Its purpose in the subsequent calculations will be to restrict all expressions to the subset W , which is a positive distance away from ∂U . This is necessary as we have no information concerning the behavior of u near ∂U . (As an interesting technical point, notice carefully in the following calculations why we put “ ζ^2 ” and not just “ ζ ” in (11) below.)

2. Now since u is a weak solution of (1), we have $B[u, v] = (f, v)$ for all $v \in H_0^1(U)$. Consequently

$$(9) \quad \sum_{i,j=1}^n \int_U a^{ij} u_{x_i} v_{x_j} dx = \int_U \tilde{f} v dx,$$

where

$$(10) \quad \tilde{f} := f - \sum_{i=1}^n b^i u_{x_i} - cu.$$

3. Now let $|h| > 0$ be small, choose $k \in \{1, \dots, n\}$, and then substitute

$$(11) \quad v := -D_k^{-h}(\zeta^2 D_k^h u)$$

into (9), where as in §5.8.2 the expression $D_k^h u$ denotes the difference quotient

$$D_k^h u(x) = \frac{u(x + he_k) - u(x)}{h} \quad (h \in \mathbb{R}, h \neq 0).$$

We write the resulting expression as

$$(12) \quad A = B,$$

for

$$(13) \quad A := \sum_{i,j=1}^n \int_U a^{ij} u_{x_i} v_{x_j} dx$$

and

$$(14) \quad B := \int_U \tilde{f} v dx.$$

4. *Estimate of A.* We have

$$(15) \quad \begin{aligned} A &= - \sum_{i,j=1}^n \int_U a^{ij} u_{x_i} \left[D_k^{-h} \left(\zeta^2 D_k^h u \right) \right]_{x_j} dx \\ &= \sum_{i,j=1}^n \int_U D_k^h (a^{ij} u_{x_i}) \left(\zeta^2 D_k^h u \right)_{x_j} dx \\ &= \sum_{i,j=1}^n \int_U a^{ij,h} \left(D_k^h u_{x_i} \right) \left(\zeta^2 D_k^h u \right)_{x_j} \\ &\quad + \left(D_k^h a^{ij} \right) u_{x_i} \left(\zeta^2 D_k^h u \right)_{x_j} dx. \end{aligned}$$

Here we used the formulas

$$(16) \quad \int_U v D_k^{-h} w dx = - \int_U w D_k^h v dx$$

and

$$(17) \quad D_k^h(vw) = v^h D_k^h w + w D_k^h v,$$

for $v^h(x) := v(x + he_k)$.

Returning now to (15), we find

$$(18) \quad \begin{aligned} A &= \sum_{i,j=1}^n \int_U a^{ij,h} D_k^h u_{x_i} D_k^h u_{x_j} \zeta^2 dx \\ &\quad + \sum_{i,j=1}^n \int_U [a^{ij,h} D_k^h u_{x_i} D_k^h u_{x_j} 2\zeta \zeta_{x_j} + \left(D_k^h a^{ij} \right) u_{x_i} D_k^h u_{x_j} \zeta^2 \\ &\quad \quad \quad + \left(D_k^h a^{ij} \right) u_{x_i} D_k^h u_{x_j} 2\zeta \zeta_{x_j}] dx \\ &=: A_1 + A_2. \end{aligned}$$

The uniform ellipticity condition implies

$$(19) \quad A_1 \geq \theta \int_U \zeta^2 |D_k^h Du|^2 dx.$$

Furthermore we see from (5) that

$$|A_2| \leq C \int_U \zeta |D_k^h Du| |D_k^h u| + \zeta |D_k^h Du| |Du| + \zeta |D_k^h u| |Du| dx,$$

for some appropriate constant C . But then Cauchy's inequality with ϵ (§B.2) yields the bound

$$|A_2| \leq \epsilon \int_U \zeta^2 |D_k^h Du|^2 dx + \frac{C}{\epsilon} \int_W |D_k^h u|^2 + |Du|^2 dx.$$

We choose $\epsilon = \frac{\theta}{2}$ and further recall from Theorem 3(i) in §5.8.2 the estimate

$$\int_W |D_k^h u|^2 dx \leq C \int_U |Du|^2 dx,$$

thereby obtaining the inequality

$$|A_2| \leq \frac{\theta}{2} \int_U \zeta^2 |D_k^h Du|^2 dx + C \int_U |Du|^2 dx.$$

This estimate, (19) and (18) imply finally

$$(20) \quad A \geq \frac{\theta}{2} \int_U \zeta^2 |D_k^h Du|^2 dx - C \int_U |Du|^2 dx.$$

5. *Estimate of B.* Recalling now (10), (11), and (14), we estimate

$$(21) \quad |B| \leq C \int_U (|f| + |Du| + |u|)|v| dx.$$

Now Theorem 3(i) in §5.8.2 implies

$$\begin{aligned} \int_U |v|^2 dx &\leq C \int_U |D(\zeta^2 D_k^h u)|^2 dx \\ &\leq C \int_W |D_k^h u|^2 + \zeta^2 |D_k^h Du|^2 dx \\ &\leq C \int_U |Du|^2 + \zeta^2 |D_k^h Du|^2 dx. \end{aligned}$$

Thus (21) and Cauchy's inequality with ϵ imply

$$|B| \leq \epsilon \int_U \zeta^2 |D_k^h Du|^2 dx + \frac{C}{\epsilon} \int_U f^2 + u^2 dx + \frac{C}{\epsilon} \int_U |Du|^2 dx.$$

Select $\epsilon = \frac{\theta}{4}$, to obtain

$$(22) \quad |B| \leq \frac{\theta}{4} \int_U \zeta^2 |D_k^h Du|^2 dx + C \int_U f^2 + u^2 + |Du|^2 dx.$$

6. We finally combine (12), (20) and (22), to discover

$$\int_V |D_k^h Du|^2 dx \leq \int_U \zeta^2 |D_k^h Du|^2 dx \leq C \int_U f^2 + u^2 + |Du|^2 dx$$

for $k = 1, \dots, n$ and all sufficiently small $|h| \neq 0$.

In view of Theorem 3(ii) in §5.8.2, we deduce $Du \in H_{\text{loc}}^1(U; \mathbb{R}^n)$, and thus $u \in H_{\text{loc}}^2(U)$, with the estimate

$$(23) \quad \|u\|_{H^2(V)} \leq C (\|f\|_{L^2(U)} + \|u\|_{H^1(U)}).$$

7. We now refine estimate (23) by noting that if $V \subset\subset W \subset\subset U$, then the same argument shows

$$(24) \quad \|u\|_{H^2(V)} \leq C (\|f\|_{L^2(W)} + \|u\|_{H^1(W)}),$$

for an appropriate constant C depending on V, W , etc. Choose a new cutoff function ζ satisfying

$$\begin{cases} \zeta \equiv 1 \text{ on } W, \text{ spt } \zeta \subset U, \\ 0 \leq \zeta \leq 1. \end{cases}$$

Now set $v = \zeta^2 u$ in identity (9) and perform elementary calculations, to discover

$$\int_U \zeta^2 |Du|^2 dx \leq C \int_U f^2 + u^2 dx.$$

Thus

$$\|u\|_{H^1(W)} \leq C (\|f\|_{L^2(U)} + \|u\|_{L^2(U)}).$$

This inequality and (24) yield (8). □

Our intention next is to iterate the argument above, thereby deducing our weak solution lies in various higher Sobolev spaces (provided the coefficients are smooth enough and the right-hand side lies in sufficiently good spaces).

THEOREM 2 (Higher interior regularity). *Let m be a nonnegative integer, and assume*

$$(25) \quad a^{ij}, b^i, c \in C^{m+1}(U) \quad (i, j = 1, \dots, n)$$

and

$$(26) \quad f \in H^m(U).$$

Suppose $u \in H^1(U)$ is a weak solution of the elliptic PDE

$$Lu = f \quad \text{in } U.$$

Then

$$(27) \quad u \in H_{\text{loc}}^{m+2}(U);$$

and for each $V \subset\subset U$ we have the estimate

$$(28) \quad \|u\|_{H^{m+2}(V)} \leq C(\|f\|_{H^m(U)} + \|u\|_{L^2(U)}),$$

the constant C depending only on m, U, V and the coefficients of L .

Proof. 1. We will establish (27), (28) by induction on m , the case $m = 0$ being Theorem 1 above.

2. Assume now assertions (27) and (28) are valid for some nonnegative integer m and all open sets U , coefficients a^{ij}, b^i, c , etc., as above. Suppose then

$$(29) \quad a^{ij}, b^i, c \in C^{m+2}(U),$$

$$(30) \quad f \in H^{m+1}(U),$$

and $u \in H^1(U)$ is a weak solution of $Lu = f$ in U . By the induction hypotheses, we have

$$(31) \quad u \in H_{\text{loc}}^{m+2}(U),$$

with the estimate

$$(32) \quad \|u\|_{H^{m+2}(W)} \leq C(\|f\|_{H^m(U)} + \|u\|_{L^2(U)}),$$

for each $W \subset\subset U$ and an appropriate constant C , depending only on W , the coefficients of L , etc. Fix $V \subset\subset W \subset\subset U$.

3. Now let α be any multiindex with

$$(33) \quad |\alpha| = m + 1,$$

and choose any test function $\tilde{v} \in C_c^\infty(W)$. Insert

$$v := (-1)^{|\alpha|} D^\alpha \tilde{v}$$

into the identity $B[u, v] = (f, v)_{L^2(U)}$, and perform some integrations by parts, eventually to discover

$$(34) \quad B[\tilde{u}, \tilde{v}] = (\tilde{f}, \tilde{v})$$

for

$$(35) \quad \tilde{u} := D^\alpha u \in H^1(W)$$

and

$$(36) \quad \tilde{f} := D^\alpha f - \sum_{\substack{\beta \leq \alpha \\ \beta \neq \alpha}} \binom{\alpha}{\beta} \left[- \sum_{i,j=1}^n (D^{\alpha-\beta} a^{ij} D^\beta u_{x_i})_{x_j} + \sum_{i=1}^n D^{\alpha-\beta} b^i D^\beta u_{x_i} + D^{\alpha-\beta} c D^\beta u \right].$$

Since the identity (34) holds for each $\tilde{v} \in C_c^\infty(W)$, we see that \tilde{u} is a weak solution of

$$L\tilde{u} = \tilde{f} \quad \text{in } W.$$

In view of (29)–(32) and (36), we have $\tilde{f} \in L^2(W)$, with

$$(37) \quad \|\tilde{f}\|_{L^2(W)} \leq C(\|f\|_{H^{m+1}(U)} + \|u\|_{L^2(U)}).$$

4. In light of Theorem 1 then, we see $\tilde{u} \in H^2(V)$, with the estimate

$$\begin{aligned} \|\tilde{u}\|_{H^2(V)} &\leq C(\|\tilde{f}\|_{L^2(W)} + \|\tilde{u}\|_{L^2(W)}) \\ &\leq C(\|f\|_{H^{m+1}(U)} + \|u\|_{L^2(U)}). \end{aligned}$$

This inequality holds for each multiindex α with $|\alpha| = m + 1$ and $\tilde{u} = D^\alpha u$ as above. Consequently $u \in H^{m+3}(V)$, and

$$\|u\|_{H^{m+3}(V)} \leq C(\|f\|_{H^{m+1}(U)} + \|u\|_{L^2(U)}). \quad \square$$

We can now repeatedly apply Theorem 2 for $m = 0, 1, 2, \dots$ to deduce the infinite differentiability of u .

THEOREM 3 (Infinite differentiability in the interior). *Assume*

$$a^{ij}, b^i, c \in C^\infty(U) \quad (i, j = 1, \dots, n)$$

and

$$f \in C^\infty(U).$$

Suppose $u \in H^1(U)$ is a weak solution of the elliptic PDE

$$Lu = f \quad \text{in } U.$$

Then

$$u \in C^\infty(U).$$

We are again making no assumptions here about the behavior of u on ∂U . Therefore, in particular, we are asserting that any possible singularities of u on the boundary do not “propagate” into the interior.

Proof. According to Theorem 2, we have $u \in H_{\text{loc}}^m(U)$ for each integer $m = 1, 2, \dots$. Hence Theorem 6 in §5.6.3 implies $u \in C^k(U)$ for each $k = 1, 2, \dots$. \square

6.3.2. Boundary regularity.

Now we extend the estimates from §6.3.1 to study the smoothness of weak solutions up to the boundary.

THEOREM 4 (Boundary H^2 -regularity). *Assume*

$$(38) \quad a^{ij} \in C^1(\bar{U}), \quad b^i, c \in L^\infty(U) \quad (i, j = 1, \dots, n)$$

and

$$(39) \quad f \in L^2(U).$$

Suppose that $u \in H_0^1(U)$ is a weak solution of the elliptic boundary-value problem

$$(40) \quad \begin{cases} Lu = f & \text{in } U \\ u = 0 & \text{on } \partial U. \end{cases}$$

Assume finally

$$(41) \quad \partial U \text{ is } C^2.$$

Then

$$u \in H^2(U),$$

and we have the estimate

$$(42) \quad \|u\|_{H^2(U)} \leq C(\|f\|_{L^2(U)} + \|u\|_{L^2(U)}),$$

the constant C depending only on U and the coefficients of L .

Remarks. (i) If $u \in H_0^1(U)$ is the *unique* weak solution of (40), estimate (42) simplifies to read

$$\|u\|_{H^2(U)} \leq C\|f\|_{L^2(U)}.$$

This follows from Theorem 6 in §6.2.

(ii) Observe also that in contrast to Theorem 1 in §6.3.1, we are now assuming $u = 0$ along ∂U (in the trace sense).

Proof. 1. We first investigate the special case that U is a half-ball:

$$(43) \quad U = B^0(0, 1) \cap \mathbb{R}_+^n.$$

Set $V := B^0(0, \frac{1}{2}) \cap \mathbb{R}_+^n$. Then select a smooth cutoff function ζ satisfying

$$\begin{cases} \zeta \equiv 1 \text{ on } B(0, \frac{1}{2}), \quad \zeta \equiv 0 \text{ on } \mathbb{R}^n - B(0, 1), \\ 0 \leq \zeta \leq 1. \end{cases}$$

So $\zeta \equiv 1$ on V and ζ vanishes near the *curved* part of ∂U .

2. Since u is a weak solution of (40), we have $B[u, v] = (f, v)$ for all $v \in H_0^1(U)$; consequently

$$(44) \quad \sum_{i,j=1}^n \int_U a^{ij} u_{x_i} v_{x_j} dx = \int_U \tilde{f} v dx,$$

for

$$(45) \quad \tilde{f} := f - \sum_{i=1}^n b^i u_{x_i} - cu.$$

3. Now let $h > 0$ be small, choose $k \in \{1, \dots, n-1\}$, and write

$$v := -D_k^{-h}(\zeta^2 D_k^h u).$$

Let us note carefully

$$\begin{aligned} v(x) &= -\frac{1}{h} D_k^{-h}(\zeta^2(x)[u(x + he_k) - u(x)]) \\ &= \frac{1}{h^2}(\zeta^2(x - he_k)[u(x) - u(x - he_k)] \\ &\quad - \zeta^2(x)[u(x + he_k) - u(x)]) \end{aligned}$$

if $x \in U$. Now since $u = 0$ along $\{x_n = 0\}$ in the trace sense and $\zeta \equiv 0$ near the curved portion of ∂U , we see $v \in H_0^1(U)$.

We may therefore substitute v into the identity (44) and write the resulting expression as

$$(46) \quad A = B,$$

for

$$(47) \quad A := \sum_{i,j=1}^n \int_U a^{ij} u_{x_i} v_{x_j} dx$$

and

$$(48) \quad B := \int_U \tilde{f} v dx.$$

4. We can now estimate the terms A and B in almost exactly the same way that we estimated their counterparts in the proof of Theorem 1. After some calculations we find

$$(49) \quad A \geq \frac{\theta}{2} \int_U \zeta^2 |D_k^h Du|^2 dx - C \int_U |Du|^2 dx$$

and

$$(50) \quad |B| \leq \frac{\theta}{4} \int_U \zeta^2 |D_k^h Du|^2 dx + C \int_U f^2 + u^2 + |Du|^2 dx,$$

for appropriate constants C . We then combine (46), (49), and (50) to discover

$$\int_V |D_k^h Du|^2 dx \leq C \int_U f^2 + u^2 + |Du|^2 dx$$

for $k = 1, \dots, n-1$. Thus recalling the remark after the proof of Theorem 3 in §5.8.2, we deduce

$$u_{x_k} \in H^1(V) \quad (k = 1, \dots, n-1),$$

with the estimate

$$(51) \quad \sum_{\substack{k,l=1 \\ k+l < 2n}}^n \|u_{x_k x_l}\|_{L^2(V)} \leq C(\|f\|_{L^2(U)} + \|u\|_{H^1(U)}).$$

5. We must now augment (51) with an estimate of the L^2 -norm of $u_{x_n x_n}$ over V . For this we recall from the Remarks after Theorem 1 that $Lu = f$

a.e. in U . Remembering the definition of L , we can rewrite this equality into nondivergence form, as

$$(52) \quad - \sum_{i,j=1}^n a^{ij} u_{x_i x_j} + \sum_{i=1}^n \tilde{b}^i u_{x_i} + cu = f,$$

for $\tilde{b}^i := b^i - \sum_{j=1}^n a_{x_j}^{ij}$ ($i = 1, \dots, n$). So we discover

$$(53) \quad a^{nn} u_{x_n x_n} = - \sum_{\substack{i,j=1 \\ i+j < 2n}}^n a^{ij} u_{x_i x_j} + \sum_{i=1}^n \tilde{b}^i u_{x_i} + cu - f.$$

Now according to the uniform ellipticity condition, $\sum_{i,j=1}^n a^{ij}(x) \xi_i \xi_j \geq \theta |\xi|^2$ for all $x \in U$, $\xi \in \mathbb{R}^n$. We set $\xi = e_n = (0, \dots, 0, 1)$, to conclude

$$(54) \quad a^{nn}(x) \geq \theta > 0$$

for all $x \in U$. But then (38), (53) and (54) imply

$$(55) \quad |u_{x_n x_n}| \leq C \left(\sum_{\substack{i,j=1 \\ i+j < 2n}}^n |u_{x_i x_j}| + |Du| + |u| + |f| \right)$$

in U . Utilizing this estimate in inequality (51), we conclude $u \in H^2(V)$ and

$$(56) \quad \|u\|_{H^2(V)} \leq C(\|f\|_{L^2(U)} + \|u\|_{L^2(U)})$$

for some appropriate constant C .

6. We now drop the assumption that U is a half-ball and so has the special form (43). In the general case we choose any point $x^0 \in \partial U$ and note that since ∂U is C^2 , we may assume—upon relabeling the coordinate axes if needs be—that

$$U \cap B(x^0, r) = \{x \in B(x^0, r) \mid x_n > \gamma(x_1, \dots, x_{n-1})\}$$

for some $r > 0$ and some C^2 function $\gamma : \mathbb{R}^{n-1} \rightarrow \mathbb{R}$. As usual, we change variables utilizing §C.1 and write

$$(57) \quad y = \Phi(x), \quad x = \Psi(y).$$

7. Choose $s > 0$ so small that the half-ball $U' := B^0(0, s) \cap \{y_n > 0\}$ lies in $\Phi(U \cap B(x^0, r))$. Set $V' := B^0(0, s/2) \cap \{y_n > 0\}$. Finally define

$$(58) \quad u'(y) := u(\Psi(y)) \quad (y \in U').$$

It is straightforward to check

$$(59) \quad u' \in H^1(U')$$

and

$$(60) \quad u' = 0 \quad \text{on } \partial U' \cap \{y_n = 0\}$$

in the trace sense.

8. We now claim u' is a weak solution of the PDE

$$(61) \quad L'u' = f' \quad \text{in } U',$$

for

$$(62) \quad f'(y) := f(\Psi(y))$$

and

$$(63) \quad L'u' := - \sum_{k,l=1}^n (a'^{kl} u'_{y_k})_{y_l} + \sum_{k=1}^n b'^k u'_{y_k} + c'u',$$

where

$$(64) \quad a'^{kl}(y) := \sum_{r,s=1}^n a^{rs}(\Psi(y)) \Phi_{x_r}^k(\Psi(y)) \Phi_{x_s}^l(\Psi(y)) \quad (k, l = 1, \dots, n),$$

$$(65) \quad b'^k(y) := \sum_{r=1}^n b^r(\Psi(y)) \Phi_{x_r}^k(\Psi(y)) \quad (k = 1, \dots, n),$$

and

$$(66) \quad c'(y) := c(\Psi(y))$$

for $y \in U'$, $k, l = 1, \dots, n$.

If $v' \in H_0^1(U')$ and $B'[\cdot, \cdot]$ denotes the bilinear form associated with the operator L' , we have

$$(67) \quad B'[u', v'] = \int_{U'} \sum_{k,l=1}^n a'^{kl} u'_{y_k} v'_{y_l} + \sum_{k=1}^n b'^k u'_{y_k} v' + c'u'v' dy.$$

Now define

$$v(x) := v'(\Phi(x)).$$

Then from (67) we calculate

$$(68) \quad \begin{aligned} B'[u', v'] &= \sum_{i,j=1}^n \sum_{k,l=1}^n \int_{U'} a'^{kl} u_{x_i} \Psi_{y_k}^i v_{x_j} \Psi_{y_l}^j dy \\ &+ \sum_{i=1}^n \sum_{k=1}^n \int_{U'} b'^k u_{x_i} \Psi_{y_k}^i v dy + \int_{U'} c' uv dy. \end{aligned}$$

Now according to (64), we find for each $i, j = 1, \dots, n$ that

$$\sum_{k,l=1}^n a'^{kl} \Psi_{y_k}^i \Psi_{y_l}^j = \sum_{r,s=1}^n \sum_{k,l=1}^n a^{rs} \Phi_{x_r}^k \Phi_{x_s}^l \Psi_{y_k}^i \Psi_{y_l}^j = a^{ij},$$

since $D\Phi = (D\Psi)^{-1}$. Similarly for $i = 1, \dots, n$, we have

$$\sum_{k=1}^n b'^k \Psi_{y_k}^i = \sum_{k=1}^n \sum_{r=1}^n b^r \Phi_{x_r}^k \Psi_{y_k}^i = b^i.$$

Substituting these calculations into (68) and changing variables yields, since $|\det D\Phi| = 1$,

$$\begin{aligned} B'[u', v'] &= \int_U \sum_{i,j=1}^n a^{ij} u_{x_i} v_{x_j} + \sum_{i=1}^n b^i u_{x_i} v + cuv dx \\ &= B[u, v] = (f, v)_{L^2(U)} = (f', v')_{L^2(U')}. \end{aligned}$$

This establishes (61).

9. We now check that *the operator L' is uniformly elliptic in U'* . Indeed if $y \in U'$ and $\xi \in \mathbb{R}^n$, we note that

$$(69) \quad \begin{aligned} \sum_{k,l=1}^n a'^{kl}(y) \xi_k \xi_l &= \sum_{r,s=1}^n \sum_{k,l=1}^n a^{rs}(\Psi(y)) \Phi_{x_r}^k \Phi_{x_s}^l \xi_k \xi_l \\ &= \sum_{r,s=1}^n a^{rs}(\Psi(y)) \eta_r \eta_s \geq \theta |\eta|^2, \end{aligned}$$

where $\eta = \xi D\Phi$; that is, $\eta_r = \sum_{k=1}^n \Phi_{x_r}^k \xi_k$ ($r = 1, \dots, n$). But then, since $D\Phi D\Psi = I$, we have $\xi = \eta D\Psi$; and so $|\xi| \leq C|\eta|$ for some constant C . This inequality and (69) imply

$$(70) \quad \sum_{k,l=1}^n a'^{kl}(y) \xi_k \xi_l \geq \theta' |\xi|^2$$

for some $\theta' > 0$ and all $y \in U'$, $\xi \in \mathbb{R}^n$.

Observe also from (64) that the coefficients a'^{kl} are C^1 , since Φ and Ψ are C^2 .

10. In view of (61) and (70), we may apply the results from steps 1–5 in the proof above to ascertain that $u' \in H^2(V')$, with the bound

$$\|u'\|_{H^2(V')} \leq C(\|f'\|_{L^2(U')} + \|u'\|_{L^2(U')}).$$

Consequently

$$(71) \quad \|u\|_{H^2(V)} \leq C(\|f\|_{L^2(U)} + \|u\|_{L^2(U)})$$

for $V := \Psi(V')$.

Since ∂U is compact, we can as usual cover ∂U with finitely many sets V_1, \dots, V_N as above. We sum the resulting estimates, along with the interior estimate, to find $u \in H^2(U)$, with the inequality (42). \square

Now we derive higher regularity for our weak solutions, all the way up to ∂U .

THEOREM 5 (Higher boundary regularity). *Let m be a nonnegative integer, and assume*

$$(72) \quad a^{ij}, b^i, c \in C^{m+1}(\bar{U}) \quad (i, j = 1, \dots, n)$$

and

$$(73) \quad f \in H^m(U).$$

Suppose that $u \in H_0^1(U)$ is a weak solution of the boundary-value problem

$$(74) \quad \begin{cases} Lu = f & \text{in } U \\ u = 0 & \text{on } \partial U. \end{cases}$$

Assume finally

$$(75) \quad \partial U \text{ is } C^{m+2}.$$

Then

$$(76) \quad u \in H^{m+2}(U),$$

and we have the estimate

$$(77) \quad \|u\|_{H^{m+2}(U)} \leq C(\|f\|_{H^m(U)} + \|u\|_{L^2(U)}),$$

the constant C depending only on m, U and the coefficients of L .

Remark. If u is the *unique* solution of (74), then estimate (77) simplifies to read

$$\|u\|_{H^{m+2}(U)} \leq C \|f\|_{H^m(U)}.$$

Proof. 1. We first investigate the special case

$$(78) \quad U := B^0(0, s) \cap \mathbb{R}_+^n$$

for some $s > 0$. Fix $0 < t < s$ and set $V := B^0(0, t) \cap \mathbb{R}_+^n$.

2. We intend to prove by induction on m that whenever $u = 0$ along $\{x_n = 0\}$ in the trace sense, (72) and (73) imply

$$(79) \quad u \in H^{m+2}(V),$$

with the estimate

$$(80) \quad \|u\|_{H^{m+2}(V)} \leq C (\|f\|_{H^m(U)} + \|u\|_{L^2(U)}),$$

for a constant C depending only on U, V and the coefficients of L . The case $m = 0$ follows as in the proof of Theorem 4 above.

Suppose then

$$(81) \quad a^{ij}, b^i, c \in C^{m+2}(\bar{U}),$$

$$(82) \quad f \in H^{m+1}(U),$$

and u is a weak solution of $Lu = f$ in U , which vanishes in the trace sense along $\{x_n = 0\}$. Fix any $0 < t < r < s$, and write $W := B^0(0, r) \cap \mathbb{R}_+^n$. By the induction assumption we have

$$(83) \quad u \in H^{m+2}(W),$$

with the estimate

$$(84) \quad \|u\|_{H^{m+2}(W)} \leq C (\|f\|_{H^m(U)} + \|u\|_{L^2(U)}).$$

Furthermore according to the interior regularity Theorem 2, $u \in H_{\text{loc}}^{m+3}(U)$.

3. Next, let α be any multiindex with

$$(85) \quad |\alpha| = m + 1$$

and

$$(86) \quad \alpha_n = 0.$$

Then

$$(87) \quad \tilde{u} := D^\alpha u$$

belongs to $H^1(U)$ and vanishes along the plane $\{x_n = 0\}$ in the trace sense. Furthermore, as in the proof of Theorem 2, \tilde{u} is a weak solution of $L\tilde{u} = \tilde{f}$ in U , for

$$\begin{aligned} \tilde{f} := D^\alpha f - \sum_{\substack{\beta \leq \alpha \\ \beta \neq \alpha}} \binom{\alpha}{\beta} \left[\sum_{i,j=1}^n - \left(D^{\alpha-\beta} a^{ij} D^\beta u_{x_i} \right)_{x_j} \right. \\ \left. + \sum_{i=1}^n D^{\alpha-\beta} b^i D^\beta u_{x_i} + D^{\alpha-\beta} c D^\beta u \right]. \end{aligned}$$

In view of (72), (73), (82) and (84), we see $\tilde{f} \in L^2(W)$, with

$$(88) \quad \|\tilde{f}\|_{L^2(W)} \leq C (\|f\|_{H^{m+1}(U)} + \|u\|_{L^2(U)}).$$

Consequently the proof of Theorem 4 shows $\tilde{u} \in H^2(V)$, with the estimate

$$\begin{aligned} \|\tilde{u}\|_{H^2(V)} &\leq C (\|\tilde{f}\|_{L^2(W)} + \|\tilde{u}\|_{L^2(W)}) \\ &\leq C (\|f\|_{H^{m+1}(U)} + \|u\|_{L^2(U)}). \end{aligned}$$

In light of (85)–(88), we thus deduce

$$(89) \quad \|D^\beta u\|_{L^2(V)} \leq C (\|f\|_{H^{m+1}(U)} + \|u\|_{L^2(U)})$$

for any multiindex β with $|\beta| = m + 3$ and

$$(90) \quad \beta_n = 0, 1, \text{ or } 2.$$

4. We must extend estimate (89) to remove the restriction (90). For this, let us suppose by induction

$$(91) \quad \|D^\beta u\|_{L^2(V)} \leq C (\|f\|_{H^{m+1}(U)} + \|u\|_{L^2(U)})$$

for any multiindex β with $|\beta| = m + 3$ and

$$(92) \quad \beta_n = 0, 1, \dots, j,$$

for some $j \in \{2, \dots, m + 2\}$. Assume then $|\beta| = m + 3$,

$$(93) \quad \beta_n = j + 1.$$

Let us write $\beta = \gamma + \delta$, for $\delta = (0, \dots, 2)$ and $|\gamma| = m + 1$. Since $u \in H_{\text{loc}}^{m+3}(U)$ and $Lu = f$ in U , we have $D^\gamma Lu = D^\gamma f$ a.e. in U . Now

$$D^\gamma Lu = a^{nn} D^\beta u + \left\{ \begin{array}{l} \text{sum of terms involving at most } j \\ \text{derivatives of } u \text{ with respect to } x_n \text{ and} \\ \text{at most } m + 3 \text{ derivatives in all } \end{array} \right\}.$$

Since $a^{nn} \geq \theta > 0$, we thus find by utilizing (91), (92) that

$$(94) \quad \|D^\beta u\|_{L^2(V)} \leq C (\|f\|_{H^{m+1}(U)} + \|u\|_{L^2(U)})$$

provided $|\beta| = m + 3$ and $\beta_n = j + 1$. By induction on j then, we have

$$\|u\|_{H^{m+3}(U)} \leq C (\|f\|_{H^{m+1}(U)} + \|u\|_{L^2(U)}).$$

This estimate in turn completes the induction on m , begun in step 2.

5. We have now shown that (72) and (73) imply (79) and (80), provided U has the form (78). The general case follows once we straighten out the boundary, using the ideas explained in the proof of Theorem 4. \square

We finally iterate the foregoing estimates to obtain

THEOREM 6 (Infinite differentiability up to the boundary). *Assume*

$$a^{ij}, b^i, c \in C^\infty(\bar{U}) \quad (i, j = 1, \dots, n)$$

and

$$f \in C^\infty(\bar{U}).$$

Suppose $u \in H_0^1(U)$ is a weak solution of the boundary-value problem

$$\begin{cases} Lu = f & \text{in } U \\ u = 0 & \text{on } \partial U. \end{cases}$$

Assume also that ∂U is C^∞ . Then

$$u \in C^\infty(\bar{U}).$$

Proof. According to Theorem 5 we have $u \in H^m(U)$ for each integer $m = 1, 2, \dots$. Thus Theorem 6 in §5.6.3 implies $u \in C^k(\bar{U})$ for each $k = 1, 2, \dots$. \square

The computations in this section have basically been repeated applications of “energy” methods to higher and higher partial derivatives. The basic tool of integration by parts has eventually taken us from weak solutions (belonging merely to $H_0^1(U)$) to smooth, classical solutions.

6.4. MAXIMUM PRINCIPLES

This section develops the *maximum principle* for second-order elliptic partial differential equations.

Maximum principle methods are based upon the observation that if a C^2 function u attains its maximum over an open set U at a point $x_0 \in U$, then

$$(1) \quad Du(x_0) = 0, \quad D^2u(x_0) \leq 0,$$

the latter inequality meaning that the symmetric matrix $D^2u = ((u_{x_i x_j}))$ is nonpositive definite at x_0 . Deductions based upon (1) are consequently *pointwise* in character and are thus utterly different from the integral-based energy methods set forth in §§6.1–6.3.

Furthermore we will need to require that our solutions u are at least C^2 , so that it makes sense to consider the pointwise values of Du, D^2u . (In view of the regularity theory from §6.3 we know however that a weak solution is this smooth, at least provided the coefficients are sufficiently regular.) As we will shortly learn, it is also most appropriate now to consider elliptic operators L having the *nondivergence form*

$$(2) \quad Lu = - \sum_{i,j=1}^n a^{ij} u_{x_i x_j} + \sum_{i=1}^n b^i u_{x_i} + cu,$$

where the coefficients a^{ij}, b^i, c are continuous and—as always—the uniform ellipticity condition (4) in §6.1 holds. We continue also to assume, without loss of generality, the symmetry condition $a^{ij} = a^{ji}$ ($i, j = 1, \dots, n$).

6.4.1. Weak maximum principle.

First, we identify circumstances under which a function must attain its maximum (or minimum) on the boundary. We always assume $U \subset \mathbb{R}^n$ is open, bounded.

THEOREM 1 (Weak maximum principle). *Assume $u \in C^2(U) \cap C(\bar{U})$ and*

$$c \equiv 0 \quad \text{in } U.$$

(i) *If*

$$(3) \quad Lu \leq 0 \quad \text{in } U,$$

then

$$\max_{\bar{U}} u = \max_{\partial U} u.$$

(ii) If

$$(4) \quad Lu \geq 0 \quad \text{in } U,$$

then

$$\min_{\bar{U}} u = \min_{\partial U} u.$$

Remark. A function satisfying (3) is called a *subsolution*. We are thus asserting that a *subsolution attains its maximum on ∂U* . Similarly, if (4) holds, u is a *supersolution* and attains its minimum on ∂U . \square

Proof. 1. Let us first suppose we have the strict inequality

$$(5) \quad Lu < 0 \quad \text{in } U,$$

and yet there exists a point $x_0 \in U$ with

$$(6) \quad u(x_0) = \max_{\bar{U}} u.$$

Now at this maximum point x_0 , we have

$$(7) \quad Du(x_0) = 0$$

and

$$(8) \quad D^2u(x_0) \leq 0.$$

2. Since the matrix $A = ((a^{ij}(x_0)))$ is symmetric and positive definite, there exists an orthogonal matrix $O = ((o_{ij}))$ so that

$$(9) \quad OAO^T = \text{diag}(d_1, \dots, d_n), \quad OO^T = I,$$

with $d_k > 0$ ($k = 1, \dots, n$). Write $y = x_0 + O(x - x_0)$. Then $x - x_0 = O^T(y - x_0)$, and so

$$u_{x_i} = \sum_{k=1}^n u_{y_k} o_{ki}, \quad u_{x_i x_j} = \sum_{k,l=1}^n u_{y_k y_l} o_{ki} o_{lj} \quad (i, j = 1, \dots, n).$$

Hence at the point x_0 ,

$$(10) \quad \begin{aligned} \sum_{i,j=1}^n a^{ij} u_{x_i x_j} &= \sum_{k,l=1}^n \sum_{i,j=1}^n a^{ij} u_{y_k y_l} o_{ki} o_{lj} \\ &= \sum_{k=1}^n d_k u_{y_k y_k} \quad \text{by (9)} \\ &\leq 0, \end{aligned}$$

since $d_k > 0$ and $u_{y_k y_k}(x_0) \leq 0$ ($k = 1, \dots, n$), according to (8).

3. Thus at x_0

$$Lu = - \sum_{i,j=1}^n a^{ij} u_{x_i x_j} + \sum_{i=1}^n b^i u_{x_i} \geq 0,$$

in light of (7) and (10). So (5) and (6) are incompatible, and we have a contradiction.

4. In the general case that (3) holds, write

$$u^\epsilon(x) := u(x) + \epsilon e^{\lambda x_1} \quad (x \in U),$$

where $\lambda > 0$ will be selected below and $\epsilon > 0$. Recall (as in the proof of Theorem 4 in §6.3.2) that the uniform ellipticity condition implies $a^{ii}(x) \geq \theta$ ($i = 1, \dots, n$, $x \in U$). Therefore

$$\begin{aligned} Lu^\epsilon &= Lu + \epsilon L(e^{\lambda x_1}) \\ &\leq \epsilon e^{\lambda x_1} [-\lambda^2 a^{11} + \lambda b^1] \\ &\leq \epsilon e^{\lambda x_1} [-\lambda^2 \theta + \|\mathbf{b}\|_{L^\infty} \lambda] \\ &< 0 \quad \text{in } U, \end{aligned}$$

provided we choose $\lambda > 0$ sufficiently large. Then according to steps 1 and 2 above $\max_{\bar{U}} u^\epsilon = \max_{\partial U} u^\epsilon$. Let $\epsilon \rightarrow 0$ to find $\max_{\bar{U}} u = \max_{\partial U} u$. This proves (i).

5. Since $-u$ is a subsolution whenever u is a supersolution, assertion (ii) follows. \square

We next modify the maximum principle to allow for a *nonnegative* zeroth-order coefficient c . Remember from §A.3 that $u^+ = \max(u, 0)$, $u^- = -\min(u, 0)$.

THEOREM 2 (Weak maximum principle for $c \geq 0$). *Assume $u \in C^2(U) \cap C(\bar{U})$ and*

$$c \geq 0 \quad \text{in } U.$$

(i) *If*

$$Lu \leq 0 \quad \text{in } U,$$

then

$$(11) \quad \max_{\bar{U}} u \leq \max_{\partial U} u^+.$$

(ii) Likewise, if

$$Lu \geq 0 \quad \text{in } U,$$

then

$$(12) \quad \min_{\bar{U}} u \geq -\max_{\partial U} u^-.$$

Remark. So in particular, if $Lu = 0$ in U , then

$$(13) \quad \max_{\bar{U}} |u| = \max_{\partial U} |u|.$$

Proof. 1. Let u be a subsolution and set $V := \{x \in U \mid u(x) > 0\}$. Then

$$\begin{aligned} Ku &:= Lu - cu \\ &\leq -cu \leq 0 \quad \text{in } V. \end{aligned}$$

The operator K has no zeroth-order term and consequently Theorem 1 implies $\max_{\bar{V}} u = \max_{\partial V} u = \max_{\partial U} u^+$. This gives (11) in the case that $V \neq \emptyset$. Otherwise $u \leq 0$ everywhere in U , and (11) likewise follows.

2. Assertion (ii) follows from (i) applied to $-u$, once we observe that $(-u)^+ = u^-$. \square

6.4.2. Strong maximum principle.

We next substantially strengthen the foregoing assertions, by demonstrating that a subsolution u cannot attain its maximum at an interior point of a connected region at all, unless u is constant. This statement is the *strong maximum principle*, which depends on the following subtle analysis of the outer normal derivative $\frac{\partial u}{\partial \nu}$ at a boundary maximum point.

LEMMA (Hopf's Lemma). Assume $u \in C^2(U) \cap C^1(\bar{U})$ and

$$c \equiv 0 \quad \text{in } U.$$

Suppose further

$$Lu \leq 0 \quad \text{in } U$$

and there exists a point $x^0 \in \partial U$ such that

$$(14) \quad u(x^0) > u(x) \quad \text{for all } x \in U.$$

Assume finally that U satisfies the interior ball condition at x^0 ; that is, there exists an open ball $B \subset U$ with $x^0 \in \partial B$.

(i) Then

$$\frac{\partial u}{\partial \nu}(x^0) > 0,$$

where ν is the outer unit normal to B at x^0 .

(ii) If

$$c \geq 0 \quad \text{in } U,$$

the same conclusion holds provided

$$u(x^0) \geq 0.$$

The importance of (i) is the *strict* inequality: that $\frac{\partial u}{\partial \nu}(x^0) \geq 0$ is obvious. Note that the interior ball condition automatically holds if ∂U is C^2 .

Proof. 1. Assume $c \geq 0$. We may as well further assume $B = B^0(0, r)$ for some radius $r > 0$. Define

$$v(x) := e^{-\lambda|x|^2} - e^{-\lambda r^2} \quad (x \in B(0, r))$$

for $\lambda > 0$ as selected below. Then using the uniform ellipticity condition, we compute

$$\begin{aligned} Lv &= - \sum_{i,j=1}^n a^{ij} v_{x_i x_j} + \sum_{i=1}^n b^i v_{x_i} + cv \\ &= e^{-\lambda|x|^2} \sum_{i,j=1}^n a^{ij} (-4\lambda^2 x_i x_j + 2\lambda \delta_{ij}) \\ &\quad - e^{-\lambda|x|^2} \sum_{i=1}^n b^i 2\lambda x_i + c(e^{-\lambda|x|^2} - e^{-\lambda r^2}) \\ &\leq e^{-\lambda|x|^2} (-4\theta\lambda^2|x|^2 + 2\lambda \operatorname{tr} \mathbf{A} + 2\lambda|\mathbf{b}||x| + c), \end{aligned}$$


for $\mathbf{A} = ((a^{ij}))$, $\mathbf{b} = (b^1, \dots, b^n)$. Consider next the open annular region $R := B^0(0, r) - B(0, r/2)$. We have

$$(15) \quad Lv \leq e^{-\lambda|x|^2} (-\theta\lambda^2 r^2 + 2\lambda \operatorname{tr} \mathbf{A} + 2\lambda|\mathbf{b}|r + c) \leq 0$$

in R , provided $\lambda > 0$ is fixed large enough.

2. In view of (14) there exists a constant $\epsilon > 0$ so small that

$$(16) \quad u(x^0) \geq u(x) + \epsilon v(x) \quad (x \in \partial B(0, r/2)).$$



In addition note

$$(17) \quad u(x^0) \geq u(x) + \epsilon v(x) \quad (x \in \partial B(0, r)),$$

since $v \equiv 0$ on $\partial B(0, r)$.

3. From (15) we see

$$L(u + \epsilon v - u(x^0)) \leq -cu(x^0) \leq 0 \quad \text{in } R,$$

and from (16), (17) we observe

$$u + \epsilon v - u(x^0) \leq 0 \quad \text{on } \partial R.$$

In view of the weak maximum principle, Theorem 2, $u + \epsilon v - u(x^0) \leq 0$ in R . But $u(x^0) + \epsilon v(x^0) - u(x^0) = 0$, and so

$$\frac{\partial u}{\partial \nu}(x^0) + \epsilon \frac{\partial v}{\partial \nu}(x^0) \geq 0.$$

Consequently

$$\frac{\partial u}{\partial \nu}(x^0) \geq -\epsilon \frac{\partial v}{\partial \nu}(x^0) = -\frac{\epsilon}{r} Dv(x^0) \cdot x^0 = 2\lambda \epsilon r e^{-\lambda r^2} > 0,$$

as required. □

Hopf's Lemma is the primary technical tool in the next proof:

THEOREM 3 (Strong maximum principle). Assume $u \in C^2(U) \cap C(\bar{U})$ and

$$c \equiv 0 \quad \text{in } U.$$

Suppose also U is connected, open and bounded.

(i) If

$$Lu \leq 0 \quad \text{in } U$$

and u attains its maximum over \bar{U} at an interior point, then

u is constant within U .

(ii) Similarly, if

$$Lu \geq 0 \quad \text{in } U$$

and u attains its minimum over \bar{U} at an interior point, then

u is constant within U .

Proof. Write $M := \max_{\bar{U}} u$ and $C := \{x \in U \mid u(x) = M\}$. Then if $u \not\equiv M$, set

$$V := \{x \in U \mid u(x) < M\}.$$

Choose a point $y \in V$ satisfying $\text{dist}(y, C) < \text{dist}(y, \partial U)$, and let B denote the largest ball with center y whose interior lies in V . Then there exists some point $x^0 \in C$, with $x^0 \in \partial B$. Clearly V satisfies the interior ball condition at x^0 , whence Hopf's Lemma, (i), implies $\frac{\partial u}{\partial \nu}(x^0) > 0$. But this is a contradiction: since u attains its maximum at $x^0 \in U$, we have $Du(x^0) = 0$. \square

If the zeroth-order term c is *nonnegative*, we have this version of the strong maximum principle:

THEOREM 4 (Strong maximum principle with $c \geq 0$). Assume $u \in C^2(U) \cap C(\bar{U})$ and

$$c \geq 0 \quad \text{in } U.$$

Suppose also U is connected.

(i) If

$$Lu \leq 0 \quad \text{in } U$$

and u attains a nonnegative maximum over \bar{U} at an interior point, then

u is constant within U .

(ii) Similarly, if

$$Lu \geq 0 \quad \text{in } U$$

and u attains a nonpositive minimum over \bar{U} at an interior point, then

u is constant within U .

The proof is like that above, except that we use statement (ii) in Hopf's Lemma.

6.4.3. Harnack's inequality.

Harnack's inequality states that the values of a nonnegative solution are comparable, at least in any subregion away from the boundary. We assume as usual that

$$Lu = - \sum_{i,j=1}^n a^{ij} u_{x_i x_j} + \sum_{i=1}^n b^i u_{x_i} + cu.$$

THEOREM 5 (Harnack's inequality). *Assume $u \geq 0$ is a C^2 solution of*

$$Lu = 0 \quad \text{in } U,$$

and suppose $V \subset\subset U$ is connected. Then there exists a constant C such that

$$(18) \quad \sup_V u \leq C \inf_V u.$$

The constant C depends only on V and the coefficients of L .

This assertion is true if the coefficients are merely bounded and measurable: see Gilbarg–Trudinger [G-T]. We will however provide a proof only for the special case that $b^i \equiv c \equiv 0$ and the a^{ij} are smooth ($i, j = 1, \dots, n$).

Proof. 1. We may assume $u > 0$ in U , for otherwise we could apply the result to $u + \epsilon$ and then let $\epsilon \rightarrow 0^+$.

Set

$$(19) \quad v := \log u.$$

Since $Lu = 0$, we compute

$$(20) \quad \sum_{i,j=1}^n a^{ij} v_{x_i x_j} + a^{ij} v_{x_i} v_{x_j} = 0 \quad \text{in } U.$$

Define

$$(21) \quad w := \sum_{i,j=1}^n a^{ij} v_{x_i} v_{x_j},$$

so that (20) says

$$(22) \quad - \sum_{i,j=1}^n a^{ij} v_{x_i x_j} = w.$$

2. We calculate for $k, l = 1, \dots, n$ that

$$w_{x_k x_l} = \sum_{i,j=1}^n 2a^{ij} v_{x_i x_k x_l} v_{x_j} + 2a^{ij} v_{x_i x_k} v_{x_j x_l} + R,$$

where the remainder term R , resulting from derivatives falling upon the coefficients, satisfies an estimate of the form

$$(23) \quad |R| \leq \epsilon |D^2 v|^2 + C(\epsilon) |Dv|^2$$

for each $\epsilon > 0$. Thus

$$(24) \quad - \sum_{k,l=1}^n a^{kl} w_{x_k x_l} = 2 \sum_{i,j=1}^n a^{ij} v_{x_j} \left(- \sum_{k,l=1}^n a^{lk} v_{x_i x_k x_l} \right) - 2 \sum_{k,l=1}^n a^{ij} a^{kl} v_{x_i x_k} v_{x_j x_l} - R.$$

Differentiating (22), we see that

$$(25) \quad - \sum_{k,l=1}^n a^{lk} v_{x_i x_k x_l} = w_{x_i} + R_i \quad (i = 1, \dots, n),$$

where R_i denotes another remainder term satisfying estimate (23). Furthermore, the uniform ellipticity condition implies

$$(26) \quad \sum_{k,l=1}^n a^{ij} a^{kl} v_{x_i x_k} v_{x_j x_l} \geq \theta^2 |D^2 v|^2.$$

Substituting (25) and (26) into (24) and then choosing $\epsilon > 0$ small enough, we derive the differential inequality

$$(27) \quad - \sum_{k,l=1}^n a^{kl} w_{x_k x_l} + \sum_{k=1}^n b^k w_{x_k} \geq \theta^2 |D^2 v|^2 - C |Dv|^2,$$

where

$$(28) \quad b^k := -2 \sum_{l=1}^n a^{kl} v_{x_l} \quad (k = 1, \dots, n).$$

3. Suppose next $V \subset\subset U$ is an open ball of radius $r > 0$. Choose a cutoff function $\zeta \in C^\infty(U)$ such that

$$0 \leq \zeta \leq 1, \quad \zeta = 0 \text{ on } \partial U, \quad \zeta \equiv 1 \text{ on } V.$$

Let

$$(29) \quad z := \zeta^4 w.$$

We now assume

$$(30) \quad z \text{ attains its maximum at some point } x_0 \in U.$$

Then at x_0 we have

$$(31) \quad \zeta w_{x_k} + 4\zeta_{x_k} w = 0 \quad (k = 1, \dots, n).$$

Furthermore, at this point we have the inequality

$$0 \leq - \sum_{k,l=1}^n a^{kl} z_{x_k x_l} + \sum_{k=1}^n b^k z_{x_k} = - \sum_{k,l=1}^n a^{kl} (\zeta^4 w)_{x_k x_l} + \sum_{k=1}^n b^k (\zeta^4 w)_{x_k}.$$

Hence

$$(32) \quad 0 \leq \zeta^4 \left(- \sum_{k,l=1}^n a^{kl} w_{x_k x_l} + \sum_{k=1}^n b^k w_{x_k} \right) + \hat{R},$$

where the remainder term \hat{R} , which comprises terms for which derivatives fall upon the cutoff function ζ , satisfies the estimate

$$|\hat{R}| \leq C(\zeta^2 w + \zeta^3 |Dw|).$$

Recalling now (31), we see that in fact

$$(33) \quad |\hat{R}| \leq C\zeta^2 w.$$

Insert this estimate and (27) into (32):

$$\zeta^4 |D^2 v|^2 \leq C\zeta^4 |Dv|^2 + C\zeta^2 w.$$

But $\theta |Dv|^2 \leq w$, and furthermore $w \leq C |D^2 v|$, according to (22). It follows that

$$\zeta^4 w^2 \leq C\zeta^2 w,$$

and therefore

$$z = \zeta^4 w \leq C \quad \text{at the point } x_0.$$

Since z attains its maximum at x_0 and since $\zeta \equiv 1$ on V , we consequently have the estimate

$$(34) \quad |Dv| \leq C \quad \text{within the region } V.$$

4. Now select $x_1, x_2 \in V$. Then

$$v(x_2) - v(x_1) \leq \sup_V |Dv| r \leq C,$$

where we recall that r is the radius of the ball V . We exponentiate to deduce

$$u(x_2) \leq u(x_1)e^C.$$

This inequality is valid for all $x_1, x_2 \in V$, and so the Harnack inequality (18) is valid if V is a ball. In the general case we cover $V \subset\subset U$ with balls and repeatedly apply the inequality above. \square

6.5. EIGENVALUES AND EIGENFUNCTIONS

We consider in this section the boundary-value problem

$$(1) \quad \begin{cases} Lw = \lambda w & \text{in } U \\ w = 0 & \text{on } \partial U, \end{cases}$$

where U is open and bounded, and recall that λ is an eigenvalue of L provided there exists a nontrivial solution w of (1). From the theory developed in §6.2 we recall that the set Σ of eigenvalues of L is at most countable.

The theorems in §6.5.1 below are analogues for elliptic PDE of the standard linear algebra assertion that a real symmetric matrix possesses real eigenvalues and an orthonormal basis of eigenvectors. Similarly, the results in §6.5.2 are PDE versions of the Perron–Frobenius theorem that a matrix with positive entries has a real, positive eigenvalue and a corresponding eigenvector with positive entries (cf. Gantmacher [Ga]).

6.5.1. Eigenvalues of symmetric elliptic operators.

For simplicity, we consider now an elliptic operator having the divergence form

$$(2) \quad Lu = - \sum_{i,j=1}^n (a^{ij} u_{x_i})_{x_j},$$

where $a^{ij} \in C^\infty(\bar{U})$ ($i, j = 1, \dots, n$). We suppose the usual uniform ellipticity condition to hold and as usual suppose

$$(3) \quad a^{ij} = a^{ji} \quad (i, j = 1, \dots, n).$$

The operator L is thus formally symmetric, and in particular the associated bilinear form $B[\ , \]$ satisfies $B[u, v] = B[v, u]$ ($u, v \in H_0^1(U)$). Assume also U is connected.

THEOREM 1 (Eigenvalues of symmetric elliptic operators).

- (i) *Each eigenvalue of L is real.*
 (ii) *Furthermore, if we repeat each eigenvalue according to its (finite) multiplicity, we have*

$$\Sigma = \{\lambda_k\}_{k=1}^{\infty},$$

where

$$0 < \lambda_1 \leq \lambda_2 \leq \lambda_3 \leq \dots$$

and

$$\lambda_k \rightarrow \infty \quad \text{as } k \rightarrow \infty.$$

- (iii) *Finally, there exists an orthonormal basis $\{w_k\}_{k=1}^{\infty}$ of $L^2(U)$, where $w_k \in H_0^1(U)$ is an eigenfunction corresponding to λ_k :*

$$(4) \quad \begin{cases} Lw_k = \lambda_k w_k & \text{in } U \\ w_k = 0 & \text{on } \partial U, \end{cases}$$

for $k = 1, 2, \dots$

Owing to the regularity theory in §6.3, $w_k \in C^\infty(U)$, and furthermore $w_k \in C^\infty(\bar{U})$ if ∂U is smooth, for $k = 1, 2, \dots$

Proof. 1. As in §6.2,

$$S := L^{-1}$$

is a bounded, linear, compact operator mapping $L^2(U)$ into itself.

2. We claim further that S is symmetric. To see this, select $f, g \in L^2(U)$. Then $Sf = u$ means $u \in H_0^1(U)$ is the weak solution of

$$\begin{cases} Lu = f & \text{in } U \\ u = 0 & \text{on } \partial U, \end{cases}$$

and likewise $Sg = v$ means $v \in H_0^1(U)$ solves

$$\begin{cases} Lv = g & \text{in } U \\ v = 0 & \text{on } \partial U \end{cases}$$

in the weak sense. Thus

$$(Sf, g) = (u, g) = B[v, u]$$

and

$$(f, Sg) = (f, v) = B[u, v].$$

Since $B[u, v] = B[v, u]$, we see $(Sf, g) = (f, Sg)$ for all $f, g \in L^2(U)$. Therefore S is symmetric.

3. Notice also

$$(Sf, f) = (u, f) = B[u, u] \geq 0 \quad (f \in L^2(U)).$$

Consequently the theory of compact, symmetric operators from §D.6 implies that all the eigenvalues of S are real, positive, and there are corresponding eigenfunctions which make up an orthonormal basis of $L^2(U)$. But observe as well that for $\eta \neq 0$, we have $Sw = \eta w$ if and only if $Lw = \lambda w$ for $\lambda = \frac{1}{\eta}$. The theorem follows. \square

Weyl's Law. The study of the distribution of the eigenvalues of elliptic operators is extremely important in mathematical physics. A landmark assertion concerning the asymptotic distribution of eigenvalues is due to H. Weyl. For the special case of the eigenvalues $\{\lambda_k\}_{k=1}^{\infty}$ of the Laplacian in the smooth, bounded open set $U \subset \mathbb{R}^n$, taken with zero boundary conditions, Weyl's Law asserts

$$\lim_{k \rightarrow \infty} \frac{\lambda_k^{\frac{n}{2}}}{k} = \frac{(2\pi)^n}{|U|\alpha(n)}.$$

Here $|U|$ denotes the volume of U .

We next scrutinize more carefully the first eigenvalue of L .

DEFINITION. We call $\lambda_1 > 0$ the principal eigenvalue of L .

THEOREM 2 (Variational principle for the principal eigenvalue).

(i) We have

$$(5) \quad \lambda_1 = \min\{B[u, u] \mid u \in H_0^1(U), \|u\|_{L^2} = 1\}.$$

(ii) Furthermore, the above minimum is attained for a function w_1 , positive within U , which solves

$$\begin{cases} Lw_1 = \lambda_1 w_1 & \text{in } U \\ w_1 = 0 & \text{on } \partial U. \end{cases}$$

(iii) Finally, if $u \in H_0^1(U)$ is any weak solution of

$$\begin{cases} Lu = \lambda_1 u & \text{in } U \\ u = 0 & \text{on } \partial U, \end{cases}$$

then u is a multiple of w_1 .

Remarks. (i) Assertion (iii) says *the principal eigenvalue λ_1 is simple*. In particular

$$0 < \lambda_1 < \lambda_2 \leq \lambda_3 \leq \dots$$

(ii) Expression (5) is *Rayleigh's formula* and is equivalent to the statement

$$\lambda_1 = \min_{\substack{u \in H_0^1(U) \\ u \neq 0}} \frac{B[u, u]}{\|u\|_{L^2(U)}^2}.$$

Proof. 1. In view of (4) we see

$$(6) \quad B[w_k, w_k] = \lambda_k \|w_k\|_{L^2(U)}^2 = \lambda_k$$

and

$$(7) \quad B[w_k, w_l] = \lambda_k (w_k, w_l) = 0$$

for $k, l = 1, 2, \dots, k \neq l$.

2. As $\{w_k\}_{k=1}^\infty$ is an orthonormal basis of $L^2(U)$, if $u \in H_0^1(U)$ and $\|u\|_{L^2(U)} = 1$, we can write

$$(8) \quad u = \sum_{k=1}^{\infty} d_k w_k$$

for $d_k = (u, w_k)_{L^2(U)}$, the series converging in $L^2(U)$: see §D.2. In addition

$$(9) \quad \sum_{k=1}^{\infty} d_k^2 = \|u\|_{L^2(U)}^2 = 1.$$

3. Furthermore from (6) and (7) we see that $\left\{ \frac{w_k}{\lambda_k^{1/2}} \right\}_{k=1}^\infty$ is an orthonormal subset of $H_0^1(U)$, endowed with the new inner product $B[\ , \]$.

We claim further that $\left\{ \frac{w_k}{\lambda_k^{1/2}} \right\}_{k=1}^\infty$ is in fact an orthonormal basis of $H_0^1(U)$, with this new inner product. To see this, it suffices to verify that

$$B[w_k, u] = 0 \quad (k = 1, 2, \dots)$$

implies $u \equiv 0$. But this assertion is clearly true, since the identities

$$B[w_k, u] = \lambda_k (w_k, u) = 0 \quad (k = 1, \dots)$$

force $u \equiv 0$, as $\{w_k\}_{k=1}^\infty$ is a basis of $L^2(U)$. Consequently

$$u = \sum_{k=1}^{\infty} \mu_k \frac{w_k}{\lambda_k^{1/2}}$$

for $\mu_k = B \left[u, \frac{w_k}{\lambda_k^{1/2}} \right]$, the series converging in $H_0^1(U)$. But then according to (8), $\mu_k = d_k \lambda_k^{1/2}$; and so the series (8) in fact converges also in $H_0^1(U)$.

4. Thus (6) and (8) imply

$$B[u, u] = \sum_{k=1}^{\infty} d_k^2 \lambda_k \geq \lambda_1 \quad \text{by (9).}$$

As equality holds for $u = w_1$, we obtain formula (5).

5. We next *claim* that if $u \in H_0^1(U)$ and $\|u\|_{L^2(U)} = 1$, then u is a weak solution of

$$(10) \quad \begin{cases} Lu = \lambda_1 u & \text{in } U \\ u = 0 & \text{on } \partial U \end{cases}$$

if and only if

$$(11) \quad B[u, u] = \lambda_1.$$

Obviously (10) implies (11). On the other hand, suppose (11) is valid. Then, writing $d_k = (u, w_k)$ as above, we have

$$(12) \quad \sum_{k=1}^{\infty} d_k^2 \lambda_1 = \lambda_1 = B[u, u] = \sum_{k=1}^{\infty} d_k^2 \lambda_k.$$

Hence

$$(13) \quad \sum_{k=1}^{\infty} (\lambda_k - \lambda_1) d_k^2 = 0.$$

Consequently

$$d_k = (u, w_k) = 0 \quad \text{if } \lambda_k > \lambda_1.$$

Since λ_1 has finite multiplicity, it follows that

$$(14) \quad u = \sum_{k=1}^m (u, w_k) w_k$$

for some m , where $Lw_k = \lambda_1 w_k$. Therefore

$$(15) \quad Lu = \sum_{k=1}^m (u, w_k) Lw_k = \lambda_1 u.$$

This proves (10).

6. Next we will show that if $u \in H_0^1(U)$ is a weak solution of (10), $u \neq 0$, then either

$$(16) \quad u > 0 \quad \text{in } U$$

or else

$$(17) \quad u < 0 \quad \text{in } U.$$

To see this, let us assume without loss of generality that $\|u\|_{L^2} = 1$ and note

$$(18) \quad \alpha + \beta = 1$$

for

$$\alpha := \int_U (u^+)^2 dx, \quad \beta := \int_U (u^-)^2 dx.$$

Furthermore since $u^\pm \in H_0^1(U)$, with

$$Du^+ = \begin{cases} Du & \text{a.e. on } \{u \geq 0\} \\ 0 & \text{a.e. on } \{u \leq 0\}, \end{cases}$$

$$Du^- = \begin{cases} 0 & \text{a.e. on } \{u \geq 0\} \\ -Du & \text{a.e. on } \{u \leq 0\} \end{cases}$$

(cf. Problem 18 in Chapter 5), we have $B[u^+, u^-] = 0$. Accordingly

$$\begin{aligned} \lambda_1 &= B[u, u] = B[u^+, u^+] + B[u^-, u^-] \\ &\geq \lambda_1 \|u^+\|_{L^2(U)}^2 + \lambda_1 \|u^-\|_{L^2(U)}^2 \quad \text{by (5)} \\ &= (\alpha + \beta)\lambda_1 = \lambda_1. \end{aligned}$$

But then we see that the inequality above must in fact be an equality, and so

$$B[u^+, u^+] = \lambda_1 \|u^+\|_{L^2(U)}^2, \quad B[u^-, u^-] = \lambda_1 \|u^-\|_{L^2(U)}^2.$$

Therefore the claim proved in step 5 asserts

$$(19) \quad \begin{cases} Lu^+ = \lambda_1 u^+ & \text{in } U \\ u^+ = 0 & \text{on } \partial U \end{cases}$$

and

$$(20) \quad \begin{cases} Lu^- = \lambda_1 u^- & \text{in } U \\ u^- = 0 & \text{on } \partial U \end{cases}$$

in the weak sense.

7. Next, since the coefficients a^{ij} are smooth, we deduce from (19) that $u^+ \in C^\infty(U)$ and

$$Lu^+ = \lambda_1 u^+ \geq 0 \quad \text{in } U.$$

The function u^+ is therefore a supersolution. Thus the strong maximum principle implies either $u^+ > 0$ in U or else $u^+ \equiv 0$ in U . Similar arguments apply to u^- , and so either (16) or (17) holds.

8. Finally assume that u and \tilde{u} are two nontrivial weak solutions of (10). In view of steps 6 and 7 above

$$\int_U \tilde{u} \, dx \neq 0,$$

and so there exists a real constant χ such that

$$(21) \quad \int_U u - \chi \tilde{u} \, dx = 0.$$

But since $u - \chi \tilde{u}$ is also a weak solution of (10), steps 6 and 7 and the equality (21) imply $u \equiv \chi \tilde{u}$ in U . Hence the eigenvalue λ_1 is simple. \square

6.5.2. Eigenvalues of nonsymmetric elliptic operators.

We will now consider a uniformly elliptic operator L in the nondivergence form:

$$Lu = - \sum_{i,j=1}^n a^{ij} u_{x_i x_j} + \sum_{i=1}^n b^i u_{x_i} + cu.$$

Let us for simplicity assume that $a^{ij}, b^i, c \in C^\infty(\bar{U})$, that U is open, bounded and connected, and that ∂U is smooth. We suppose also $a^{ij} = a^{ji}$ ($i, j = 1, \dots, n$) and

$$(22) \quad c \geq 0 \quad \text{in } U.$$

Notice however that in general the operator L will not equal its formal adjoint. We therefore *cannot* invoke as above the abstract theory from §D.6. And in fact L will in general have complex eigenvalues and eigenfunctions.

Remarkably, however, the principal eigenvalue of L is real, and the corresponding eigenfunction is of one sign within U .

THEOREM 3 (Principal eigenvalue for nonsymmetric elliptic operators).

(i) *There exists a real eigenvalue λ_1 for the operator L , taken with zero boundary conditions, such that if $\lambda \in \mathbb{C}$ is any other eigenvalue, we have*

$$\operatorname{Re}(\lambda) \geq \lambda_1.$$

(ii) *There exists a corresponding eigenfunction w_1 , which is positive within U .*

(iii) *The eigenvalue λ_1 is simple; that is, if u is any solution of (1), then u is a multiple of w_1 .*

Proof*. 1. Choose $m = \lceil \frac{n}{2} \rceil + 3$ and consider the Banach space $X = H^m(U) \cap H_0^1(U)$. According to Theorem 6 in §5.6.3, $X \subset C^2(\bar{U})$. We define the linear, compact operator $A : X \rightarrow X$ by setting $Af = u$, where u is the unique solution of

$$(23) \quad \begin{cases} Lu = f & \text{in } U \\ u = 0 & \text{on } \partial U. \end{cases}$$

Next define the cone

$$C = \{u \in X \mid u \geq 0 \text{ in } U\}.$$

According to the maximum principle, $A : C \rightarrow C$.

2. Hereafter fix any function $w \in C$, $w \not\equiv 0$. Employing the strong maximum principle and Hopf's Lemma, we deduce

$$(24) \quad v > 0 \text{ in } U, \quad \frac{\partial v}{\partial \nu} < 0 \text{ on } \partial U$$

for $v = A(w)$.

Remember that $w = 0$ on ∂U . So in view of (24) there exists a constant $\mu > 0$ so that

$$(25) \quad \mu v \geq w \quad \text{in } U.$$

3. Fix $\epsilon > 0$, $\eta > 0$, and consider then the equation

$$(26) \quad u = \eta A[u + \epsilon w]$$

for the unknown $u \in C$. We claim that

$$(27) \quad \text{if (26) has a solution } u, \text{ then } \eta \leq \mu.$$

*Omit on first reading.

To verify this assertion, suppose in fact $u \in C$ solves (26). We compute

$$u \geq \eta A[\epsilon w] = \eta \epsilon v \geq \frac{\eta}{\mu} \epsilon w,$$

according to (25). Hence

$$u \geq \eta A u \geq \frac{\eta^2 \epsilon}{\mu} A w = \frac{\eta^2 \epsilon}{\mu} v \geq \left(\frac{\eta}{\mu}\right)^2 \epsilon w.$$

Continuing, we deduce

$$u \geq \left(\frac{\eta}{\mu}\right)^k \epsilon w \quad (k = 1, \dots),$$

a contradiction unless $\eta \leq \mu$. This observation confirms the assertion (27).

4. Define

$$S_\epsilon := \{u \in C \mid \text{there exists } 0 \leq \eta \leq 2\mu \text{ such that } u = \eta A[u + \epsilon w]\}.$$

We assert that

$$(28) \quad S_\epsilon \text{ is unbounded in } X.$$

Indeed, if this were not so, it would follow that the equation

$$u = 2\mu A[u + \epsilon w]$$

has a solution, in contradiction to (27). To draw this conclusion we have applied the variant for mappings on convex sets of Schaefer's fixed point theorem, stated in §9.2.2.

5. Owing to (28), there exist

$$(29) \quad 0 \leq \eta_\epsilon \leq 2\mu$$

and $v_\epsilon \in C$, with $\|v_\epsilon\|_X \geq \frac{1}{\epsilon}$, satisfying

$$(30) \quad v_\epsilon = \eta_\epsilon A[v_\epsilon + \epsilon w].$$

Renormalize by setting

$$(31) \quad u_\epsilon := \frac{v_\epsilon}{\|v_\epsilon\|_X}.$$

Using (29)–(31) and the compactness of the operator A , we obtain a subsequence $\epsilon_k \rightarrow 0$ so that

$$\eta_{\epsilon_k} \rightarrow \eta \text{ and } u_{\epsilon_k} \rightarrow u \text{ in } X.$$

Then (31) implies

$$(32) \quad \|u\|_X = 1, \quad u \in C.$$

Since $u_\epsilon = \eta_\epsilon A \left[u_\epsilon + \frac{\epsilon w}{\|v_\epsilon\|_X} \right]$, we deduce in the limit that $u = \eta Au$. In view of (32), $\eta > 0$. We may consequently rewrite the above to read

$$\begin{cases} Lw_1 = \lambda_1 w_1 & \text{in } U \\ w_1 = 0 & \text{on } \partial U, \end{cases}$$

for $\lambda_1 = \eta$, $u = w_1$. Thus λ_1 is a real eigenvalue for the operator L , taken with zero boundary conditions, and $w_1 \geq 0$ is a corresponding eigenfunction. In view of the strong maximum principle and Hopf's Lemma, we have

$$(33) \quad w_1 > 0 \text{ in } U, \quad \frac{\partial w_1}{\partial \nu} < 0 \text{ on } \partial U.$$

Additionally, we know w_1 is smooth, owing to the regularity theory in §6.3.

6. All expressions occurring in steps 1–5 above are real. Suppose now $\lambda \in \mathbb{C}$ and u is a complex-valued solution of

$$(34) \quad \begin{cases} Lu = \lambda u & \text{in } U \\ u = 0 & \text{on } \partial U. \end{cases}$$

Now choose any smooth function $w : U \rightarrow \mathbb{R}$, with $w > 0$ in U , and set $v := \frac{u}{w}$. We compute

$$(35) \quad \begin{aligned} \lambda v &= \frac{1}{w} L(vw) \quad \text{by (34)} \\ &= Lv - cv - \frac{2}{w} \sum_{i,j=1}^n a^{ij} w_{x_j} v_{x_i} + \frac{v}{w} Lw. \end{aligned}$$

Writing

$$Kv := - \sum_{i,j=1}^n a^{ij} v_{x_i x_j} + \sum_{i=1}^n b'^i v_{x_i}$$

for $b'^i := b^i - \frac{2}{w} \sum_{j=1}^n a^{ij} w_{x_j}$ ($i = 1, \dots, n$), we deduce from (35) that

$$(36) \quad Kv + \left(\frac{Lw}{w} - \lambda \right) v = 0 \quad \text{in } U.$$

Take complex conjugates:

$$(37) \quad K\bar{v} + \left(\frac{Lw}{w} - \bar{\lambda} \right) \bar{v} = 0 \quad \text{in } U.$$

Next we compute

$$(38) \quad K(|v|^2) = K(v\bar{v}) = \bar{v}Kv + vK\bar{v} - 2 \sum_{i,j=1}^n a^{ij} v_{x_i} \bar{v}_{x_j} \leq \bar{v}Kv + vK\bar{v},$$

since

$$\sum_{i,j=1}^n a^{ij} \xi_i \bar{\xi}_j = \sum_{i,j=1}^n a^{ij} (\operatorname{Re}(\xi_i) \operatorname{Re}(\xi_j) + \operatorname{Im}(\xi_i) \operatorname{Im}(\xi_j)) \geq 0$$

for $\xi \in \mathbb{C}^n$. Combining (36)–(38), we discover

$$K(|v|^2) \leq 2 \left(\operatorname{Re} \lambda - \frac{Lw}{w} \right) |v|^2.$$

Now choose

$$(39) \quad w := w_1^{1-\epsilon}$$

for $0 < \epsilon < 1$. Then

$$Lw = \frac{(1-\epsilon)}{w_1^\epsilon} Lw_1 + \frac{\epsilon(1-\epsilon)}{w_1^{1+\epsilon}} \sum_{i,j=1}^n a^{ij} w_{1,x_i} w_{1,x_j} + \epsilon c w_1^{1-\epsilon} \geq (1-\epsilon) \lambda_1 w.$$

Consequently

$$K(|v|^2) \leq 2(\operatorname{Re} \lambda - (1-\epsilon)\lambda_1) |v|^2 \quad \text{in } U.$$

Thus if $\operatorname{Re}(\lambda) \leq (1-\epsilon)\lambda_1$, then $K(|v|^2) \leq 0$ in U . As $v = 0$ on ∂U , according to (33) and (39), we deduce from the maximum principle that $v = \frac{u}{w} = 0$ in U . Thus $u \equiv 0$ in U and so λ cannot be an eigenvalue. This conclusion obtains for each $\epsilon > 0$, and so $\operatorname{Re} \lambda \geq \lambda_1$ if λ is any complex eigenvalue.

7. Finally, let u be any (possibly complex-valued) solution of

$$(40) \quad \begin{cases} Lu = \lambda_1 u & \text{in } U \\ u = 0 & \text{on } \partial U. \end{cases}$$

Since $\operatorname{Re}(u)$ and $\operatorname{Im}(u)$ also solve (40), we may as well suppose from the outset u is real-valued. Replacing u by $-u$ if needs be, we may also suppose $u > 0$ somewhere in U . Now set

$$(41) \quad \chi := \sup\{\mu > 0 \mid w_1 - \mu u \geq 0 \text{ in } U\}.$$

Then $0 < \chi < \infty$. Write $v = w_1 - \chi u$; so that $v \geq 0$ in U and

$$\begin{cases} Lv = \lambda_1 v \geq 0 & \text{in } U \\ v = 0 & \text{on } \partial U. \end{cases}$$

Now if v is not identically zero, the strong maximum principle and Hopf's Lemma imply

$$v > 0 \text{ in } U, \quad \frac{\partial v}{\partial \nu} < 0 \text{ on } \partial U.$$

Thus

$$v - \epsilon u \geq 0 \quad \text{in } U \text{ for some } \epsilon > 0,$$

and so

$$w_1 - (\chi + \epsilon)u \geq 0 \quad \text{in } U,$$

a contradiction to (41). Hence $v \equiv 0$ in U , and so u is a multiple of w_1 . \square

6.6. PROBLEMS

In the following exercises we assume the coefficients of the various PDE are smooth and satisfy the uniform ellipticity condition. Also $U \subset \mathbb{R}^n$ is always an open, bounded set, with smooth boundary ∂U .

1. Consider Laplace's equation with potential function c :

$$(*) \quad -\Delta u + cu = 0,$$

and the divergence structure equation:

$$(**) \quad -\operatorname{div}(aDv) = 0,$$

where the function a is positive.

(a) Show that if u solves (*) and $w > 0$ also solves (*), then $v := u/w$ solves (**) for $a := w^2$.

(b) Conversely, show that if v solves (**), then $u := va^{1/2}$ solves (*) for some potential c .

2. Let

$$Lu = - \sum_{i,j=1}^n (a^{ij}u_{x_i})_{x_j} + cu.$$

Prove that there exists a constant $\mu > 0$ such that the corresponding bilinear form $B[\cdot, \cdot]$ satisfies the hypotheses of the Lax–Milgram Theorem, provided

$$c(x) \geq -\mu \quad (x \in U).$$

3. A function $u \in H_0^2(U)$ is a weak solution of this boundary-value problem for the *biharmonic equation*

$$(*) \quad \begin{cases} \Delta^2 u = f & \text{in } U \\ u = \frac{\partial u}{\partial \nu} = 0 & \text{on } \partial U \end{cases}$$

provided

$$\int_U \Delta u \Delta v \, dx = \int_U f v \, dx$$

for all $v \in H_0^2(U)$. Given $f \in L^2(U)$, prove that there exists a unique weak solution of (*).

4. Assume U is connected. A function $u \in H^1(U)$ is a weak solution of *Neumann's problem*

$$(*) \quad \begin{cases} -\Delta u = f & \text{in } U \\ \frac{\partial u}{\partial \nu} = 0 & \text{on } \partial U \end{cases}$$

if

$$\int_U Du \cdot Dv \, dx = \int_U f v \, dx$$

for all $v \in H^1(U)$. Suppose $f \in L^2(U)$. Prove (*) has a weak solution if and only if

$$\int_U f \, dx = 0.$$

5. Explain how to define $u \in H^1(U)$ to be a weak solution of Poisson's equation with *Robin boundary conditions*:

$$\begin{cases} -\Delta u = f & \text{in } U \\ u + \frac{\partial u}{\partial \nu} = 0 & \text{on } \partial U. \end{cases}$$

Discuss the existence and uniqueness of a weak solution for a given $f \in L^2(U)$.

6. Suppose U is connected and ∂U consists of two disjoint, closed sets Γ_1 and Γ_2 . Define what it means for u to be a weak solution of Poisson's equation with *mixed Dirichlet-Neumann boundary conditions*:

$$\begin{cases} -\Delta u = f & \text{in } U \\ u = 0 & \text{on } \Gamma_1 \\ \frac{\partial u}{\partial \nu} = 0 & \text{on } \Gamma_2. \end{cases}$$

Discuss the existence and uniqueness of weak solutions.

7. Let $u \in H^1(\mathbb{R}^n)$ have compact support and be a weak solution of the semilinear PDE

$$-\Delta u + c(u) = f \quad \text{in } \mathbb{R}^n,$$

where $f \in L^2(\mathbb{R}^n)$ and $c : \mathbb{R} \rightarrow \mathbb{R}$ is smooth, with $c(0) = 0$ and $c' \geq 0$. Prove $u \in H^2(\mathbb{R}^n)$.

(Hint: Mimic the proof of Theorem 1 in §6.3.1, but without the cutoff function ζ .)

8. Let u be a smooth solution of the uniformly elliptic equation $Lu = -\sum_{i,j=1}^n a^{ij}(x)u_{x_i x_j} = 0$ in U . Assume that the coefficients have bounded derivatives.

Set $v := |Du|^2 + \lambda u^2$ and show that

$$Lv \leq 0 \quad \text{in } U$$

if λ is large enough. Deduce

$$\|Du\|_{L^\infty(U)} \leq C(\|Du\|_{L^\infty(\partial U)} + \|u\|_{L^\infty(\partial U)}).$$

9. Assume u is a smooth solution of $Lu = -\sum_{i,j=1}^n a^{ij}u_{x_i x_j} = f$ in U , $u = 0$ on ∂U , where f is bounded. Fix $x^0 \in \partial U$. A *barrier* at x^0 is a C^2 function w such that

$$Lw \geq 1 \text{ in } U, \quad w(x^0) = 0, \quad w \geq 0 \text{ on } \partial U.$$

Show that if w is a barrier at x^0 , there exists a constant C such that

$$|Du(x^0)| \leq C \left| \frac{\partial w}{\partial \nu}(x^0) \right|.$$

10. Assume U is connected. Use (a) energy methods and (b) the maximum principle to show that the only smooth solutions of the Neumann boundary-value problem

$$\begin{cases} -\Delta u = 0 & \text{in } U \\ \frac{\partial u}{\partial \nu} = 0 & \text{on } \partial U \end{cases}$$

are $u \equiv C$, for some constant C .

11. Assume $u \in H^1(U)$ is a bounded weak solution of

$$-\sum_{i,j=1}^n (a^{ij}u_{x_i})_{x_j} = 0 \quad \text{in } U.$$

Let $\phi : \mathbb{R} \rightarrow \mathbb{R}$ be convex and smooth, and set $w = \phi(u)$. Show w is a weak subsolution; that is, $B[w, v] \leq 0$ for all $v \in H_0^1(U)$, $v \geq 0$.

12. We say that the uniformly elliptic operator

$$Lu = - \sum_{i,j=1}^n a^{ij} u_{x_i x_j} + \sum_{i=1}^n b^i u_{x_i} + cu$$

satisfies the *weak maximum principle* if for all $u \in C^2(U) \cap C(\bar{U})$

$$\begin{cases} Lu \leq 0 & \text{in } U \\ u \leq 0 & \text{on } \partial U \end{cases}$$

implies that $u \leq 0$ in U .

Suppose that there exists a function $v \in C^2(U) \cap C(\bar{U})$ such that $Lv \geq 0$ in U and $v > 0$ on \bar{U} . Show that L satisfies the weak maximum principle.

(Hint: Find an elliptic operator M with no zeroth-order term such that $w := u/v$ satisfies $Mw \leq 0$ in the region $\{u > 0\}$. To do this, first compute $(v^2 w_{x_i})_{x_j}$.)

13. (Courant minimax principle) Let $L = - \sum_{i,j=1}^n (a^{ij} u_{x_i})_{x_j}$, where $((a^{ij}))$ is symmetric. Assume the operator L , with zero boundary conditions, has eigenvalues $0 < \lambda_1 < \lambda_2 \leq \dots$. Show

$$\lambda_k = \max_{S \in \Sigma_{k-1}} \min_{\substack{u \in S^\perp \\ \|u\|_{L^2} = 1}} B[u, u] \quad (k = 1, 2, \dots).$$

Here Σ_{k-1} denotes the collection of $(k-1)$ -dimensional subspaces of $H_0^1(U)$.

14. Let λ_1 be the principal eigenvalue of the uniformly elliptic, nonsymmetric operator

$$Lu = - \sum_{i,j=1}^n a^{ij} u_{x_i x_j} + \sum_{i=1}^n b^i u_{x_i} + cu,$$

taken with zero boundary conditions. Prove the “max-min” representation formula:

$$\lambda_1 = \sup_u \inf_x \frac{Lu(x)}{u(x)},$$

the “sup” taken over functions $u \in C^\infty(\bar{U})$ with $u > 0$ in U , $u = 0$ on ∂U , and the “inf” taken over points $x \in U$.

(Hint: Consider the eigenfunction w_1^* corresponding to λ_1 for the adjoint operator L^* .)

15. (Eigenvalues and domain variations) Consider a family of smooth, bounded domains $U(\tau) \subset \mathbb{R}^n$ that depend smoothly upon the parameter $\tau \in \mathbb{R}$. As τ changes, each point on $\partial U(\tau)$ moves with velocity \mathbf{v} .

For each τ , we consider eigenvalues $\lambda = \lambda(\tau)$ and corresponding eigenfunctions $w = w(x, \tau)$:

$$\begin{cases} -\Delta w = \lambda w & \text{in } U(\tau) \\ w = 0 & \text{on } \partial U(\tau), \end{cases}$$

normalized so that $\|w\|_{L^2(U(\tau))} = 1$. Suppose that λ and w are smooth functions of τ and x .

Prove *Hadamard's variational formula*

$$\dot{\lambda} = - \int_{\partial U(\tau)} \left| \frac{\partial w}{\partial \nu} \right|^2 \mathbf{v} \cdot \nu \, dS,$$

where $\dot{\cdot} = \frac{d}{d\tau}$ and $\mathbf{v} \cdot \nu$ is the normal velocity of $\partial U(\tau)$.

(Hint: Use the calculus formula from §C.4.)

16. (Radiation condition) If we separate variables to look for a complex-valued solution of the wave equation having the form $u = e^{-i\sigma t} w$ for $w = w(x)$ and $\sigma \in \mathbb{R}, \sigma \neq 0$, we are led to the eigenvalue problem

$$(*) \quad -\Delta w = \lambda w \quad \text{in } \mathbb{R}^n$$

where $\lambda := \sigma^2$.

(a) Show that $w = e^{i\sigma\omega \cdot x}$ solves (*), provided $|\omega| = 1$. Then $u = e^{i\sigma(\omega \cdot x - t)}$ is a traveling wave solution of the wave equation.

(b) Show that for $n = 3$, the function $\Phi := \frac{e^{i\sigma|x|}}{4\pi|x|}$ solves

$$-\Delta \Phi = \lambda \Phi + \delta_0 \quad \text{in } \mathbb{R}^3.$$

(c) The *Sommerfeld radiation condition* requires for a solution of (*) that

$$\lim_{r \rightarrow \infty} r(w_r - i\sigma w) = 0,$$

for $w_r := Dw \cdot \frac{x}{|x|}$. Prove that the solution w from (a) does not satisfy this condition but that Φ from (b) does.

17. (Continuation) Prove that if w is a complex-valued solution of eigenvalue problem (*) in \mathbb{R}^3 and if w satisfies the radiation condition, then $w \equiv 0$.

(Hints: First observe that

$$0 = \int_{B(0,R)} \bar{w} \Delta w - w \Delta \bar{w} \, dx = \int_{\partial B(0,R)} \bar{w} w_r - w \bar{w}_r \, dS.$$

Use this and the radiation condition to show

$$\int_{\partial B(0,R)} |w_r|^2 + \sigma^2 |w|^2 \, dS = \int_{\partial B(0,R)} |w_r - i\sigma w|^2 \, dS \rightarrow 0$$

as $R \rightarrow \infty$. Given now a point $x_0 \in \mathbb{R}^3$, select $R > |x_0|$. Then

$$w(x_0) = \int_{\partial B(0,R)} \Phi w_r - w \Phi_r \, dS,$$

where $\Phi = \Phi(x - x_0)$. Show the integral goes to zero as $R \rightarrow \infty$.)

6.7. REFERENCES

- Section 6.1 See Gilbarg–Trudinger [**G-T**, Chapters 5, 8].
- Section 6.3 Consult Gilbarg–Trudinger [**G-T**], Krylov [**Kr1**], [**Kr2**] and Ladyzhenskaya–Uraltseva [**L-U**] for more on regularity theory for elliptic PDE.
- Section 6.4 Gilbarg–Trudinger [**G-T**, Chapter 3]. Protter–Weinberger [**P-W**] and Pucci–Serrin [**P-S**] are good references for further maximum principle methods.
- Section 6.5 See Smoller [**S**, pp. 122–125]. The last part of the proof of Theorem 3 is modified from Protter–Weinberger [**P-W**, §2.8]. O. Hald simplified my proof of Theorem 2.
- Section 6.6 S. Armstrong contributed Problem 12. Problem 15: see Garabedian [**G**, Chapter 15] for more domain variation formulas. Colton [**C1**] has more on the Sommerfeld radiation condition.

LINEAR EVOLUTION EQUATIONS

- 7.1 Second-order parabolic equations
- 7.2 Second-order hyperbolic equations
- 7.3 Hyperbolic systems of first-order equations
- 7.4 Semigroup theory
- 7.5 Problems
- 7.6 References

This long chapter studies various linear partial differential equations that involve time. We often call such PDE *evolution equations*, the idea being that the solution evolves in time from a given initial configuration. We will study by energy methods general second-order parabolic and hyperbolic equations and also certain first-order hyperbolic systems. The Fourier transform, utilized in §7.3.3, and the semigroup technique, discussed in §7.4, provide alternative approaches.

7.1. SECOND-ORDER PARABOLIC EQUATIONS

Second-order parabolic PDE are natural generalizations of the heat equation (§2.3). We will study in this section the existence and uniqueness of appropriately defined weak solutions, their smoothness and other properties.

7.1.1. Definitions.

a. Parabolic equations. For this chapter we assume U to be an open, bounded subset of \mathbb{R}^n and as before set $U_T = U \times (0, T]$ for some fixed time $T > 0$.

We will first study the initial/boundary-value problem

$$(1) \quad \begin{cases} u_t + Lu = f & \text{in } U_T \\ u = 0 & \text{on } \partial U \times [0, T] \\ u = g & \text{on } U \times \{t = 0\}, \end{cases}$$

where $f : U_T \rightarrow \mathbb{R}$ and $g : U \rightarrow \mathbb{R}$ are given and $u : \bar{U}_T \rightarrow \mathbb{R}$ is the unknown, $u = u(x, t)$. The letter L denotes for each time t a second-order partial differential operator, having either the divergence form

$$(2) \quad Lu = - \sum_{i,j=1}^n (a^{ij}(x, t)u_{x_i})_{x_j} + \sum_{i=1}^n b^i(x, t)u_{x_i} + c(x, t)u$$

or else the nondivergence form

$$(3) \quad Lu = - \sum_{i,j=1}^n a^{ij}(x, t)u_{x_i x_j} + \sum_{i=1}^n b^i(x, t)u_{x_i} + c(x, t)u,$$

for given coefficients a^{ij}, b^i, c ($i, j = 1, \dots, n$).

DEFINITION. We say that the partial differential operator $\frac{\partial}{\partial t} + L$ is (uniformly) parabolic if there exists a constant $\theta > 0$ such that

$$(4) \quad \sum_{i,j=1}^n a^{ij}(x, t)\xi_i \xi_j \geq \theta |\xi|^2$$

for all $(x, t) \in U_T$, $\xi \in \mathbb{R}^n$.

Remark. Note in particular that for each fixed time $0 \leq t \leq T$ the operator L is a uniformly elliptic operator in the spatial variable x .

An obvious example is $a^{ij} \equiv \delta_{ij}$, $b^i \equiv c \equiv f \equiv 0$, in which case $L = -\Delta$ and the PDE $\frac{\partial u}{\partial t} + Lu = 0$ becomes the heat equation. We will see in fact that solutions of the general second-order parabolic PDE are similar in many ways to solutions of the heat equation.

General second-order parabolic equations describe in physical applications the time-evolution of the density of some quantity u , say a chemical

concentration, within the region U . As noted for the equilibrium setting (i.e., second-order elliptic PDE, in §6.1.1), the second-order term $\sum_{i,j=1}^n a^{ij} u_{x_i x_j}$ describes diffusion, the first-order term $\sum_{i=1}^n b^i u_{x_i}$ describes transport, and the zeroth-order term cu describes creation or depletion.

The Fokker–Planck and Kolmogorov equations from the probabilistic study of diffusion processes are also second-order parabolic equations.

b. Weak solutions. Mimicking the developments in §6.1.2 for elliptic equations, we consider first the case that L has the divergence form (2) and try to find an appropriate notion of weak solution for the initial/boundary-value problem (1). We assume for now that

$$(5) \quad a^{ij}, b^i, c \in L^\infty(U_T) \quad (i, j = 1, \dots, n),$$

$$(6) \quad f \in L^2(U_T),$$

$$(7) \quad g \in L^2(U).$$

We will also always suppose $a^{ij} = a^{ji}$ ($i, j = 1, \dots, n$).

Let us now define, by analogy with the notation introduced in Chapter 6, the time-dependent bilinear form

$$(8) \quad B[u, v; t] := \int_U \sum_{i,j=1}^n a^{ij}(\cdot, t) u_{x_i} v_{x_j} + \sum_{i=1}^n b^i(\cdot, t) u_{x_i} v + c(\cdot, t) uv \, dx$$

for $u, v \in H_0^1(U)$ and a.e. $0 \leq t \leq T$.

Motivation for definition of weak solution. To make plausible the following definition of weak solution, let us first temporarily suppose that $u = u(x, t)$ is in fact a smooth solution of our parabolic problem (1). We now switch our viewpoint, by associating with u a mapping

$$\mathbf{u} : [0, T] \rightarrow H_0^1(U)$$

defined by

$$[\mathbf{u}(t)](x) := u(x, t) \quad (x \in U, 0 \leq t \leq T).$$

In other words, we are going to consider u not as a function of x and t together, but rather as a mapping \mathbf{u} of t into the space $H_0^1(U)$ of functions of x . This point of view will greatly clarify the following presentation.

Returning to problem (1), let us similarly define

$$\mathbf{f} : [0, T] \rightarrow L^2(U)$$

by

$$[\mathbf{f}(t)](x) := f(x, t) \quad (x \in U, 0 \leq t \leq T).$$

Then if we fix a function $v \in H_0^1(U)$, we can multiply the PDE $\frac{\partial u}{\partial t} + Lu = f$ by v and integrate by parts, to find

$$(9) \quad (\mathbf{u}', v) + B[\mathbf{u}, v; t] = (\mathbf{f}, v) \quad \left(' = \frac{d}{dt} \right)$$

for each $0 \leq t \leq T$, the pairing (\cdot, \cdot) denoting inner product in $L^2(U)$.

Next, observe that

$$(10) \quad u_t = g^0 + \sum_{j=1}^n g_{x_j}^j \quad \text{in } U_T$$

for $g^0 := f - \sum_{i=1}^n b^i u_{x_i} - cu$ and $g^j := \sum_{i=1}^n a^{ij} u_{x_i}$ ($j = 1, \dots, n$). Consequently (10) and the definitions from §5.9.1 imply the right-hand side of (10) lies in the Sobolev space $H^{-1}(U)$, with

$$\|u_t\|_{H^{-1}(U)} \leq \left(\sum_{j=0}^n \|g^j\|_{L^2(U)}^2 \right)^{1/2} \leq C \left(\|u\|_{H_0^1(U)} + \|f\|_{L^2(U)} \right).$$

This estimate suggests it may be reasonable to look for a weak solution with $\mathbf{u}' \in H^{-1}(U)$ for a.e. time $0 \leq t \leq T$, in which case the first term in (9) can be reexpressed as $\langle \mathbf{u}', v \rangle$, $\langle \cdot, \cdot \rangle$ being the pairing of $H^{-1}(U)$ and $H_0^1(U)$. \square

All these considerations motivate the following

DEFINITION. We say a function

$$\mathbf{u} \in L^2(0, T; H_0^1(U)), \quad \text{with } \mathbf{u}' \in L^2(0, T; H^{-1}(U)),$$

is a weak solution of the parabolic initial/boundary-value problem (1) provided

$$(i) \quad \langle \mathbf{u}', v \rangle + B[\mathbf{u}, v; t] = (\mathbf{f}, v)$$

for each $v \in H_0^1(U)$ and a.e. time $0 \leq t \leq T$ and

$$(ii) \quad \mathbf{u}(0) = g.$$

Remark. In view of Theorem 3 in §5.9.2, we see $\mathbf{u} \in C([0, T]; L^2(U))$, and thus the equality (ii) makes sense.

7.1.2. Existence of weak solutions.

a. Galerkin approximations. We intend to build a weak solution of the parabolic problem

$$(11) \quad \begin{cases} u_t + Lu = f & \text{in } U_T \\ u = 0 & \text{on } \partial U \times [0, T] \\ u = g & \text{on } U \times \{t = 0\} \end{cases}$$

by first constructing solutions of certain finite-dimensional approximations to (11) and then passing to limits. This is called *Galerkin's method*.

More precisely, assume the functions $w_k = w_k(x)$ ($k = 1, \dots$) are smooth,

$$(12) \quad \{w_k\}_{k=1}^{\infty} \text{ is an orthogonal basis of } H_0^1(U),$$

and

$$(13) \quad \{w_k\}_{k=1}^{\infty} \text{ is an orthonormal basis of } L^2(U).$$

(For instance, we could take $\{w_k\}_{k=1}^{\infty}$ to be the complete set of appropriately normalized eigenfunctions for $L = -\Delta$ in $H_0^1(U)$: see §6.5.1.)

Fix now a positive integer m . We will look for a function $\mathbf{u}_m : [0, T] \rightarrow H_0^1(U)$ of the form

$$(14) \quad \mathbf{u}_m(t) := \sum_{k=1}^m d_m^k(t) w_k,$$

where we hope to select the coefficients $d_m^k(t)$ ($0 \leq t \leq T$, $k = 1, \dots, m$) so that

$$(15) \quad d_m^k(0) = (g, w_k) \quad (k = 1, \dots, m)$$

and

$$(16) \quad (\mathbf{u}'_m, w_k) + B[\mathbf{u}_m, w_k; t] = (\mathbf{f}, w_k) \quad (0 \leq t \leq T, k = 1, \dots, m).$$

(Here, as before, $(\ , \)$ denotes the inner product in $L^2(U)$.)

Thus we seek a function \mathbf{u}_m of the form (14) that satisfies the “projection” (16) of problem (11) onto the finite-dimensional subspace spanned by $\{w_k\}_{k=1}^m$.

THEOREM 1 (Construction of approximate solutions). *For each integer $m = 1, 2, \dots$ there exists a unique function \mathbf{u}_m of the form (14) satisfying (15), (16).*

Proof. Assuming \mathbf{u}_m has the structure (14), we first note from (13) that

$$(17) \quad (\mathbf{u}'_m(t), w_k) = d_m^{k'}(t).$$

Furthermore

$$(18) \quad B[\mathbf{u}_m, w_k; t] = \sum_{l=1}^m e^{kl}(t) d_m^l(t),$$

for $e^{kl}(t) := B[w_l, w_k; t]$ ($k, l = 1, \dots, m$). Let us further write $f^k(t) := (\mathbf{f}(t), w_k)$ ($k = 1, \dots, m$). Then (16) becomes the linear system of ODE

$$(19) \quad d_m^{k'}(t) + \sum_{l=1}^m e^{kl}(t) d_m^l(t) = f^k(t) \quad (k = 1, \dots, m),$$

subject to the initial conditions (15). According to standard existence theory for ordinary differential equations, there exists a unique absolutely continuous function $\mathbf{d}_m(t) = (d_m^1(t), \dots, d_m^m(t))$ satisfying (15) and (19) for a.e. $0 \leq t \leq T$. And then \mathbf{u}_m defined by (14) solves (16) for a.e. $0 \leq t \leq T$. \square

b. Energy estimates. We propose now to send m to infinity and to show a subsequence of our solutions \mathbf{u}_m of the approximate problems (15), (16) converges to a weak solution of (11). For this we will need some uniform estimates.

THEOREM 2 (Energy estimates). *There exists a constant C , depending only on U, T and the coefficients of L , such that*

$$(20) \quad \max_{0 \leq t \leq T} \|\mathbf{u}_m(t)\|_{L^2(U)} + \|\mathbf{u}_m\|_{L^2(0, T; H_0^1(U))} + \|\mathbf{u}'_m\|_{L^2(0, T; H^{-1}(U))} \\ \leq C(\|\mathbf{f}\|_{L^2(0, T; L^2(U))} + \|g\|_{L^2(U)})$$

for $m = 1, 2, \dots$

Proof. 1. Multiply equation (16) by $d_m^k(t)$, sum for $k = 1, \dots, m$, and then recall (14) to find

$$(21) \quad (\mathbf{u}'_m, \mathbf{u}_m) + B[\mathbf{u}_m, \mathbf{u}_m; t] = (\mathbf{f}, \mathbf{u}_m)$$

for a.e. $0 \leq t \leq T$. We proved in §6.2.2 that there exist constants $\beta > 0$, $\gamma \geq 0$ such that

$$(22) \quad \beta \|\mathbf{u}_m\|_{H_0^1(U)}^2 \leq B[\mathbf{u}_m, \mathbf{u}_m; t] + \gamma \|\mathbf{u}_m\|_{L^2(U)}^2$$

for all $0 \leq t \leq T$, $m = 1, \dots$. Furthermore $|(\mathbf{f}, \mathbf{u}_m)| \leq \frac{1}{2}\|\mathbf{f}\|_{L^2(U)}^2 + \frac{1}{2}\|\mathbf{u}_m\|_{L^2(U)}^2$, and $(\mathbf{u}'_m, \mathbf{u}_m) = \frac{d}{dt}(\frac{1}{2}\|\mathbf{u}_m\|_{L^2(U)}^2)$ for a.e. $0 \leq t \leq T$. Consequently (21) yields the inequality

$$(23) \quad \frac{d}{dt} \left(\|\mathbf{u}_m\|_{L^2(U)}^2 \right) + 2\beta\|\mathbf{u}_m\|_{H_0^1(U)}^2 \leq C_1\|\mathbf{u}_m\|_{L^2(U)}^2 + C_2\|\mathbf{f}\|_{L^2(U)}^2$$

for a.e. $0 \leq t \leq T$ and appropriate constants C_1 and C_2 .

2. Now write

$$(24) \quad \eta(t) := \|\mathbf{u}_m(t)\|_{L^2(U)}^2$$

and

$$(25) \quad \xi(t) := \|\mathbf{f}(t)\|_{L^2(U)}^2.$$

Then (23) implies

$$\eta'(t) \leq C_1\eta(t) + C_2\xi(t)$$

for a.e. $0 \leq t \leq T$. Thus the differential form of Gronwall's inequality (§B.2) yields the estimate

$$(26) \quad \eta(t) \leq e^{C_1 t} \left(\eta(0) + C_2 \int_0^t \xi(s) ds \right) \quad (0 \leq t \leq T).$$

Since $\eta(0) = \|\mathbf{u}_m(0)\|_{L^2(U)}^2 \leq \|g\|_{L^2(U)}^2$ by (15), we obtain from (24)–(26) the estimate

$$\max_{0 \leq t \leq T} \|\mathbf{u}_m(t)\|_{L^2(U)}^2 \leq C \left(\|g\|_{L^2(U)}^2 + \|\mathbf{f}\|_{L^2(0,T;L^2(U))}^2 \right).$$

3. Returning once more to inequality (23), we integrate from 0 to T and employ the inequality above to find

$$\begin{aligned} \|\mathbf{u}_m\|_{L^2(0,T;H_0^1(U))}^2 &= \int_0^T \|\mathbf{u}_m\|_{H_0^1(U)}^2 dt \\ &\leq C \left(\|g\|_{L^2(U)}^2 + \|\mathbf{f}\|_{L^2(0,T;L^2(U))}^2 \right). \end{aligned}$$

4. Fix any $v \in H_0^1(U)$, with $\|v\|_{H_0^1(U)} \leq 1$, and write $v = v^1 + v^2$, where $v^1 \in \text{span}\{w_k\}_{k=1}^m$ and $(v^2, w_k) = 0$ ($k = 1, \dots, m$). Since the functions $\{w_k\}_{k=0}^\infty$ are orthogonal in $H_0^1(U)$, $\|v^1\|_{H_0^1(U)} \leq \|v\|_{H_0^1(U)} \leq 1$. Utilizing (16), we deduce for a.e. $0 \leq t \leq T$ that

$$(\mathbf{u}'_m, v^1) + B[\mathbf{u}_m, v^1; t] = (\mathbf{f}, v^1).$$

Then (14) implies

$$\langle \mathbf{u}'_m, v \rangle = (\mathbf{u}'_m, v) = (\mathbf{u}'_m, v^1) = (\mathbf{f}, v^1) - B[\mathbf{u}_m, v^1; t].$$

Consequently

$$|\langle \mathbf{u}'_m, v \rangle| \leq C(\|\mathbf{f}\|_{L^2(U)} + \|\mathbf{u}_m\|_{H_0^1(U)}),$$

since $\|v^1\|_{H_0^1(U)} \leq 1$. Thus

$$\|\mathbf{u}'_m\|_{H^{-1}(U)} \leq C(\|\mathbf{f}\|_{L^2(U)} + \|\mathbf{u}_m\|_{H_0^1(U)}),$$

and therefore

$$\begin{aligned} \int_0^T \|\mathbf{u}'_m\|_{H^{-1}(U)}^2 dt &\leq C \int_0^T \|\mathbf{f}\|_{L^2(U)}^2 + \|\mathbf{u}_m\|_{H_0^1(U)}^2 dt \\ &\leq C(\|g\|_{L^2(U)}^2 + \|\mathbf{f}\|_{L^2(0,T;L^2(U))}^2). \end{aligned} \quad \square$$

c. Existence and uniqueness. Next we pass to limits as $m \rightarrow \infty$, to build a weak solution of our initial/boundary-value problem (11).

THEOREM 3 (Existence of weak solution). *There exists a weak solution of (11).*

Proof. 1. According to the energy estimates (20), we see that the sequence $\{\mathbf{u}_m\}_{m=1}^\infty$ is bounded in $L^2(0, T; H_0^1(U))$ and $\{\mathbf{u}'_m\}_{m=1}^\infty$ is bounded in $L^2(0, T; H^{-1}(U))$.

Consequently there exists a subsequence $\{\mathbf{u}_{m_l}\}_{l=1}^\infty \subset \{\mathbf{u}_m\}_{m=1}^\infty$ and a function $\mathbf{u} \in L^2(0, T; H_0^1(U))$, with $\mathbf{u}' \in L^2(0, T; H^{-1}(U))$, such that

$$(27) \quad \begin{cases} \mathbf{u}_{m_l} \rightharpoonup \mathbf{u} & \text{weakly in } L^2(0, T; H_0^1(U)) \\ \mathbf{u}'_{m_l} \rightharpoonup \mathbf{u}' & \text{weakly in } L^2(0, T; H^{-1}(U)). \end{cases}$$

(See §D.4 and Problem 5.)

2. Next fix an integer N and choose a function $\mathbf{v} \in C^1([0, T]; H_0^1(U))$ having the form

$$(28) \quad \mathbf{v}(t) = \sum_{k=1}^N d^k(t) w_k,$$

where $\{d^k\}_{k=1}^N$ are given smooth functions. We choose $m \geq N$, multiply (16) by $d^k(t)$, sum $k = 1, \dots, N$, and then integrate with respect to t to find

$$(29) \quad \int_0^T \langle \mathbf{u}'_m, \mathbf{v} \rangle + B[\mathbf{u}_m, \mathbf{v}; t] dt = \int_0^T (\mathbf{f}, \mathbf{v}) dt.$$

We set $m = m_l$ and recall (27), to find upon passing to weak limits that

$$(30) \quad \int_0^T \langle \mathbf{u}', \mathbf{v} \rangle + B[\mathbf{u}, \mathbf{v}; t] dt = \int_0^T (\mathbf{f}, \mathbf{v}) dt.$$

This equality then holds for all functions $\mathbf{v} \in L^2(0, T; H_0^1(U))$, as functions of the form (28) are dense in this space. Hence in particular

$$(31) \quad \langle \mathbf{u}', v \rangle + B[\mathbf{u}, v; t] = (\mathbf{f}, v)$$

for each $v \in H_0^1(U)$ and a.e. $0 \leq t \leq T$. From Theorem 3 in §5.9.2 we see that furthermore $\mathbf{u} \in C([0, T]; L^2(U))$.

3. In order to prove $\mathbf{u}(0) = g$, we first note from (30) that

$$(32) \quad \int_0^T -\langle \mathbf{v}', \mathbf{u} \rangle + B[\mathbf{u}, \mathbf{v}; t] dt = \int_0^T (\mathbf{f}, \mathbf{v}) dt + (\mathbf{u}(0), \mathbf{v}(0))$$

for each $\mathbf{v} \in C^1([0, T]; H_0^1(U))$ with $\mathbf{v}(T) = 0$. Similarly, from (29) we deduce

$$(33) \quad \int_0^T -\langle \mathbf{v}', \mathbf{u}_m \rangle + B[\mathbf{u}_m, \mathbf{v}; t] dt = \int_0^T (\mathbf{f}, \mathbf{v}) dt + (\mathbf{u}_m(0), \mathbf{v}(0)).$$

We set $m = m_l$ and once again employ (27) to find

$$(34) \quad \int_0^T -\langle \mathbf{v}', \mathbf{u} \rangle + B[\mathbf{u}, \mathbf{v}; t] dt = \int_0^T (\mathbf{f}, \mathbf{v}) dt + (g, \mathbf{v}(0)),$$

since $\mathbf{u}_{m_l}(0) \rightarrow g$ in $L^2(U)$. As $\mathbf{v}(0)$ is arbitrary, comparing (32) and (34), we conclude $\mathbf{u}(0) = g$. \square

THEOREM 4 (Uniqueness of weak solutions). *A weak solution of (11) is unique.*

Proof. It suffices to check that the only weak solution of (11) with $\mathbf{f} \equiv g \equiv 0$ is

$$(35) \quad \mathbf{u} \equiv 0.$$

To prove this, observe that by setting $\mathbf{v} = \mathbf{u}$ in identity (31) (for $\mathbf{f} \equiv 0$) we learn, using Theorem 3 in §5.9.2, that

$$(36) \quad \frac{d}{dt} \left(\frac{1}{2} \|\mathbf{u}\|_{L^2(U)}^2 \right) + B[\mathbf{u}, \mathbf{u}; t] = \langle \mathbf{u}', \mathbf{u} \rangle + B[\mathbf{u}, \mathbf{u}; t] = 0.$$

Since

$$B[\mathbf{u}, \mathbf{u}; t] \geq \beta \|\mathbf{u}\|_{H_0^1(U)}^2 - \gamma \|\mathbf{u}\|_{L^2(U)}^2 \geq -\gamma \|\mathbf{u}\|_{L^2(U)}^2,$$

Gronwall's inequality and (36) imply (35). \square

7.1.3. Regularity.

In this section we discuss the regularity of our weak solutions u to the initial/boundary-value problem for second-order parabolic equations. Our eventual goal is to prove that u is smooth, provided the coefficients of the PDE, the boundary of the domain, etc. are smooth. The following presentation mirrors that from §6.3.

Motivation: formal derivation of estimates. (i) To gain some insight as to what regularity assertions could possibly be valid, let us temporarily suppose $u = u(x, t)$ is a smooth solution of this initial-value problem for the heat equation:

$$(37) \quad \begin{cases} u_t - \Delta u = f & \text{in } \mathbb{R}^n \times (0, T] \\ u = g & \text{on } \mathbb{R}^n \times \{t = 0\}, \end{cases}$$

and assume also u goes to zero as $|x| \rightarrow \infty$ sufficiently rapidly to justify the following computations. We then calculate for $0 \leq t \leq T$

$$(38) \quad \begin{aligned} \int_{\mathbb{R}^n} f^2 dx &= \int_{\mathbb{R}^n} (u_t - \Delta u)^2 dx \\ &= \int_{\mathbb{R}^n} u_t^2 - 2\Delta u u_t + (\Delta u)^2 dx \\ &= \int_{\mathbb{R}^n} u_t^2 + 2Du \cdot Du_t + (\Delta u)^2 dx. \end{aligned}$$

Now $2Du \cdot Du_t = \frac{d}{dt}(|Du|^2)$, and consequently

$$\int_0^t \int_{\mathbb{R}^n} 2Du \cdot Du_t dx ds = \int_{\mathbb{R}^n} |Du|^2 dx \Big|_{s=0}^{s=t}.$$

Furthermore, as demonstrated in §6.3,

$$\int_{\mathbb{R}^n} (\Delta u)^2 dx = \int_{\mathbb{R}^n} |D^2u|^2 dx.$$

We utilize the two equalities above in (38) and integrate in time, to obtain

$$(39) \quad \begin{aligned} \sup_{0 \leq t \leq T} \int_{\mathbb{R}^n} |Du|^2 dx + \int_0^T \int_{\mathbb{R}^n} u_t^2 + |D^2u|^2 dx dt \\ \leq C \left(\int_0^T \int_{\mathbb{R}^n} f^2 dx dt + \int_{\mathbb{R}^n} |Dg|^2 dx \right). \end{aligned}$$

We therefore see that we can estimate the L^2 -norms of u_t and D^2u within $\mathbb{R}^n \times (0, T)$, in terms of the L^2 -norm of f on $\mathbb{R}^n \times (0, T)$ and the L^2 -norm of Dg on \mathbb{R}^n .

(ii) Next differentiate the PDE with respect to t and set $\tilde{u} := u_t$. Then

$$(40) \quad \begin{cases} \tilde{u}_t - \Delta \tilde{u} = \tilde{f} & \text{in } \mathbb{R}^n \times (0, T] \\ \tilde{u} = \tilde{g} & \text{on } \mathbb{R}^n \times \{t = 0\}, \end{cases}$$

for $\tilde{f} := f_t$, $\tilde{g} := u_t(\cdot, 0) = f(\cdot, 0) + \Delta g$. Multiplying by \tilde{u} , integrating by parts and invoking Gronwall's inequality, we deduce

$$(41) \quad \sup_{0 \leq t \leq T} \int_{\mathbb{R}^n} |u_t|^2 dx + \int_0^T \int_{\mathbb{R}^n} |Du_t|^2 dx dt \leq C \left(\int_0^T \int_{\mathbb{R}^n} f_t^2 dx dt + \int_{\mathbb{R}^n} |D^2 g|^2 + f(\cdot, 0)^2 dx \right).$$

But

$$(42) \quad \max_{0 \leq t \leq T} \|f(\cdot, t)\|_{L^2(\mathbb{R}^n)} \leq C(\|f\|_{L^2(\mathbb{R}^n \times (0, T))} + \|f_t\|_{L^2(\mathbb{R}^n \times (0, T))}),$$

according to Theorem 2(iii) of §5.9.2. Furthermore, writing $-\Delta u = f - u_t$, we find as in §6.3 that

$$(43) \quad \int_{\mathbb{R}^n} |D^2 u|^2 dx \leq C \int_{\mathbb{R}^n} f^2 + u_t^2 dx.$$

Combining (41)–(43) leads us to the estimate

$$(44) \quad \sup_{0 \leq t \leq T} \int_{\mathbb{R}^n} |u_t|^2 + |D^2 u|^2 dx + \int_0^T \int_{\mathbb{R}^n} |Du_t|^2 dx dt \leq C \left(\int_0^T \int_{\mathbb{R}^n} f_t^2 + f^2 dx dt + \int_{\mathbb{R}^n} |D^2 g|^2 dx \right),$$

for some constant C . □

The foregoing formal computations suggest that we have estimates corresponding to (39) and (44) for our weak solution to a general second-order parabolic PDE. These calculations do not constitute a proof, however, since our weak solution of (11), constructed in §7.1.2, is not smooth enough to justify the foregoing computations.

We will instead calculate using the Galerkin approximations. To streamline the presentation, we hereafter assume that $\{w_k\}_{k=1}^\infty$ is the complete collection of eigenfunctions for $-\Delta$ on $H_0^1(U)$ and that U is bounded, open, with ∂U smooth. We furthermore suppose that

$$(45) \quad \begin{cases} \text{the coefficients } a^{ij}, b^i, c \text{ (} i, j = 1, \dots, n \text{) are smooth on} \\ \bar{U} \text{ and do not depend on } t. \end{cases}$$

THEOREM 5 (Improved regularity).

(i) Assume

$$g \in H_0^1(U), \mathbf{f} \in L^2(0, T; L^2(U)).$$

Suppose also $\mathbf{u} \in L^2(0, T; H_0^1(U))$, with $\mathbf{u}' \in L^2(0, T; H^{-1}(U))$, is the weak solution of

$$\begin{cases} u_t + Lu = f & \text{in } U_T \\ u = 0 & \text{on } \partial U \times [0, T] \\ u = g & \text{on } U \times \{t = 0\}. \end{cases}$$

Then in fact

$$\mathbf{u} \in L^2(0, T; H^2(U)) \cap L^\infty(0, T; H_0^1(U)), \mathbf{u}' \in L^2(0, T; L^2(U)),$$

and we have the estimate

$$(46) \quad \operatorname{ess\,sup}_{0 \leq t \leq T} \|\mathbf{u}(t)\|_{H_0^1(U)} + \|\mathbf{u}\|_{L^2(0, T; H^2(U))} + \|\mathbf{u}'\|_{L^2(0, T; L^2(U))} \leq C \left(\|\mathbf{f}\|_{L^2(0, T; L^2(U))} + \|g\|_{H_0^1(U)} \right),$$

the constant C depending only on U, T and the coefficients of L .

(ii) If, in addition,

$$g \in H^2(U), \mathbf{f}' \in L^2(0, T; L^2(U)),$$

then

$$\begin{aligned} \mathbf{u} &\in L^\infty(0, T; H^2(U)), \mathbf{u}' \in L^\infty(0, T; L^2(U)) \cap L^2(0, T; H_0^1(U)), \\ \mathbf{u}'' &\in L^2(0, T; H^{-1}(U)), \end{aligned}$$

with the estimate

$$(47) \quad \operatorname{ess\,sup}_{0 \leq t \leq T} (\|\mathbf{u}(t)\|_{H^2(U)} + \|\mathbf{u}'(t)\|_{L^2(U)}) + \|\mathbf{u}'\|_{L^2(0, T; H_0^1(U))} + \|\mathbf{u}''\|_{L^2(0, T; H^{-1}(U))} \leq C (\|\mathbf{f}\|_{H^1(0, T; L^2(U))} + \|g\|_{H^2(U)}).$$

Assertions (i), (ii) of Theorem 5 are precise versions of the formal estimates (39), (44) (for the heat equation on $U = \mathbb{R}^n$).

Proof. 1. Fixing $m \geq 1$, we multiply equation (16) in §7.1.2 by $d_m^k(t)$ and sum $k = 1, \dots, m$, to discover

$$(\mathbf{u}'_m, \mathbf{u}'_m) + B[\mathbf{u}_m, \mathbf{u}'_m] = (\mathbf{f}, \mathbf{u}'_m)$$

for a.e. $0 \leq t \leq T$. Now

$$\begin{aligned} B[\mathbf{u}_m, \mathbf{u}'_m] &= \int_U \sum_{i,j=1}^n a^{ij} \mathbf{u}_{m,x_i} \mathbf{u}'_{m,x_j} dx \\ &\quad + \int_U \sum_{i=1}^n b^i \mathbf{u}_{m,x_i} \mathbf{u}'_m + c \mathbf{u}_m \mathbf{u}'_m dx \\ &=: A + B. \end{aligned}$$

Since $a^{ij} = a^{ji}$ ($i, j = 1, \dots, n$) and these coefficients do not depend on t , we see $A = \frac{d}{dt} \left(\frac{1}{2} A[\mathbf{u}_m, \mathbf{u}_m] \right)$, for the symmetric bilinear form

$$A[u, v] := \int_U \sum_{i,j=1}^n a^{ij} u_{x_i} v_{x_j} dx \quad (u, v \in H_0^1(U)).$$

Furthermore,

$$|B| \leq \frac{C}{\epsilon} \|\mathbf{u}_m\|_{H_0^1(U)}^2 + \epsilon \|\mathbf{u}'_m\|_{L^2(U)}^2, \quad |(\mathbf{f}, \mathbf{u}'_m)| \leq \frac{C}{\epsilon} \|\mathbf{f}\|_{L^2(U)}^2 + \epsilon \|\mathbf{u}'_m\|_{L^2(U)}^2$$

for each $\epsilon > 0$.

2. Combining the above inequalities, we deduce

$$\begin{aligned} \|\mathbf{u}'_m\|_{L^2(U)}^2 + \frac{d}{dt} \left(\frac{1}{2} A[\mathbf{u}_m, \mathbf{u}_m] \right) \\ \leq \frac{C}{\epsilon} (\|\mathbf{u}_m\|_{H_0^1(U)}^2 + \|\mathbf{f}\|_{L^2(U)}^2) + 2\epsilon \|\mathbf{u}'_m\|_{L^2(U)}^2. \end{aligned}$$

Choosing $\epsilon = \frac{1}{4}$ and integrating, we find

$$\begin{aligned} \int_0^T \|\mathbf{u}'_m\|_{L^2(U)}^2 dt + \sup_{0 \leq t \leq T} A[\mathbf{u}_m(t), \mathbf{u}_m(t)] \\ \leq C \left(A[\mathbf{u}_m(0), \mathbf{u}_m(0)] + \int_0^T \|\mathbf{u}_m\|_{H_0^1(U)}^2 + \|\mathbf{f}\|_{L^2(U)}^2 dt \right) \\ \leq C (\|g\|_{H_0^1(U)}^2 + \|\mathbf{f}\|_{L^2(0,T;L^2(U))}^2), \end{aligned}$$

according to Theorem 2 in §7.1.2, where we estimated $\|\mathbf{u}_m(0)\|_{H_0^1(U)} \leq \|g\|_{H_0^1(U)}$. As $A[u, u] \geq \theta \int_U |Du|^2 dx$ for each $u \in H_0^1(U)$, we find that

$$(48) \quad \sup_{0 \leq t \leq T} \|\mathbf{u}_m(t)\|_{H_0^1(U)}^2 \leq C (\|g\|_{H_0^1(U)}^2 + \|\mathbf{f}\|_{L^2(0,T;L^2(U))}^2).$$

Passing to limits as $m = m_l \rightarrow \infty$, we deduce $\mathbf{u} \in L^\infty(0, T; H_0^1(U))$, $\mathbf{u}' \in L^2(0, T; L^2(U))$, with the stated bounds; cf. Problem 6.

3. In particular, for a.e. t we have the identity

$$(\mathbf{u}', v) + B[\mathbf{u}, v] = (\mathbf{f}, v)$$

for each $v \in H_0^1(U)$. This equality we rewrite to read

$$B[\mathbf{u}, v] = (\mathbf{h}, v)$$

for $\mathbf{h} := \mathbf{f} - \mathbf{u}'$. Since $\mathbf{h}(t) \in L^2(U)$ for a.e. $0 \leq t \leq T$, we deduce from the elliptic regularity Theorem 4 in §6.3.2 that $\mathbf{u}(t) \in H^2(U)$ for a.e. $0 \leq t \leq T$, with the estimate

$$\begin{aligned} \|\mathbf{u}\|_{H^2(U)}^2 &\leq C(\|\mathbf{h}\|_{L^2(U)}^2 + \|\mathbf{u}\|_{L^2(U)}^2) \\ (49) \quad &\leq C(\|\mathbf{f}\|_{L^2(U)}^2 + \|\mathbf{u}'\|_{L^2(U)}^2 + \|\mathbf{u}\|_{L^2(U)}^2). \end{aligned}$$

Integrating and utilizing the estimates from step 2, we complete the proof of (i).

4. The goal next is to establish higher regularity for our weak solution. So now suppose $g \in H^2(U) \cap H_0^1(U)$, $\mathbf{f} \in H^1(0, T; L^2(U))$. Fix $m \geq 1$ and differentiate equation (16) in §7.1.2 with respect to t . Owing to (45), we find

$$(50) \quad (\tilde{\mathbf{u}}'_m, w_k) + B[\tilde{\mathbf{u}}_m, w_k] = (\mathbf{f}', w_k) \quad (k = 1, \dots, m),$$

where $\tilde{\mathbf{u}}_m := \mathbf{u}'_m$. Multiply (50) by $d_m^k(t)$ and sum $k = 1, \dots, m$:

$$(\tilde{\mathbf{u}}'_m, \tilde{\mathbf{u}}_m) + B[\tilde{\mathbf{u}}_m, \tilde{\mathbf{u}}_m] = (\mathbf{f}', \tilde{\mathbf{u}}_m).$$

Employing Gronwall's inequality, we deduce

$$\begin{aligned} (51) \quad \sup_{0 \leq t \leq T} \|\mathbf{u}'_m(t)\|_{L^2(U)}^2 + \int_0^T \|\mathbf{u}'_m\|_{H_0^1(U)}^2 dt \\ \leq C(\|\mathbf{u}'_m(0)\|_{L^2(U)}^2 + \|\mathbf{f}'\|_{L^2(0, T; L^2(U))}^2) \\ \leq C(\|\mathbf{f}\|_{H^1(0, T; L^2(U))}^2 + \|\mathbf{u}_m(0)\|_{H^2(U)}^2). \end{aligned}$$

We employed (16) in the last inequality.

5. We must estimate the last term in (51). Remember that we have taken $\{w_k\}_{k=1}^\infty$ to be the complete collection of (smooth) eigenfunctions for $-\Delta$ on $H_0^1(U)$. In particular, $\Delta \mathbf{u}_m = 0$ on ∂U . Thus

$$\|\mathbf{u}_m(0)\|_{H^2(U)}^2 \leq C\|\Delta \mathbf{u}_m(0)\|_{L^2(U)}^2 = C(\mathbf{u}_m(0), \Delta^2 \mathbf{u}_m(0)).$$

Since $\Delta^2 \mathbf{u}_m(0) \in \text{span}\{w_k\}_{k=1}^m$ and $(\mathbf{u}_m(0), w_k) = (g, w_k)$ for $k = 1, \dots, m$, we have

$$\begin{aligned} \|\mathbf{u}_m(0)\|_{H^2(U)}^2 &\leq C(g, \Delta^2 \mathbf{u}_m(0)) = C(\Delta g, \Delta \mathbf{u}_m(0)) \\ &\leq \frac{1}{2} \|\mathbf{u}_m(0)\|_{H^2(U)}^2 + C\|g\|_{H^2(U)}^2. \end{aligned}$$

Hence $\|\mathbf{u}_m(0)\|_{H^2(U)} \leq C\|g\|_{H^2(U)}$. Therefore (51) implies

$$(52) \quad \sup_{0 \leq t \leq T} \|\mathbf{u}'_m(t)\|_{L^2(U)}^2 + \int_0^T \|\mathbf{u}'_m\|_{H_0^1(U)}^2 dt \leq C(\|\mathbf{f}\|_{H^1(0,T;L^2(U))}^2 + \|g\|_{H^2(U)}^2).$$

6. Now

$$B[\mathbf{u}_m, w_k] = (\mathbf{f} - \mathbf{u}'_m, w_k) \quad (k = 1, \dots, m).$$

Let λ_k denote the k^{th} eigenvalue of $-\Delta$ on $H_0^1(U)$. Multiplying the above identity by $\lambda_k d_m^k(t)$ and summing $k = 1, \dots, m$, we deduce for $0 \leq t \leq T$ that

$$(53) \quad B[\mathbf{u}_m, -\Delta \mathbf{u}_m] = (\mathbf{f} - \mathbf{u}'_m, -\Delta \mathbf{u}_m).$$

Since $\Delta \mathbf{u}_m = 0$ on ∂U , we see $B[\mathbf{u}_m, -\Delta \mathbf{u}_m] = (L\mathbf{u}_m, -\Delta \mathbf{u}_m)$. Next we invoke the inequality

$$(54) \quad \beta \|u\|_{H^2(U)}^2 \leq (Lu, -\Delta u) + \gamma \|u\|_{L^2(U)}^2 \quad (u \in H^2(U) \cap H_0^1(U))$$

for constants $\beta > 0$, $\gamma \geq 0$. (See Problem 9 and also the remark following the proof.)

We thereupon conclude from (53) that

$$\|\mathbf{u}_m\|_{H^2(U)} \leq C(\|\mathbf{f}\|_{L^2(U)} + \|\mathbf{u}'_m\|_{L^2(U)} + \|\mathbf{u}_m\|_{L^2(U)}).$$

This inequality, (52), (48) and Theorem 2 in §5.9.2 imply

$$\begin{aligned} \sup_{0 \leq t \leq T} (\|\mathbf{u}'_m(t)\|_{L^2(U)}^2 + \|\mathbf{u}_m(t)\|_{H^2(U)}^2) + \int_0^T \|\mathbf{u}'_m\|_{H_0^1(U)}^2 dt \\ \leq C(\|\mathbf{f}\|_{H^1(0,T;L^2(U))}^2 + \|g\|_{H^2(U)}^2). \end{aligned}$$

Passing to limits as $m = m_l \rightarrow \infty$, we deduce the same bound for \mathbf{u} .

7. It remains to show $\mathbf{u}'' \in L^2(0, T; H^{-1}(U))$. To do so, take $v \in H_0^1(U)$, with $\|v\|_{H_0^1(U)} \leq 1$, and write $v = v^1 + v^2$, as in the proof of Theorem 2 in §7.1.2. Then for a.e. time $0 \leq t \leq T$

$$\langle \mathbf{u}''_m, v \rangle = (\mathbf{u}''_m, v) = (\mathbf{u}''_m, v^1) = (\mathbf{f}', v^1) - B[\mathbf{u}'_m, v^1]$$

according to (50), since $\mathbf{u}''_m = \tilde{\mathbf{u}}'_m$. Consequently

$$|\langle \mathbf{u}''_m, v \rangle| \leq C(\|\mathbf{f}'\|_{L^2(U)} + \|\mathbf{u}'_m\|_{H^1_0(U)}),$$

since $\|v^1\|_{H^1_0(U)} \leq 1$. Thus

$$\|\mathbf{u}''_m\|_{H^{-1}(U)} \leq C(\|\mathbf{f}'\|_{L^2(U)} + \|\mathbf{u}'_m\|_{H^1_0(U)}),$$

and so \mathbf{u}''_m is bounded in $L^2(0, T; H^{-1}(U))$. Passing to limits, we deduce that $\mathbf{u}'' \in L^2(0, T; H^{-1}(U))$, with the stated estimate. \square

Remark. If L were symmetric, we could alternatively have taken $\{w_k\}_{k=1}^\infty$ to be a basis of eigenfunctions of L on $H^1_0(U)$ and so avoided the need for inequality (54).

Let us now build upon the previous regularity assertion:

THEOREM 6 (Higher regularity). *Assume*

$$g \in H^{2m+1}(U), \quad \frac{d^k \mathbf{f}}{dt^k} \in L^2(0, T; H^{2m-2k}(U)) \quad (k = 0, \dots, m).$$

Suppose also that the following m^{th} -order compatibility conditions hold:

$$\begin{cases} g_0 := g \in H^1_0(U), & g_1 := \mathbf{f}(0) - Lg_0 \in H^1_0(U), \\ \dots, & g_m := \frac{d^{m-1} \mathbf{f}}{dt^{m-1}}(0) - Lg_{m-1} \in H^1_0(U). \end{cases}$$

Then

$$\frac{d^k \mathbf{u}}{dt^k} \in L^2(0, T; H^{2m+2-2k}(U)) \quad (k = 0, \dots, m+1);$$

and we have the estimate

$$(55) \quad \sum_{k=0}^{m+1} \left\| \frac{d^k \mathbf{u}}{dt^k} \right\|_{L^2(0, T; H^{2m+2-2k}(U))} \leq C \left(\sum_{k=0}^m \left\| \frac{d^k \mathbf{f}}{dt^k} \right\|_{L^2(0, T; H^{2m-2k}(U))} + \|g\|_{H^{2m+1}(U)} \right),$$

the constant C depending only on m , U , T and the coefficients of L .

Remark. Taking into account Theorem 4 in §5.9.2, we see that

$$\mathbf{f}(0) \in H^{2m-1}(U), \quad \mathbf{f}'(0) \in H^{2m-3}(U), \quad \dots, \quad \mathbf{f}^{(m-1)}(0) \in H^1(U),$$

and consequently

$$g_0 \in H^{2m+1}(U), \quad g_1 \in H^{2m-1}(U), \quad \dots, \quad g_m \in H^1(U).$$

The compatibility conditions are consequently the requirements that in addition each of these functions equals 0 on ∂U , in the trace sense.

Proof. 1. The proof is an induction on m , the case $m = 0$ being Theorem 5(i) above.

Assume now the theorem is valid for some nonnegative integer m , and suppose then

$$(56) \quad g \in H^{2m+3}(U), \quad \frac{d^k \mathbf{f}}{dt^k} \in L^2(0, T; H^{2m+2-2k}(U)) \quad (k = 0, \dots, m+1)$$

and the $(m+1)$ th-order compatibility conditions hold. Now set $\tilde{\mathbf{u}} := \mathbf{u}'$. Differentiating the PDE with respect to t , we check that $\tilde{\mathbf{u}}$ is the unique weak solution of

$$(57) \quad \begin{cases} \tilde{\mathbf{u}}_t + L\tilde{\mathbf{u}} = \tilde{\mathbf{f}} & \text{in } U_T \\ \tilde{\mathbf{u}} = 0 & \text{on } \partial U \times [0, T] \\ \tilde{\mathbf{u}} = \tilde{\mathbf{g}} & \text{on } U \times \{t = 0\}, \end{cases}$$

for $\tilde{\mathbf{f}} := \mathbf{f}_t$, $\tilde{\mathbf{g}} := \mathbf{f}(\cdot, 0) - Lg$. In particular, for $m = 0$ we rely upon Theorem 5(ii) to be sure that $\tilde{\mathbf{u}} \in L^2(0, T; H_0^1(U))$, $\tilde{\mathbf{u}}' \in L^2(0, T; H^{-1}(U))$.

Since f and g satisfy the $(m+1)$ th-order compatibility conditions, it follows that $\tilde{\mathbf{f}}$ and $\tilde{\mathbf{g}}$ satisfy the m th-order compatibility condition. Thus applying the induction assumption, we deduce

$$\frac{d^k \tilde{\mathbf{u}}}{dt^k} \in L^2(0, T; H^{2m+2-2k}(U)) \quad (k = 0, \dots, m+1)$$

and

$$\begin{aligned} \sum_{k=0}^{m+1} \left\| \frac{d^k \tilde{\mathbf{u}}}{dt^k} \right\|_{L^2(0, T; H^{2m+2-2k}(U))} \\ \leq C \left(\sum_{k=0}^m \left\| \frac{d^k \tilde{\mathbf{f}}}{dt^k} \right\|_{L^2(0, T; H^{2m-2k}(U))} + \|\tilde{\mathbf{g}}\|_{H^{2m+1}(U)} \right) \end{aligned}$$

for $\tilde{\mathbf{f}} := \mathbf{f}'$. Since $\tilde{\mathbf{u}} = \mathbf{u}'$, we can rewrite the foregoing:

$$\frac{d^k \mathbf{u}}{dt^k} \in L^2(0, T; H^{2m+4-2k}(U)) \quad (k = 1, \dots, m+2),$$

$$(58) \quad \begin{aligned} \sum_{k=1}^{m+2} \left\| \frac{d^k \mathbf{u}}{dt^k} \right\|_{L^2(0, T; H^{2m+4-2k}(U))} \\ \leq C \left(\sum_{k=1}^{m+1} \left\| \frac{d^k \mathbf{f}}{dt^k} \right\|_{L^2(0, T; H^{2m+2-2k}(U))} + \|\mathbf{f}(0)\|_{H^{2m+1}(U)} + \|Lg\|_{H^{2m+1}(U)} \right) \\ \leq C \left(\sum_{k=0}^{m+1} \left\| \frac{d^k \mathbf{f}}{dt^k} \right\|_{L^2(0, T; H^{2m+2-2k}(U))} + \|g\|_{H^{2m+3}(U)} \right). \end{aligned}$$

Here we used the estimate

$$(59) \quad \|\mathbf{f}(0)\|_{H^{2m+1}(U)} \leq C(\|\mathbf{f}\|_{L^2(0,T;H^{2m+2}(U))} + \|\mathbf{f}'\|_{L^2(0,T;H^{2m}(U))}),$$

which follows from Theorem 4 in §5.9.2.

2. Now write for a.e. $0 \leq t \leq T$: $Lu = \mathbf{f} - \mathbf{u}' =: \mathbf{h}$. According to Theorem 5 in §6.3.2, we have

$$\begin{aligned} \|\mathbf{u}\|_{H^{2m+4}(U)} &\leq C(\|\mathbf{h}\|_{H^{2m+2}(U)} + \|\mathbf{u}\|_{L^2(U)}) \\ &\leq C(\|\mathbf{f}\|_{H^{2m+2}(U)} + \|\mathbf{u}'\|_{H^{2m+2}(U)} + \|\mathbf{u}\|_{L^2(U)}). \end{aligned}$$

Integrating with respect to t from 0 to T and adding the resulting expression to (58), we deduce

$$(60) \quad \begin{aligned} &\sum_{k=0}^{m+2} \left\| \frac{d^k \mathbf{u}}{dt^k} \right\|_{L^2(0,T;H^{2m+4-2k}(U))} \\ &\leq C \left(\sum_{k=0}^{m+1} \left\| \frac{d^k \mathbf{f}}{dt^k} \right\|_{L^2(0,T;H^{2m+2-2k}(U))} + \|g\|_{H^{2m+3}(U)} + \|\mathbf{u}\|_{L^2(0,T;L^2(U))} \right). \end{aligned}$$

Since

$$\|\mathbf{u}\|_{L^2(0,T;L^2(U))} \leq C(\|\mathbf{f}\|_{L^2(0,T;L^2(U))} + \|g\|_{L^2(U)}),$$

we thereby obtain the assertion of the theorem for $m + 1$. \square

THEOREM 7 (Infinite differentiability). *Assume*

$$g \in C^\infty(\bar{U}), \quad f \in C^\infty(\bar{U}_T),$$

and the m^{th} -order compatibility conditions hold for $m = 0, 1, \dots$. Then the parabolic initial/boundary-value problem (11) has a unique solution

$$u \in C^\infty(\bar{U}_T).$$

Proof. Apply Theorem 6 for $m = 0, 1, \dots$ \square

As we did for elliptic operators in Chapter 6, we have succeeded in repeatedly applying fairly straightforward “energy” estimates to produce a smooth solution of our parabolic initial/boundary-value problem (1). This assertion requires that the compatibility conditions (53) hold for all m , and it is easy to see that these conditions are necessary for the existence of a solution smooth on all of \bar{U}_T .

Interior estimates, analogous to those developed for elliptic PDE in §6.3.1, can also be derived, and these in particular do not require the compatibility conditions.

7.1.4. Maximum principles.

This section develops the maximum principle and Harnack's inequality for second-order parabolic operators.

a. Weak maximum principle. We will from now on assume that the operator L has the nondivergence form

$$(61) \quad Lu = - \sum_{i,j=1}^n a^{ij} u_{x_i x_j} + \sum_{i=1}^n b^i u_{x_i} + cu,$$

where the coefficients a^{ij}, b^i, c are continuous. We will always suppose the uniform parabolicity condition from §7.1.1 and also that $a^{ij} = a^{ji}$ ($i, j = 1, \dots, n$). Recall also that the parabolic boundary of U_T is $\Gamma_T = \bar{U}_T - U_T$.

THEOREM 8 (Weak maximum principle). *Assume $u \in C_1^2(U_T) \cap C(\bar{U}_T)$ and*

$$(62) \quad c \equiv 0 \quad \text{in } U_T.$$

(i) *If*

$$(63) \quad u_t + Lu \leq 0 \quad \text{in } U_T,$$

then

$$\max_{\bar{U}_T} u = \max_{\Gamma_T} u.$$

(ii) *Likewise, if*

$$(64) \quad u_t + Lu \geq 0 \quad \text{in } U_T,$$

then

$$\min_{\bar{U}_T} u = \min_{\Gamma_T} u.$$

A function u satisfying the inequality (63) is called a *subsolution*, and so we are asserting that a subsolution attains its maximum on the parabolic boundary Γ_T . Similarly, u is a *supersolution* provided (64) holds, in which case u attains its minimum on Γ_T .

Proof. 1. Let us first suppose we have the strict inequality

$$(65) \quad u_t + Lu < 0 \quad \text{in } U_T,$$

but there exists a point $(x_0, t_0) \in U_T$ with $u(x_0, t_0) = \max_{\bar{U}_T} u$.

2. If $0 < t_0 < T$, then (x_0, t_0) belongs to the interior of U_T and consequently

$$(66) \quad u_t = 0 \quad \text{at } (x_0, t_0),$$

since u attains its maximum at this point. On the other hand $Lu \geq 0$ at (x_0, t_0) , as explained in the proof of Theorem 1 in §6.4. Thus $u_t + Lu \geq 0$ at (x_0, t_0) , a contradiction to (65).

3. Now suppose $t_0 = T$. Then since u attains its maximum over \bar{U}_T at (x_0, t_0) , we see

$$u_t \geq 0 \quad \text{at } (x_0, t_0).$$

Since we still have the inequality $Lu \geq 0$ at (x_0, t_0) , we once more deduce the contradiction

$$u_t + Lu \geq 0 \quad \text{at } (x_0, t_0).$$

4. In the general case that (63) holds, write $u^\epsilon(x, t) := u(x, t) - \epsilon t$ where $\epsilon > 0$. Then

$$u_t^\epsilon + Lu^\epsilon = u_t + Lu - \epsilon < 0 \quad \text{in } U_T,$$

and so $\max_{\bar{U}_T} u^\epsilon = \max_{\Gamma_T} u^\epsilon$. Let $\epsilon \rightarrow 0$ to find $\max_{\bar{U}_T} u = \max_{\Gamma_T} u$. This proves assertion (i).

5. As $-u$ is a subsolution whenever u is a supersolution, assertion (ii) follows at once. \square

Next we allow for zeroth-order terms.

THEOREM 9 (Weak maximum principle for $c \geq 0$). *Assume $u \in C_1^2(U_T) \cap C(\bar{U}_T)$ and*

$$c \geq 0 \quad \text{in } U_T.$$

(i) *If*

$$u_t + Lu \leq 0 \quad \text{in } U_T,$$

then

$$\max_{\bar{U}_T} u \leq \max_{\Gamma_T} u^+.$$

(ii) *If*

$$u_t + Lu \geq 0 \quad \text{in } U_T,$$

then

$$\min_{\bar{U}_T} u \geq -\max_{\Gamma_T} u^-.$$

Remark. In particular, if $u_t + Lu = 0$ within U_T , then

$$\max_{\bar{U}_T} |u| = \max_{\Gamma_T} |u|.$$

Proof. 1. Assume u satisfies

$$(67) \quad u_t + Lu < 0 \quad \text{in } U_T$$

and attains a *positive* maximum at a point $(x_0, t_0) \in U_T$. Since $u(x_0, t_0) > 0$ and $c \geq 0$, we as above derive the contradiction

$$u_t + Lu \geq 0 \quad \text{at } (x_0, t_0).$$

2. If instead of (67), we have only

$$u_t + Lu \leq 0 \quad \text{in } U_T,$$

then as before $u^\epsilon(x, t) := u(x, t) - \epsilon t$ satisfies

$$u_t^\epsilon + Lu^\epsilon < 0 \quad \text{in } U_T.$$

Furthermore if u attains a positive maximum at some point in U_T , then u^ϵ also attains a positive maximum at some point belonging to U_T , provided $\epsilon > 0$ is small enough. But then, as in the previous proof, we obtain a contradiction.

3. Assertion (ii) follows similarly. □

Remark. Unlike the situation for elliptic PDE, various versions of the maximum principle obtain for parabolic PDE, even if the zeroth-order coefficient is negative: see Problem 8.

b. Harnack's inequality. Harnack's inequality states that if u is a non-negative solution of our parabolic PDE, then the maximum of u in some interior region at a positive time can be estimated by the minimum of u in the same region *at a later time*.

THEOREM 10 (Parabolic Harnack inequality). *Assume $u \in C_1^2(U_T)$ solves*

$$(68) \quad u_t + Lu = 0 \quad \text{in } U_T$$

and

$$u \geq 0 \quad \text{in } U_T.$$

Suppose $V \subset\subset U$ is connected. Then for each $0 < t_1 < t_2 \leq T$, there exists a constant C such that

$$(69) \quad \sup_V u(\cdot, t_1) \leq C \inf_V u(\cdot, t_2).$$

The constant C depends only on V , t_1, t_2 , and the coefficients of L .

This is true if the coefficients are continuous, or even merely bounded and measurable; see Lieberman [Lb]. We will however provide a proof only for the special case that $b^i \equiv c \equiv 0$ and the a^{ij} are smooth ($i, j = 1, \dots, n$). The following computations are elementary but tricky.

Proof*. 1. We may assume $u > 0$ in U_T , for otherwise we could apply the result to $u + \epsilon$ and then let $\epsilon \rightarrow 0^+$.

Set

$$(70) \quad v := \log u \quad \text{in } U_T.$$

Using (68), we compute

$$(71) \quad v_t = \sum_{i,j=1}^n a^{ij} v_{x_i x_j} + a^{ij} v_{x_i} v_{x_j} \quad \text{in } U_T.$$

Define

$$(72) \quad w := \sum_{i,j=1}^n a^{ij} v_{x_i x_j}, \quad \tilde{w} := \sum_{i,j=1}^n a^{ij} v_{x_i} v_{x_j},$$

so that (71) reads

$$(73) \quad v_t = w + \tilde{w}.$$

2. We calculate using (72), (73) for $k, l = 1, \dots, n$ that

$$v_{x_k x_l t} = w_{x_k x_l} + \sum_{i,j=1}^n 2a^{ij} v_{x_i x_k x_l} v_{x_j} + 2a^{ij} v_{x_i x_k} v_{x_j x_l} + R,$$

where the remainder term R satisfies an estimate of the form

$$(74) \quad |R| \leq \epsilon |D^2 v|^2 + C(\epsilon) |Dv|^2 + C$$

*Omit on first reading.

for each $\epsilon > 0$. Thus

$$\begin{aligned} w_t &= \sum_{k,l=1}^n a^{kl} v_{x_k x_l t} + a_t^{kl} v_{x_k x_l} \\ &= \sum_{k,l=1}^n a^{kl} w_{x_k x_l} + 2 \sum_{i,j=1}^n a^{ij} v_{x_j} w_{x_i} \\ &\quad + 2 \sum_{i,j,k,l=1}^n a^{ij} a^{kl} v_{x_i x_k} v_{x_j x_l} + R, \end{aligned}$$

where R now denotes another remainder term satisfying estimate (74). Therefore choosing $\epsilon > 0$ small enough in (74) and remembering the uniform parabolicity condition, we discover

$$(75) \quad w_t - \sum_{k,l=1}^n a^{kl} w_{x_k x_l} + \sum_{k=1}^n b^k w_{x_k} \geq \theta^2 |D^2 v|^2 - C |Dv|^2 - C,$$

where

$$(76) \quad b^k := -2 \sum_{l=1}^n a^{kl} v_{x_l} \quad (k = 1, \dots, n).$$

3. Estimate (75) is a differential inequality for w , and our task next is to obtain a similar inequality for \tilde{w} . Indeed, using (72) and (71), we compute

$$\begin{aligned} \tilde{w}_t - \sum_{k,l=1}^n a^{kl} \tilde{w}_{x_k x_l} &= 2 \sum_{i,j=1}^n a^{ij} v_{x_i} \left(v_{t x_j} - \sum_{k,l=1}^n a^{kl} v_{x_k x_l x_j} \right) \\ &\quad - 2 \sum_{i,j,k,l=1}^n a^{ij} a^{kl} v_{x_i x_k} v_{x_j x_l} + R, \end{aligned}$$

R yet another remainder term satisfying (74) for each $\epsilon > 0$. Recalling (71) and (76), we simplify to discover that

$$(77) \quad \begin{aligned} \tilde{w}_t - \sum_{k,l=1}^n a^{kl} \tilde{w}_{x_k x_l} + \sum_{k=1}^n b^k \tilde{w}_{x_k} \\ \geq -C |D^2 v|^2 - C |Dv|^2 - C \quad \text{in } U_T. \end{aligned}$$

4. Next set

$$(78) \quad \hat{w} := w + \kappa \tilde{w},$$

$\kappa > 0$ to be selected below. Combining (75) and (77), we deduce

$$(79) \quad \hat{w}_t - \sum_{k,l=1}^n a^{kl} \hat{w}_{x_k x_l} + \sum_{k=1}^n b^k \hat{w}_{x_k} \geq \frac{\theta^2}{2} |D^2 v|^2 - C |Dv|^2 - C$$

provided $0 < \kappa \leq \frac{1}{2}$ is now fixed to be sufficiently small.

5. Suppose next $V \subset\subset U$ is an open ball and $0 < t_1 < t_2 \leq T$. Choose a cutoff function $\zeta \in C^\infty(U_T)$ such that

$$(80) \quad \begin{cases} 0 \leq \zeta \leq 1, \zeta = 0 \text{ on } \Gamma_T, \\ \zeta \equiv 1 \text{ on } V \times [t_1, t_2]. \end{cases}$$

Note that ζ vanishes along $\{t = 0\}$.

Let μ be a positive constant (to be adjusted below), and assume then

$$(81) \quad \begin{cases} \zeta^4 \hat{w} + \mu t \text{ attains a negative minimum} \\ \text{at some point } (x_0, t_0) \in U \times (0, T]. \end{cases}$$

Consequently

$$(82) \quad \zeta \hat{w}_{x_k} + 4\zeta_{x_k} \hat{w} = 0 \quad \text{at } (x_0, t_0) \quad (k = 1, \dots, n).$$

In addition

$$0 \geq (\zeta^4 \hat{w} + \mu t)_t - \sum_{k,l=1}^n a^{kl} (\zeta^4 \hat{w} + \mu t)_{x_k x_l} \quad \text{at } (x_0, t_0).$$

Hence at the point (x_0, t_0) ,

$$(83) \quad 0 \geq \mu + \zeta^4 \left(\hat{w}_t - \sum_{k,l=1}^n a^{kl} \hat{w}_{x_k x_l} \right) - 2 \sum_{k,l=1}^n a^{kl} (\zeta^4)_{x_l} \hat{w}_{x_k} + \hat{R},$$

where

$$(84) \quad |\hat{R}| \leq C \zeta^2 |\hat{w}|.$$

Recall now (79) and (82):

$$0 \geq \mu + \zeta^4 \left(\frac{\theta^2}{2} |D^2 v|^2 - C |Dv|^2 - C - \sum_{k=1}^n b^k \hat{w}_{x_k} \right) + \hat{R},$$

\hat{R} another remainder term satisfying estimate (84). Utilizing (82) and (76), we deduce

$$(85) \quad 0 \geq \mu + \zeta^4 \left(\frac{\theta^2}{2} |D^2 v|^2 - C |Dv|^2 - C \right) + \tilde{R},$$

where now

$$(86) \quad |\tilde{R}| \leq C\zeta^2|\hat{w}| + C\zeta^3|Dv||\hat{w}|.$$

Remember that (85), (86) are valid at the point (x_0, t_0) where $\zeta^4\hat{w} + \mu t$ attains a negative minimum. In particular, at this point $\hat{w} = w + \kappa\tilde{w} < 0$. Recalling the definition (72) of w, \tilde{w} , we deduce

$$(87) \quad |Dv|^2 \leq C|D^2v|,$$

and so

$$|\hat{w}| \leq C|D^2v| \quad \text{at } (x_0, t_0).$$

Consequently (86) implies the estimate

$$(88) \quad |\tilde{R}| \leq C\zeta^2|D^2v| + C\zeta^3|D^2v|^{3/2} \leq \epsilon\zeta^4|D^2v|^2 + C(\epsilon),$$

where we employed Young's inequality with ϵ from §B.2. Making use of (85), (87), (88), we at last discover a contradiction to (81), provided μ is large enough.

6. Therefore $\zeta^4\hat{w} + \mu t \geq 0$ in U_T , and so in particular

$$\hat{w} + \mu t \geq 0 \quad \text{in } V \times [t_1, t_2].$$

Using (73), we deduce then that

$$(89) \quad v_t \geq \alpha|Dv|^2 - \beta \quad \text{in } V \times [t_1, t_2]$$

for appropriate constants $\alpha, \beta > 0$.

7. The differential inequality (89) for $v = \log u$ now leads us to the Harnack inequality, as follows. Fix $x_1, x_2 \in V, t_2 > t_1$. Then

$$\begin{aligned} v(x_2, t_2) - v(x_1, t_1) &= \int_0^1 \frac{d}{ds} v(sx_2 + (1-s)x_1, st_2 + (1-s)t_1) ds \\ &= \int_0^1 Dv \cdot (x_2 - x_1) + v_t(t_2 - t_1) ds \\ &\geq \int_0^1 -|Dv||x_2 - x_1| + (t_2 - t_1)[\alpha|Dv|^2 - \beta] ds \text{ by (89)} \\ &\geq -\gamma, \end{aligned}$$


where γ depends only upon $\alpha, \beta, |x_1 - x_2|, |t_1 - t_2|$. Thus (70) implies

$$\log u(x_2, t_2) \geq \log u(x_1, t_1) - \gamma,$$

and so

$$u(x_2, t_2) \geq e^{-\gamma}u(x_1, t_1).$$

This inequality is valid for each $x_1, x_2 \in V$, and so (69) is valid in the case V is a ball. In the general case we cover $V \subset\subset U$ with balls and repeatedly apply the estimate above. \square



Parabolic strong maximum principle

c. Strong maximum principle. Now we employ Harnack's inequality to establish

THEOREM 11 (Strong maximum principle). Assume $u \in C_1^2(U_T) \cap C(\bar{U}_T)$ and

$$c \equiv 0 \quad \text{in } U_T.$$

Suppose also U is connected.

(i) If

$$u_t + Lu \leq 0 \quad \text{in } U_T$$

and u attains its maximum over \bar{U}_T at a point $(x_0, t_0) \in U_T$, then

u is constant on U_{t_0} .

(ii) Likewise, if

$$u_t + Lu \geq 0 \quad \text{in } U_T$$

and u attains its minimum over \bar{U}_T at a point $(x_0, t_0) \in U_T$, then

u is constant on U_{t_0} .

Thus our uniformly parabolic partial differential equations support “infinite propagation speed of disturbances”.

We will for the following proofs assume that u and the coefficients of L are in fact smooth.

Proof. 1. Assume $u_t + Lu \leq 0$ in U_T and u attains its maximum at some point $(x_0, t_0) \in U_T$.

Select a smooth, open set $W \subset\subset U$, with $x_0 \in W$. Let v solve

$$\begin{cases} v_t + Lv = 0 & \text{in } W_T \\ v = u & \text{on } \Delta_T, \end{cases}$$

where Δ_T denotes the parabolic boundary of W_T .

Then by the weak maximum principle $u \leq v$. Since

$$u \leq v \leq M,$$

for $M := \max_{\bar{U}_T} u$, we deduce that $v = M$ at (x_0, t_0) .

2. Now write $\tilde{v} := M - v$. Then, since $c \equiv 0$, we have

$$(90) \quad \tilde{v}_t + L\tilde{v} = 0, \quad \tilde{v} \geq 0 \quad \text{in } W_T.$$

Choose any $V \subset\subset W$ with $x_0 \in V$, V connected. Let $0 < t < t_0$. Then owing to the Harnack inequality,

$$(91) \quad \max_V \tilde{v}(\cdot, t) \leq C \inf_V \tilde{v}(\cdot, t_0).$$

But $\inf_V \tilde{v}(\cdot, t_0) \leq \tilde{v}(x_0, t_0) = 0$. As $\tilde{v} \geq 0$, (91) therefore implies $\tilde{v} \equiv 0$ on $V \times \{t\}$, for each $0 < t < t_0$. This deduction holds for each set V as above, and so $\tilde{v} \equiv 0$ in W_{t_0} . But therefore $v \equiv M$ in W_{t_0} . As $v = u$ on Δ_T , we conclude $u \equiv M$ on $\partial W \times [0, t_0]$.

This conclusion holds for all sets W as above, and therefore $u \equiv M$ on U_{t_0} . \square

THEOREM 12 (Strong maximum principle for $c \geq 0$). Assume $u \in C_1^2(U_T) \cap C(\bar{U}_T)$ and

$$c \geq 0 \quad \text{in } U_T.$$

Suppose also U is connected.

(i) If

$$u_t + Lu \leq 0 \quad \text{in } U_T$$

and u attains a nonnegative maximum over \bar{U}_T at a point $(x_0, t_0) \in U_T$, then

$$u \text{ is constant on } U_{t_0}.$$

(ii) Similarly, if

$$u_t + Lu \geq 0 \quad \text{in } U_T$$

and u attains a nonpositive minimum over \bar{U}_T at a point $(x_0, t_0) \in U_T$, then

$$u \text{ is constant on } U_{t_0}.$$

Proof. 1. As above, set $M := \max_{\bar{U}_T} u$. Assume $M \geq 0$, $u_t + Lu \leq 0$ in U_T , and u attains this maximum of M at some point $(x_0, t_0) \in U_T$.

If $M = 0$, the foregoing proof directly applies, as then

$$\tilde{v}_t + L\tilde{v} = 0, \quad \tilde{v} \geq 0 \quad \text{in } W_T.$$

2. Assume instead that $M > 0$. Select as in the previous proof a smooth, open set $W \subset\subset U$, with $x_0 \in W$. Now let v solve

$$\begin{cases} v_t + Kv = 0 & \text{in } W_T \\ v = u^+ & \text{on } \Delta_T, \end{cases}$$

where

$$Kv := Lv - cv.$$

Note $0 \leq v \leq M$. Since $u_t + Ku \leq -cu \leq 0$ on $\{u \geq 0\}$, we deduce from the weak maximum principle that $u \leq v$. As before it follows that $v = M$ at (x_0, t_0) .

3. Now write $\tilde{v} := M - v$. Then, since the operator K has no zeroth-order term, we have

$$\tilde{v}_t + K\tilde{v} = 0, \quad \tilde{v} \geq 0 \quad \text{in } W_T.$$

Select any $V \subset\subset W$ with $x_0 \in V$, V connected. Let $0 < t < t_0$. Then the Harnack inequality implies as above that $v \equiv u^+ \equiv M$ on $\partial W \times [0, t_0]$. Since $M > 0$, we conclude that $u \equiv M$ on $\partial W \times [0, t_0]$.

This deduction is valid for all sets W as above, and therefore $u \equiv M$ on U_{t_0} . \square

7.2. SECOND-ORDER HYPERBOLIC EQUATIONS

Second-order hyperbolic equations are natural generalizations of the wave equation (§2.4). We will build in this section appropriately defined weak solutions and study their uniqueness, smoothness and other properties. It is interesting, given the utterly different physical character of second-order parabolic and hyperbolic PDE, that we can provide rather similar functional analytic constructions of weak solutions.

7.2.1. Definitions.

a. Hyperbolic equations. As in §7.1 we write $U_T = U \times (0, T]$, where $T > 0$ and $U \subset \mathbb{R}^n$ is an open, bounded set.

We will next study the initial/boundary-value problem

$$(1) \quad \begin{cases} u_{tt} + Lu = f & \text{in } U_T \\ u = 0 & \text{on } \partial U \times [0, T] \\ u = g, u_t = h & \text{on } U \times \{t = 0\}, \end{cases}$$

where $f : U_T \rightarrow \mathbb{R}$, $g, h : U \rightarrow \mathbb{R}$ are given, and $u : \bar{U}_T \rightarrow \mathbb{R}$ is the unknown, $u = u(x, t)$. The symbol L denotes for each time t a second-order partial differential operator, having either the divergence form

$$(2) \quad Lu = - \sum_{i,j=1}^n (a^{ij}(x, t)u_{x_i})_{x_j} + \sum_{i=1}^n b^i(x, t)u_{x_i} + c(x, t)u$$

or else the nondivergence form

$$(3) \quad Lu = - \sum_{i,j=1}^n a^{ij}(x, t)u_{x_i x_j} + \sum_{i=1}^n b^i(x, t)u_{x_i} + c(x, t)u$$

for given coefficients a^{ij}, b^i, c ($i, j = 1, \dots, n$).

DEFINITION. We say the partial differential operator $\frac{\partial^2}{\partial t^2} + L$ is (uniformly) hyperbolic if there exists a constant $\theta > 0$ such that

$$(4) \quad \sum_{i,j=1}^n a^{ij}(x, t)\xi_i \xi_j \geq \theta |\xi|^2$$

for all $(x, t) \in U_T$, $\xi \in \mathbb{R}^n$.

If $a^{ij} \equiv \delta_{ij}$, $b^i \equiv c \equiv f \equiv 0$, then $L = -\Delta$ and the partial differential equation becomes the wave equation, already studied in Chapter 2. General second-order hyperbolic PDE model wave transmission in heterogeneous, nonisotropic media.

b. Weak solutions. As before, in §6.1.2 and §7.1.1, we first assume L has the divergence form (2) and look for an appropriate notion of weak solution for problem (1). We will suppose initially that

$$(5) \quad a^{ij}, b^i, c \in C^1(\bar{U}_T) \quad (i, j = 1, \dots, n),$$

$$(6) \quad f \in L^2(U_T),$$

$$(7) \quad g \in H_0^1(U), \quad h \in L^2(U)$$

and always assume $a^{ij} = a^{ji}$ ($i, j = 1, \dots, n$).

As in §7.1.1, let us also introduce the time-dependent bilinear form

$$(8) \quad B[u, v; t] := \int_U \sum_{i,j=1}^n a^{ij}(\cdot, t)u_{x_i}v_{x_j} + \sum_{i=1}^n b^i(\cdot, t)u_{x_i}v + c(\cdot, t)uv \, dx$$

for $u, v \in H_0^1(U)$ and $0 \leq t \leq T$.

Motivation for definition of weak solution. We begin by supposing $u = u(x, t)$ to be a smooth solution of (1) and defining the associated mapping

$$\mathbf{u} : [0, T] \rightarrow H_0^1(U),$$

by

$$[\mathbf{u}(t)](x) := u(x, t) \quad (x \in U, 0 \leq t \leq T).$$

We similarly introduce the function

$$\mathbf{f} : [0, T] \rightarrow L^2(U)$$

defined by

$$[\mathbf{f}(t)](x) := f(x, t) \quad (x \in U, 0 \leq t \leq T).$$

Now fix any function $v \in H_0^1(U)$, multiply the PDE $u_{tt} + Lu = f$ by v , and integrate by parts, to obtain the identity

$$(9) \quad (\mathbf{u}'', v) + B[\mathbf{u}, v; t] = (\mathbf{f}, v)$$

for $0 \leq t \leq T$, where (\cdot, \cdot) denotes the inner product in $L^2(U)$. Almost exactly as in the parallel discussion for parabolic PDE in §7.1.1, we see from the PDE $u_{tt} + Lu = f$ that

$$u_{tt} = g^0 + \sum_{j=1}^n g_{x_j}^j$$

for $g^0 := f - \sum_{i=1}^n b^i u_{x_i} - cu$ and $g^j := \sum_{i=1}^n a^{ij} u_{x_i}$ ($j = 1, \dots, n$). This suggests that we should look for a weak solution \mathbf{u} with $\mathbf{u}'' \in H^{-1}(U)$ for a.e. $0 \leq t \leq T$ and then reinterpret the first term of (9) as $\langle \mathbf{u}'', v \rangle$, $\langle \cdot, \cdot \rangle$ denoting as usual the pairing between $H^{-1}(U)$ and $H_0^1(U)$. \square

DEFINITION. We say a function

$$\mathbf{u} \in L^2(0, T; H_0^1(U)), \text{ with } \mathbf{u}' \in L^2(0, T; L^2(U)), \mathbf{u}'' \in L^2(0, T; H^{-1}(U)),$$

is a weak solution of the hyperbolic initial/boundary-value problem (1) provided

$$(i) \quad \langle \mathbf{u}'', v \rangle + B[\mathbf{u}, v; t] = (\mathbf{f}, v)$$

for each $v \in H_0^1(U)$ and a.e. time $0 \leq t \leq T$ and

$$(ii) \quad \mathbf{u}(0) = g, \mathbf{u}'(0) = h.$$

Remark. In view of Theorem 2 in §5.9.2, we know $\mathbf{u} \in C([0, T]; L^2(U))$ and $\mathbf{u}' \in C([0, T]; H^{-1}(U))$. Consequently the equalities (ii) above make sense.

7.2.2. Existence of weak solutions.

a. Galerkin approximations. By analogy with the approach taken in §7.1.2 we will construct our weak solution of the hyperbolic initial/boundary-value problem

$$(10) \quad \begin{cases} u_{tt} + Lu = f & \text{in } U_T \\ u = 0 & \text{on } \partial U \times [0, T] \\ u = g, u_t = h & \text{on } U \times \{t = 0\} \end{cases}$$

by first solving a finite-dimensional approximation. We thus once more employ *Galerkin's method* by selecting smooth functions $w_k = w_k(x)$ ($k = 1, \dots$) such that

$$(11) \quad \{w_k\}_{k=1}^{\infty} \text{ is an orthogonal basis of } H_0^1(U)$$

and

$$(12) \quad \{w_k\}_{k=1}^{\infty} \text{ is an orthonormal basis of } L^2(U).$$

Fix a positive integer m , and write

$$(13) \quad \mathbf{u}_m(t) := \sum_{k=1}^m d_m^k(t) w_k,$$

where we intend to select the coefficients $d_m^k(t)$ ($0 \leq t \leq T$, $k = 1, \dots, m$) to satisfy

$$(14) \quad d_m^k(0) = (g, w_k) \quad (k = 1, \dots, m),$$

$$(15) \quad d_m^{k'}(0) = (h, w_k) \quad (k = 1, \dots, m),$$

and

$$(16) \quad (\mathbf{u}_m'', w_k) + B[\mathbf{u}_m, w_k; t] = (\mathbf{f}, w_k) \quad (0 \leq t \leq T, k = 1, \dots, m).$$

THEOREM 1 (Construction of approximate solutions). *For each integer $m = 1, 2, \dots$, there exists a unique function \mathbf{u}_m of the form (13) satisfying (14)–(16).*

Proof. Assuming \mathbf{u}_m to be given by (13), we observe using (12)

$$(17) \quad (\mathbf{u}_m''(t), w_k) = d_m^{k''}(t).$$

Furthermore, exactly as in the proof of Theorem 1 in §7.1.2, we have

$$B[\mathbf{u}_m, w_k; t] = \sum_{l=1}^m e^{kl}(t) d_m^l(t)$$

for $e^{kl}(t) := B[w_l, w_k; t]$ ($k, l = 1, \dots, m$). We also write $f^k(t) := (\mathbf{f}(t), w_k)$ ($k = 1, \dots, m$). Consequently (16) becomes the linear system of ODE

$$(18) \quad d_m^{k''}(t) + \sum_{l=1}^m e^{kl}(t) d_m^l(t) = f^k(t) \quad (0 \leq t \leq T, k = 1, \dots, m),$$

subject to the initial conditions (14), (15). According to standard theory for ordinary differential equations, there exists a unique function $\mathbf{d}_m(t) = (d_m^1(t), \dots, d_m^m(t))$, satisfying (14), (15) and solving (18) for $0 \leq t \leq T$. \square

b. Energy estimates. Our plan is hereafter to send $m \rightarrow \infty$, and so we will need some estimates, uniform in m .

THEOREM 2 (Energy estimates). *There exists a constant C , depending only on U, T and the coefficients of L , such that*

$$(19) \quad \max_{0 \leq t \leq T} \left(\|\mathbf{u}_m(t)\|_{H_0^1(U)} + \|\mathbf{u}_m'(t)\|_{L^2(U)} \right) + \|\mathbf{u}_m''\|_{L^2(0, T; H^{-1}(U))} \leq C \left(\|\mathbf{f}\|_{L^2(0, T; L^2(U))} + \|g\|_{H_0^1(U)} + \|h\|_{L^2(U)} \right),$$

for $m = 1, 2, \dots$

Proof. 1. Multiply equality (16) by $d_m^{k'}(t)$, sum $k = 1, \dots, m$, and recall (13) to discover

$$(20) \quad (\mathbf{u}_m'', \mathbf{u}_m') + B[\mathbf{u}_m, \mathbf{u}_m'; t] = (\mathbf{f}, \mathbf{u}_m')$$

for a.e. $0 \leq t \leq T$. Observe next $(\mathbf{u}_m'', \mathbf{u}_m') = \frac{d}{dt} \left(\frac{1}{2} \|\mathbf{u}_m'\|_{L^2(U)}^2 \right)$. Furthermore, we can write

$$(21) \quad \begin{aligned} B[\mathbf{u}_m, \mathbf{u}_m'; t] &= \int_U \sum_{i,j=1}^n a^{ij} \mathbf{u}_{m,x_i} \mathbf{u}'_{m,x_j} dx \\ &\quad + \int_U \sum_{i=1}^n b^i \mathbf{u}_{m,x_i} \mathbf{u}'_m + c \mathbf{u}_m \mathbf{u}'_m dx \\ &=: B_1 + B_2. \end{aligned}$$

Since $a^{ij} = a^{ji}$ ($i, j = 1, \dots, n$), we see

$$(22) \quad B_1 = \frac{d}{dt} \left(\frac{1}{2} A[\mathbf{u}_m, \mathbf{u}_m; t] \right) - \frac{1}{2} \int_U \sum_{i,j=1}^n a_t^{ij} \mathbf{u}_{m,x_i} \mathbf{u}_{m,x_j} dx,$$

for the symmetric bilinear form

$$A[u, v; t] := \int_U \sum_{i,j=1}^n a^{ij} u_{x_i} v_{x_j} dx \quad (u, v \in H_0^1(U)).$$

The equality (22) implies

$$(23) \quad B_1 \geq \frac{d}{dt} \left(\frac{1}{2} A[\mathbf{u}_m, \mathbf{u}_m; t] \right) - C \|\mathbf{u}_m\|_{H_0^1(U)}^2,$$

and we note also

$$(24) \quad |B_2| \leq C (\|\mathbf{u}_m\|_{H_0^1(U)}^2 + \|\mathbf{u}'_m\|_{L^2(U)}^2).$$

Combining estimates (20)–(24), we discover

$$(25) \quad \begin{aligned} \frac{d}{dt} \left(\|\mathbf{u}'_m\|_{L^2(U)}^2 + A[\mathbf{u}_m, \mathbf{u}_m; t] \right) &\leq C \left(\|\mathbf{u}'_m\|_{L^2(U)}^2 + \|\mathbf{u}_m\|_{H_0^1(U)}^2 + \|\mathbf{f}\|_{L^2(U)}^2 \right) \\ &\leq C \left(\|\mathbf{u}'_m\|_{L^2(U)}^2 + A[\mathbf{u}_m, \mathbf{u}_m; t] + \|\mathbf{f}\|_{L^2(U)}^2 \right), \end{aligned}$$

where we used the inequality

$$(26) \quad \theta \int_U |Du|^2 dx \leq A[u, u; t] \quad (u \in H_0^1(U)),$$

which follows from the uniform hyperbolicity condition.

2. Now write

$$(27) \quad \eta(t) := \|\mathbf{u}'_m(t)\|_{L^2(U)}^2 + A[\mathbf{u}_m(t), \mathbf{u}_m(t); t]$$

and

$$(28) \quad \xi(t) := \|\mathbf{f}(t)\|_{L^2(U)}^2.$$

Then inequality (25) reads

$$\eta'(t) \leq C_1 \eta(t) + C_2 \xi(t)$$

for $0 \leq t \leq T$ and appropriate constants C_1, C_2 . Thus Gronwall's inequality (§B.2) yields the estimate

$$(29) \quad \eta(t) \leq e^{C_1 t} \left(\eta(0) + C_2 \int_0^t \xi(s) ds \right) \quad (0 \leq t \leq T).$$

However,

$$\begin{aligned} \eta(0) &= \|\mathbf{u}'_m(0)\|_{L^2(U)}^2 + A[\mathbf{u}_m(0), \mathbf{u}_m(0); 0] \\ &\leq C \left(\|h\|_{L^2(U)}^2 + \|g\|_{H_0^1(U)}^2 \right), \end{aligned}$$

according to (14) and (15) and the estimate $\|\mathbf{u}_m(0)\|_{H_0^1(U)} \leq \|g\|_{H_0^1(U)}$. Thus formulas (27)–(29) provide the bound

$$\begin{aligned} &\|\mathbf{u}'_m(t)\|_{L^2(U)}^2 + A[\mathbf{u}_m(t), \mathbf{u}_m(t); t] \\ &\leq C \left(\|g\|_{H_0^1(U)}^2 + \|h\|_{L^2(U)}^2 + \|\mathbf{f}\|_{L^2(0,T;L^2(U))}^2 \right). \end{aligned}$$

Since $0 \leq t \leq T$ was arbitrary, we see from this estimate and (26) that

$$\begin{aligned} &\max_{0 \leq t \leq T} \left(\|\mathbf{u}_m(t)\|_{H_0^1(U)}^2 + \|\mathbf{u}'_m(t)\|_{L^2(U)}^2 \right) \\ &\leq C \left(\|g\|_{H_0^1(U)}^2 + \|h\|_{L^2(U)}^2 + \|\mathbf{f}\|_{L^2(0,T;L^2(U))}^2 \right). \end{aligned}$$

3. Fix any $v \in H_0^1(U)$, $\|v\|_{H_0^1(U)} \leq 1$, and write $v = v^1 + v^2$, where $v^1 \in \text{span}\{w_k\}_{k=1}^m$ and $(v^2, w_k) = 0$ ($k = 1, \dots, m$). Note $\|v^1\|_{H_0^1(U)} \leq 1$. Then (13) and (16) imply

$$\langle \mathbf{u}''_m, v \rangle = (\mathbf{u}''_m, v) = (\mathbf{u}''_m, v^1) = (\mathbf{f}, v^1) - B[\mathbf{u}_m, v^1; t].$$

Thus

$$|\langle \mathbf{u}''_m, v \rangle| \leq C(\|\mathbf{f}\|_{L^2(U)} + \|\mathbf{u}_m\|_{H_0^1(U)}),$$

since $\|v^1\|_{H_0^1(U)} \leq 1$. Consequently

$$\begin{aligned} \int_0^T \|\mathbf{u}''_m\|_{H^{-1}(U)}^2 dt &\leq C \int_0^T \|\mathbf{f}\|_{L^2(U)}^2 + \|\mathbf{u}_m\|_{H_0^1(U)}^2 dt \\ &\leq C \left(\|g\|_{H_0^1(U)}^2 + \|h\|_{L^2(U)}^2 + \|\mathbf{f}\|_{L^2(0,T;L^2(U))}^2 \right). \quad \square \end{aligned}$$

c. Existence and uniqueness. Now we pass to limits in our Galerkin approximations.

THEOREM 3 (Existence of weak solution). *There exists a weak solution of (1).*

Proof. 1. According to the energy estimates (19), we see that the sequence $\{\mathbf{u}_m\}_{m=1}^\infty$ is bounded in $L^2(0, T; H_0^1(U))$, $\{\mathbf{u}'_m\}_{m=1}^\infty$ is bounded in $L^2(0, T; L^2(U))$, and $\{\mathbf{u}''_m\}_{m=1}^\infty$ is bounded in $L^2(0, T; H^{-1}(U))$.

As a consequence there exists a subsequence $\{\mathbf{u}_{m_l}\}_{l=1}^\infty \subset \{\mathbf{u}_m\}_{m=1}^\infty$ and $\mathbf{u} \in L^2(0, T; H_0^1(U))$, with $\mathbf{u}' \in L^2(0, T; L^2(U))$, $\mathbf{u}'' \in L^2(0, T; H^{-1}(U))$, such that

$$(30) \quad \begin{cases} \mathbf{u}_{m_l} \rightharpoonup \mathbf{u} & \text{weakly in } L^2(0, T; H_0^1(U)) \\ \mathbf{u}'_{m_l} \rightharpoonup \mathbf{u}' & \text{weakly in } L^2(0, T; L^2(U)) \\ \mathbf{u}''_{m_l} \rightharpoonup \mathbf{u}'' & \text{weakly in } L^2(0, T; H^{-1}(U)). \end{cases}$$

2. Next fix an integer N and choose a function $\mathbf{v} \in C^1([0, T]; H_0^1(U))$ of the form

$$(31) \quad \mathbf{v}(t) = \sum_{k=1}^N d^k(t) w_k,$$

where $\{d^k\}_{k=1}^N$ are smooth functions. We select $m \geq N$, multiply (16) by $d^k(t)$, sum $k = 1, \dots, N$, and then integrate with respect to t , to discover

$$(32) \quad \int_0^T \langle \mathbf{u}''_m, \mathbf{v} \rangle + B[\mathbf{u}_m, \mathbf{v}; t] dt = \int_0^T (\mathbf{f}, \mathbf{v}) dt.$$

We set $m = m_l$ and recall (30), to find in the limit that

$$(33) \quad \int_0^T \langle \mathbf{u}'', \mathbf{v} \rangle + B[\mathbf{u}, \mathbf{v}; t] dt = \int_0^T (\mathbf{f}, \mathbf{v}) dt.$$

This equality then holds for all functions $\mathbf{v} \in L^2(0, T; H_0^1(U))$, since functions of the form (31) are dense in this space. From (33) it follows further that

$$\langle \mathbf{u}'', v \rangle + B[\mathbf{u}, v; t] = (\mathbf{f}, v)$$

for all $v \in H_0^1(U)$ and a.e. $0 \leq t \leq T$. Furthermore, $\mathbf{u} \in C([0, T]; L^2(U))$ and $\mathbf{u}' \in C([0, T]; H^{-1}(U))$.

3. We must now verify

$$(34) \quad \mathbf{u}(0) = g,$$

$$(35) \quad \mathbf{u}'(0) = h.$$

For this, choose any function $\mathbf{v} \in C^2([0, T]; H_0^1(U))$, with $\mathbf{v}(T) = \mathbf{v}'(T) = 0$. Then integrating by parts twice with respect to t in (33), we find

$$(36) \quad \int_0^T (\mathbf{v}'', \mathbf{u}) + B[\mathbf{u}, \mathbf{v}; t] dt = \int_0^T (\mathbf{f}, \mathbf{v}) dt - (\mathbf{u}(0), \mathbf{v}'(0)) + \langle \mathbf{u}'(0), \mathbf{v}(0) \rangle.$$

Similarly from (32) we deduce

$$\int_0^T (\mathbf{v}'', \mathbf{u}_m) + B[\mathbf{u}_m, \mathbf{v}; t] dt = \int_0^T (\mathbf{f}, \mathbf{v}) dt - (\mathbf{u}_m(0), \mathbf{v}'(0)) + (\mathbf{u}'_m(0), \mathbf{v}(0)).$$

We set $m = m_l$ and recall (14), (15) and (30), to deduce

$$(37) \quad \int_0^T (\mathbf{v}'', \mathbf{u}) + B[\mathbf{u}, \mathbf{v}; t] dt = \int_0^T (\mathbf{f}, \mathbf{v}) dt - (g, \mathbf{v}'(0)) + (h, \mathbf{v}(0)).$$

Comparing identities (36) and (37), we conclude (34), (35), since $\mathbf{v}(0)$, $\mathbf{v}'(0)$ are arbitrary. Hence \mathbf{u} is a weak solution of (1). \square

Recalling the energy estimates from Theorem 2, we observe that in fact $\mathbf{u} \in L^\infty(0, T; H_0^1(U))$, $\mathbf{u}' \in L^\infty(0, T; L^2(U))$, $\mathbf{u}'' \in L^2(0, T; H^{-1}(U))$: see Theorem 5 below.

THEOREM 4 (Uniqueness of weak solution). *A weak solution of (1) is unique.*

The following tricky demonstration would be greatly simplified if we knew $\mathbf{u}'(t)$ itself were smooth enough to insert in place of v in the definition of weak solution. This is not so, however.

Proof. 1. It suffices to show that the only weak solution of (1) with $\mathbf{f} \equiv g \equiv h \equiv 0$ is

$$(38) \quad \mathbf{u} \equiv \mathbf{0}.$$

To verify this, fix $0 \leq s \leq T$ and set

$$\mathbf{v}(t) := \begin{cases} \int_t^s \mathbf{u}(\tau) d\tau & \text{if } 0 \leq t \leq s \\ \mathbf{0} & \text{if } s \leq t \leq T. \end{cases}$$

Then $\mathbf{v}(t) \in H_0^1(U)$ for each $0 \leq t \leq T$, and so

$$\int_0^s \langle \mathbf{u}'', \mathbf{v} \rangle + B[\mathbf{u}, \mathbf{v}; t] dt = 0.$$

Since $\mathbf{u}'(0) = \mathbf{v}(s) = \mathbf{0}$, we obtain after integrating by parts in the first term above:

$$(39) \quad \int_0^s -(\mathbf{u}', \mathbf{v}') + B[\mathbf{u}, \mathbf{v}; t] dt = 0.$$

Now $\mathbf{v}' = -\mathbf{u}$ ($0 \leq t \leq s$), and so

$$\int_0^s \langle \mathbf{u}', \mathbf{u} \rangle - B[\mathbf{v}', \mathbf{v}; t] dt = 0.$$

Thus

$$\int_0^s \frac{d}{dt} \left(\frac{1}{2} \|\mathbf{u}\|_{L^2(U)}^2 - \frac{1}{2} B[\mathbf{v}, \mathbf{v}; t] \right) dt = - \int_0^s C[\mathbf{u}, \mathbf{v}; t] + D[\mathbf{v}, \mathbf{v}; t] dt,$$

where

$$C[u, v; t] := - \int_U \sum_{i=1}^n b^i v_{x_i} u + \frac{1}{2} b_{x_i}^i u v dx$$

and

$$D[u, v; t] := \frac{1}{2} \int_U \sum_{i,j=1}^n a_t^{ij} u_{x_i} v_{x_j} + \sum_{i=1}^n b_t^i u_{x_i} v + c_t u v dx,$$

for $u, v \in H_0^1(U)$. Hence

$$\frac{1}{2} \|\mathbf{u}(s)\|_{L^2(U)}^2 + \frac{1}{2} B[\mathbf{v}(0), \mathbf{v}(0); t] = - \int_0^s C[\mathbf{u}, \mathbf{v}; t] + D[\mathbf{v}, \mathbf{v}; t] dt,$$

and consequently

$$(40) \quad \begin{aligned} & \|\mathbf{u}(s)\|_{L^2(U)}^2 + \|\mathbf{v}(0)\|_{H_0^1(U)}^2 \\ & \leq C \left(\int_0^s \|\mathbf{v}\|_{H_0^1(U)}^2 + \|\mathbf{u}\|_{L^2(U)}^2 dt + \|\mathbf{v}(0)\|_{L^2(U)}^2 \right). \end{aligned}$$

2. Now let us write

$$\mathbf{w}(t) := \int_0^t \mathbf{u}(\tau) d\tau \quad (0 \leq t \leq T),$$

whereupon (40) becomes

$$(41) \quad \begin{aligned} & \|\mathbf{u}(s)\|_{L^2(U)}^2 + \|\mathbf{w}(s)\|_{H_0^1(U)}^2 \\ & \leq C \left(\int_0^s \|\mathbf{w}(t) - \mathbf{w}(s)\|_{H_0^1(U)}^2 + \|\mathbf{u}(t)\|_{L^2(U)}^2 dt + \|\mathbf{w}(s)\|_{L^2(U)}^2 \right). \end{aligned}$$

But $\|\mathbf{w}(t) - \mathbf{w}(s)\|_{H_0^1(U)}^2 \leq 2\|\mathbf{w}(t)\|_{H_0^1(U)}^2 + 2\|\mathbf{w}(s)\|_{H_0^1(U)}^2$, and $\|\mathbf{w}(s)\|_{L^2(U)} \leq \int_0^s \|\mathbf{u}(t)\|_{L^2(U)} dt$. Therefore (41) implies

$$\|\mathbf{u}(s)\|_{L^2(U)}^2 + (1 - 2sC_1)\|\mathbf{w}(s)\|_{H_0^1(U)}^2 \leq C_1 \int_0^s \|\mathbf{w}\|_{H_0^1(U)}^2 + \|\mathbf{u}\|_{L^2(U)}^2 dt.$$

Choose T_1 so small that

$$1 - 2T_1C_1 \geq \frac{1}{2}.$$

Then if $0 \leq s \leq T_1$, we have

$$\|\mathbf{u}(s)\|_{L^2(U)}^2 + \|\mathbf{w}(s)\|_{H_0^1(U)}^2 \leq C \int_0^s \|\mathbf{u}\|_{L^2(U)}^2 + \|\mathbf{w}\|_{H_0^1(U)}^2 dt.$$

Consequently the integral form of Gronwall's inequality (§B.2) implies $\mathbf{u} \equiv \mathbf{0}$ on $[0, T_1]$.

3. We apply the same argument on the intervals $[T_1, 2T_1]$, $[2T_1, 3T_1]$, etc., eventually to deduce (38). \square

7.2.3. Regularity.

As in our earlier treatments of second-order elliptic and parabolic PDE, the next task is to study the smoothness of our weak solutions.

Motivation: formal derivation of estimates. (i) Suppose for the moment $u = u(x, t)$ is a smooth solution of this initial-value problem for the wave equation:

$$\begin{cases} u_{tt} - \Delta u = f & \text{in } \mathbb{R}^n \times (0, T] \\ u = g, u_t = h & \text{on } \mathbb{R}^n \times \{t = 0\} \end{cases}$$

and assume also u goes to zero as $|x| \rightarrow \infty$ sufficiently rapidly to justify the following calculations. Then as in §2.4.3, we compute

$$\begin{aligned} \frac{d}{dt} \left(\int_{\mathbb{R}^n} |Du|^2 + u_t^2 dx \right) &= 2 \int_{\mathbb{R}^n} Du \cdot Du_t + u_t u_{tt} dx \\ &= 2 \int_{\mathbb{R}^n} u_t (u_{tt} - \Delta u) dx = 2 \int_{\mathbb{R}^n} u_t f dx \\ &\leq \int_{\mathbb{R}^n} u_t^2 dx + \int_{\mathbb{R}^n} f^2 dx. \end{aligned}$$

Applying Gronwall's inequality, we deduce

$$(42) \quad \sup_{0 \leq t \leq T} \int_{\mathbb{R}^n} |Du|^2 + u_t^2 dx \leq C \left(\int_0^T \int_{\mathbb{R}^n} f^2 dx dt + \int_{\mathbb{R}^n} |Dg|^2 + h^2 dx \right),$$

with the constant C depending only on T .

(ii) Next differentiate the PDE with respect to t and set $\tilde{u} := u_t$. Then

$$(43) \quad \begin{cases} \tilde{u}_{tt} - \Delta \tilde{u} = \tilde{f} & \text{in } \mathbb{R}^n \times (0, T] \\ \tilde{u} = \tilde{g}, \tilde{u}_t = \tilde{h} & \text{on } \mathbb{R}^n \times \{t = 0\}, \end{cases}$$

for $\tilde{f} := f_t$, $\tilde{g} := h$, $\tilde{h} := u_{tt}(\cdot, 0) = f(\cdot, 0) + \Delta g$. Applying estimate (42) to \tilde{u} , we discover

$$(44) \quad \begin{aligned} & \sup_{0 \leq t \leq T} \int_{\mathbb{R}^n} |Du_t|^2 + u_{tt}^2 dx \\ & \leq C \left(\int_0^T \int_{\mathbb{R}^n} f_t^2 dx dt + \int_{\mathbb{R}^n} |D^2g|^2 + |Dh|^2 + f(\cdot, 0)^2 dx \right). \end{aligned}$$

Now

$$(45) \quad \max_{0 \leq t \leq T} \|f(\cdot, t)\|_{L^2(\mathbb{R}^n)} \leq C(\|f\|_{L^2(\mathbb{R}^n \times (0, T))} + \|f_t\|_{L^2(\mathbb{R}^n \times (0, T))}),$$

according to Theorem 2 in §5.9.2. Furthermore, writing $-\Delta u = f - u_{tt}$, we deduce as in §6.3 that

$$(46) \quad \int_{\mathbb{R}^n} |D^2u|^2 dx \leq C \int_{\mathbb{R}^n} f^2 + u_{tt}^2 dx$$

for each $0 \leq t \leq T$. Combining (44)–(46), we conclude

$$(47) \quad \begin{aligned} & \sup_{0 \leq t \leq T} \int_{\mathbb{R}^n} |D^2u|^2 + |Du_t|^2 + u_{tt}^2 dx \\ & \leq C \left(\int_0^T \int_{\mathbb{R}^n} f_t^2 + f^2 dx dt + \int_{\mathbb{R}^n} |D^2g|^2 + |Dh|^2 dx \right), \end{aligned}$$

the constant C depending only on T . □

This estimate suggests that bounds similar to (42) and (47) should be valid for our weak solution of a general second-order hyperbolic PDE.

We will calculate using the Galerkin approximations. To simplify the presentation, we hereafter assume that $\{w_k\}_{k=1}^\infty$ is the complete collection of eigenfunctions for $-\Delta$ on $H_0^1(U)$ and also that U is bounded, open, with ∂U smooth. In addition we suppose

$$(48) \quad \begin{cases} \text{the coefficients } a^{ij}, b^i, c \text{ (} i, j = 1, \dots, n \text{) are smooth on} \\ \bar{U} \text{ and do not depend on } t. \end{cases}$$

THEOREM 5 (Improved regularity).

(i) Assume

$$g \in H_0^1(U), \quad h \in L^2(U), \quad \mathbf{f} \in L^2(0, T; L^2(U)),$$

and suppose also $\mathbf{u} \in L^2(0, T; H_0^1(U))$, with $\mathbf{u}' \in L^2(0, T; L^2(U))$, $\mathbf{u}'' \in L^2(0, T; H^{-1}(U))$, is the weak solution of the problem

$$(49) \quad \begin{cases} u_{tt} + Lu = f & \text{in } U_T \\ u = 0 & \text{on } \partial U \times [0, T] \\ u = g, u_t = h & \text{on } U \times \{t = 0\}. \end{cases}$$

Then in fact

$$\mathbf{u} \in L^\infty(0, T; H_0^1(U)), \quad \mathbf{u}' \in L^\infty(0, T; L^2(U)),$$

and we have the estimate

$$(50) \quad \begin{aligned} \operatorname{ess\,sup}_{0 \leq t \leq T} (\|\mathbf{u}(t)\|_{H_0^1(U)} + \|\mathbf{u}'(t)\|_{L^2(U)}) \\ \leq C(\|\mathbf{f}\|_{L^2(0, T; L^2(U))} + \|g\|_{H_0^1(U)} + \|h\|_{L^2(U)}). \end{aligned}$$

(ii) If, in addition,

$$g \in H^2(U), \quad h \in H_0^1(U), \quad \mathbf{f}' \in L^2(0, T; L^2(U)),$$

then

$$\begin{aligned} \mathbf{u} \in L^\infty(0, T; H^2(U)), \quad \mathbf{u}' \in L^\infty(0, T; H_0^1(U)), \\ \mathbf{u}'' \in L^\infty(0, T; L^2(U)), \quad \mathbf{u}''' \in L^2(0, T; H^{-1}(U)), \end{aligned}$$

with the estimate

$$(51) \quad \begin{aligned} \operatorname{ess\,sup}_{0 \leq t \leq T} (\|\mathbf{u}(t)\|_{H^2(U)} + \|\mathbf{u}'(t)\|_{H_0^1(U)} + \|\mathbf{u}''(t)\|_{L^2(U)}) \\ + \|\mathbf{u}'''\|_{L^2(0, T; H^{-1}(U))} \leq C(\|\mathbf{f}'\|_{L^2(0, T; L^2(U))} + \|g\|_{H^2(U)} + \|h\|_{H^1(U)}). \end{aligned}$$

Assertions (i), (ii) of this theorem are precise versions of the formal estimates (42), (47) (for the wave equation in $U = \mathbb{R}^n$).

Proof. 1. In the proof of Theorem 2, we have already derived the bounds

$$(52) \quad \begin{aligned} \sup_{0 \leq t \leq T} (\|\mathbf{u}_m(t)\|_{H_0^1(U)} + \|\mathbf{u}'_m(t)\|_{L^2(U)}) \\ \leq C(\|\mathbf{f}\|_{L^2(0, T; L^2(U))} + \|g\|_{H_0^1(U)} + \|h\|_{L^2(U)}). \end{aligned}$$

Passing to limits as $m = m_l \rightarrow \infty$, we deduce (50).

2. Assume now the hypotheses of assertion (ii). Fix a positive integer m , and next differentiate the identity (16) with respect to t . Writing $\tilde{\mathbf{u}}_m := \mathbf{u}'_m$, we obtain

$$(53) \quad (\tilde{\mathbf{u}}''_m, w_k) + B[\tilde{\mathbf{u}}_m, w_k] = (\mathbf{f}', w_k) \quad (k = 1, \dots, m).$$

Multiplying by $d_m^k(t)$ and adding for $k = 1, \dots, m$, we discover

$$(54) \quad (\tilde{\mathbf{u}}''_m, \tilde{\mathbf{u}}'_m) + B[\tilde{\mathbf{u}}_m, \tilde{\mathbf{u}}'_m] = (\mathbf{f}', \tilde{\mathbf{u}}'_m).$$

Arguing as in the proof of the energy estimates, we observe

$$(55) \quad \frac{d}{dt} (\|\tilde{\mathbf{u}}'_m\|_{L^2(U)}^2 + A[\tilde{\mathbf{u}}_m, \tilde{\mathbf{u}}_m]) \\ \leq C (\|\tilde{\mathbf{u}}'_m\|_{L^2(U)}^2 + A[\tilde{\mathbf{u}}_m, \tilde{\mathbf{u}}_m] + \|\mathbf{f}'\|_{L^2(U)}^2),$$

the bilinear form $A[\ , \]$ defined as before.

3. Now

$$(56) \quad B[\mathbf{u}_m, w_k] = (\mathbf{f} - \mathbf{u}''_m, w_k) \quad (k = 1, \dots, m).$$

Recall we are taking $\{w_k\}_{k=1}^\infty$ to be the complete collection of eigenfunctions for $-\Delta$ on $H_0^1(U)$. Multiplying (56) by $\lambda_k d_m^k(t)$ and summing $k = 1, \dots, m$, we deduce

$$(57) \quad B[\mathbf{u}_m, -\Delta \mathbf{u}_m] = (\mathbf{f} - \mathbf{u}''_m, -\Delta \mathbf{u}_m).$$

Since $\Delta \mathbf{u}_m = 0$ on ∂U , we have

$$(58) \quad B[\mathbf{u}_m, -\Delta \mathbf{u}_m] = (L\mathbf{u}_m, -\Delta \mathbf{u}_m).$$

Next we employ the inequality

$$(59) \quad \beta \|u\|_{H^2(U)}^2 \leq (Lu, -\Delta u) + \gamma \|u\|_{L^2(U)}^2 \quad (u \in H^2(U) \cap H_0^1(U));$$

see Problem 9. We deduce from (56)–(59) that

$$(60) \quad \|\mathbf{u}_m\|_{H^2(U)}^2 \leq C (\|\mathbf{f}\|_{L^2(U)}^2 + \|\mathbf{u}''_m\|_{L^2(U)}^2 + \|\mathbf{u}_m\|_{L^2(U)}^2).$$

Using this estimate in (55), recalling $\tilde{\mathbf{u}}_m = \mathbf{u}'_m$, and applying Gronwall's inequality, we deduce

$$(61) \quad \sup_{0 \leq t \leq T} (\|\mathbf{u}_m(t)\|_{H^2(U)}^2 + \|\mathbf{u}'_m(t)\|_{H^1(U)}^2 + \|\mathbf{u}''_m(t)\|_{L^2(U)}^2) \\ \leq C (\|\mathbf{f}\|_{H^1(0,T;L^2(U))}^2 + \|g\|_{H^2(U)}^2 + \|h\|_{H^1(U)}^2).$$

Here we estimated $\|\mathbf{u}_m(0)\|_{H^2(U)} \leq C \|g\|_{H^2(U)}$, as in the proof of Theorem 5 in §7.1.3.

Passing to limits as $m = m_l \rightarrow \infty$, we derive the same bound for \mathbf{u} .

4. As in the earlier proof of Theorem 5 in §7.1.3, we likewise deduce $\mathbf{u}''' \in L^2(0, T; H^{-1}(U))$, with the stated estimate. \square

Remark. If L were symmetric, we could alternatively have taken $\{w_k\}_{k=1}^{\infty}$ to be a basis of eigenfunctions of L on $H_0^1(U)$ and so avoided the need for inequality (59).

Now let m be a nonnegative integer.

THEOREM 6 (Higher regularity). *Assume*

$$\begin{cases} g \in H^{m+1}(U), h \in H^m(U), \\ \frac{d^k \mathbf{f}}{dt^k} \in L^2(0, T; H^{m-k}(U)) \quad (k = 0, \dots, m). \end{cases}$$

Suppose also that the following m^{th} -order compatibility conditions hold:

$$(62) \quad \begin{cases} g_0 := g \in H_0^1(U), h_1 := h \in H_0^1(U), \dots, \\ g_{2l} := \frac{d^{2l-2} \mathbf{f}}{dt^{2l-2}}(\cdot, 0) - Lg_{2l-2} \in H_0^1(U) \quad (\text{if } m = 2l) \\ h_{2l+1} := \frac{d^{2l-1} \mathbf{f}}{dt^{2l-1}}(\cdot, 0) - Lh_{2l-1} \in H_0^1(U) \quad (\text{if } m = 2l + 1). \end{cases}$$

Then

$$(63) \quad \frac{d^k \mathbf{u}}{dt^k} \in L^\infty(0, T; H^{m+1-k}(U)) \quad (k = 0, \dots, m + 1),$$

and we have the estimate

$$(64) \quad \begin{aligned} & \text{ess sup}_{0 \leq t \leq T} \sum_{k=0}^{m+1} \left\| \frac{d^k \mathbf{u}}{dt^k} \right\|_{H^{m+1-k}(U)} \\ & \leq C \left(\sum_{k=0}^m \left\| \frac{d^k \mathbf{f}}{dt^k} \right\|_{L^2(0, T; H^{m-k}(U))} + \|g\|_{H^{m+1}(U)} + \|h\|_{H^m(U)} \right). \end{aligned}$$

Remark. In view of Theorem 2 in §5.9.2, we see that

$$(65) \quad \mathbf{f}(0) \in H^{m-1}(U), \mathbf{f}'(0) \in H^{m-2}(U), \dots, \mathbf{f}^{(m-2)}(0) \in H^1(U),$$

and consequently

$$(66) \quad \begin{aligned} & g_0 \in H^{m+1}(U), h_1 \in H^m(U), g_2 \in H^{m-1}(U), h_3 \in H^{m-2}(U), \\ & \dots, g_{2l} \in H^1(U) \quad (\text{if } m = 2l), h_{2l+1} \in H^1(U) \quad (\text{if } m = 2l + 1). \end{aligned}$$

The compatibility conditions are consequently the requirements that, in addition, each of these functions equals 0 on ∂U , in the trace sense.

Proof. 1. The proof is by an induction, the case $m = 0$ following from Theorem 5(i) above.

2. Assume next the theorem is valid for some nonnegative integer m , and suppose

$$(67) \quad \begin{cases} g \in H^{m+2}(U), h \in H^{m+1}(U), \\ \frac{d^k \mathbf{f}}{dt^k} \in L^2(0, T; H^{m+1-k}(U)) \quad (k = 0, \dots, m+1). \end{cases}$$

Suppose also the $(m+1)^{\text{th}}$ -order compatibility conditions obtain. Now set $\tilde{\mathbf{u}} := \mathbf{u}'$. Differentiating the PDE with respect to t , we check that $\tilde{\mathbf{u}}$ is the unique, weak solution of

$$(68) \quad \begin{cases} \tilde{u}_{tt} + L\tilde{u} = \tilde{f} & \text{in } U_T \\ \tilde{u} = 0 & \text{on } \partial U \times [0, T] \\ \tilde{u} = \tilde{g}, \tilde{u}_t = \tilde{h} & \text{on } U \times \{t = 0\}, \end{cases}$$

for

$$(69) \quad \tilde{f} := f_t, \tilde{g} := h, \tilde{h} := f(\cdot, 0) - Lg.$$

In particular, for $m = 0$ we rely upon Theorem 5(ii) to be sure that $\tilde{\mathbf{u}} \in L^2(0, T; H_0^1(U))$, $\tilde{\mathbf{u}}' \in L^2(0, T; L^2(U))$, $\tilde{\mathbf{u}}'' \in L^2(0, T; H^{-1}(U))$.

Since f, g and h satisfy the $(m+1)^{\text{th}}$ -order compatibility conditions, \tilde{f}, \tilde{g} and \tilde{h} satisfy the m^{th} -order compatibility conditions. Thus applying the induction assumption, we see

$$\frac{d^k \tilde{\mathbf{u}}}{dt^k} \in L^\infty(0, T; H^{m+1-k}(U)) \quad (k = 0, \dots, m+1),$$

with the estimate

$$\begin{aligned} & \text{ess sup}_{0 \leq t \leq T} \sum_{k=0}^{m+1} \left\| \frac{d^k \tilde{\mathbf{u}}}{dt^k} \right\|_{H^{m+1-k}(U)} \\ & \leq C \left(\sum_{k=0}^m \left\| \frac{d^k \tilde{\mathbf{f}}}{dt^k} \right\|_{L^2(0, T; H^{m-k}(U))} + \|\tilde{g}\|_{H^{m+1}(U)} + \|\tilde{h}\|_{H^m(U)} \right). \end{aligned}$$

Since $\tilde{\mathbf{u}} = \mathbf{u}'$, we can rewrite:

$$(70) \quad \begin{aligned} & \text{ess sup}_{0 \leq t \leq T} \sum_{k=1}^{m+2} \left\| \frac{d^k \mathbf{u}}{dt^k} \right\|_{H^{m+2-k}(U)} \\ & \leq C \left(\sum_{k=1}^{m+1} \left\| \frac{d^k \mathbf{f}}{dt^k} \right\|_{L^2(0, T; H^{m+1-k}(U))} + \|h\|_{H^{m+1}(U)} + \|Lg\|_{H^m(U)} + \|\mathbf{f}(0)\|_{H^m(U)} \right) \\ & \leq C \left(\sum_{k=0}^{m+1} \left\| \frac{d^k \mathbf{f}}{dt^k} \right\|_{L^2(0, T; H^{m+1-k}(U))} + \|g\|_{H^{m+2}(U)} + \|h\|_{H^{m+1}(U)} \right). \end{aligned}$$

Here we used the inequality

$$\|\mathbf{f}\|_{C([0,T];H^m(U))} \leq C(\|\mathbf{f}\|_{L^2(0,T;H^m(U))} + \|\mathbf{f}'\|_{L^2(0,T;H^m(U))}),$$

which follows from Theorem 2 in §5.9.2.

3. Now write for a.e. $0 \leq t \leq T$ that $Lu = \mathbf{f} - \mathbf{u}'' =: \mathbf{h}$. We have

$$\begin{aligned} \|\mathbf{u}\|_{H^{m+2}(U)} &\leq C(\|\mathbf{h}\|_{H^m(U)} + \|\mathbf{u}\|_{L^2(U)}) \\ &\leq C(\|\mathbf{f}\|_{H^m(U)} + \|\mathbf{u}''\|_{H^m(U)} + \|\mathbf{u}\|_{L^2(U)}). \end{aligned}$$

Taking the essential supremum with respect to t , adding this inequality to (70) and making standard estimates, we deduce

$$\begin{aligned} \operatorname{ess\,sup}_{0 \leq t \leq T} \sum_{k=0}^{m+2} \left\| \frac{d^k \mathbf{u}}{dt^k} \right\|_{H^{m+2-k}(U)} \\ \leq C \left(\sum_{k=0}^{m+1} \left\| \frac{d^k \mathbf{f}}{dt^k} \right\|_{L^2(0,T;H^{m+1-k}(U))} + \|g\|_{H^{m+2}(U)} + \|h\|_{H^{m+1}(U)} \right). \end{aligned}$$

This is the assertion of the theorem for $m + 1$. □

THEOREM 7 (Infinite differentiability). *Assume*

$$g, h \in C^\infty(\bar{U}), \quad f \in C^\infty(\bar{U}_T),$$

and the m^{th} -order compatibility conditions hold for $m = 0, 1, \dots$


Then the hyperbolic initial/boundary-value problem (1) has a unique solution

$$u \in C^\infty(\bar{U}_T).$$

Proof. Apply Theorem 6 for $m = 0, 1, \dots$ □

7.2.4. Propagation of disturbances.

Our study of second-order hyperbolic equations has thus far pretty much paralleled our treatment of second-order parabolic PDE, in §7.1. In the corresponding earlier section §7.1.4, we discussed maximum principles for second-order parabolic equations and noted in particular that the strong maximum principle implies an “infinite propagation speed” of initial disturbances for such PDE. Now strong maximum principles are false for second-order hyperbolic partial differential equations, and we will instead address here the opposite phenomenon, namely the “finite propagation speed” of



Domain of dependence

initial disturbances. This study extends some ideas already introduced in §2.4.3.

For simplicity we will consider in this subsection the case that $U = \mathbb{R}^n$ and L has the simple nondivergence form

$$(71) \quad Lu = - \sum_{i,j=1}^n a^{ij} u_{x_i x_j},$$

where the coefficients are smooth, independent of time, and there are no lower-order terms. We as usual require the uniform hyperbolicity condition.

Let us assume now u is a smooth solution of the PDE

$$(72) \quad u_{tt} + Lu = 0 \quad \text{in } \mathbb{R}^n \times (0, \infty).$$

We wish to prove a uniqueness/finite propagation speed assertion analogous to that obtained for the wave equation in §2.4.3. For this, we fix a point $(x_0, t_0) \in \mathbb{R}^n \times (0, \infty)$ and then try to find some sort of a curved “cone-like” region C , with vertex (x_0, t_0) , such that $u \equiv 0$ within C if $u \equiv u_t \equiv 0$ on $C_0 = C \cap \{t = 0\}$.

Motivated by the geometric optics computation in Example 3 of §4.5.3, let us guess that the boundary of such a region C is given as a level set $\{p = 0\}$, where p solves the Hamilton–Jacobi PDE

$$(73) \quad p_t - \left(\sum_{i,j=1}^n a^{ij}(x) p_{x_i} p_{x_j} \right)^{1/2} = 0 \quad \text{in } \mathbb{R}^n \times (0, \infty).$$

We will simplify (73) by separating variables, to write

$$(74) \quad p(x, t) = q(x) + t - t_0 \quad (x \in \mathbb{R}^n, 0 \leq t \leq t_0),$$

where q solves

$$(75) \quad \begin{cases} \sum_{i,j=1}^n a^{ij} q_{x_i} q_{x_j} = 1, & q > 0 \quad \text{in } \mathbb{R}^n - \{x_0\} \\ q(x_0) = 0. \end{cases}$$

We henceforth assume that q is a smooth solution of (75) on $\mathbb{R}^n - \{x_0\}$. (In fact $q(x)$ is the distance of x to x_0 , in the Riemannian metric determined by $((a^{ij}))$.) We introduce the “curved” backwards wave cone

$$K := \{(x, t) \mid p(x, t) < 0\} = \{(x, t) \mid q(x) < t_0 - t\}.$$

For each $t > 0$, we further define

$$(76) \quad K_t := \{x \mid q(x) < t_0 - t\} = \text{cross section of } K \text{ at time } t.$$

Since (75) implies $Dq \neq 0$ in $\mathbb{R}^n - \{x_0\}$, ∂K_t is a smooth, $(n-1)$ -dimensional surface for $0 \leq t < t_0$.

THEOREM 8 (Finite propagation speed). *Assume u is a smooth solution of the hyperbolic equation (72). If $u \equiv u_t \equiv 0$ on K_0 , then $u \equiv 0$ within K .*

We see in particular that if u is a solution of (72) with the initial conditions

$$(77) \quad u = g, \quad u_t = h \quad \text{on } \mathbb{R}^n \times \{t = 0\},$$

then $u(x_0, t_0)$ depends only upon the values of g and h within K_0 .

Proof. 1. We modify a proof from §2.4.3 and so define the energy

$$e(t) := \frac{1}{2} \int_{K_t} u_t^2 + \sum_{i,j=1}^n a^{ij} u_{x_i} u_{x_j} dx \quad (0 \leq t \leq t_0).$$

2. In order to compute $\dot{e}(t)$, we first note that if f is a continuous function of x , then

$$\frac{d}{dt} \left(\int_{K_t} f dx \right) = - \int_{\partial K_t} \frac{f}{|Dq|} dS$$

according to the coarea formula from §C.3. Thus

$$(78) \quad \begin{aligned} \dot{e}(t) &= \int_{K_t} u_t u_{tt} + \sum_{i,j=1}^n a^{ij} u_{x_i} u_{x_j t} dx \\ &\quad - \frac{1}{2} \int_{\partial K_t} \left(u_t^2 + \sum_{i,j=1}^n a^{ij} u_{x_i} u_{x_j} \right) \frac{1}{|Dq|} dS \\ &=: A - B. \end{aligned}$$

Integrating by parts, we calculate

$$\begin{aligned}
 (79) \quad A &= \int_{K_t} u_t \left(u_{tt} - \sum_{i,j=1}^n (a^{ij} u_{x_i})_{x_j} \right) dx + \int_{\partial K_t} \sum_{i,j=1}^n a^{ij} u_{x_i} \nu^j u_t dS \\
 &= - \int_{K_t} u_t \sum_{i,j=1}^n a_{x_j}^{ij} u_{x_i} dx + \int_{\partial K_t} \sum_{i,j=1}^n a^{ij} u_{x_i} \nu^j u_t dS,
 \end{aligned}$$

with $\nu = (\nu^1, \dots, \nu^n)$ being as usual the outer unit normal to ∂K_t . But according to the generalized Cauchy–Schwarz inequality (§B.2)

$$(80) \quad \left| \sum_{i,j=1}^n a^{ij} u_{x_i} \nu^j \right| \leq \left(\sum_{i,j=1}^n a^{ij} u_{x_i} u_{x_j} \right)^{1/2} \left(\sum_{i,j=1}^n a^{ij} \nu^i \nu^j \right)^{1/2}.$$

In addition, since $q = t_0 - t$ on ∂K_t , we have $\nu = \frac{Dq}{|Dq|}$ on ∂K_t . Hence

$$\sum_{i,j=1}^n a^{ij} \nu^i \nu^j = \sum_{i,j=1}^n \frac{a^{ij} q_{x_i} q_{x_j}}{|Dq|^2} = \frac{1}{|Dq|^2}$$

by (75). Consequently inequality (80) reads

$$(81) \quad \left| \sum_{i,j=1}^n a^{ij} u_{x_i} \nu^j \right| \leq \left(\sum_{i,j=1}^n a^{ij} u_{x_i} u_{x_j} \right)^{1/2} \frac{1}{|Dq|}.$$

Then returning to (79), we estimate using (81) and Cauchy's inequality:

$$\begin{aligned}
 |A| &\leq Ce(t) + \int_{\partial K_t} \left(\sum_{i,j=1}^n a^{ij} u_{x_i} u_{x_j} \right)^{1/2} |u_t| \frac{1}{|Dq|} dS \\
 &\leq Ce(t) + \frac{1}{2} \int_{\partial C_t} \left(u_t^2 + \sum_{i,j=1}^n a^{ij} u_{x_i} u_{x_j} \right) \frac{1}{|Dq|} dS \\
 &= Ce(t) + B.
 \end{aligned}$$

3. Therefore inequality (78) gives

$$\dot{e}(t) \leq Ce(t).$$

Since hypothesis (76) implies $e(0) = 0$, we deduce using Gronwall's inequality that

$$e(t) = 0 \quad \text{for all } 0 \leq t \leq t_0.$$

Hence $u_t \equiv Du \equiv 0$ in K , and consequently $u \equiv 0$ in K .

□

7.2.5. Equations in two variables.

In this subsection we briefly consider second-order hyperbolic partial differential equations involving only two variables and demonstrate that in this setting rather more precise information can be obtained. The very rough idea is that since a function of two variables has “only” three second partial derivatives, algebraic and analytic simplifications in the structure of the PDE may be possible, which are unavailable for more than two variables.

We begin by considering a general linear second-order PDE in two variables

$$(82) \quad \sum_{i,j=1}^2 a^{ij} u_{x_i x_j} + \sum_{i=1}^2 b^i u_{x_i} + cu = 0,$$

where the coefficients a^{ij}, b^i, c ($i, j = 1, 2$), with $a^{ij} = a^{ji}$, and the unknown u are functions of the two variables x_1 and x_2 in some region $U \subset \mathbb{R}^2$. Note that for the moment, and in contrast to the theory developed above, we do *not* identify either x_1 or x_2 with the variable t denoting time.

We now pose the following basic question: *is it possible to simplify the structure of the PDE (82) by introducing new independent variables?* In other words, can we expect to convert the PDE into some “nicer” form by rewriting in terms of new variables $y = \Phi(x)$?

More precisely, let us set

$$(83) \quad \begin{cases} y_1 = \Phi^1(x_1, x_2) \\ y_2 = \Phi^2(x_1, x_2) \end{cases}$$

for some appropriate function $\Phi = (\Phi^1, \Phi^2)$. To investigate this possibility let us now write

$$(84) \quad u(x) = v(\Phi(x)).$$

That is, we define $v(y) := u(\Psi(y))$, where $\Psi = \Phi^{-1}$.

From (84), we compute

$$\begin{cases} u_{x_i} = \sum_{k=1}^2 v_{y_k} \Phi_{x_i}^k \\ u_{x_i x_j} = \sum_{k,l=1}^2 v_{y_k y_l} \Phi_{x_i}^k \Phi_{x_j}^l + \sum_{k=1}^2 v_{y_k} \Phi_{x_i x_j}^k \end{cases}$$

for $i, j = 1, 2$. Substituting into (82), we discover that v solves the PDE

$$(85) \quad \sum_{k,l=1}^n \tilde{a}^{kl} v_{y_k y_l} + \cdots = 0,$$

for

$$(86) \quad \tilde{a}^{kl} := \sum_{i,j=1}^2 a^{ij} \Phi_{x_i}^k \Phi_{x_j}^l \quad (k, l = 1, 2),$$

where the dots in (85) represent terms of lower order.

We examine the first term in the PDE (85) in the hope we can perhaps choose the transformation $\Phi = (\Phi^1, \Phi^2)$ so this expression is particularly simple. Let us try to achieve

$$(87) \quad \tilde{a}^{11} \equiv \tilde{a}^{22} \equiv 0.$$

In view of formula (86) this will be possible provided we can choose both Φ^1 and Φ^2 to solve the nonlinear first-order PDE

$$(88) \quad a^{11}(v_{x_1})^2 + 2a^{12}v_{x_1}v_{x_2} + a^{22}(v_{x_2})^2 = 0 \quad \text{in } U.$$

Observe this is the *characteristic equation* associated with the partial differential equation (82), as discussed in §4.6.2.

To proceed further, let us suppose

$$(89) \quad \det \mathbf{A} = a^{11}a^{22} - (a^{12})^2 < 0 \quad \text{in } U,$$

in which case we say the PDE (82) is *hyperbolic*.

Utilizing condition (89), we can then factor equation (88) as follows:

$$(90) \quad \begin{aligned} & \left(a^{11}v_{x_1} + \left[a^{12} + \left((a^{12})^2 - a^{11}a^{22} \right)^{1/2} \right] v_{x_2} \right) \\ & \cdot \left(a^{11}v_{x_1} + \left[a^{12} - \left((a^{12})^2 - a^{11}a^{22} \right)^{1/2} \right] v_{x_2} \right) \\ & = a^{11} \left(a^{11}(v_{x_1})^2 + 2a^{12}v_{x_1}v_{x_2} + a^{22}(v_{x_2})^2 \right) = 0. \end{aligned}$$

Now the left-hand side of (90) is the product of two linear first-order PDE:

$$(91_1) \quad a^{11}v_{x_1} + \left[a^{12} + \left((a^{12})^2 - a^{11}a^{22} \right)^{1/2} \right] v_{x_2} = 0 \quad \text{in } U$$

and

$$(91_2) \quad a^{11}v_{x_1} + \left[a^{12} - \left((a^{12})^2 - a^{11}a^{22} \right)^{1/2} \right] v_{x_2} = 0 \quad \text{in } U.$$

We now assume that we can choose Φ^1 to be a smooth solution of the PDE (91₁), with $D\Phi^1 \neq 0$ in U . Then Φ^1 is constant along trajectories $\mathbf{x} = (x^1, x^2)$ of the ODE

$$(92) \quad \begin{cases} \dot{x}^1 = a^{11} \\ \dot{x}^2 = \left[a^{12} + ((a^{12})^2 - a^{11}a^{22})^{1/2} \right]. \end{cases}$$

Similarly, suppose Φ^2 is a smooth solution of (91₂), with $D\Phi^2 \neq 0$ in U ; then Φ^2 is constant along trajectories $\mathbf{x} = (x^1, x^2)$ of the ODE

$$(93) \quad \begin{cases} \dot{x}^1 = a^{11} \\ \dot{x}^2 = \left[a^{12} - ((a^{12})^2 - a^{11}a^{22})^{1/2} \right]. \end{cases}$$

Curves which are trajectories of either the ODE (92) or (93) are called *characteristics* of the original partial differential equation (82). Returning now to (83), we see that trajectories of solutions of the characteristic ODE (92) and (93) provide our new coordinate lines.

Additionally we can verify using (89) that

$$(94) \quad \tilde{a}^{12} = \sum_{i,j=1}^2 a^{ij} \Phi_{x_i}^1 \Phi_{x_j}^2 \neq 0 \quad \text{in } U.$$

Combining then (85), (86), (87) and (94), we see that our PDE (82) becomes in the y coordinates

$$(95) \quad v_{y_1 y_2} + \cdots = 0,$$

the dots as before denoting terms of lower order. Let us call equation (95) the *first canonical form* for the hyperbolic PDE (82).

If we change variables again by setting $z_1 = y_1 + y_2$, $z_2 = y_1 - y_2$, then (95) becomes

$$(96) \quad w_{z_1 z_1} - w_{z_2 z_2} + \cdots = 0.$$

If we then further rename the variables $t = z_1$, $x = z_2$, then (96) reads

$$(97) \quad w_{tt} - w_{xx} + \cdots = 0,$$

the second-order term of which is the one-dimensional wave operator. Equation (97) is the *second canonical form*.

Hence any hyperbolic PDE in two variables of the form (82) can be converted by a change of variables into the wave equation plus a lower-order term, assuming we can find the mapping Φ as above.

7.3. HYPERBOLIC SYSTEMS OF FIRST-ORDER EQUATIONS

We next broaden our study of hyperbolic PDE (which we may informally interpret as equations supporting “wave-like” solutions) to the case of first-order systems. We continue in the manner of §§7.1 and 7.2 by first employing energy bounds to construct weak solutions for symmetric hyperbolic systems. For nonsymmetric, constant coefficient hyperbolic systems, however, we will instead employ Fourier transform methods.

7.3.1. Definitions.

We investigate in this section *systems* of linear first-order partial differential equations having the form

$$(1) \quad \mathbf{u}_t + \sum_{j=1}^n \mathbf{B}_j \mathbf{u}_{x_j} = \mathbf{f} \quad \text{in } \mathbb{R}^n \times (0, \infty),$$

subject to the initial condition

$$(2) \quad \mathbf{u} = \mathbf{g} \quad \text{on } \mathbb{R}^n \times \{t = 0\}.$$

The unknown is $\mathbf{u} : \mathbb{R}^n \times [0, \infty) \rightarrow \mathbb{R}^m$, $\mathbf{u} = (u^1, \dots, u^m)$, and the functions $\mathbf{B}_j : \mathbb{R}^n \times [0, \infty) \rightarrow \mathbb{M}^{m \times m}$ ($j = 1, \dots, n$), $\mathbf{f} : \mathbb{R}^n \times [0, \infty) \rightarrow \mathbb{R}^m$, $\mathbf{g} : \mathbb{R}^n \rightarrow \mathbb{R}^m$ are given.

NOTATION. For each $\mathbf{y} \in \mathbb{R}^n$, set

$$\mathbf{B}(\mathbf{x}, t; \mathbf{y}) := \sum_{j=1}^n y_j \mathbf{B}_j(\mathbf{x}, t) \quad (\mathbf{x} \in \mathbb{R}^n, t \geq 0).$$

DEFINITION. *The system of PDE (1) is called hyperbolic if the $m \times m$ matrix $\mathbf{B}(\mathbf{x}, t; \mathbf{y})$ is diagonalizable for each $\mathbf{x}, \mathbf{y} \in \mathbb{R}^n$, $t \geq 0$.*

In other words, (1) is hyperbolic provided for each $\mathbf{x}, \mathbf{y}, t$, the matrix $\mathbf{B}(\mathbf{x}, t; \mathbf{y})$ has m real eigenvalues

$$\lambda_1(\mathbf{x}, t; \mathbf{y}) \leq \lambda_2(\mathbf{x}, t; \mathbf{y}) \leq \dots \leq \lambda_m(\mathbf{x}, t; \mathbf{y})$$

and corresponding eigenvectors $\{\mathbf{r}_k(\mathbf{x}, t; \mathbf{y})\}_{k=1}^m$ that form a basis of \mathbb{R}^m .

There are two important special cases:

DEFINITIONS. (i) We say (1) is a symmetric hyperbolic system if $\mathbf{B}_j(x, t)$ is a symmetric $m \times m$ matrix for each $x \in \mathbb{R}^n$, $t \geq 0$ ($j = 1, \dots, m$).

(ii) The system (1) is strictly hyperbolic if for each $x, y \in \mathbb{R}^n$, $y \neq 0$, and each $t \geq 0$, the matrix $\mathbf{B}(x, t; y)$ has m distinct real eigenvalues:

$$\lambda_1(x, t; y) < \lambda_2(x, t; y) < \dots < \lambda_m(x, t; y).$$

Motivation for the definition of hyperbolicity. We justify the hyperbolicity condition as follows. Assume $\mathbf{f} \equiv 0$ and, further, the matrices B_j are constant ($j = 1, \dots, n$). Thus

$$(3) \quad \sum_{j=1}^n y_j B_j = \mathbf{B}(y)$$

depends only on $y \in \mathbb{R}^n$.

As in §4.2 let us look for a *plane wave* solution of (1), (2). That is, we seek a solution \mathbf{u} having the form

$$(4) \quad \mathbf{u}(x, t) = \mathbf{v}(y \cdot x - \sigma t) \quad (x \in \mathbb{R}^n, t \geq 0)$$

for some direction $y \in \mathbb{R}^n$, velocity $\frac{\sigma}{|y|}$ ($\sigma \in \mathbb{R}$), and profile $\mathbf{v} : \mathbb{R} \rightarrow \mathbb{R}^m$. Plugging (4) into (1), we compute

$$\left(-\sigma I + \sum_{j=1}^n y_j B_j \right) \mathbf{v}' = 0.$$

This equality asserts \mathbf{v}' is an eigenvector of the matrix $\mathbf{B}(y)$ corresponding to the eigenvalue σ .

The hyperbolicity condition requires that there are m distinct plane wave solutions of (1) for each direction y . These are

$$(y \cdot x - \lambda_k(y)t) \mathbf{r}_k(y) \quad (k = 1, \dots, m),$$

where

$$\lambda_1(y) \leq \lambda_2(y) \leq \dots \leq \lambda_m(y)$$

are the eigenvalues of $\mathbf{B}(y)$ and $\{\mathbf{r}_k(y)\}_{k=1}^m$ the corresponding eigenvectors. The eigenvalues for $|y| = 1$ are the *wave speeds*.

7.3.2. Symmetric hyperbolic systems.

In this section we apply energy methods and the vanishing viscosity technique to build a solution to the hyperbolic initial-value problem

$$(5) \quad \begin{cases} \mathbf{u}_t + \sum_{j=1}^n \mathbf{B}_j \mathbf{u}_{x_j} = \mathbf{f} & \text{in } \mathbb{R}^n \times (0, T] \\ \mathbf{u} = \mathbf{g} & \text{on } \mathbb{R}^n \times \{t = 0\}, \end{cases}$$

where $T > 0$, under the fundamental assumption that

$$(6) \quad \text{the matrices } \mathbf{B}_j(x, t) \text{ are symmetric } (j = 1, \dots, n),$$

for $x \in \mathbb{R}^n$, $0 \leq t \leq T$. We will further assume $\mathbf{B}_j \in C^2(\mathbb{R}^n \times [0, T]; \mathbb{M}^{m \times m})$, with

$$(7) \quad \sup_{\mathbb{R}^n \times [0, T]} (|\mathbf{B}_j|, |D_{x,t} \mathbf{B}_j|, |D_{x,t}^2 \mathbf{B}_j|) < \infty \quad (j = 1, \dots, n)$$

and

$$(8) \quad \mathbf{g} \in H^1(\mathbb{R}^n; \mathbb{R}^m), \quad \mathbf{f} \in H^1(\mathbb{R}^n \times (0, T); \mathbb{R}^m).$$

Remark. More general systems having the form

$$(9) \quad \mathbf{B}_0 \mathbf{u}_t + \sum_{j=1}^n \mathbf{B}_j \mathbf{u}_{x_j} = \mathbf{f}$$

are also called *symmetric*, provided the matrices \mathbf{B}_j are symmetric for $j = 0, \dots, n$. The theory set forth below easily extends to such systems, provided \mathbf{B}_0 is positive definite.

Symmetric hyperbolic systems of the type (9) generalize the second-order hyperbolic PDE studied in §7.2. For suppose v is a smooth solution of the scalar equation

$$(10) \quad v_{tt} - \sum_{i,j=1}^n a^{ij} v_{x_i x_j} = 0,$$

where without loss of generality we may take $a^{ij} = a^{ji}$ ($i, j = 1, \dots, n$). Writing

$$\mathbf{u} = (u^1, \dots, u^{n+1}) := (v_{x_1}, \dots, v_{x_n}, v_t),$$

we discover \mathbf{u} solves a system of the form (9), for $m = n + 1$, $\mathbf{f} \equiv 0$,

$$\mathbf{B}_j = \begin{pmatrix} 0 & \dots & 0 & -a^{1j} \\ & \ddots & & \vdots \\ 0 & \dots & 0 & -a^{nj} \\ -a^{1j} & \dots & -a^{nj} & 0 \end{pmatrix}_{(n+1) \times (n+1)} \quad (j = 1, \dots, n),$$

$$\mathbf{B}_0 = \begin{pmatrix} a^{11} & \dots & a^{1n} & 0 \\ & \ddots & & \vdots \\ a^{1n} & \dots & a^{nn} & 0 \\ 0 & \dots & 0 & 1 \end{pmatrix}_{(n+1) \times (n+1)}.$$

Observe that the uniform hyperbolicity condition for (10) implies that the matrix \mathbf{B}_0 is positive definite.

a. Weak solutions. To ease notation, let us define the bilinear form

$$B[\mathbf{u}, \mathbf{v}; t] := \int_{\mathbb{R}^n} \sum_{j=1}^n (\mathbf{B}_j(\cdot, t) \mathbf{u}_{x_j}) \cdot \mathbf{v} \, dx$$

for $0 \leq t \leq T$, $\mathbf{u}, \mathbf{v} \in H^1(\mathbb{R}^n; \mathbb{R}^m)$.

DEFINITION. We say

$$\mathbf{u} \in L^2(0, T; H^1(\mathbb{R}^n; \mathbb{R}^m)), \text{ with } \mathbf{u}' \in L^2(0, T; L^2(\mathbb{R}^n; \mathbb{R}^m)),$$

is a weak solution of the initial-value problem (5) for the symmetric hyperbolic system provided

$$(i) \quad (\mathbf{u}', \mathbf{v}) + B[\mathbf{u}, \mathbf{v}; t] = (\mathbf{f}, \mathbf{v})$$

for each $\mathbf{v} \in H^1(\mathbb{R}^n; \mathbb{R}^m)$ and a.e. $0 \leq t \leq T$ and

$$(ii) \quad \mathbf{u}(0) = \mathbf{g}.$$

Here and afterwards (\cdot, \cdot) denotes the inner product in $L^2(\mathbb{R}^n; \mathbb{R}^m)$.

Remark. According to Theorem 2 in §5.9.2, $\mathbf{u} \in C([0, T]; L^2(\mathbb{R}^n; \mathbb{R}^m))$ and so the initial condition (ii) makes sense.

b. Vanishing viscosity method. We will approximate problem (5) by the parabolic initial-value problem

$$(11) \quad \begin{cases} \mathbf{u}_t^\epsilon - \epsilon \Delta \mathbf{u}^\epsilon + \sum_{j=1}^n \mathbf{B}_j \mathbf{u}_{x_j}^\epsilon = \mathbf{f} & \text{in } \mathbb{R}^n \times (0, T] \\ \mathbf{u}^\epsilon = \mathbf{g}^\epsilon & \text{on } \mathbb{R}^n \times \{t = 0\} \end{cases}$$

for $0 < \epsilon \leq 1$, $\mathbf{g}^\epsilon := \eta_\epsilon * \mathbf{g}$. The idea is that for each $\epsilon > 0$, problem (11) has a unique smooth solution \mathbf{u}^ϵ , which converges to zero as $|x| \rightarrow \infty$. The plan is to show that as $\epsilon \rightarrow 0$, the \mathbf{u}^ϵ converge to a limit function \mathbf{u} , which is a weak solution of (5).

THEOREM 1 (Existence of approximate solutions). *For each $\epsilon > 0$, there exists a unique solution \mathbf{u}^ϵ of (11), with*

$$(12) \quad \mathbf{u}^\epsilon \in L^2(0, T; H^3(\mathbb{R}^n; \mathbb{R}^m)), \quad \mathbf{u}^{\epsilon'} \in L^2(0, T; H^1(\mathbb{R}^n; \mathbb{R}^m)).$$

Proof. 1. Set $X = L^\infty((0, T); H^1(\mathbb{R}^n; \mathbb{R}^m))$. For each $\mathbf{v} \in X$, consider the linear system

$$\begin{cases} \mathbf{u}_t - \epsilon \Delta \mathbf{u} = \mathbf{f} - \sum_{j=1}^n \mathbf{B}_j \mathbf{v}_{x_j} & \text{in } \mathbb{R}^n \times (0, T] \\ \mathbf{u} = \mathbf{g}^\epsilon & \text{on } \mathbb{R}^n \times \{t = 0\}. \end{cases}$$

As the right-hand side is bounded in L^2 , there exists a unique solution $\mathbf{u} \in L^2(0, T; H^2(\mathbb{R}^n; \mathbb{R}^m))$, $\mathbf{u}' \in L^2(0, T; L^2(\mathbb{R}^n; \mathbb{R}^m))$. Indeed, we can utilize the fundamental solution Φ of the heat equation (§2.3.1) to represent \mathbf{u}^ϵ in terms of \mathbf{g}^ϵ and $\mathbf{f} - \sum_{j=1}^n \mathbf{B}_j \mathbf{v}_{x_j}$.

Similarly, take $\tilde{\mathbf{v}} \in X$ and let $\tilde{\mathbf{u}}$ solve

$$\begin{cases} \tilde{\mathbf{u}}_t - \epsilon \Delta \tilde{\mathbf{u}} = \mathbf{f} - \sum_{j=1}^n \mathbf{B}_j \tilde{\mathbf{v}}_{x_j} & \text{in } \mathbb{R}^n \times (0, T] \\ \tilde{\mathbf{u}} = \mathbf{g}^\epsilon & \text{on } \mathbb{R}^n \times \{t = 0\}. \end{cases}$$

2. Subtracting, we find $\hat{\mathbf{u}} := \mathbf{u} - \tilde{\mathbf{u}}$ satisfies

$$(13) \quad \begin{cases} \hat{\mathbf{u}}_t - \epsilon \Delta \hat{\mathbf{u}} = - \sum_{j=1}^n \mathbf{B}_j \hat{\mathbf{v}}_{x_j} & \text{in } \mathbb{R}^n \times (0, T] \\ \hat{\mathbf{u}} = \mathbf{0} & \text{on } \mathbb{R}^n \times \{t = 0\}, \end{cases}$$

for $\hat{\mathbf{v}} := \mathbf{v} - \tilde{\mathbf{v}}$. From the representation formula of $\hat{\mathbf{u}}$ in terms of the fundamental solution Φ and $\sum_{j=1}^n \mathbf{B}_j \hat{\mathbf{v}}_{x_j}$, we obtain the estimate

$$(14) \quad \begin{aligned} \operatorname{ess\,sup}_{0 \leq t \leq T} \|\hat{\mathbf{u}}(t)\|_{H^1(\mathbb{R}^n; \mathbb{R}^m)} &\leq C(\epsilon) \left\| \sum_{j=1}^n \mathbf{B}_j \hat{\mathbf{v}}_{x_j} \right\|_{L^2(0, T; L^2(\mathbb{R}^n; \mathbb{R}^m))} \\ &\leq C(\epsilon) \|\hat{\mathbf{v}}\|_{L^2(0, T; H^1(\mathbb{R}^n; \mathbb{R}^m))} \\ &\leq C(\epsilon) T^{1/2} \operatorname{ess\,sup}_{0 \leq t \leq T} \|\hat{\mathbf{v}}(t)\|_{H^1(\mathbb{R}^n; \mathbb{R}^m)}. \end{aligned}$$

Thus

$$(15) \quad \|\hat{\mathbf{u}}\| \leq C(\epsilon) T^{1/2} \|\hat{\mathbf{v}}\|.$$

3. If T is so small that

$$(16) \quad C(\epsilon) T^{1/2} \leq 1/2,$$

then (15) reads $\|\mathbf{u} - \tilde{\mathbf{u}}\| \leq \frac{1}{2} \|\mathbf{v} - \tilde{\mathbf{v}}\|$. According to Banach's Fixed Point Theorem (§9.2.1) the mapping $\mathbf{v} \mapsto \mathbf{u}$ has a unique fixed point. Then $\mathbf{u} = \mathbf{u}^\epsilon$ solves (11), provided (16) holds.

If (16) fails, we choose $0 < T_1 < T$ so that $CT_1^{1/2} = 1/2$ and repeat the above argument on the time intervals $[0, T_1]$, $[T_1, 2T_1]$, etc.

Assertion (12) follows from parabolic regularity theory (cf. §7.1.3). \square

c. Energy estimates. We want to send $\varepsilon \rightarrow 0$ in (11), and for this as usual need some uniform estimates.

THEOREM 2 (Energy estimates). *There exists a constant C , depending only on n and the coefficients, such that*

$$(17) \quad \begin{aligned} & \max_{0 \leq t \leq T} (\|\mathbf{u}^\varepsilon(t)\|_{H^1(\mathbb{R}^n; \mathbb{R}^m)} + \|\mathbf{u}^{\varepsilon'}(t)\|_{L^2(\mathbb{R}^n; \mathbb{R}^m)}) \\ & \leq C(\|g\|_{H^1(\mathbb{R}^n; \mathbb{R}^m)} + \|\mathbf{f}\|_{L^2(0, T; H^1(\mathbb{R}^n; \mathbb{R}^m))} + \|\mathbf{f}'\|_{L^2(0, T; L^2(\mathbb{R}^n; \mathbb{R}^m))}) \end{aligned}$$

for each $0 < \varepsilon \leq 1$.

Proof. 1. We compute

$$(18) \quad \frac{d}{dt} \left(\frac{1}{2} \|\mathbf{u}^\varepsilon\|_{L^2(\mathbb{R}^n; \mathbb{R}^m)}^2 \right) = (\mathbf{u}^\varepsilon, \mathbf{u}^{\varepsilon'}) = \left(\mathbf{u}^\varepsilon, \mathbf{f} - \sum_{j=1}^n B_j \mathbf{u}_{x_j}^\varepsilon + \varepsilon \Delta \mathbf{u}^\varepsilon \right).$$

Now

$$(19) \quad |(\mathbf{u}^\varepsilon, \mathbf{f})| \leq \|\mathbf{u}^\varepsilon\|_{L^2(\mathbb{R}^n; \mathbb{R}^m)}^2 + \|\mathbf{f}\|_{L^2(\mathbb{R}^n; \mathbb{R}^m)}^2$$

and

$$(20) \quad (\mathbf{u}^\varepsilon, \varepsilon \Delta \mathbf{u}^\varepsilon) = -\varepsilon \|D\mathbf{u}^\varepsilon\|_{L^2(\mathbb{R}^n; \mathbb{M}^{m \times n})}^2 \leq 0.$$

2. Suppose $\mathbf{v} \in C_c^\infty(\mathbb{R}^n; \mathbb{R}^m)$. Then

$$\begin{aligned} \left(\mathbf{v}, \sum_{j=1}^n \mathbf{B}_j \mathbf{v}_{x_j} \right) &= \sum_{j=1}^n \int_{\mathbb{R}^n} (\mathbf{B}_j \mathbf{v}_{x_j}) \cdot \mathbf{v} \, dx \\ &= \frac{1}{2} \sum_{j=1}^n \int_{\mathbb{R}^n} ((\mathbf{B}_j \mathbf{v}) \cdot \mathbf{v})_{x_j} \, dx - \frac{1}{2} \sum_{j=1}^n \int_U (\mathbf{B}_{j, x_j} \mathbf{v}) \cdot \mathbf{v} \, dx, \end{aligned}$$

the last equality following from the symmetry assumption (6). As \mathbf{v} has compact support, we deduce using (7) that

$$\left| \left(\mathbf{v}, \sum_{j=1}^n \mathbf{B}_j \mathbf{v}_{x_j} \right) \right| \leq \frac{1}{2} \sum_{j=1}^n \left| \int_{\mathbb{R}^n} (\mathbf{B}_{j, x_j} \mathbf{v}) \cdot \mathbf{v} \, dx \right| \leq C \|\mathbf{v}\|_{L^2(\mathbb{R}^n; \mathbb{R}^m)}^2.$$

By approximation therefore

$$\left| \left(\mathbf{u}^\varepsilon, \sum_{j=1}^n \mathbf{B}_j \mathbf{u}_{x_j}^\varepsilon \right) \right| \leq C \|\mathbf{u}^\varepsilon\|_{L^2(\mathbb{R}^n; \mathbb{R}^m)}^2.$$

Utilizing this bound, (19) and (20) in (18), we obtain the estimate

$$\frac{d}{dt} (\|\mathbf{u}^\epsilon\|_{L^2(\mathbb{R}^n; \mathbb{R}^m)}^2) \leq C (\|\mathbf{u}^\epsilon\|_{L^2(\mathbb{R}^n; \mathbb{R}^m)}^2 + \|\mathbf{f}\|_{L^2(\mathbb{R}^n; \mathbb{R}^m)}^2).$$

We next apply Gronwall's inequality, to deduce

$$(21) \quad \max_{0 \leq t \leq T} \|\mathbf{u}^\epsilon(t)\|_{L^2(\mathbb{R}^n; \mathbb{R}^m)}^2 \leq C \left(\|\mathbf{g}\|_{L^2(\mathbb{R}^n; \mathbb{R}^m)}^2 + \|\mathbf{f}\|_{L^2(0, T; L^2(\mathbb{R}^n; \mathbb{R}^m))}^2 \right),$$

since $\|\mathbf{g}^\epsilon\|_{L^2(\mathbb{R}^n; \mathbb{R}^m)} \leq \|\mathbf{g}\|_{L^2(\mathbb{R}^n; \mathbb{R}^m)}$.

3. Fix $k \in \{1, \dots, n\}$ and write $\mathbf{v}^k := \mathbf{u}_{x_k}^\epsilon$. Differentiating (11) with respect to x_k , we find

$$\begin{cases} \mathbf{v}_t^k - \epsilon \Delta \mathbf{v}^k + \sum_{j=1}^n \mathbf{B}_j \mathbf{v}_{x_j}^k = \mathbf{f}_{x_k} - \sum_{j=1}^n \mathbf{B}_{j, x_k} \mathbf{u}_{x_j}^\epsilon & \text{in } \mathbb{R}^n \times (0, T] \\ \mathbf{v}^k = \mathbf{g}_{x_k}^\epsilon & \text{on } \mathbb{R}^n \times \{t = 0\}. \end{cases}$$

Reasoning as above, we find

$$(22) \quad \frac{d}{dt} (\|\mathbf{v}^k\|_{L^2(\mathbb{R}^n; \mathbb{R}^m)}^2) \leq C (\|D\mathbf{u}^\epsilon\|_{L^2(\mathbb{R}^n; \mathbb{M}^{m \times n})}^2 + \|D\mathbf{f}\|_{L^2(\mathbb{R}^n; \mathbb{M}^{m \times n})}^2).$$

Sum the previous inequalities for $k = 1, \dots, n$, to deduce

$$\frac{d}{dt} (\|D\mathbf{u}^\epsilon\|_{L^2(\mathbb{R}^n; \mathbb{M}^{m \times n})}^2) \leq C (\|D\mathbf{u}^\epsilon\|_{L^2(\mathbb{R}^n; \mathbb{M}^{m \times n})}^2 + \|D\mathbf{f}\|_{L^2(\mathbb{R}^n; \mathbb{M}^{m \times n})}^2).$$

Gronwall's inequality now provides the bound

$$(23) \quad \begin{aligned} \max_{0 \leq t \leq T} \|D\mathbf{u}^\epsilon(t)\|_{L^2(\mathbb{R}^n; \mathbb{M}^{m \times n})}^2 \\ \leq C (\|D\mathbf{g}\|_{L^2(\mathbb{R}^n; \mathbb{M}^{m \times n})}^2 + \|\mathbf{f}\|_{L^2(0, T; H^1(\mathbb{R}^n; \mathbb{R}^m))}^2), \end{aligned}$$

since $\|D\mathbf{g}^\epsilon\|_{L^2(\mathbb{R}^n; \mathbb{M}^{m \times n})} \leq \|D\mathbf{g}\|_{L^2(\mathbb{R}^n; \mathbb{M}^{m \times n})}$.

4. Next set $\mathbf{v} := \mathbf{u}^\epsilon$ and differentiate (11) with respect to t , to discover

$$(24) \quad \begin{cases} \mathbf{v}_t - \epsilon \Delta \mathbf{v} + \sum_{j=1}^n \mathbf{B}_j \mathbf{v}_{x_j} = \mathbf{f}' - \sum_{j=1}^n \mathbf{B}_{j, t} \mathbf{u}_{x_j}^\epsilon & \text{in } \mathbb{R}^n \times (0, T] \\ \mathbf{v} = \mathbf{f} - \sum_{j=1}^n \mathbf{B}_j \mathbf{g}_{x_j}^\epsilon + \epsilon \Delta \mathbf{g}^\epsilon & \text{on } \mathbb{R}^n \times \{t = 0\}. \end{cases}$$

Reasoning as before, we compute

$$(25) \quad \begin{aligned} \max_{0 \leq t \leq T} \|\mathbf{u}^{\epsilon'}(t)\|_{L^2(\mathbb{R}^n; \mathbb{R}^m)}^2 \leq C (\|D\mathbf{g}\|_{L^2(\mathbb{R}^n; \mathbb{M}^{m \times n})}^2 + \epsilon^2 \|\Delta \mathbf{g}^\epsilon\|_{L^2(\mathbb{R}^n; \mathbb{R}^m)}^2 \\ + \|\mathbf{f}(0)\|_{L^2(\mathbb{R}^n; \mathbb{R}^m)}^2 + \|\mathbf{f}\|_{L^2(0, T; H^1(\mathbb{R}^n; \mathbb{R}^m))}^2 + \|\mathbf{f}'\|_{L^2(0, T; L^2(\mathbb{R}^n; \mathbb{R}^m))}^2). \end{aligned}$$

Now

$$(26) \quad \|\Delta \mathbf{g}^\epsilon\|_{L^2(\mathbb{R}^n; \mathbb{R}^m)}^2 \leq \frac{C}{\epsilon^2} \|D\mathbf{g}\|_{L^2(\mathbb{R}^n; \mathbb{M}^{m \times n})}^2,$$

since $\mathbf{g}^\epsilon = \eta_\epsilon * \mathbf{g}$. Furthermore

$$(27) \quad \|\mathbf{f}(0)\|_{L^2(\mathbb{R}^n; \mathbb{R}^m)}^2 \leq C (\|\mathbf{f}\|_{L^2(0, T; L^2(\mathbb{R}^n; \mathbb{R}^m))}^2 + \|\mathbf{f}'\|_{L^2(0, T; L^2(\mathbb{R}^n; \mathbb{R}^m))}^2).$$

This bound, together with (21) and (23), completes the proof. \square

d. Existence and uniqueness.

THEOREM 3 (Existence of weak solution). *There exists a weak solution of the initial value problem (5).*

Proof. 1. According to the energy estimates (17) there exists a subsequence $\epsilon_k \rightarrow 0$ and a function $\mathbf{u} \in L^2(0, T; H^1(\mathbb{R}^n; \mathbb{R}^m))$, such that $\mathbf{u}' \in L^2(0, T; L^2(\mathbb{R}^n; \mathbb{R}^m))$, with

$$(28) \quad \begin{cases} \mathbf{u}^{\epsilon_k} \rightharpoonup \mathbf{u} & \text{weakly in } L^2(0, T; H^1(\mathbb{R}^n; \mathbb{R}^m)) \\ \mathbf{u}'^{\epsilon_k} \rightharpoonup \mathbf{u}' & \text{weakly in } L^2(0, T; L^2(\mathbb{R}^n; \mathbb{R}^m)). \end{cases}$$

2. Choose a function $\mathbf{v} \in C^1([0, T]; H^1(\mathbb{R}^n; \mathbb{R}^m))$. Then from (11) we deduce

$$(29) \quad \int_0^T (\mathbf{u}'^{\epsilon}, \mathbf{v}) + \epsilon D\mathbf{u}^{\epsilon} : D\mathbf{v} + B[\mathbf{u}^{\epsilon}, \mathbf{v}; t] dt = \int_0^T (\mathbf{f}, \mathbf{v}) dt.$$

Let $\epsilon = \epsilon_k \rightarrow 0$:

$$(30) \quad \int_0^T (\mathbf{u}', \mathbf{v}) + B[\mathbf{u}, \mathbf{v}; t] dt = \int_0^T (\mathbf{f}, \mathbf{v}) dt.$$

This identity is valid for all $\mathbf{v} \in C([0, T]; H^1(\mathbb{R}^n; \mathbb{R}^m))$, and so

$$(\mathbf{u}', \mathbf{v}) + B[\mathbf{u}, \mathbf{v}; t] = (\mathbf{f}, \mathbf{v})$$

for a.e. t and each $\mathbf{v} \in H^1(\mathbb{R}^n; \mathbb{R}^m)$.

3. Assume now $\mathbf{v}(T) = 0$. Then (29) implies

$$\int_0^T -(\mathbf{u}^{\epsilon}, \mathbf{v}') + \epsilon D\mathbf{u}^{\epsilon} : D\mathbf{v} + B[\mathbf{u}^{\epsilon}, \mathbf{v}; t] dt = \int_0^T (\mathbf{f}, \mathbf{v}) dt + (\mathbf{g}^{\epsilon}, \mathbf{v}(0)).$$

Upon sending $\epsilon = \epsilon_k \rightarrow 0$, we obtain

$$\int_0^T -(\mathbf{u}, \mathbf{v}') + B[\mathbf{u}, \mathbf{v}; t] dt = \int_0^T (\mathbf{f}, \mathbf{v}) dt + (\mathbf{g}, \mathbf{v}(0)).$$

Integrating by parts in (30) gives us the identity

$$\int_0^T -(\mathbf{u}, \mathbf{v}') + B[\mathbf{u}, \mathbf{v}; t] dt = \int_0^T (\mathbf{f}, \mathbf{v}) dt + (\mathbf{u}(0), \mathbf{v}(0)).$$

Consequently $\mathbf{u}(0) = \mathbf{g}$, as $\mathbf{v}(0)$ is arbitrary. □

THEOREM 4 (Uniqueness of weak solution). *A weak solution of (5) is unique.*

Proof. It suffices to show the only weak solution of (5) with $\mathbf{f} \equiv \mathbf{g} \equiv 0$ is $\mathbf{u} \equiv 0$.

To verify this, note

$$(31) \quad (\mathbf{u}', \mathbf{u}) + B[\mathbf{u}, \mathbf{u}; t] = 0 \quad \text{for a.e. } 0 \leq t \leq T.$$

Since $|B[\mathbf{u}, \mathbf{u}; t]| \leq C\|\mathbf{u}\|_{L^2(\mathbb{R}^n; \mathbb{R}^m)}^2$, we as usual compute from (31) that

$$\frac{d}{dt} (\|\mathbf{u}(t)\|_{L^2(\mathbb{R}^n; \mathbb{R}^m)}^2) \leq C\|\mathbf{u}(t)\|_{L^2(\mathbb{R}^n; \mathbb{R}^m)}^2,$$

whence Gronwall's inequality forces $\|\mathbf{u}(t)\|_{L^2(\mathbb{R}^n; \mathbb{R}^m)}^2 = 0$ ($0 \leq t \leq T$), since $\mathbf{u}(0) = 0$. \square

7.3.3. Systems with constant coefficients.

In this subsection we apply the Fourier transform (§4.3) to solve the *constant coefficient* system

$$(32) \quad \mathbf{u}_t + \sum_{j=1}^n B_j \mathbf{u}_{x_j} = \mathbf{0} \quad \text{in } \mathbb{R}^n \times (0, \infty),$$

with the initial condition

$$(33) \quad \mathbf{u} = \mathbf{g} \quad \text{on } \mathbb{R}^n \times \{t = 0\}.$$

We assume that the $\{B_j\}_{j=1}^n$ are constant $m \times m$ matrices and that the $m \times m$ matrix

$$(34) \quad \mathbf{B}(y) := \sum_{j=1}^n y_j B_j$$

has for each $y \in \mathbb{R}^n$ m real eigenvalues

$$(35) \quad \lambda_1(y) \leq \lambda_2(y) \leq \cdots \leq \lambda_m(y).$$

There is no hypothesis concerning the eigenvectors, and so we are supposing only a very weak sort of hyperbolicity here. We also make no assumption of symmetry for the matrices $\{B_j\}_{j=1}^n$. Consequently the foregoing energy estimates do not apply. We need a new tool, which we discover in the Fourier transform.

THEOREM 5 (Existence of solution). *Assume*

$$\mathbf{g} \in H^s(\mathbb{R}^n; \mathbb{R}^m) \quad \left(s > \frac{n}{2} + m \right).$$

Then there is a unique solution $\mathbf{u} \in C^1([0, \infty); \mathbb{R}^m)$ of the initial-value problem (32), (33).

See §5.8.4 for the definition of the fractional Sobolev spaces H^s .

Proof. 1. We apply the Fourier transform (§4.3.1), as follows. First, temporarily assume $\mathbf{u} = (u^1, \dots, u^m)$ is a smooth solution. Then set

$$\hat{\mathbf{u}} = (\hat{u}^1, \dots, \hat{u}^m),$$

where $\hat{\cdot}$ denotes the Fourier transform in the variable x : we do not transform with respect to the time variable t . Equation (32) becomes

$$\hat{\mathbf{u}}_t + i \sum_{j=1}^n y_j B_j \hat{\mathbf{u}} = \mathbf{0};$$

that is,

$$(36) \quad \hat{\mathbf{u}}_t + i\mathbf{B}(y)\hat{\mathbf{u}} = \mathbf{0} \quad \text{in } \mathbb{R}^n \times (0, \infty).$$

In addition

$$(37) \quad \hat{\mathbf{u}} = \hat{\mathbf{g}} \quad \text{on } \mathbb{R}^n \times \{t = 0\}.$$

For each fixed $y \in \mathbb{R}^n$ we solve (36), (37) by integrating in time, to find

$$(38) \quad \hat{\mathbf{u}}(y, t) = e^{-it\mathbf{B}(y)} \hat{\mathbf{g}}(y) \quad (y \in \mathbb{R}^n, t \geq 0).$$

Consequently $\mathbf{u} = (e^{-it\mathbf{B}(y)} \hat{\mathbf{g}})^\vee$, so that

$$(39) \quad \mathbf{u}(x, t) = \frac{1}{(2\pi)^{n/2}} \int_{\mathbb{R}^n} e^{ix \cdot y} e^{-it\mathbf{B}(y)} \hat{\mathbf{g}}(y) dy \quad (x \in \mathbb{R}^n, t \geq 0).$$

2. We have derived formula (39) assuming \mathbf{u} to be a smooth solution of (32), (33). We now verify that the function \mathbf{u} defined by (39) is in truth a solution, and so must first check that the integral in (39) converges.

Since $\mathbf{g} \in H^s(\mathbb{R}^n; \mathbb{R}^m)$, we know according to §5.8.4 that there exists $\mathbf{f} \in L^2(\mathbb{R}^n; \mathbb{R}^m)$ such that

$$(40) \quad |\hat{\mathbf{g}}(y)| \leq C(1 + |y|^s)^{-1} |\mathbf{f}(y)| \quad (y \in \mathbb{R}^n).$$

So in order to investigate the convergence of the integral (39), we must estimate $\|e^{-it\mathbf{B}(y)}\|$.

3. For a fixed y , let Γ denote the path $\partial B(0, r)$ in the complex plane, traversed counterclockwise, the radius r selected so large that the eigenvalues $\lambda_1(y), \dots, \lambda_m(y)$ lie within Γ .

We have the formula

$$(41) \quad e^{-it\mathbf{B}(y)} = \frac{1}{2\pi i} \int_{\Gamma} e^{-itz} (zI - \mathbf{B}(y))^{-1} dz.$$

To verify this, let $\mathbf{A}(t, y)$ denote the right-hand side of (41) and fix $x \in \mathbb{R}^m$. Then

$$\begin{aligned} \mathbf{B}(y)\mathbf{A}(t, y)x &= \frac{1}{2\pi i} \int_{\Gamma} e^{-itz} \mathbf{B}(y)(zI - \mathbf{B}(y))^{-1} x dz \\ &= \frac{1}{2\pi i} \int_{\Gamma} e^{-itz} (z(zI - \mathbf{B}(y))^{-1} x - x) dz \\ &= -\frac{1}{i} \frac{d}{dt} \mathbf{A}(t, y)x, \end{aligned}$$

since $\int_{\Gamma} e^{-itz} dz = 0$. Consequently

$$(42) \quad \left(\frac{d}{dt} + i\mathbf{B}(y) \right) \mathbf{A}(t, y) = 0.$$

In addition

$$\begin{aligned} \mathbf{A}(0, y)x &= \frac{1}{2\pi i} \int_{\Gamma} (zI - \mathbf{B}(y))^{-1} x dz \\ (43) \quad &= \frac{1}{2\pi i} \int_{\Gamma} \frac{x + \mathbf{B}(y)(zI - \mathbf{B}(y))^{-1} x}{z} dz \\ &= x + \frac{1}{2\pi i} \int_{\Gamma} \frac{\mathbf{B}(y)(zI - \mathbf{B}(y))^{-1} x}{z} dz. \end{aligned}$$

Now set

$$(44) \quad w := (zI - \mathbf{B}(y))^{-1} x,$$

so that $zw - \mathbf{B}(y)w = x$. Taking the product with \bar{w} , we deduce $|w| \leq \frac{C}{|z|}$ for some constant C . Using this estimate and letting r go to infinity, we conclude from (43) that $\mathbf{A}(0, y)x = x$. This equality and (42) verify the representation formula (41).

4. Define a new path Δ in the complex plane as follows. For fixed y , draw circles $B_k = B(\lambda_k(y), 1)$ of radius 1, centered at $\lambda_k(y)$ ($k = 1, \dots, m$). Then take Δ to be the boundary of $\bigcup_{k=1}^m B_k$, traversed counterclockwise.

Deforming the path Γ into Δ , we deduce from (41) that

$$(45) \quad e^{-it\mathbf{B}(y)} = \frac{1}{2\pi i} \int_{\Delta} e^{-itz} (zI - \mathbf{B}(y))^{-1} dz.$$

Now

$$(46) \quad |e^{-itz}| \leq e^t \quad (z \in \Delta).$$

Furthermore

$$\det(zI - \mathbf{B}(y)) = \prod_{k=1}^m (z - \lambda_k(y)),$$

whence

$$(47) \quad |\det(zI - \mathbf{B}(y))| \geq 1 \quad \text{if } z \in \Delta.$$

Now

$$(zI - \mathbf{B}(y))^{-1} = \frac{\text{cof}(zI - \mathbf{B}(y))^T}{\det(zI - \mathbf{B}(y))},$$

where “cof” denotes the cofactor matrix (see §8.1.4). We deduce

$$(48) \quad \begin{aligned} \|(zI - \mathbf{B}(y))^{-1}\| &\leq \|\text{cof}(zI - \mathbf{B}(y))\| \\ &\leq C(1 + |z|^{m-1} + \|\mathbf{B}(y)\|^{m-1}) \\ &\leq C(1 + |y|^{m-1}) \quad \text{if } z \in \Delta. \end{aligned}$$

We have utilized in this calculation the elementary inequality

$$|\lambda_k(y)| \leq C|y| \quad (k = 1, \dots, m).$$

Combining (45)–(48), we derive the estimate

$$(49) \quad \|e^{-it\mathbf{B}(y)}\| \leq Ce^t(1 + |y|^{m-1}) \quad (y \in \mathbb{R}^n).$$

5. Return now to (37). We deduce using (40), (49) that

$$\begin{aligned} \int_{\mathbb{R}^n} |e^{ix \cdot y} e^{-it\mathbf{B}(y)} \hat{\mathbf{g}}(y)| dy &\leq C \int_{\mathbb{R}^n} \|e^{-it\mathbf{B}(y)}\| (1 + |y|^s)^{-1} |\mathbf{f}(y)| dy \\ &\leq Ce^t \int_{\mathbb{R}^n} |\mathbf{f}(y)| (1 + |y|^{m-1}) (1 + |y|^s)^{-1} dy \\ &\leq C \left(\int_{\mathbb{R}^n} |\mathbf{f}|^2 dy \right)^{1/2} \left(\int_{\mathbb{R}^n} \frac{dy}{1 + |y|^{2(s-m+1)}} \right)^{1/2} \\ &< \infty, \end{aligned}$$

since $s > \frac{n}{2} + m - 1$. Hence the integral in (39) converges, and it follows easily that the function

$$\mathbf{u}(x, t) = \frac{1}{(2\pi)^{n/2}} \int_{\mathbb{R}^n} e^{ix \cdot y} e^{-it\mathbf{B}(y)} \hat{\mathbf{g}}(y) dy$$

is continuous on $\mathbb{R}^n \times [0, \infty)$.

6. To show \mathbf{u} is C^1 , observe for $0 < |h| \leq 1$ that

$$\frac{\mathbf{u}(x, t+h) - \mathbf{u}(x, t)}{h} = \frac{1}{(2\pi)^{n/2}h} \int_{\mathbb{R}^n} e^{ix \cdot y} (e^{-i(t+h)\mathbf{B}(y)} - e^{-it\mathbf{B}(y)}) \hat{\mathbf{g}}(y) dy.$$

Since

$$e^{-i(t+h)\mathbf{B}(y)} - e^{-it\mathbf{B}(y)} = -i \int_t^{t+h} \mathbf{B}(y) e^{-is\mathbf{B}(y)} ds,$$

we can estimate as above that

$$\left| \frac{1}{h} (e^{-i(t+h)\mathbf{B}(y)} - e^{-it\mathbf{B}(y)}) \right| \leq C e^{t+1} (1 + |y|^m).$$

Therefore

$$\left| \frac{\mathbf{u}(x, t+h) - \mathbf{u}(x, t)}{h} \right| \leq C e^{t+1} \int_{\mathbb{R}^n} |\mathbf{f}(y)| (1 + |y|^m) (1 + |y|^s)^{-1} dy,$$

and the integrand is summable since $s > \frac{n}{2} + m$. Thus \mathbf{u}_t exists and is continuous on $\mathbb{R}^n \times [0, \infty)$. A similar argument shows \mathbf{u}_{x_i} exists and is continuous ($i = 1, \dots, n$). According to the Dominated Convergence Theorem, we can furthermore differentiate under the integral sign in (39), to confirm that \mathbf{u} solves the system $\mathbf{u}_t + \sum_{j=1}^n B_j \mathbf{u}_{x_j} = \mathbf{0}$. \square

In Chapter 11 we will encounter nonlinear first-order systems of hyperbolic equations.

7.4. SEMIGROUP THEORY

Semigroup theory is the abstract study of first-order ordinary differential equations with values in Banach spaces, driven by linear, but possibly unbounded, operators. In this section we outline the basics of the theory and present as well two applications to linear PDE. This approach provides an elegant alternative to some of the existence theory for evolution equations set forth in §§7.1–7.3.

7.4.1. Definitions, elementary properties.

We begin in an abstract setting. Let X denote a real Banach space, and consider then the ordinary differential equation

$$(1) \quad \begin{cases} \mathbf{u}'(t) = A\mathbf{u}(t) & (t \geq 0) \\ \mathbf{u}(0) = u, \end{cases}$$

where $' = \frac{d}{dt}$, $u \in X$ is given, and A is a linear operator. More precisely, suppose $D(A)$, the *domain* of A , is a linear subspace of X and we are given a possibly unbounded linear operator

$$(2) \quad A : D(A) \rightarrow X.$$

We investigate the existence and uniqueness of a solution

$$\mathbf{u} : [0, \infty) \rightarrow X$$

of the ODE (1). The key problem is to ascertain reasonable conditions on the operator A so that (a) the ODE has a unique solution \mathbf{u} for each initial point $u \in X$ and (b) many interesting PDE can be cast into the abstract form (1). (We have in mind the situation that X is an L^p space of functions and A is a linear partial differential operator involving variables other than t . In this case A is necessarily an unbounded operator.)

a. Semigroups. Let us for the moment informally assume $\mathbf{u} : [0, \infty) \rightarrow X$ is a solution of the differential equation (1) and that (1) in fact has a unique solution for each initial point $u \in X$.

NOTATION. We will write

$$(3) \quad \mathbf{u}(t) := S(t)u \quad (t \geq 0)$$

to display explicitly the dependence of $\mathbf{u}(t)$ on the initial value $u \in X$. For each time $t \geq 0$ we may therefore regard $S(t)$ as a mapping from X into X .

What properties does the family of operators $\{S(t)\}_{t \geq 0}$ satisfy? Clearly $S(t) : X \rightarrow X$ is linear. Furthermore

$$(4) \quad S(0)u = u \quad (u \in X)$$

and

$$(5) \quad S(t+s)u = S(t)S(s)u = S(s)S(t)u \quad (t, s \geq 0, u \in X).$$

Condition (5) is simply our assumption that the ODE (1) has a unique solution for each initial point. Finally, it seems reasonable to suppose that for each $u \in X$

$$(6) \quad \text{the mapping } t \mapsto S(t)u \text{ is continuous from } [0, \infty) \text{ into } X.$$

DEFINITIONS. (i) A family $\{S(t)\}_{t \geq 0}$ of bounded linear operators mapping X into X is called a semigroup if conditions (4)–(6) are satisfied.

(ii) We say $\{S(t)\}_{t \geq 0}$ is a contraction semigroup if in addition

$$(7) \quad \|S(t)\| \leq 1 \quad (t \geq 0),$$

$\| \cdot \|$ here denoting the operator norm. Thus

$$\|S(t)u\| \leq \|u\| \quad (t \geq 0, u \in X).$$

The notion of contraction semigroup captures many properties of a nice flow on X generated by the ODE (1).

b. Elementary properties, generators. The real problem now is to determine which operators A generate contraction semigroups. We will answer this in §7.4.2, after recording in this section some further general facts.

Henceforth assume $\{S(t)\}_{t \geq 0}$ is a contraction semigroup on X .

DEFINITIONS. Write

$$(8) \quad D(A) := \left\{ u \in X \mid \lim_{t \rightarrow 0^+} \frac{S(t)u - u}{t} \text{ exists in } X \right\}$$

and

$$(9) \quad Au := \lim_{t \rightarrow 0^+} \frac{S(t)u - u}{t} \quad (u \in D(A)).$$

We call $A : D(A) \rightarrow X$ the (infinitesimal) generator of the semigroup $\{S(t)\}_{t \geq 0}$; $D(A)$ is the domain of A .

THEOREM 1 (Differential properties of semigroups). Assume $u \in D(A)$. Then

- (i) $S(t)u \in D(A)$ for each $t \geq 0$.
- (ii) $AS(t)u = S(t)Au$ for each $t \geq 0$.
- (iii) The mapping $t \mapsto S(t)u$ is differentiable for each $t > 0$.
- (iv) $\frac{d}{dt}S(t)u = AS(t)u \quad (t > 0)$.

Proof. 1. Let $u \in D(A)$. Then

$$\begin{aligned} & \lim_{s \rightarrow 0^+} \frac{S(s)S(t)u - S(t)u}{s} \\ &= \lim_{s \rightarrow 0^+} \frac{S(t)S(s)u - S(t)u}{s} \quad \text{by the semigroup property (5)} \\ &= S(t) \lim_{s \rightarrow 0^+} \frac{S(s)u - u}{s} = S(t)Au. \end{aligned}$$

Thus $S(t)u \in D(A)$ and $AS(t)u = S(t)Au$. Assertions (i) and (ii) are proved.

2. Let $u \in D(A)$, $h > 0$. Then if $t > 0$,

$$\begin{aligned} & \lim_{h \rightarrow 0^+} \left\{ \frac{S(t)u - S(t-h)u}{h} - S(t)Au \right\} \\ &= \lim_{h \rightarrow 0^+} \left\{ S(t-h) \left(\frac{S(h)u - u}{h} \right) - S(t)Au \right\} \\ &= \lim_{h \rightarrow 0^+} \left\{ S(t-h) \left(\frac{S(h)u - u}{h} - Au \right) + (S(t-h) - S(t))Au \right\} = 0, \end{aligned}$$

since $\frac{S(h)u - u}{h} \rightarrow Au$ and $\|S(t-h)\| \leq 1$. Consequently

$$\lim_{h \rightarrow 0^+} \frac{S(t)u - S(t-h)u}{h} = S(t)Au.$$

Similarly

$$\lim_{h \rightarrow 0^+} \frac{S(t+h)u - S(t)u}{h} = S(t) \lim_{h \rightarrow 0^+} \frac{S(h)u - u}{h} = S(t)Au.$$

Thus $\frac{d}{dt}S(t)u$ exists for each time $t > 0$ and equals $S(t)Au = AS(t)u$. \square

Remark. Since $t \mapsto AS(t)u = S(t)Au$ is continuous, the mapping $t \mapsto S(t)u$ is C^1 in $(0, \infty)$, if $u \in D(A)$.

THEOREM 2 (Properties of generators).

(i) The domain $D(A)$ is dense in X

and

(ii) A is a closed operator.

Remark. To say A is closed means that whenever $u_k \in D(A)$ ($k = 1, \dots$) and $u_k \rightarrow u$, $Au_k \rightarrow v$ as $k \rightarrow \infty$, then

$$u \in D(A), \quad v = Au.$$

Proof. 1. Fix any $u \in X$ and define then $u^t := \int_0^t S(s)u ds$. In view of (6), $\frac{u^t}{t} \rightarrow u$ in X , as $t \rightarrow 0^+$.

2. We claim

$$(10) \quad u^t \in D(A) \quad (t > 0).$$

Indeed if $r > 0$, we have

$$\begin{aligned} \frac{S(r)u^t - u^t}{r} &= \frac{1}{r} \left[S(r) \left(\int_0^t S(s)u \, ds \right) - \left(\int_0^t S(s)u \, ds \right) \right] \\ &= \frac{1}{r} \int_0^t S(r+s)u - S(s)u \, ds, \end{aligned}$$

where we used the semigroup property (5). Thus

$$\begin{aligned} \frac{S(r)u^t - u^t}{r} &= \frac{1}{r} \int_t^{t+r} S(s)u \, ds - \frac{1}{r} \int_0^r S(s)u \, ds \\ &\rightarrow S(t)u - u, \quad \text{as } r \rightarrow 0+. \end{aligned}$$

Hence $u^t \in D(A)$, with $Au^t = S(t)u - u$. This proves (10) and completes the proof of assertion (i).

3. To prove A is closed, let $u_k \in D(A)$ ($k = 1, \dots$) and suppose

$$(11) \quad u_k \rightarrow u, \quad Au_k \rightarrow v \quad \text{in } X.$$

We must prove $u \in D(A)$, $v = Au$. According to Theorem 1

$$S(t)u_k - u_k = \int_0^t S(s)Au_k \, ds.$$

Let $k \rightarrow \infty$ and recall (11):

$$S(t)u - u = \int_0^t S(s)v \, ds.$$

Hence we have

$$\lim_{t \rightarrow 0+} \frac{S(t)u - u}{t} = \lim_{t \rightarrow 0+} \frac{1}{t} \int_0^t S(s)v \, ds = v.$$

But then by definition $u \in D(A)$, $v = Au$. □

c. Resolvents. Let A be a closed linear operator on X , with domain $D(A)$.

DEFINITIONS. (i) We say a real number λ belongs to $\rho(A)$, the resolvent set of A , provided the operator

$$\lambda I - A : D(A) \rightarrow X$$

is one-to-one and onto.

(ii) If $\lambda \in \rho(A)$, the resolvent operator $R_\lambda : X \rightarrow X$ is defined by

$$R_\lambda u := (\lambda I - A)^{-1}u.$$

According to the Closed Graph Theorem (§D.3), $R_\lambda : X \rightarrow D(A) \subseteq X$ is a bounded linear operator. Furthermore,

$$AR_\lambda u = R_\lambda Au \quad \text{if } u \in D(A).$$

THEOREM 3 (Properties of resolvent operators).

(i) If $\lambda, \mu \in \rho(A)$, we have

$$(12) \quad R_\lambda - R_\mu = (\mu - \lambda)R_\lambda R_\mu \quad (\text{resolvent identity})$$

and

$$(13) \quad R_\lambda R_\mu = R_\mu R_\lambda.$$

(ii) If $\lambda > 0$, then $\lambda \in \rho(A)$,

$$(14) \quad R_\lambda u = \int_0^\infty e^{-\lambda t} S(t)u \, dt \quad (u \in X),$$

and so $\|R_\lambda\| \leq \frac{1}{\lambda}$.

Thus the resolvent operator is the Laplace transform of the semigroup (cf. Example 8 in §4.3.3).

Proof. 1. Verification of the identities (12), (13) is left to the reader (Problem 12).

2. Note first that since $\lambda > 0$ and $\|S(t)\| \leq 1$, the integral on the right-hand side of (14) is defined. Let $\tilde{R}_\lambda u$ denote this integral. Then for $h > 0$ and $u \in X$,

$$\begin{aligned} \frac{S(h)\tilde{R}_\lambda u - \tilde{R}_\lambda u}{h} &= \frac{1}{h} \left\{ \int_0^\infty e^{-\lambda t} [S(t+h)u - S(t)u] \, dt \right\} \\ &= -\frac{1}{h} \int_0^h e^{-\lambda(t-h)} S(t)u \, dt \\ &\quad + \frac{1}{h} \int_0^\infty (e^{-\lambda(t-h)} - e^{-\lambda t}) S(t)u \, dt \\ &= -e^{\lambda h} \frac{1}{h} \int_0^h e^{-\lambda t} S(t)u \, dt \\ &\quad + \left(\frac{e^{\lambda h} - 1}{h} \right) \int_0^\infty e^{-\lambda t} S(t)u \, dt. \end{aligned}$$

Hence

$$\lim_{h \rightarrow 0^+} \frac{S(h)\tilde{R}_\lambda u - \tilde{R}_\lambda u}{h} = -u + \lambda \tilde{R}_\lambda u.$$

Thus $A\tilde{R}_\lambda u = -u + \lambda \tilde{R}_\lambda u$; that is,

$$(15) \quad (\lambda I - A)\tilde{R}_\lambda u = u \quad (u \in X).$$

On the other hand if $u \in D(A)$,

$$(16) \quad \begin{aligned} A\tilde{R}_\lambda u &= A \int_0^\infty e^{-\lambda t} S(t)u \, dt = \int_0^\infty e^{-\lambda t} AS(t)u \, dt \\ &= \int_0^\infty e^{-\lambda t} S(t)Au \, dt = \tilde{R}_\lambda Au. \end{aligned}$$

Our passing A under the integral sign is justified since A is a closed operator: see Problem 13. Thus

$$\tilde{R}_\lambda(\lambda I - A)u = u \quad (u \in D(A)).$$

In view of (15) and the formula above $\lambda I - A$ is one-to-one and onto. Consequently $\lambda \in \rho(A)$, $\tilde{R}_\lambda = (\lambda I - A)^{-1} = R_\lambda$. \square

7.4.2. Generating contraction semigroups.

We now characterize the generators of contraction semigroups:

THEOREM 4 (Hille–Yosida Theorem). *Let A be a closed, densely-defined linear operator on X . Then A is the generator of a contraction semigroup $\{S(t)\}_{t \geq 0}$ if and only if*

$$(17) \quad (0, \infty) \subset \rho(A) \quad \text{and} \quad \|R_\lambda\| \leq \frac{1}{\lambda} \quad \text{for } \lambda > 0.$$

Proof. 1. If A is a generator, then from Theorem 3(ii) we immediately deduce (17).

2. Conversely, suppose A is closed, densely-defined, and satisfies (17). We must build a contraction semigroup with A as its generator. For this, fix $\lambda > 0$ and define

$$(18) \quad A_\lambda := -\lambda I + \lambda^2 R_\lambda = \lambda A R_\lambda.$$

The operator A_λ is a kind of regularized approximation to A .

3. We first claim

$$(19) \quad A_\lambda u \rightarrow Au \quad \text{as } \lambda \rightarrow \infty \quad (u \in D(A)).$$

Indeed, since $\lambda R_\lambda u - u = A R_\lambda u = R_\lambda A u$, $\|\lambda R_\lambda u - u\| \leq \|R_\lambda\| \|A u\| \leq \frac{1}{\lambda} \|A u\| \rightarrow 0$. Thus $\lambda R_\lambda u \rightarrow u$ as $\lambda \rightarrow \infty$ if $u \in D(A)$. But since $\|\lambda R_\lambda\| \leq 1$ and $D(A)$ is dense, we deduce then as well

$$(20) \quad \lambda R_\lambda u \rightarrow u \quad \text{as } \lambda \rightarrow \infty, \quad \text{for all } u \in X.$$

Now if $u \in D(A)$, then

$$A_\lambda u = \lambda A R_\lambda u = \lambda R_\lambda A u.$$

In view of (20), our claim (19) is proved.

4. Next, define

$$S_\lambda(t) := e^{tA_\lambda} = e^{-\lambda t} e^{\lambda^2 t R_\lambda} = e^{-\lambda t} \sum_{k=0}^{\infty} \frac{(\lambda^2 t)^k}{k!} R_\lambda^k.$$

Observe that since $\|R_\lambda\| \leq \lambda^{-1}$,

$$\|S_\lambda(t)\| \leq e^{-\lambda t} \sum_{k=0}^{\infty} \frac{\lambda^{2k} t^k}{k!} \|R_\lambda\|^k \leq e^{-\lambda t} \sum_{k=0}^{\infty} \frac{\lambda^k t^k}{k!} = 1.$$

Consequently $\{S_\lambda(t)\}_{t \geq 0}$ is a contraction semigroup, and it is easy to check that its generator is A_λ , with $D(A_\lambda) = X$.

5. Let $\lambda, \mu > 0$. Since $R_\lambda R_\mu = R_\mu R_\lambda$, we see $A_\lambda A_\mu = A_\mu A_\lambda$, and so

$$A_\mu S_\lambda(t) = S_\lambda(t) A_\mu \quad \text{for each } t > 0.$$

Thus we can compute

$$\begin{aligned} S_\lambda(t)u - S_\mu(t)u &= \int_0^t \frac{d}{ds} [S_\mu(t-s)S_\lambda(s)u] ds \\ &= \int_0^t S_\mu(t-s)S_\lambda(s)(A_\lambda u - A_\mu u) ds, \end{aligned}$$

because $\frac{d}{dt} S_\lambda(t)u = A_\lambda S_\lambda(t)u = S_\lambda(t)A_\lambda u$. Consequently (19) implies that if $u \in D(A)$, then $\|S_\lambda(t)u - S_\mu(t)u\| \leq t\|A_\lambda u - A_\mu u\| \rightarrow 0$ as $\lambda, \mu \rightarrow \infty$. Hence

$$(21) \quad S(t)u := \lim_{\lambda \rightarrow \infty} S_\lambda(t)u \quad \text{exists for each } t \geq 0, u \in D(A).$$

As $\|S_\lambda(t)\| \leq 1$, the limit (21) in fact exists for all $u \in X$, uniformly for t in compact subsets of $[0, \infty)$. It is now straightforward to verify $\{S(t)\}_{t \geq 0}$ is a contraction semigroup on X .

6. It remains to show A is the generator of $\{S(t)\}_{t \geq 0}$. Write B to denote this generator. Now

$$(22) \quad S_\lambda(t)u - u = \int_0^t S_\lambda(s)A_\lambda u ds.$$

In addition

$$\|S_\lambda(s)A_\lambda u - S(s)Au\| \leq \|S_\lambda(s)\| \|A_\lambda u - Au\| + \|(S_\lambda(s) - S(s))Au\| \rightarrow 0$$

as $\lambda \rightarrow \infty$, if $u \in D(A)$. Passing therefore to limits in (22), we deduce

$$S(t)u - u = \int_0^t S(s)Au \, ds$$

if $u \in D(A)$. Thus $D(A) \subseteq D(B)$ and

$$Bu = \lim_{t \rightarrow 0^+} \frac{S(t)u - u}{t} = Au \quad (u \in D(A)).$$

Now if $\lambda > 0$, $\lambda \in \rho(A) \cap \rho(B)$. Also $(\lambda I - B)(D(A)) = (\lambda I - A)(D(A)) = X$, according to (17). Hence $(\lambda I - B)|_{D(A)}$ is one-to-one and onto, whence $D(A) = D(B)$. Therefore $A = B$, and so A is indeed the generator of $\{S(t)\}_{t \geq 0}$. \square

Remark. Let $\omega \in \mathbb{R}$. A semigroup $\{S(t)\}_{t \geq 0}$ is called ω -contractive if $\|S(t)\| \leq e^{\omega t}$ ($t \geq 0$). An easy variant of Theorem 4 asserts that a closed, densely defined linear operator A generates an ω -contractive semigroup if and only if

$$(23) \quad (\omega, \infty) \subset \rho(A) \quad \text{and} \quad \|R_\lambda\| \leq \frac{1}{\lambda - \omega} \quad \text{for all } \lambda > \omega.$$

This version of the Hille–Yosida Theorem will be required for our first example below.

7.4.3. Applications.

We demonstrate in this section that certain second-order parabolic and hyperbolic PDE can be realized within the semigroup framework.

a. Second-order parabolic PDE. We consider the initial/boundary-value problem

$$(24) \quad \begin{cases} u_t + Lu = 0 & \text{in } U_T \\ u = 0 & \text{on } \partial U \times [0, T] \\ u = g & \text{on } U \times \{t = 0\}, \end{cases}$$

a special case (corresponding to $f \equiv 0$) of (1) in §7.1.1. We assume L has the divergence structure (2) from §7.1.1, satisfies the usual strong ellipticity condition, and has smooth coefficients, *which do not depend on t* . We additionally suppose that the bounded open set U has a smooth boundary.

We propose to reinterpret (24) as the flow determined by a semigroup on $X = L^2(U)$. For this, we set

$$(25) \quad D(A) := H_0^1(U) \cap H^2(U)$$

and define

$$(26) \quad Au := -Lu \quad \text{if } u \in D(A).$$

Clearly then A is an unbounded linear operator on X . Recall from §6.2.2 the energy estimate

$$(27) \quad \beta \|u\|_{H_0^1(U)}^2 \leq B[u, u] + \gamma \|u\|_{L^2(U)}^2,$$

for constants $\beta > 0$, $\gamma \geq 0$, where $B[\ , \]$ is the bilinear form associated with L .

THEOREM 5 (Second-order parabolic PDE as semigroups). *The operator A generates a γ -contraction semigroup $\{S(t)\}_{t \geq 0}$ on $L^2(U)$.*

Proof. 1. We must verify the hypotheses of the variant of the Hille–Yosida Theorem mentioned in the concluding Remark in §7.4.2, with γ replacing ω .

First, $D(A)$ given by (25) is clearly dense in $L^2(U)$.

2. We claim now that the operator A is closed. Indeed, let $\{u_k\}_{k=1}^\infty \subset D(A)$ with

$$(28) \quad u_k \rightarrow u, \quad Au_k \rightarrow f \quad \text{in } L^2(U).$$

According to the regularity Theorem 4 in §6.3.2,

$$\|u_k - u_l\|_{H^2(U)} \leq C(\|Au_k - Au_l\|_{L^2(U)} + \|u_k - u_l\|_{L^2(U)})$$

for all k and l . Thus (28) implies $\{u_k\}_{k=1}^\infty$ is a Cauchy sequence in $H^2(U)$ and so

$$(29) \quad u_k \rightarrow u \quad \text{in } H^2(U).$$

Therefore $u \in D(A)$. Furthermore (29) implies $Au_k \rightarrow Au$ in $L^2(U)$, and consequently $f = Au$.

3. Next we check the resolvent conditions (23), with γ replacing ω . According to Theorem 3 in §6.2.2, for each $\lambda \geq \gamma$ the boundary-value problem

$$(30) \quad \begin{cases} Lu + \lambda u = f & \text{in } U \\ u = 0 & \text{in } \partial U \end{cases}$$

has a unique weak solution $u \in H_0^1(U)$ for each $f \in L^2(U)$. Owing to elliptic regularity theory, in fact $u \in H^2(U) \cap H_0^1(U)$. Thus $u \in D(A)$. We may now rewrite (30), using (26), and find

$$(31) \quad \lambda u - Au = f.$$

Thus $(\lambda I - A) : D(A) \rightarrow X$ is one-to-one and onto, provided $\lambda \geq \gamma$. Hence $\rho(A) \supset [\gamma, \infty)$.

4. Consider the weak form of the boundary-value problem (30):

$$B[u, v] + \lambda(u, v) = (f, v)$$

for each $v \in H_0^1(U)$, where (\cdot, \cdot) is the inner product in $L^2(U)$. Set $v = u$ and recall (27) to compute for $\lambda > \gamma$

$$(\lambda - \gamma)\|u\|_{L^2(U)}^2 \leq \|f\|_{L^2(U)}\|u\|_{L^2(U)}.$$

Hence, since $u = R_\lambda f$, we have the estimate

$$\|R_\lambda f\|_{L^2(U)} \leq \frac{1}{\lambda - \gamma} \|f\|_{L^2(U)}.$$

This bound is valid for all $f \in L^2(U)$ and so

$$(32) \quad \|R_\lambda\| \leq \frac{1}{\lambda - \gamma} \quad (\lambda > \gamma),$$

as required. □

Semigroup theory provides an elegant method for constructing a solution to the initial/boundary-value problem (24). It is worth noting however that this technique requires that the coefficients a^{ij}, b^i, c ($i, j = 1, \dots, n$) of L be independent of t . The Galerkin method in §7.1 works without this restriction. On the other hand, semigroup theory constructs at the outset a more regular solution than the weak solution produced by the Galerkin technique. But we can also apply the regularity theory in §7.1.3 to demonstrate that this weak solution is in fact more regular.

b. Second-order hyperbolic PDE. We turn our attention next to the hyperbolic initial/boundary-value problem

$$(33) \quad \begin{cases} u_{tt} + Lu = 0 & \text{in } U_T \\ u = 0 & \text{on } \partial U \times [0, T] \\ u = g, u_t = h & \text{on } U \times \{t = 0\}, \end{cases}$$

for the operator L and open set U as above. We recast (33) as a first-order system by setting $v := u_t$. Then the foregoing reads

$$\begin{cases} u_t = v, & v_t + Lu = 0 & \text{in } U_T \\ & u = 0 & \text{on } \partial U \times [0, T] \\ & u = g, v = h & \text{on } U \times \{t = 0\}. \end{cases}$$

We will further assume L has the symmetric form

$$Lu = - \sum_{i,j=1}^n (a^{ij} u_{x_i})_{x_j} + cu,$$

where

$$(34) \quad c \geq 0, \quad a^{ij} = a^{ji} \quad (i, j = 1, \dots, n).$$

Thus for some constant $\beta > 0$

$$(35) \quad \beta \|u\|_{H_0^1(U)}^2 \leq B[u, u] \quad \text{for all } u \in H_0^1(U).$$

Now take

$$X = H_0^1(U) \times L^2(U),$$

with the norm

$$(36) \quad \|(u, v)\| := (B[u, u] + \|v\|_{L^2(U)}^2)^{1/2}.$$

Define

$$D(A) := [H^2(U) \cap H_0^1(U)] \times H_0^1(U)$$

and set

$$(37) \quad A(u, v) := (v, -Lu) \quad \text{for } (u, v) \in D(A).$$

We will show A verifies the hypothesis of the Hille–Yosida Theorem.

THEOREM 6 (Second-order hyperbolic PDE as semigroups). *The operator A generates a contraction semigroup $\{S(t)\}_{t \geq 0}$ on $H_0^1(U) \times L^2(U)$.*

Proof. 1. The domain of A is clearly dense in $H_0^1(U) \times L^2(U)$.

2. To see A is closed, let $\{(u_k, v_k)\}_{k=1}^\infty \subset D(A)$, with

$$(u_k, v_k) \rightarrow (u, v), \quad A(u_k, v_k) \rightarrow (f, g) \quad \text{in } H_0^1(U) \times L^2(U).$$

Since $A(u_k, v_k) = (v_k, -Lu_k)$, we conclude $f = v$ and $Lu_k \rightarrow -g$ in $L^2(U)$. As in the previous proof, it follows that $u_k \rightarrow u$ in $H^2(U)$ and $g = -Lu$. Thus $(u, v) \in D(A)$, $A(u, v) = (v, -Lu) = (f, g)$.

3. Now let $\lambda > 0$, $(f, g) \in X := H_0^1(U) \times L^2(U)$ and consider the operator equation

$$(38) \quad \lambda(u, v) - A(u, v) = (f, g).$$

This is equivalent to the two scalar equations

$$(39) \quad \begin{cases} \lambda u - v = f & (u \in H^2(U) \cap H_0^1(U)) \\ \lambda v + Lu = g & (v \in H_0^1(U)). \end{cases}$$

But (39) implies

$$(40) \quad \lambda^2 u + Lu = \lambda f + g \quad (u \in H^2(U) \cap H_0^1(U)).$$

Since $\lambda^2 > 0$, estimate (35) and the regularity theory imply there exists a unique solution u of (40). Defining then $v := \lambda u - f \in H_0^1(U)$, we have shown that (38) has a unique solution (u, v) . Consequently $\rho(A) \supset (0, \infty)$.

4. Whenever (39) holds, we write $(u, v) = R_\lambda(f, g)$. Now from the second equation in (39), we deduce

$$\lambda \|v\|_{L^2}^2 + B[u, v] = (g, v)_{L^2}.$$

Substituting $v = \lambda u - f$, we obtain

$$\begin{aligned} \lambda (\|v\|_{L^2}^2 + B[u, u]) &= (g, v)_{L^2} + B[u, f] \\ &\leq (\|g\|_{L^2}^2 + B[f, f])^{1/2} (\|v\|_{L^2}^2 + B[u, u])^{1/2}. \end{aligned}$$

Here we used the generalized Cauchy–Schwarz inequality (§B.2), which holds owing to the symmetry condition (34). In light of our definition (36),

$$\|(u, v)\| \leq \frac{1}{\lambda} \|(f, g)\|;$$

and so

$$\|R_\lambda\| \leq \frac{1}{\lambda} \quad (\lambda > 0),$$

as required. □

See Friedman [Fr1] or Yosida [Y] for the theory of *analytic semigroups*. Aspects of *nonlinear* semigroup theory will be developed later, in §9.6.

7.5. PROBLEMS

In the following exercises we assume $U \subset \mathbb{R}^n$ is an open, bounded set, with smooth boundary, and $T > 0$.

1. Prove there is at most one smooth solution of this initial/boundary-value problem for the heat equation with Neumann boundary conditions

$$\begin{cases} u_t - \Delta u = f & \text{in } U_T \\ \frac{\partial u}{\partial \nu} = 0 & \text{on } \partial U \times [0, T] \\ u = g & \text{on } U \times \{t = 0\}. \end{cases}$$

2. Assume u is a smooth solution of

$$\begin{cases} u_t - \Delta u = 0 & \text{in } U \times (0, \infty) \\ u = 0 & \text{on } \partial U \times [0, \infty) \\ u = g & \text{on } U \times \{t = 0\}. \end{cases}$$

Prove the exponential decay estimate:

$$\|u(\cdot, t)\|_{L^2(U)} \leq e^{-\lambda_1 t} \|g\|_{L^2(U)} \quad (t \geq 0),$$

where $\lambda_1 > 0$ is the principal eigenvalue of $-\Delta$ (with zero boundary conditions) on U .

3. (Adjoint dynamics) Suppose that u is a smooth solution of

$$\begin{cases} u_t + Lu = 0 & \text{in } U_T \\ u = 0 & \text{on } \partial U \times [0, T] \\ u = g & \text{on } U \times \{t = 0\}, \end{cases}$$

where L denotes a second-order elliptic operator, and that v is a smooth solution of the adjoint problem

$$\begin{cases} v_t - L^*v = 0 & \text{in } U_T \\ v = 0 & \text{on } \partial U \times [0, T] \\ v = h & \text{on } U \times \{t = T\}. \end{cases}$$

Show

$$\int_U g(x)v(x, 0) dx = \int_U u(x, T)h(x) dx.$$

4. (Galerkin's method for Poisson's equation) Suppose $f \in L^2(U)$ and assume that $u_m = \sum_{k=1}^m d_m^k w_k$ solves

$$\int_U Du_m \cdot Dw_k dx = \int_U f \cdot w_k dx$$

for $k = 1, \dots, m$. Show that a subsequence of $\{u_m\}_{m=1}^\infty$ converges weakly in $H_0^1(U)$ to the weak solution u of

$$\begin{cases} -\Delta u = f & \text{in } U \\ u = 0 & \text{on } \partial U. \end{cases}$$

5. Assume

$$\begin{cases} \mathbf{u}_k \rightharpoonup \mathbf{u} & \text{in } L^2(0, T; H_0^1(U)) \\ \mathbf{u}'_k \rightharpoonup \mathbf{v} & \text{in } L^2(0, T; H^{-1}(U)). \end{cases}$$

Prove that $\mathbf{v} = \mathbf{u}'$. (Hint: Let $\phi \in C_c^1(0, T)$, $w \in H_0^1(U)$. Then

$$\int_0^T \langle \mathbf{u}'_k, \phi w \rangle dt = - \int_0^T \langle \mathbf{u}_k, \phi' w \rangle dt.)$$

6. Suppose H is a Hilbert space and $\mathbf{u}_k \rightharpoonup \mathbf{u}$ in $L^2(0, T; H)$. Assume further we have the uniform bounds

$$\operatorname{ess\,sup}_{0 \leq t \leq T} \|\mathbf{u}_k(t)\| \leq C \quad (k = 1, \dots)$$

for some constant C . Prove

$$\operatorname{ess\,sup}_{0 \leq t \leq T} \|\mathbf{u}(t)\| \leq C.$$

(Hint: We have $\int_a^b (v, \mathbf{u}_k(t)) dt \leq C \|v\| |b - a|$ for $0 \leq a \leq b \leq T$ and $v \in H$.)

7. Suppose u is a smooth solution of

$$\begin{cases} u_t - \Delta u + cu = 0 & \text{in } U \times (0, \infty) \\ u = 0 & \text{on } \partial U \times [0, \infty) \\ u = g & \text{on } U \times \{t = 0\} \end{cases}$$

and the function c satisfies $c \geq \gamma > 0$.

Prove the exponential decay estimate

$$|u(x, t)| \leq C e^{-\gamma t} \quad ((x, t) \in U_T).$$

8. Assume that u is a smooth solution of the PDE from Problem 7, that $g \geq 0$, and that c is bounded (but *not* necessarily nonnegative). Show $u \geq 0$. (Hint: What PDE does $v := e^{-\lambda t} u$ solve?)

9. Prove inequality (54) in §7.1.3, (59) in §7.2.3.

(Hints: Assume u is smooth, $u = 0$ on ∂U . Transform the term $(Lu, -\Delta u)$ by integrating by parts twice, and then estimate the boundary terms. After changing variables locally and using cutoff functions, you may assume the boundary is flat. This problem is difficult.)

10. Show there exists at most one smooth solution of this initial/boundary-value problem for the *telegraph equation*

$$\begin{cases} u_{tt} + du_t - u_{xx} = f & \text{in } (0, 1) \times (0, T) \\ u = 0 & \text{on } (\{0\} \times [0, T]) \cup (\{1\} \times [0, T]) \\ u = g, u_t = h & \text{on } (0, 1) \times \{t = 0\}. \end{cases}$$

Here d is a constant.

11. Prove there exists at most one smooth solution u of this problem for the *beam equation*

$$\begin{cases} u_{tt} + u_{xxxx} = 0 & \text{in } (0, 1) \times (0, T) \\ u = u_x = 0 & \text{on } (\{0\} \times [0, T]) \cup (\{1\} \times [0, T]) \\ u = g, u_t = h & \text{on } [0, 1] \times \{t = 0\}. \end{cases}$$

12. Prove the resolvent identities (12) and (13) in §7.4.1.

13. Justify the equality

$$A \int_0^\infty e^{-\lambda t} S(t)u dt = \int_0^\infty e^{-\lambda t} AS(t)u dt$$

used in (16) of §7.4.1. (Hint: Approximate the integral by a Riemann sum and recall A is a closed operator.)

14. Define for $t > 0$

$$[S(t)g](x) = \int_{\mathbb{R}^n} \Phi(x - y, t)g(y) dy \quad (x \in \mathbb{R}^n),$$

where $g : \mathbb{R}^n \rightarrow \mathbb{R}$ and Φ is the fundamental solution of the heat equation. Also set $S(0)g = g$.

- (a) Prove $\{S(t)\}_{t \geq 0}$ is a contraction semigroup on $L^2(\mathbb{R}^n)$.
 (b) Show $\{S(t)\}_{t \geq 0}$ is *not* a contraction semigroup on $L^\infty(\mathbb{R}^n)$.
15. Let $\{S(t)\}_{t \geq 0}$ be a contraction semigroup on X , with generator A . Inductively define $D(A^k) := \{u \in D(A^{k-1}) \mid A^{k-1}u \in D(A)\}$ ($k = 2, \dots$). Show that if $u \in D(A^k)$ for some k , then $S(t)u \in D(A^k)$ for each $t \geq 0$.
16. Use Problem 15 to prove that if u is the semigroup solution in $X = L^2(U)$ of

$$\begin{cases} u_t - \Delta u = 0 & \text{in } U_T \\ u = 0 & \text{on } \partial U \times [0, T] \\ u = g & \text{on } U \times \{t = 0\}, \end{cases}$$

with $g \in C_c^\infty(U)$, then $u(\cdot, t) \in C^\infty(U)$ for each $0 \leq t \leq T$.

7.6. REFERENCES

- Section 7.1 See Ladyzhenskaya [L, Chapter 3], J.-L. Lions [L1], Lions–Magenes [L-M], Wloka [W1]. W. Schlag helped me with the proof of Theorems 5, 6. The proof of the parabolic Harnack inequality is similar to calculations found in Davies [Da]. For more on regularity theory, see the books of Krylov [Kr1], [Kr2], Ladyzhenskaya–Solonnikov–Uraltseva [L-S-U] and Lieberman [Lb].
- Section 7.2 See Ladyzhenskaya [L, Chapter 4], J.-L. Lions [L1], Lions–Magenes [L-M] and Wloka [W1].
- Section 7.3 The Fourier transform argument is taken from John [J2]; cf. also Treves [T, §15]. D. Serre has shown me a much more precise version of Theorem 5, under the further assumption of strict hyperbolicity.
- Section 7.4 See Friedman [Fr1, Part 2, §1] and Yosida [Y, Chapter IX].
- Section 7.5 Problem 9: see Brezis–Evans (Arch. Rational Mech. Analysis 71 (1979), 1–13) and also Ladyzhenskaya–Uraltseva [L-U, p. 182].

This page is intentionally left blank.

Part III

**THEORY FOR
NONLINEAR
PARTIAL
DIFFERENTIAL
EQUATIONS**

This page is intentionally left blank.

THE CALCULUS OF VARIATIONS

- 8.1 Introduction
- 8.2 Existence of minimizers
- 8.3 Regularity
- 8.4 Constraints
- 8.5 Critical points
- 8.6 Invariance, Noether's Theorem
- 8.7 Problems
- 8.8 References

8.1. INTRODUCTION

8.1.1. Basic ideas.

We introduce some new ideas by supposing that we wish to solve a particular partial differential equation, which for simplicity we write in the abstract form

$$(1) \quad A[u] = 0.$$

In this formula $A[\cdot]$ denotes a given, possibly nonlinear partial differential operator and u is the unknown. There is, of course, no general theory for solving all such PDE.

The *calculus of variations* identifies an important class of such nonlinear problems that can be solved using relatively simple techniques from nonlinear functional analysis. This is the class of *variational problems*, that is,

PDE of the form (1), where the nonlinear operator $A[\cdot]$ is the “derivative” of an appropriate “energy” functional $I[\cdot]$. Symbolically we write

$$(2) \quad A[\cdot] = I'[\cdot].$$

Then problem (1) reads

$$(3) \quad I'[u] = 0.$$

The advantage of this new formulation is that we now can recognize solutions of (1) as being critical points of $I[\cdot]$. These in certain circumstances may be relatively easy to find: if, for instance, the functional $I[\cdot]$ has a minimum at u , then presumably (3) is valid and thus u is a solution of the original PDE (1). *The point is that whereas it is usually extremely difficult to solve (1) directly, it may be much easier to discover minimum (or maximum or other critical) points of the functional $I[\cdot]$.*

In addition of course, many of the laws of physics and other scientific disciplines arise directly as variational principles.

8.1.2. First variation, Euler–Lagrange equation.

Suppose now $U \subset \mathbb{R}^n$ is a bounded, open set with smooth boundary ∂U and we are given a smooth function

$$L : \mathbb{R}^n \times \mathbb{R} \times \bar{U} \rightarrow \mathbb{R}.$$

We call L the *Lagrangian*.

NOTATION. We will write

$$L = L(p, z, x) = L(p_1, \dots, p_n, z, x_1, \dots, x_n)$$

for $p \in \mathbb{R}^n$, $z \in \mathbb{R}$, and $x \in U$. Thus “ p ” is the name of the variable for which we substitute $Dw(x)$ below, and “ z ” is the variable for which we substitute $w(x)$. We also set

$$\begin{cases} D_p L = (L_{p_1}, \dots, L_{p_n}) \\ D_z L = L_z \\ D_x L = (L_{x_1}, \dots, L_{x_n}). \end{cases}$$

This notation will clarify the theory to follow.

We make the vague ideas in §8.1.1 more precise by now assuming $I[\cdot]$ to have the explicit form

$$(4) \quad I[w] := \int_U L(Dw(x), w(x), x) dx,$$

for smooth functions $w : \bar{U} \rightarrow \mathbb{R}$ satisfying, say, the boundary condition

$$(5) \quad w = g \quad \text{on } \partial U.$$

Let us now additionally suppose some particular smooth function u , satisfying the requisite boundary condition $u = g$ on ∂U , happens to be a minimizer of $I[\cdot]$ among all functions w satisfying (5). *We will demonstrate that u is then automatically a solution of a certain nonlinear partial differential equation.*

To confirm this, first choose any smooth function $v \in C_c^\infty(U)$ and consider the real-valued function

$$(6) \quad i(\tau) := I[u + \tau v] \quad (\tau \in \mathbb{R}).$$

Since u is a minimizer of $I[\cdot]$ and $u + \tau v = u = g$ on ∂U , we observe that $i(\cdot)$ has a minimum at $\tau = 0$. Therefore

$$(7) \quad i'(0) = 0.$$

We explicitly compute this derivative (called the *first variation*) by writing out

$$(8) \quad i(\tau) = \int_U L(Du + \tau Dv, u + \tau v, x) dx.$$

Thus

$$i'(\tau) = \int_U \sum_{i=1}^n L_{p_i}(Du + \tau Dv, u + \tau v, x) v_{x_i} + L_z(Du + \tau Dv, u + \tau v, x) v dx.$$

Let $\tau = 0$, to deduce from (7) that

$$0 = i'(0) = \int_U \sum_{i=1}^n L_{p_i}(Du, u, x) v_{x_i} + L_z(Du, u, x) v dx.$$

Finally, since v has compact support, we can integrate by parts and obtain

$$0 = \int_U \left[- \sum_{i=1}^n (L_{p_i}(Du, u, x))_{x_i} + L_z(Du, u, x) \right] v dx.$$

As this equality holds for all test functions v , we conclude u solves the

nonlinear PDE

$$(9) \quad - \sum_{i=1}^n (L_{p_i}(Du, u, x))_{x_i} + L_z(Du, u, x) = 0 \quad \text{in } U.$$

This is the *Euler–Lagrange equation* associated with the energy functional $I[\cdot]$ defined by (4). Observe that (9) is a quasilinear, second-order PDE in divergence form.

In summary, any smooth minimizer of $I[\cdot]$ is a solution of the Euler–Lagrange partial differential equation (9), and thus—conversely—we can try to find a solution of (9) by searching for minimizers of (4).

Example 1 (Dirichlet’s principle). Take

$$L(p, z, x) = \frac{1}{2}|p|^2.$$

Then $L_{p_i} = p_i$ ($i = 1, \dots, n$), $L_z = 0$; and so the Euler–Lagrange equation associated with the functional

$$I[w] := \frac{1}{2} \int_U |Dw|^2 dx$$

is

$$\Delta u = 0 \quad \text{in } U.$$

This fact is *Dirichlet’s principle*, previously introduced in §2.2.5. □

Example 2 (Generalized Dirichlet’s principle). Write

$$L(p, z, x) = \frac{1}{2} \sum_{i,j=1}^n a^{ij}(x)p_i p_j - z f(x),$$

where $a^{ij} = a^{ji}$ ($i, j = 1, \dots, n$). Then $L_{p_i} = \sum_{j=1}^n a^{ij}(x)p_j$ ($i = 1, \dots, n$), $L_z = -f(x)$. Hence the Euler–Lagrange equation associated with the functional

$$I[w] := \int_U \frac{1}{2} \sum_{i,j=1}^n a^{ij} w_{x_i} w_{x_j} - w f dx$$

is the divergence structure linear equation

$$- \sum_{i,j=1}^n (a^{ij} u_{x_j})_{x_i} = f \quad \text{in } U.$$

We will see later (in §8.1.3 and §8.2) that the uniform ellipticity condition on the a^{ij} ($i, j = 1, \dots, n$) is a natural further assumption, required to prove the existence of a minimizer. Consequently from the nonlinear viewpoint of the calculus of variations, the divergence structure form of a linear second-order elliptic PDE is completely natural. This observation provides some much belated motivation for the bilinear form techniques utilized in Chapter 6. \square

Example 3 (Nonlinear Poisson equation). Assume we are given a smooth function $f : \mathbb{R} \rightarrow \mathbb{R}$, and define its antiderivative $F(z) = \int_0^z f(y) dy$. Then the Euler–Lagrange equation associated with the functional

$$I[w] := \int_U \frac{1}{2} |Dw|^2 - F(w) dx$$

is the nonlinear Poisson equation

$$-\Delta u = f(u) \quad \text{in } U. \quad \square$$

Example 4 (Minimal surfaces). Let

$$L(p, z, x) = (1 + |p|^2)^{1/2},$$


so that

$$I[w] = \int_U (1 + |Dw|^2)^{1/2} dx$$

is the area of the graph of the function $w : U \rightarrow \mathbb{R}$. The associated Euler–Lagrange equation is

$$(10) \quad \sum_{i=1}^n \left(\frac{u_{x_i}}{(1 + |Du|^2)^{1/2}} \right)_{x_i} = 0 \quad \text{in } U.$$

This partial differential equation is the *minimal surface equation*. The expression $\operatorname{div} \left(\frac{Du}{(1 + |Du|^2)^{1/2}} \right)$ on the left side of (10) is n times the *mean curvature* of the graph of u . Thus a *minimal surface has zero mean curvature*. \square



A minimal surface

8.1.3. Second variation.

We continue in the spirit of the calculations from §8.1.2 by computing now the *second variation* of $I[\cdot]$ at the function u . This we find by observing that since u gives a minimum for $I[\cdot]$, we must have

$$i''(0) \geq 0,$$

$i(\cdot)$ defined as above by (6). In view of (8) we can calculate

$$\begin{aligned} i''(\tau) &= \int_U \sum_{i,j=1}^n L_{p_i p_j}(Du + \tau Dv, u + \tau v, x) v_{x_i} v_{x_j} \\ &\quad + 2 \sum_{i=1}^n L_{p_i z}(Du + \tau Dv, u + \tau v, x) v_{x_i} v \\ &\quad + L_{zz}(Du + \tau Dv, u + \tau v, x) v^2 dx. \end{aligned}$$

Setting $\tau = 0$, we derive the inequality

$$(11) \quad \begin{aligned} 0 \leq i''(0) &= \int_U \sum_{i,j=1}^n L_{p_i p_j}(Du, u, x) v_{x_i} v_{x_j} \\ &\quad + 2 \sum_{i=1}^n L_{p_i z}(Du, u, x) v_{x_i} v + L_{zz}(Du, u, x) v^2 dx, \end{aligned}$$

holding for all test functions $v \in C_c^\infty(U)$.

We can extract useful information from inequality (11), as follows. First, note after a routine approximation argument that estimate (11) is valid for

any Lipschitz continuous function v vanishing on ∂U . We then fix $\xi \in \mathbb{R}^n$ and define

$$(12) \quad v(x) := \epsilon \rho\left(\frac{x \cdot \xi}{\epsilon}\right) \zeta(x) \quad (x \in U),$$

where $\zeta \in C_c^\infty(U)$ and $\rho : \mathbb{R} \rightarrow \mathbb{R}$ is the periodic “zig-zag” function defined by

$$\rho(x) = \begin{cases} x & \text{if } 0 \leq x \leq \frac{1}{2} \\ 1 - x & \text{if } \frac{1}{2} \leq x \leq 1 \end{cases} \quad \rho(x+1) = \rho(x) \quad (x \in \mathbb{R}).$$

Thus

$$(13) \quad |\rho'| = 1 \quad \text{a.e.}$$

Observe further that $v_{x_i}(x) = \rho'\left(\frac{x \cdot \xi}{\epsilon}\right) \xi_i \zeta + O(\epsilon)$ as $\epsilon \rightarrow 0$, and so our substituting (12) into (11) yields

$$0 \leq \int_U \sum_{i,j=1}^n L_{p_i p_j}(Du, u, x) (\rho')^2 \xi_i \xi_j \zeta^2 dx + O(\epsilon).$$

We recall (13) and send $\epsilon \rightarrow 0$, thereby obtaining the inequality

$$0 \leq \int_U \sum_{i,j=1}^n L_{p_i p_j}(Du, u, x) \xi_i \xi_j \zeta^2 dx.$$

Since this estimate holds for all $\zeta \in C_c^\infty(U)$, we deduce

$$(14) \quad \sum_{i,j=1}^n L_{p_i p_j}(Du, u, x) \xi_i \xi_j \geq 0 \quad (\xi \in \mathbb{R}^n, x \in U).$$

We will see later in §8.2 that this necessary condition contains a clue as to the basic convexity assumption on the Lagrangian L required for the existence theory.

8.1.4. Systems.

a. Euler–Lagrange equations. The foregoing considerations generalize quite easily to the case of systems, the only new complications being largely notational. Recall from §A.1 that $\mathbb{M}^{m \times n}$ denotes the space of real $m \times n$ matrices, and assume the smooth Lagrangian function

$$L : \mathbb{M}^{m \times n} \times \mathbb{R}^m \times \bar{U} \rightarrow \mathbb{R}$$

is given.

NOTATION. We will write

$$L = L(P, z, x) = L(p_1^1, \dots, p_n^m, z^1, \dots, z^m, x_1, \dots, x_n)$$

for $P \in \mathbb{M}^{m \times n}$, $z \in \mathbb{R}^m$, and $x \in U$, where

$$P = \begin{pmatrix} p_1^1 & \cdots & p_n^1 \\ & \ddots & \\ p_1^m & \cdots & p_n^m \end{pmatrix}_{m \times n}.$$

(We are now employing superscripts to denote rows, since this notational convention simplifies the following formulas.)

As in §8.1.2 we associate with L the functional

$$(15) \quad I[\mathbf{w}] := \int_U L(D\mathbf{w}(x), \mathbf{w}(x), x) dx,$$

defined for smooth functions $\mathbf{w} : \bar{U} \rightarrow \mathbb{R}^m$, $\mathbf{w} = (w^1, \dots, w^m)$, satisfying the boundary condition $\mathbf{w} = \mathbf{g}$ on ∂U , $\mathbf{g} : \partial U \rightarrow \mathbb{R}^m$ being given. Here

$$D\mathbf{w}(x) = \begin{pmatrix} w_{x_1}^1 & \cdots & w_{x_n}^1 \\ & \ddots & \\ w_{x_1}^m & \cdots & w_{x_n}^m \end{pmatrix}_{m \times n}$$

is the gradient matrix of \mathbf{w} at x .

Let us now show that any smooth minimizer $\mathbf{u} = (u^1, \dots, u^m)$ of $I[\cdot]$, taken among functions equal to \mathbf{g} on ∂U , must solve a certain *system* of nonlinear partial differential equations. We therefore fix $\mathbf{v} = (v^1, \dots, v^m) \in C_c^\infty(U; \mathbb{R}^m)$ and write

$$i(\tau) := I[\mathbf{u} + \tau\mathbf{v}].$$

As before,

$$i'(0) = 0.$$

From this we readily deduce as above the equality

$$0 = i'(0) = \int_U \sum_{i=1}^n \sum_{k=1}^m L_{p_i^k}(D\mathbf{u}, \mathbf{u}, x) v_{x_i}^k + \sum_{k=1}^m L_{z^k}(D\mathbf{u}, \mathbf{u}, x) v^k dx.$$

As this identity is valid for all choices of v^1, \dots, v^m , we conclude after integrating by parts that

$$(16) \quad - \sum_{i=1}^n \left(L_{p_i^k}(D\mathbf{u}, \mathbf{u}, x) \right)_{x_i} + L_{z^k}(D\mathbf{u}, \mathbf{u}, x) = 0 \quad \text{in } U \quad (k = 1, \dots, m).$$

This coupled, quasilinear *system* of PDE comprises the *Euler–Lagrange equations* for the functional $I[\cdot]$ defined by (15).

b. Null Lagrangians. Surprisingly, it turns out to be interesting to study certain systems of nonlinear partial differential equations for which *every* smooth function is a solution.

DEFINITION. The function L is called a null Lagrangian if the system of Euler–Lagrange equations

$$(17) \quad - \sum_{i=1}^n \left(L_{p_i^k}(D\mathbf{u}, \mathbf{u}, x) \right)_{x_i} + L_{z^k}(D\mathbf{u}, \mathbf{u}, x) = 0 \quad \text{in } U \quad (k = 1, \dots, m)$$

is automatically solved by all smooth functions $\mathbf{u} : U \rightarrow \mathbb{R}^m$.

The importance of null Lagrangians is that the corresponding energy

$$I[\mathbf{w}] = \int_U L(D\mathbf{w}, \mathbf{w}, x) dx$$

depends only on the boundary conditions:

THEOREM 1 (Null Lagrangians and boundary conditions). Let L be a null Lagrangian. Assume $\mathbf{u}, \tilde{\mathbf{u}}$ are two functions in $C^2(\bar{U}, \mathbb{R}^m)$ such that

$$(18) \quad \mathbf{u} \equiv \tilde{\mathbf{u}} \quad \text{on } \partial U.$$

Then

$$(19) \quad I[\mathbf{u}] = I[\tilde{\mathbf{u}}].$$

Proof. Define

$$i(\tau) := I[\tau\mathbf{u} + (1 - \tau)\tilde{\mathbf{u}}] \quad (0 \leq \tau \leq 1).$$

Then

$$\begin{aligned} i'(\tau) &= \int_U \sum_{i=1}^n \sum_{k=1}^m L_{p_i^k}(\tau D\mathbf{u} + (1 - \tau)D\tilde{\mathbf{u}}, \tau\mathbf{u} + (1 - \tau)\tilde{\mathbf{u}}, x)(u_{x_i}^k - \tilde{u}_{x_i}^k) \\ &\quad + \sum_{k=1}^m L_{z^k}(\tau D\mathbf{u} + (1 - \tau)D\tilde{\mathbf{u}}, \tau\mathbf{u} + (1 - \tau)\tilde{\mathbf{u}}, x)(u^k - \tilde{u}^k) dx \\ &= \sum_{k=1}^m \int_U \left[- \sum_{i=1}^n (L_{p_i^k}(\tau D\mathbf{u} + (1 - \tau)D\tilde{\mathbf{u}}, \tau\mathbf{u} + (1 - \tau)\tilde{\mathbf{u}}, x))_{x_i} \right. \\ &\quad \left. + L_{z^k}(\tau D\mathbf{u} + (1 - \tau)D\tilde{\mathbf{u}}, \tau\mathbf{u} + (1 - \tau)\tilde{\mathbf{u}}, x) \right] (u^k - \tilde{u}^k) dx \\ &= 0, \end{aligned}$$

the last equality holding since the Euler–Lagrange system of PDE is satisfied by $\tau \mathbf{u} + (1 - \tau)\tilde{\mathbf{u}}$. The identity (19) follows. \square

In the scalar case that $m = 1$ the only null Lagrangians are the boring examples where L is linear in the variable p . For the case of systems ($m > 1$), however, there are certain nontrivial examples, which will turn out to be important for us later.

NOTATION. If A is an $n \times n$ matrix, we denote by

$$\text{cof } A$$

the *cofactor* matrix, whose $(k, i)^{\text{th}}$ entry is $(\text{cof } A)_i^k = (-1)^{i+k} d(A)_i^k$, where $d(A)_i^k$ is the determinant of the $(n-1) \times (n-1)$ matrix obtained by deleting the k^{th} row and i^{th} column from A .

LEMMA (Divergence-free rows). *Let $\mathbf{u} : \mathbb{R}^n \rightarrow \mathbb{R}^n$ be a smooth function. Then*

$$(20) \quad \sum_{i=1}^n (\text{cof } D\mathbf{u})_{i,x_i}^k = 0 \quad (k = 1, \dots, n).$$

Proof. 1. From linear algebra we recall the identity

$$(21) \quad (\det P)I = P^T(\text{cof } P) \quad (P \in \mathbb{M}^{n \times n});$$

that is,

$$(22) \quad (\det P)\delta_{ij} = \sum_{k=1}^n p_i^k (\text{cof } P)_j^k \quad (i, j = 1, \dots, n).$$

Thus in particular

$$(23) \quad \frac{\partial \det P}{\partial p_m^k} = (\text{cof } P)_m^k \quad (k, m = 1, \dots, n).$$

2. Now set $P = Du$ in (22), differentiate with respect to x_j , and sum $j = 1, \dots, n$, to find

$$\sum_{j,k,m=1}^n \delta_{ij} (\text{cof } D\mathbf{u})_m^k u_{x_m x_j}^k = \sum_{k,j=1}^n u_{x_i x_j}^k (\text{cof } D\mathbf{u})_j^k + u_{x_i}^k (\text{cof } D\mathbf{u})_{j,x_j}^k$$

for $i = 1, \dots, n$. This identity simplifies to read

$$(24) \quad \sum_{k=1}^n u_{x_i}^k \left(\sum_{j=1}^n (\operatorname{cof} D\mathbf{u})_{j,x_j}^k \right) = 0 \quad (i = 1, \dots, n).$$

3. Now if $\det D\mathbf{u}(x_0) \neq 0$, we deduce from (24) that

$$\sum_{j=1}^n (\operatorname{cof} D\mathbf{u})_{j,x_j}^k = 0 \quad (k = 1, \dots, n)$$

at x_0 . If instead $\det D\mathbf{u}(x_0) = 0$, we choose a number $\epsilon > 0$ so small that $\det(D\mathbf{u}(x_0) + \epsilon I) \neq 0$, apply steps 1–2 to $\tilde{\mathbf{u}} := \mathbf{u} + \epsilon x$, and send $\epsilon \rightarrow 0$. \square

THEOREM 2 (Determinants as null Lagrangians). *The determinant function*

$$L(P) = \det P \quad (P \in \mathbb{M}^{n \times n})$$

is a null Lagrangian.

Proof. We must show that for any smooth function $\mathbf{u} : U \rightarrow \mathbb{R}^n$,

$$\sum_{i=1}^n \left(L_{p_i^k}(D\mathbf{u}) \right)_{x_i} = 0 \quad (k = 1, \dots, n).$$

According to (23) we have $L_{p_i^k} = (\operatorname{cof} P)_i^k$ ($i, k = 1, \dots, n$). But then employing the notation and conclusion of the lemma, we see

$$\sum_{i=1}^n \left(L_{p_i^k}(D\mathbf{u}) \right)_{x_i} = \sum_{i=1}^n (\operatorname{cof} D\mathbf{u})_{i,x_i}^k = 0 \quad (k = 1, \dots, n). \quad \square$$

Some other interesting null Lagrangians are introduced in the exercises.

c. Application. A nice application is a quick analytic proof of a topological fixed point theorem.

THEOREM 3 (Brouwer's Fixed Point Theorem). *Assume*

$$\mathbf{u} : B(0, 1) \rightarrow B(0, 1)$$

is continuous, where $B(0, 1)$ denotes the closed unit ball in \mathbb{R}^n . Then \mathbf{u} has a fixed point; that is, there exists a point $x \in B(0, 1)$ with

$$\mathbf{u}(x) = x.$$

Proof. 1. Write $B = B(0, 1)$. We first of all claim that there does not exist a smooth function

$$(25) \quad \mathbf{w} : B \rightarrow \partial B$$

such that

$$(26) \quad \mathbf{w}(x) = x \quad \text{for all } x \in \partial B.$$

2. Suppose to the contrary that such a function \mathbf{w} exists. Let us temporarily write $\tilde{\mathbf{w}}$ for the identity function, so that $\tilde{\mathbf{w}}(x) = x$ for all $x \in B$. In view of (26), $\mathbf{w} \equiv \tilde{\mathbf{w}}$ on ∂B . Since the determinant is a null Lagrangian, Theorem 1 implies

$$(27) \quad \int_B \det D\mathbf{w} \, dx = \int_B \det D\tilde{\mathbf{w}} \, dx = |B| \neq 0.$$

On the other hand, (25) implies $|\mathbf{w}|^2 \equiv 1$; and so differentiating, we find

$$(28) \quad (D\mathbf{w})^T \mathbf{w} = \mathbf{0}.$$

Since $|\mathbf{w}| = 1$, (28) says 0 is an eigenvalue of $D\mathbf{w}^T$ for each $x \in B$. Therefore $\det D\mathbf{w} \equiv 0$ in B . This contradicts (27) and thereby proves no smooth function \mathbf{w} satisfying (25), (26) can exist.

3. Next we show there does not exist any continuous function \mathbf{w} verifying (25), (26). Indeed if \mathbf{w} were such a function, we continuously extend \mathbf{w} by setting $\mathbf{w}(x) = x$ if $x \in \mathbb{R}^n - B$. Observe that $\mathbf{w}(x) \neq 0$ ($x \in \mathbb{R}^n$). Fix $\epsilon > 0$ so small that $\mathbf{w}_1 := \eta_\epsilon * \mathbf{w}$ satisfies $\mathbf{w}_1(x) \neq 0$ ($x \in \mathbb{R}^n$). Note also that since η_ϵ is radial, we have $\mathbf{w}_1(x) = x$ if $x \in \mathbb{R}^n - B(0, 2)$, for $\epsilon > 0$ sufficiently small. Then

$$\mathbf{w}_2 := \frac{2\mathbf{w}_1}{|\mathbf{w}_1|}$$

would be a smooth mapping satisfying (25), (26) (with the ball $B(0, 2)$ replacing $B = B(0, 1)$), in contradiction to step 1.

4. Finally suppose $\mathbf{u} : B \rightarrow B$ is continuous but has no fixed point. Define now the mapping $\mathbf{w} : B \rightarrow \partial B$ by setting $\mathbf{w}(x)$ to be the point on ∂B hit by the ray emanating from $\mathbf{u}(x)$ and passing through x . This mapping is well defined since $\mathbf{u}(x) \neq x$ for all $x \in B$. In addition \mathbf{w} is continuous and satisfies (25), (26).

But this in turn is a contradiction to step 2. □

We will employ Brouwer's Fixed Point Theorem several times in Chapter 9.

8.2. EXISTENCE OF MINIMIZERS

In this section we will identify some conditions on the Lagrangian L which ensure that the functional $I[\cdot]$ does indeed have a minimizer, at least within an appropriate Sobolev space.

8.2.1. Coercivity, lower semicontinuity.

Let us start with some largely heuristic insights as to when the functional

$$(1) \quad I[w] := \int_U L(Dw(x), w(x), x) dx,$$

defined for appropriate functions $w : U \rightarrow \mathbb{R}$ satisfying

$$(2) \quad w = g \quad \text{on } \partial U,$$

should have a minimizer.

a. Coercivity. We first of all note that even a smooth function f mapping \mathbb{R} to \mathbb{R} and bounded below need not attain its infimum. Consider, for instance, $f = e^x$ or $(1 + x^2)^{-1}$. These examples suggest that we in general will need some hypothesis controlling $I[w]$ for “large” functions w . Certainly the most effective way to ensure this would be to hypothesize that $I[w]$ “grows rapidly as $|w| \rightarrow \infty$ ”.

More specifically, let us assume

$$(3) \quad 1 < q < \infty$$

is fixed. We will then suppose

$$(4) \quad \begin{cases} \text{there exist constants } \alpha > 0, \beta \geq 0 \text{ such that} \\ L(p, z, x) \geq \alpha|p|^q - \beta \\ \text{for all } p \in \mathbb{R}^n, z \in \mathbb{R}, x \in U. \end{cases}$$

Therefore

$$(5) \quad I[w] \geq \delta \|Dw\|_{L^q(U)}^q - \gamma$$

for $\gamma := \beta|U|$ and some constant $\delta > 0$. Thus $I[w] \rightarrow \infty$ as $\|Dw\|_{L^q} \rightarrow \infty$. It is customary to call (5) a *coercivity condition* on $I[\cdot]$.

Turning once more to our basic task of finding minimizers for the functional $I[\cdot]$, we observe from inequality (5) that it seems reasonable to define $I[w]$ not only for smooth functions w , but also for functions w in the Sobolev

space $W^{1,q}(U)$ that satisfy the boundary condition (2) in the trace sense. After all, the wider the class of functions w for which $I[w]$ is defined, the more candidates we will have for a minimizer.

We will henceforth write

$$\mathcal{A} := \{w \in W^{1,q}(U) \mid w = g \text{ on } \partial U \text{ in the trace sense}\}$$

to denote this class of *admissible* functions w . Note in view of (4) that $I[w]$ is defined (but may equal $+\infty$) for each $w \in \mathcal{A}$.

b. Lower semicontinuity. Next, let us observe that although a continuous function $f : \mathbb{R} \rightarrow \mathbb{R}$ satisfying a coercivity condition does indeed attain its infimum, our integral functional $I[\cdot]$ in general will not. To understand the problem, set

$$(6) \quad m := \inf_{w \in \mathcal{A}} I[w]$$

and choose functions $u_k \in \mathcal{A}$ ($k = 1, \dots$) so that

$$(7) \quad I[u_k] \rightarrow m \quad \text{as } k \rightarrow \infty.$$

We call $\{u_k\}_{k=1}^{\infty}$ a *minimizing sequence*.

We would now like to show that some subsequence of $\{u_k\}_{k=1}^{\infty}$ converges to an actual minimizer. For this, however, we need some kind of compactness, and this is definitely a problem since the space $W^{1,q}(U)$ is infinite dimensional. Indeed, if we utilize the coercivity inequality (5), it turns out (cf. §8.2.2) that we can only conclude that the minimizing sequence lies in a bounded subset of $W^{1,q}(U)$. But this does *not* imply that there exists any subsequence which converges in $W^{1,q}(U)$.

We therefore turn our attention to the *weak topology* (cf. §D.4). Since we are assuming $1 < q < \infty$, so that $L^q(U)$ is reflexive, we conclude that there exists a subsequence $\{u_{k_j}\}_{j=1}^{\infty} \subset \{u_k\}_{k=1}^{\infty}$ and a function $u \in W^{1,q}(U)$ so that

$$(8) \quad \begin{cases} u_{k_j} \rightharpoonup u \text{ weakly in } L^q(U) \\ Du_{k_j} \rightharpoonup Du \text{ weakly in } L^q(U; \mathbb{R}^n). \end{cases}$$

We will hereafter abbreviate (8) by saying

$$(9) \quad u_{k_j} \rightharpoonup u \text{ weakly in } W^{1,q}(U).$$

Furthermore, it will be true that $u = g$ on ∂U in the trace sense, and so $u \in \mathcal{A}$.

Consequently by shifting to the weak topology we have recovered enough compactness from the coercivity inequality (5) to deduce (9) for an appropriate subsequence. But now another difficulty arises, for in essentially all cases of interest the functional $I[\cdot]$ is not continuous with respect to weak convergence. In other words, we cannot deduce from (7) and (9) that

$$(10) \quad I[u] = \lim_{j \rightarrow \infty} I[u_{k_j}],$$

and thus u is a minimizer. The problem is that $Du_{k_j} \rightharpoonup Du$ does not imply $Du_{k_j} \rightarrow Du$ a.e.: it is quite possible for instance that the gradients Du_{k_j} , although bounded in L^q , are oscillating more and more wildly as $k_j \rightarrow \infty$.

What saves us is the final, key observation that we do not really need the full strength of (10). It would suffice instead to know only

$$(11) \quad I[u] \leq \liminf_{j \rightarrow \infty} I[u_{k_j}].$$

Then from (7) we could deduce $I[u] \leq m$. But owing to (6), $m \leq I[u]$. Consequently u is indeed a minimizer.

DEFINITION. We say that a function $I[\cdot]$ is (sequentially) weakly lower semicontinuous on $W^{1,q}(U)$, provided

$$I[u] \leq \liminf_{k \rightarrow \infty} I[u_k]$$

whenever

$$u_k \rightharpoonup u \quad \text{weakly in } W^{1,q}(U).$$

Our goal therefore is now to identify reasonable conditions on the non-linear term L that ensure $I[\cdot]$ is weakly lower semicontinuous.

8.2.2. Convexity.

We next look back to our second variation analysis in §8.1.3 and recall we derived there the inequality

$$\sum_{i,j=1}^n L_{p_i p_j}(Du(x), u(x), x) \xi_i \xi_j \geq 0 \quad (\xi \in \mathbb{R}^n, x \in U)$$

holding as a necessary condition, whenever u is a smooth minimizer. This inequality strongly suggests that it is reasonable to assume that L is convex in its first argument.

THEOREM 1 (Weak lower semicontinuity). *Assume that L is smooth, bounded below and in addition*

the mapping $p \mapsto L(p, z, x)$ is convex,

for each $z \in \mathbb{R}$, $x \in U$. Then

$I[\cdot]$ is weakly lower semicontinuous on $W^{1,q}(U)$.

Proof. 1. Choose any sequence $\{u_k\}_{k=1}^\infty$ with

$$(12) \quad u_k \rightharpoonup u \quad \text{weakly in } W^{1,q}(U),$$

and set $l := \liminf_{k \rightarrow \infty} I[u_k]$. We must show

$$(13) \quad I[u] \leq l.$$

2. Note first from (12) and §D.4 that

$$(14) \quad \sup_k \|u_k\|_{W^{1,q}(U)} < \infty.$$

Upon passing to a subsequence if necessary, we may as well also suppose

$$(15) \quad l = \lim_{k \rightarrow \infty} I[u_k].$$

Furthermore we see from the compactness theorem in §5.7 that $u_k \rightarrow u$ strongly in $L^q(U)$; and thus, passing if necessary to yet another subsequence, we have

$$(16) \quad u_k \rightarrow u \quad \text{a.e. in } U.$$

3. Fix $\epsilon > 0$. Then (16) and Egoroff's Theorem (§E.2) assert

$$(17) \quad u_k \rightarrow u \quad \text{uniformly on } E_\epsilon,$$

where E_ϵ is a measurable set with

$$(18) \quad |U - E_\epsilon| \leq \epsilon.$$

We may assume $E_\epsilon \subseteq E_{\epsilon'}$ for $0 < \epsilon' < \epsilon$. Now write

$$(19) \quad F_\epsilon := \left\{ x \in U \mid |u(x)| + |Du(x)| \leq \frac{1}{\epsilon} \right\}.$$

Then

$$(20) \quad |U - F_\epsilon| \rightarrow 0 \quad \text{as } \epsilon \rightarrow 0.$$

We finally set

$$(21) \quad G_\epsilon := E_\epsilon \cap F_\epsilon$$

and notice from (18), (20) that $|U - G_\epsilon| \rightarrow 0$ as $\epsilon \rightarrow 0$.

4. Now let us observe that since L is bounded below, we may as well assume

$$(22) \quad L \geq 0$$

(for otherwise we could apply the following arguments to $\tilde{L} = L + \beta \geq 0$ for some appropriate constant β). Consequently

$$(23) \quad \begin{aligned} I[u_k] &= \int_U L(Du_k, u_k, x) \, dx \geq \int_{G_\epsilon} L(Du_k, u_k, x) \, dx \\ &\geq \int_{G_\epsilon} L(Du, u_k, x) \, dx + \int_{G_\epsilon} D_p L(Du, u_k, x) \cdot (Du_k - Du) \, dx, \end{aligned}$$

the last inequality following from the convexity of L in its first argument; see §B.1. Now in view of (17), (19) and (21)

$$(24) \quad \lim_{k \rightarrow \infty} \int_{G_\epsilon} L(Du, u_k, x) \, dx = \int_{G_\epsilon} L(Du, u, x) \, dx.$$

In addition, since $D_p L(Du, u_k, x) \rightarrow D_p L(Du, u, x)$ uniformly on G_ϵ and $Du_k \rightharpoonup Du$ weakly in $L^q(U; \mathbb{R}^n)$, we have

$$(25) \quad \lim_{k \rightarrow \infty} \int_{G_\epsilon} D_p L(Du, u_k, x) \cdot (Du_k - Du) \, dx = 0.$$

Owing now to (24), (25), we deduce from (23) that

$$l = \lim_{k \rightarrow \infty} I[u_k] \geq \int_{G_\epsilon} L(Du, u, x) \, dx.$$

This inequality holds for each $\epsilon > 0$. We now let ϵ tend to zero and recall (22) and the Monotone Convergence Theorem (§E.3) to conclude

$$l \geq \int_U L(Du, u, x) \, dx = I[u],$$

as required. □

Remark. It is very important to understand how the foregoing proof deals with the weak convergence $Du_k \rightharpoonup Du$. The key is the convexity inequality (23), on the right-hand side of which Du_k appears linearly. Weak convergence is, by its very definition, compatible with linear expressions; and so the limit (25) holds. Remember that it is not in general true that $Du_k \rightarrow Du$ a.e., even if we pass to a subsequence.

The convergence of u_k to u in L^q is much stronger, and so we do not need any convexity assumption concerning $z \mapsto L(p, z, x)$.

We can at last establish that $I[\cdot]$ has a minimizer among the functions in \mathcal{A} .

THEOREM 2 (Existence of minimizer). *Assume that L satisfies the coercivity inequality (4) and is convex in the variable p . Suppose also the admissible set \mathcal{A} is nonempty.*

Then there exists at least one function $u \in \mathcal{A}$ solving

$$I[u] = \min_{w \in \mathcal{A}} I[w] .$$

Proof. 1. Set $m := \inf_{w \in \mathcal{A}} I[w]$. If $m = +\infty$, we are done, and so we henceforth assume m is finite. Select a minimizing sequence $\{u_k\}_{k=1}^{\infty}$. Then

$$(26) \quad I[u_k] \rightarrow m.$$

2. We may as well take $\beta = 0$ in inequality (4), since we could otherwise just as well consider $\tilde{L} := L + \beta$. Thus $L \geq \alpha|p|^q$, and so

$$(27) \quad I[w] \geq \alpha \int_U |Dw|^q dx.$$

Since m is finite, we conclude from (26) and (27) that

$$(28) \quad \sup_k \|Du_k\|_{L^q(U)} < \infty.$$

3. Now fix any function $w \in \mathcal{A}$. Since u_k and w both equal g on ∂U in the trace sense, we have $u_k - w \in W_0^{1,q}(U)$. Therefore Poincaré's inequality implies

$$\begin{aligned} \|u_k\|_{L^q(U)} &\leq \|u_k - w\|_{L^q(U)} + \|w\|_{L^q(U)} \\ &\leq C\|Du_k - Dw\|_{L^q(U)} + C \leq C, \end{aligned}$$

by (28). Hence $\sup_k \|u_k\|_{L^q(U)} < \infty$. This estimate and (28) imply $\{u_k\}_{k=1}^\infty$ is bounded in $W^{1,q}(U)$.

4. Consequently there exist a subsequence $\{u_{k_j}\}_{j=1}^\infty \subset \{u_k\}_{k=1}^\infty$ and a function $u \in W^{1,q}(U)$ such that

$$u_{k_j} \rightharpoonup u \quad \text{weakly in } W^{1,q}(U).$$

We assert next that $u \in \mathcal{A}$. To see this, note that for $w \in \mathcal{A}$ as above, $u_k - w \in W_0^{1,q}(U)$. Now $W_0^{1,q}(U)$ is a closed, linear subspace of $W^{1,q}(U)$ and so, by Mazur's Theorem (§D.4), is weakly closed. Hence $u - w \in W_0^{1,q}(U)$. Consequently the trace of u on ∂U is g .

In view of Theorem 1 then, $I[u] \leq \liminf_{j \rightarrow \infty} I[u_{k_j}] = m$. But since $u \in \mathcal{A}$, it follows that

$$I[u] = m = \min_{w \in \mathcal{A}} I[w]. \quad \square$$

We turn next to the problem of uniqueness. In general there can be many minimizers, and so to ensure uniqueness we require further assumptions. Suppose for instance

$$(29) \quad L = L(p, x) \text{ does not depend on } z$$

and

$$(30) \quad \begin{cases} \text{there exists } \theta > 0 \text{ such that} \\ \sum_{i,j=1}^n L_{p_i p_j}(p, x) \xi_i \xi_j \geq \theta |\xi|^2 \quad (p, \xi \in \mathbb{R}^n; x \in U). \end{cases}$$

Condition (30) says the mapping $p \mapsto L(p, x)$ is uniformly convex for each x .

THEOREM 3 (Uniqueness of minimizer). *Suppose (29), (30) hold. Then a minimizer $u \in \mathcal{A}$ of $I[\cdot]$ is unique.*

Proof. 1. Assume $u, \tilde{u} \in \mathcal{A}$ are both minimizers of $I[\cdot]$ over \mathcal{A} . Then $v := \frac{u + \tilde{u}}{2} \in \mathcal{A}$. We claim

$$(31) \quad I[v] \leq \frac{I[u] + I[\tilde{u}]}{2},$$

with a strict inequality, unless $u = \tilde{u}$ a.e.

2. To see this, note from the uniform convexity assumption that we have

$$(32) \quad L(p, x) \geq L(q, x) + D_p L(q, x) \cdot (p - q) + \frac{\theta}{2} |p - q|^2 \quad (x \in U, p, q \in \mathbb{R}^n).$$

Set $q = \frac{Du + D\tilde{u}}{2}$, $p = Du$, and integrate over U :

$$(33) \quad I[v] + \int_U D_p L\left(\frac{Du + D\tilde{u}}{2}, x\right) \cdot \left(\frac{Du - D\tilde{u}}{2}\right) dx + \frac{\theta}{8} \int_U |Du - D\tilde{u}|^2 dx \leq I[u].$$

Similarly, set $q = \frac{Du + D\tilde{u}}{2}$, $p = D\tilde{u}$ in (32) and integrate:

$$(34) \quad I[v] + \int_U D_p L\left(\frac{Du + D\tilde{u}}{2}, x\right) \cdot \left(\frac{D\tilde{u} - Du}{2}\right) dx + \frac{\theta}{8} \int_U |Du - D\tilde{u}|^2 dx \leq I[\tilde{u}].$$

Add and divide by 2, to deduce

$$I[v] + \frac{\theta}{8} \int_U |Du - D\tilde{u}|^2 dx \leq \frac{I[u] + I[\tilde{u}]}{2}.$$

This proves (31).

3. As $I[u] = I[\tilde{u}] = \min_{w \in \mathcal{A}} I[w] \leq I[v]$, we deduce $Du = D\tilde{u}$ a.e. in U . Since $u = \tilde{u} = g$ on ∂U in the trace sense, it follows that $u = \tilde{u}$ a.e. \square

8.2.3. Weak solutions of Euler–Lagrange equation.

We wish next to demonstrate that any minimizer $u \in \mathcal{A}$ of $I[\cdot]$ solves the Euler–Lagrange equation in some suitable sense. This does not follow from the calculations in §8.1 since we do not know u is smooth, only $u \in W^{1,q}(U)$. And in fact we will need some growth conditions on L and its derivatives. Let us hereafter suppose

$$(35) \quad |L(p, z, x)| \leq C(|p|^q + |z|^q + 1)$$

and also

$$(36) \quad \begin{cases} |D_p L(p, z, x)| \leq C(|p|^{q-1} + |z|^{q-1} + 1) \\ |D_z L(p, z, x)| \leq C(|p|^{q-1} + |z|^{q-1} + 1) \end{cases}$$

for some constant C and all $p \in \mathbb{R}^n$, $z \in \mathbb{R}$, $x \in U$.

Motivation for definition of weak solution. We now turn our attention to the boundary-value problem for the Euler–Lagrange PDE associated with our functional L , which for a smooth minimizer u reads

$$(37) \quad \begin{cases} -\sum_{i=1}^n (L_{p_i}(Du, u, x))_{x_i} + L_z(Du, u, x) = 0 & \text{in } U \\ u = g & \text{on } \partial U. \end{cases}$$

If we multiply (37) by a test function $v \in C_c^\infty(U)$ and integrate by parts, we arrive at the equality

$$(38) \quad \int_U \sum_{i=1}^n L_{p_i}(Du, u, x)v_{x_i} + L_z(Du, u, x)v \, dx = 0.$$

Of course this is the identity we first obtained in our derivation of (37) in §8.1.2.

Now assume $u \in W^{1,q}(U)$. Then using (36) we see

$$|D_p L(Du, u, x)| \leq C(|Du|^{q-1} + |u|^{q-1} + 1) \in L^{q'}(U),$$

where $q' = \frac{q}{q-1}$, $\frac{1}{q} + \frac{1}{q'} = 1$. Similarly

$$(39) \quad |D_z L(Du, u, x)| \leq C(|Du|^{q-1} + |u|^{q-1} + 1) \in L^{q'}(U).$$

Consequently we see using a standard approximation argument that the equality (38) is valid for any $v \in W_0^{1,q}(U)$. This motivates the following

DEFINITION. We say $u \in \mathcal{A}$ is a weak solution of the boundary-value problem (37) for the Euler–Lagrange equation provided

$$\int_U \sum_{i=1}^n L_{p_i}(Du, u, x)v_{x_i} + L_z(Du, u, x)v \, dx = 0$$

for all $v \in W_0^{1,q}(U)$.

THEOREM 4 (Solution of Euler–Lagrange equation). Assume L verifies the growth conditions (35), (36) and $u \in \mathcal{A}$ satisfies

$$I[u] = \min_{w \in \mathcal{A}} I[w].$$

Then u is a weak solution of (37).

Proof. We proceed as in §8.1.2, taking care about differentiating inside the integrals. Fix any $v \in W_0^{1,q}(U)$ and set

$$i(\tau) := I[u + \tau v] \quad (\tau \in \mathbb{R}).$$

In view of (35) we see that $i(\tau)$ is finite for all τ .

Let $\tau \neq 0$ and write the difference quotient

$$(40) \quad \begin{aligned} \frac{i(\tau) - i(0)}{\tau} &= \int_U \frac{L(Du + \tau Dv, u + \tau v, x) - L(Du, u, x)}{\tau} \, dx \\ &= \int_U L^\tau(x) \, dx, \end{aligned}$$

where

$$L^\tau(x) := \frac{1}{\tau} [L(Du(x) + \tau Dv(x), u(x) + \tau v(x), x) - L(Du(x), u(x), x)]$$

for a.e. $x \in U$. Clearly

$$(41) \quad L^\tau(x) \rightarrow \sum_{i=1}^n L_{p_i}(Du, u, x)v_{x_i} + L_z(Du, u, x)v \quad \text{a.e.}$$

as $\tau \rightarrow 0$. Furthermore

$$\begin{aligned} L^\tau(x) &= \frac{1}{\tau} \int_0^\tau \frac{d}{ds} L(Du + sDv, u + sv, x) ds \\ &= \frac{1}{\tau} \int_0^\tau \sum_{i=1}^n L_{p_i}(Du + sDv, u + sv, x)v_{x_i} \\ &\quad + L_z(Du + sDv, u + sv, x)v ds. \end{aligned}$$

Next recall from §B.2 Young's inequality: $ab \leq \frac{a^q}{q} + \frac{b^{q'}}{q'}$, where $\frac{1}{q} + \frac{1}{q'} = 1$. Then since $u, v \in W^{1,q}(U)$, inequalities (36) and Young's inequality imply after some elementary calculations that

$$|L^\tau(x)| \leq C(|Du|^q + |u|^q + |Dv|^q + |v|^q + 1) \in L^1(U)$$

for each $\tau \neq 0$. Consequently we may invoke the Dominated Convergence Theorem to conclude from (40), (41) that $i'(0)$ exists and equals

$$\int_U \sum_{i=1}^n L_{p_i}(Du, u, x)v_{x_i} + L_z(Du, u, x)v dx.$$

But then since $i(\cdot)$ has a minimum for $\tau = 0$, we know $i'(0) = 0$; and thus u is a weak solution. \square

Remark. In general, the Euler–Lagrange equation (37) will have other solutions which do not correspond to minima of $I[\cdot]$; see §8.2.5 and §8.5. However, *in the special case that the joint mapping $(p, z) \mapsto L(p, z, x)$ is convex for each x , then each weak solution is in fact a minimizer.*

To see this, suppose $u \in \mathcal{A}$ solves

$$(42) \quad \begin{cases} -\sum_{i=1}^n (L_{p_i}(Du, u, x))_{x_i} + L_z(Du, u, x) = 0 & \text{in } U \\ u = g & \text{on } \partial U \end{cases}$$

in the weak sense and select any $w \in \mathcal{A}$. Utilizing the convexity of the mapping $(p, z) \mapsto L(p, z, x)$, we have

$$L(p, z, x) + D_p L(p, z, x) \cdot (q - p) + D_z L(p, z, x) \cdot (w - z) \leq L(q, w, x).$$

Let $p = Du(x)$, $q = Dw(x)$, $z = u(x)$, $w = w(x)$ and integrate over U :

$$I[u] + \int_U D_p L(Du, u, x) \cdot (Dw - Du) + D_z L(Du, u, x)(w - u) dx \leq I[w].$$

In view of (42) the second term on the left is zero, and therefore $I[u] \leq I[w]$ for each $w \in \mathcal{A}$.

8.2.4. Systems.

a. Convexity. We now adopt again the notation for systems set forth in §8.1.4 and consider the existence question for minimizers of the functional

$$I[\mathbf{w}] := \int_U L(D\mathbf{w}(x), \mathbf{w}(x), x) dx,$$

defined for appropriate functions $\mathbf{w} : U \rightarrow \mathbb{R}^m$, where now $L : \mathbb{M}^{m \times n} \times \mathbb{R}^m \times \bar{U} \rightarrow \mathbb{R}$ is given.

It turns out the theory developed in §8.2.2 extends with no difficulty to the case at hand. Let us therefore assume the coercivity inequality

$$(43) \quad L(P, z, x) \geq \alpha |P|^q - \beta \quad (P \in \mathbb{M}^{m \times n}, z \in \mathbb{R}^m, x \in U)$$

for constants $\alpha > 0$, $\beta \geq 0$ and set also

$$\mathcal{A} := \{\mathbf{w} \in W^{1,q}(U; \mathbb{R}^m) \mid \mathbf{w} = \mathbf{g} \text{ on } \partial U \text{ in the trace sense}\},$$

where $\mathbf{g} : \partial U \rightarrow \mathbb{R}^m$ is given.

THEOREM 5 (Existence of minimizer). *Assume that L satisfies the coercivity inequality (43) and is convex in the variable P . Suppose also the admissible set \mathcal{A} is nonempty.*

Then there exists $\mathbf{u} \in \mathcal{A}$ solving

$$I[\mathbf{u}] = \min_{\mathbf{w} \in \mathcal{A}} I[\mathbf{w}].$$

The proof follows almost exactly the proofs of Theorems 1 and 2 in §8.2.2. Similarly to Theorem 3 above we have

THEOREM 6 (Uniqueness of minimizer). *Assume L does not depend on z and the mapping $P \mapsto L(P, x)$ is uniformly convex. Then a minimizer $\mathbf{u} \in \mathcal{A}$ of $I[\cdot]$ is unique.*

Now suppose additionally

$$(44) \quad \begin{cases} |L(P, z, x)| \leq C(|P|^q + |z|^q + 1) \\ |D_P L(P, z, x)| \leq C(|P|^{q-1} + |z|^{q-1} + 1) \\ |D_z L(P, z, x)| \leq C(|P|^{q-1} + |z|^{q-1} + 1) \end{cases}$$

for some constant C and all $P \in \mathbb{M}^{m \times n}$, $z \in \mathbb{R}^m$, $x \in U$.

We consider now the system of Euler–Lagrange equations

$$(45) \quad \begin{cases} -\sum_{i=1}^n (L_{p_i^k}(D\mathbf{u}, \mathbf{u}, x))_{x_i} + L_{z^k}(D\mathbf{u}, \mathbf{u}, x) = 0 & \text{in } U \\ u^k = g^k & \text{on } \partial U \end{cases}$$

for $k = 1, \dots, m$ and define $u \in \mathcal{A}$ to be a *weak solution* provided

$$\sum_{k=1}^m \int_U \sum_{i=1}^n L_{p_i^k}(D\mathbf{u}, \mathbf{u}, x) w_{x_i}^k + L_{z^k}(D\mathbf{u}, \mathbf{u}, x) w^k dx = 0$$

for all $\mathbf{w} \in W_0^{1,q}(U; \mathbb{R}^m)$, $\mathbf{w} = (w^1, \dots, w^m)$.

THEOREM 7 (Solution of Euler–Lagrange system). *Assume L verifies the growth conditions (44) and $\mathbf{u} \in \mathcal{A}$ satisfies*

$$I[\mathbf{u}] = \min_{\mathbf{w} \in \mathcal{A}} I[\mathbf{w}].$$

Then \mathbf{u} is a weak solution of (45).

The proof is almost precisely like that of Theorem 4.

b. Polyconvexity. It is rather surprising that there are some mathematically and physically interesting systems that are not covered by Theorem 5 above but can still be studied using the calculus of variations. These include certain problems where the Lagrangian L is *not* convex in P , but $I[\cdot]$ is nonetheless weakly lower semicontinuous.

LEMMA (Weak continuity of determinants). *Assume $n < q < \infty$ and*

$$\mathbf{u}_k \rightharpoonup \mathbf{u} \quad \text{weakly in } W^{1,q}(U; \mathbb{R}^n).$$

Then

$$\det D\mathbf{u}_k \rightharpoonup \det D\mathbf{u} \quad \text{weakly in } L^{q/n}(U).$$

Proof. 1. First we recall the matrix identity $(\det P)I = P(\operatorname{cof} P)^T$; consequently

$$\det P = \sum_{j=1}^n p_j^i (\operatorname{cof} P)_j^i \quad (i = 1, \dots, n).$$

2. Now let $\mathbf{w} \in C^\infty(U; \mathbb{R}^n)$, $\mathbf{w} = (w^1, \dots, w^n)$. Then

$$(46) \quad \det D\mathbf{w} = \sum_{j=1}^n w_{x_j}^i (\operatorname{cof} D\mathbf{w})_j^i \quad (i = 1, \dots, n).$$

But the lemma in §8.1.4 asserts $\sum_{j=1}^n (\operatorname{cof} D\mathbf{w})_{j,x_j}^i = 0$. Thus formula (46) says

$$\det D\mathbf{w} = \sum_{j=1}^n (w^i (\operatorname{cof} D\mathbf{w})_j^i)_{x_j}.$$

Consequently *the determinant of the gradient matrix can be written as a divergence*. Therefore if $v \in C_c^\infty(U)$, we have

$$(47) \quad \int_U v \det D\mathbf{w} \, dx = - \sum_{j=1}^n \int_U v_{x_j} w^i (\operatorname{cof} D\mathbf{w})_j^i \, dx \quad (i = 1, \dots, n).$$

3. We have established the identity (47) for a smooth function w , and so a standard approximation argument yields

$$(48) \quad \int_U v \det D\mathbf{u}_k \, dx = - \sum_{j=1}^n \int_U v_{x_j} u_k^i (\operatorname{cof} D\mathbf{u}_k)_j^i \, dx$$

for $k = 1, 2, \dots$. Now since $n < q < \infty$ and $\mathbf{u}_k \rightharpoonup \mathbf{u}$ in $W^{1,q}(U; \mathbb{R}^n)$, we know from Morrey's inequality that $\{\mathbf{u}_k\}_{k=1}^\infty$ is bounded in $C^{0,1-n/q}(U; \mathbb{R}^n)$. Thus using the Arzela–Ascoli compactness criterion, §C.7, we deduce $\mathbf{u}_k \rightarrow \mathbf{u}$ uniformly in U . Returning then to identity (48), we see that we could conclude

$$(49) \quad \lim_{k \rightarrow \infty} \int_U v \det D\mathbf{u}_k \, dx = - \sum_{j=1}^n \int_U v_{x_j} u^i (\operatorname{cof} D\mathbf{u})_j^i \, dx = \int_U v \det D\mathbf{u} \, dx,$$

if we knew

$$(50) \quad \lim_{k \rightarrow \infty} \int_U \psi (\operatorname{cof} D\mathbf{u}_k)_j^i \, dx = \int_U \psi (\operatorname{cof} D\mathbf{u})_j^i \, dx$$

for $i, j = 1, \dots, n$ and each $\psi \in C_c^\infty(U)$. However $(\operatorname{cof} D\mathbf{u}_k)_j^i$ is the determinant of an $(n-1) \times (n-1)$ matrix, which can be analyzed as above by being

written as a sum of determinants of appropriate $(n-2) \times (n-2)$ submatrices, times uniformly convergent factors. We continue and eventually must show only the obvious fact that the entries of the matrices $D\mathbf{u}_k$ converge weakly to the corresponding entries of $D\mathbf{u}$. In this way we verify (50) and thus (49).

4. Finally, since $\{\mathbf{u}_k\}_{k=1}^\infty$ is bounded in $W^{1,q}(U; \mathbb{R}^n)$ and $|\det D\mathbf{u}_k| \leq C|D\mathbf{u}_k|^n$, we see that $\{\det D\mathbf{u}_k\}_{k=1}^\infty$ is bounded in $L^{q/n}(U)$. Hence any subsequence has a weakly convergent subsequence in $L^{q/n}(U)$, which—owing to (49)—can only converge to $\det D\mathbf{u}$. \square

We next utilize this lemma to establish a weak lower semicontinuity assertion analogous to Theorem 1, except that we will not assume that the Lagrangian L is necessarily convex in P . Instead let us suppose that $m = n$ and L has the form

$$(51) \quad L(P, z, x) = F(P, \det P, z, x) \quad (P \in \mathbb{M}^{n \times n}, z \in \mathbb{R}^n, x \in U)$$

where $F : \mathbb{M}^{n \times n} \times \mathbb{R} \times \mathbb{R}^n \times \bar{U} \rightarrow \mathbb{R}$ is smooth. We additionally hypothesize that

$$(52) \quad \begin{cases} \text{for each fixed } z \in \mathbb{R}^m, x \in \mathbb{R}^n, \text{ the joint mapping} \\ (P, r) \mapsto F(P, r, z, x) \text{ is convex.} \end{cases}$$

A Lagrangian L of the form (51) is called *polyconvex* provided (52) holds.

THEOREM 8 (Lower semicontinuity of polyconvex functionals). *Suppose $n < q < \infty$. Assume also L is bounded below and is polyconvex. Then*

$$I[\cdot] \text{ is weakly lower semicontinuous on } W^{1,q}(U; \mathbb{R}^n).$$

Proof. Choose any sequence $\{\mathbf{u}_k\}_{k=1}^\infty$ with

$$(53) \quad \mathbf{u}_k \rightharpoonup \mathbf{u} \quad \text{weakly in } W^{1,q}(U; \mathbb{R}^n).$$

According to the lemma,

$$(54) \quad \det D\mathbf{u}_k \rightharpoonup \det D\mathbf{u} \quad \text{weakly in } L^{q/n}(U).$$

We can now argue almost exactly as in the proof of Theorem 1. Indeed, using the notation from that proof, we have

$$\begin{aligned} I[\mathbf{u}_k] &= \int_U L(D\mathbf{u}_k, \mathbf{u}_k, x) dx \geq \int_{G_\epsilon} L(D\mathbf{u}_k, \mathbf{u}_k, x) dx \\ &= \int_{G_\epsilon} F(D\mathbf{u}_k, \det D\mathbf{u}_k, \mathbf{u}_k, x) dx \\ &\geq \int_{G_\epsilon} F(D\mathbf{u}, \det D\mathbf{u}, \mathbf{u}_k, x) dx \\ &\quad + \int_{G_\epsilon} F_p(D\mathbf{u}, \det D\mathbf{u}, \mathbf{u}_k, x) \cdot (D\mathbf{u}_k - D\mathbf{u}) \\ &\quad + F_r(D\mathbf{u}, \det D\mathbf{u}, \mathbf{u}_k, x)(\det D\mathbf{u}_k - \det D\mathbf{u}) dx, \end{aligned}$$

in view of (52). Reasoning as in the proof of Theorem 1, we deduce from (53), (54) that the limit of the last term is zero as $k \rightarrow \infty$. \square

As before, we immediately deduce

THEOREM 9 (Existence of minimizers, polyconvex functionals). *Assume that $n < q < \infty$ and that L satisfies the coercivity inequality (43) and is polyconvex. Suppose also the admissible set \mathcal{A} is nonempty.*

Then there exists $\mathbf{u} \in \mathcal{A}$ solving

$$I[\mathbf{u}] = \min_{\mathbf{w} \in \mathcal{A}} I[\mathbf{w}].$$

Example: elasticity. Much of the interest in polyconvexity comes from nonlinear elasticity theory, where $n = 3$. We consider an elastic body, which initially has the reference configuration U . We then displace each point $x \in \partial U$ to a new position $\mathbf{g}(x)$ and wish to determine the new displacement $\mathbf{u}(x)$ of each internal point $x \in U$.

If the material is *hyperelastic*, there exists by definition an associated energy density L such that the physical displacement \mathbf{u} minimizes the internal energy functional

$$I[\mathbf{w}] := \int_U L(D\mathbf{w}, x) dx$$

over all admissible displacements $\mathbf{w} \in \mathcal{A}$. Now it seems reasonable physically that L , which represents the internal energy density from stretching and compression, may explicitly depend on the local change in volume, that is, on $\det D\mathbf{w}$. In other words, it is physically appropriate to suppose that L has the form $L(P, x) = F(P, \det P, x)$. Then F describes in its first argument changes in internal energy due to changes in line elements, and in

its second argument changes in internal energy due to changes in volume elements. See Ball (Arch. Rational Mech. Analysis 63 (1977), 337–403) for more explanation.

8.2.5. Local minimizers.

We pose in this subsection a basic question: *under what circumstances is a critical point of an energy functional $I[\cdot]$ in fact a minimizer or a local minimizer?* To be specific, let us assume that u is a smooth solution of the Euler–Lagrange PDE

$$(55) \quad \begin{cases} -\sum_{i=1}^n (L_{p_i}(Du, u, x))_{x_i} + L_z(Du, u, x) = 0 & \text{in } U \\ u = g & \text{on } \partial U \end{cases}$$

and is therefore a critical point of the functional

$$I[w] = \int_U L(Dw, w, x) dx$$

among functions w satisfying the boundary condition $w = g$ on ∂U . We as usual assume

$$p \mapsto L(p, z, x) \text{ is convex.}$$

We will show that if the graph of $x \mapsto u(x)$ lies within a region R generated by a one-parameter family of graphs $x \mapsto u(x, \lambda)$ corresponding to other critical points, then in fact u is a minimizer of $I[\cdot]$ as compared with admissible variations w taking values within R . More precisely, suppose that $I \subseteq \mathbb{R}$ is an open interval containing 0 and $\{u(\cdot, \lambda) \mid \lambda \in I\}$ is a smooth one-parameter family of solutions of the Euler–Lagrange PDE

$$(56) \quad -\sum_{i=1}^n (L_{p_i}(Du(x, \lambda), u(x, \lambda), x))_{x_i} + L_z(Du(x, \lambda), u(x, \lambda), x) = 0$$

within U , such that

$$(57) \quad u(x) = u(x, 0) \quad (x \in U).$$

We as follows construct an admissible function w taking values in the region R , the union of the graphs of the functions $u(\cdot, \lambda)$ for $\lambda \in I$. Take $\theta : \bar{U} \rightarrow I$ to be a smooth function satisfying

$$(58) \quad \theta = 0 \quad \text{on } \partial U.$$

Define then

$$(59) \quad w(x) := u(x, \theta(x)),$$

and note that $w = u = g$ on ∂U .

THEOREM 10 (Local minimizers). *The function u is a local minimizer within the region R , in the sense that*

$$(60) \quad I[u] \leq I[w]$$

for any function w constructed as above.

Thus if u is a solution of the Euler–Lagrange PDE and is embedded within a family of other solutions, then u is a minimizer of $I[\cdot]$ among functions w having the form (59). If say $u_\lambda > 0$ for λ small, we can write any w that is sufficiently close to u pointwise in this form. Notice that Dw need not be close to Du .

Proof. 1. We first observe that

$$w_{x_i}(x) = u_{x_i}(x, \theta(x)) + u_\lambda(x, \theta(x))\theta_{x_i} \quad (i = 1, \dots, n).$$

Hence the convexity of L in its first argument implies

$$(61) \quad \begin{aligned} I[w] &= \int_U L(Dw, w, x) dx \\ &= \int_U L(Du + u_\lambda D\theta, w, x) dx \\ &\geq \int_U L(Du, w, x) + u_\lambda D_p L(Du, w, x) \cdot D\theta dx, \end{aligned}$$

where u is evaluated at $(x, \theta(x))$ and $D = D_x$.

2. We now introduce the vector field $\mathbf{b} = (b^1, \dots, b^n)$, defined by

$$(62) \quad b^i := \int_0^{\theta(x)} u_\lambda(x, \lambda) L_{p_i}(Du(x, \lambda), u(x, \lambda), x) d\lambda \quad (i = 1, \dots, n).$$

Then

$$\begin{aligned} \operatorname{div} \mathbf{b} &= \sum_{i=1}^n \theta_{x_i} u_\lambda(x, \theta(x)) L_{p_i}(Du(x, \theta(x)), u(x, \theta(x)), x) \\ &\quad + \sum_{i=1}^n \int_0^{\theta(x)} u_{\lambda x_i}(x, \lambda) L_{p_i}(Du(x, \lambda), u(x, \lambda), x) \\ &\quad \quad \quad + u_\lambda(x, \lambda) (L_{p_i}(Du(x, \lambda), u(x, \lambda), x))_{x_i} d\lambda \\ &= u_\lambda(x, \theta) D_p L(Du(x, \theta(x)), w(x), x) \cdot D\theta \\ &\quad + \int_0^{\theta(x)} \sum_{i=1}^n u_{\lambda x_i}(x, \lambda) L_{p_i}(Du(x, \lambda), u(x, \lambda), x) \\ &\quad \quad \quad + u_\lambda(x, \lambda) L_z(Du(x, \lambda), u(x, \lambda), x) d\lambda. \end{aligned}$$

We invoked the Euler–Lagrange equations (56) for the last equality in this calculation. Observe next that

$$(L(Du(x, \lambda), u(x, \lambda), x))_\lambda = \sum_{i=1}^n L_{p_i}(Du, u, x)u_{x_i\lambda} + L_z(Du, u, x)u_\lambda.$$

Hence the foregoing calculation implies

$$\begin{aligned} \operatorname{div} \mathbf{b} &= u_\lambda(x, \theta)D_p L(Du, w, x) \cdot D\theta + \int_0^{\theta(x)} L(Du, u, x)_\lambda d\lambda \\ &= u_\lambda(x, \theta)D_p L(Du, w, x) \cdot D\theta \\ &\quad + L(Du(x, \theta(x)), w, x) - L(Du, u, x), \end{aligned}$$

in view of (57).

3. We employ this computation in (61), to deduce using the Gauss–Green Theorem that

$$\begin{aligned} I[w] &\geq \int_U L(Du, u, x) + \operatorname{div} \mathbf{b} dx \\ &= I[u] + \int_{\partial U} \mathbf{b} \cdot \nu dS \\ &= I[u], \end{aligned}$$

since $\theta = 0$ on ∂U and consequently $\mathbf{b} = 0$ on ∂U . We have proved (60). \square

This calculation provides a glimpse into the deep classical theories as to when critical points are minimizers or local minimizers: see Morrey [Mo] and Giaquinta–Hildebrandt [G-H] for more.

8.3. REGULARITY

We discuss in this section the smoothness of minimizers to our energy functionals. This is generally a quite difficult topic, and so we will make a number of strong simplifying assumptions. Thus we henceforth suppose our functional $I[\cdot]$ to have the form

$$(1) \quad I[w] := \int_U L(Dw) - wf dx,$$

for $f \in L^2(U)$. We will also take $q = 2$ and suppose as well the growth condition

$$(2) \quad |D_p L(p)| \leq C(|p| + 1) \quad (p \in \mathbb{R}^n).$$

Then any minimizer $u \in \mathcal{A}$ is a weak solution of the Euler–Lagrange PDE

$$(3) \quad - \sum_{i=1}^n (L_{p_i}(Du))_{x_i} = f \quad \text{in } U;$$

that is,

$$(4) \quad \int_U \sum_{i=1}^n L_{p_i}(Du) v_{x_i} dx = \int_U f v dx$$

for all $v \in H_0^1(U)$.

8.3.1. Second derivative estimates.

We now intend to show that if $u \in H^1(U)$ is a weak solution of the nonlinear PDE (3), then in fact $u \in H_{\text{loc}}^2(U)$. But to establish this we will need to strengthen our growth conditions on L . Let us first of all suppose

$$(5) \quad |D^2L(p)| \leq C \quad (p \in \mathbb{R}^n).$$

In addition let us assume that L is uniformly convex, and so there exists a constant $\theta > 0$ such that

$$(6) \quad \sum_{i,j=1}^n L_{p_i p_j}(p) \xi_i \xi_j \geq \theta |\xi|^2 \quad (p, \xi \in \mathbb{R}^n).$$

Clearly this is some sort of nonlinear analogue of our uniform ellipticity condition for linear PDE in Chapter 6. The idea will therefore be to try to utilize, or at least mimic, some of the calculations from that chapter.

THEOREM 1 (Second derivatives for minimizers).

(i) *Let $u \in H^1(U)$ be a weak solution of the nonlinear partial differential equation (3), where L satisfies (5), (6). Then*

$$u \in H_{\text{loc}}^2(U).$$

(ii) *If in addition $u \in H_0^1(U)$ and ∂U is C^2 , then*

$$u \in H^2(U),$$

with the estimate

$$\|u\|_{H^2(U)} \leq C \|f\|_{L^2(U)}.$$

Proof. 1. We will largely follow the proof of Theorem 1 in §6.3.1, the corresponding assertion of local H^2 regularity for solutions of linear second-order elliptic PDE.

Fix any open set $V \subset\subset U$ and choose then an open set W so that $V \subset\subset W \subset\subset U$. Select a smooth cutoff function ζ satisfying

$$\zeta \equiv 1 \text{ on } V, \quad \zeta \equiv 0 \text{ in } \mathbb{R}^n - W, \quad 0 \leq \zeta \leq 1.$$

Let $|h| > 0$ be small, choose $k \in \{1, \dots, n\}$, and substitute

$$v := -D_k^{-h}(\zeta^2 D_k^h u)$$

into (4). We are employing here the notation from §5.8.2:

$$D_k^h u(x) = \frac{u(x + he_k) - u(x)}{h} \quad (x \in W).$$

Using the identity $\int_U u D_k^{-h} v \, dx = - \int_U v D_k^h u \, dx$, we deduce

$$(7) \quad \sum_{i=1}^n \int_U D_k^h (L_{p_i}(Du)) (\zeta^2 D_k^h u)_{x_i} \, dx = - \int_U f D_k^{-h} (\zeta^2 D_k^h u) \, dx.$$

Now

$$(8) \quad \begin{aligned} D_k^h L_{p_i}(Du(x)) &= \frac{L_{p_i}(Du(x + he_k)) - L_{p_i}(Du(x))}{h} \\ &= \frac{1}{h} \int_0^1 \frac{d}{ds} L_{p_i}(sDu(x + he_k) + (1-s)Du(x)) \, ds \\ &= \frac{1}{h} \int_0^1 \sum_{j=1}^n L_{p_i p_j}(sDu(x + he_k) + (1-s)Du(x)) \\ &\quad (u_{x_j}(x + he_k) - u_{x_j}(x)) \, ds \\ &= \sum_{j=1}^n a^{ij,h}(x) D_k^h u_{x_j}(x), \end{aligned}$$

for

$$(9) \quad a^{ij,h}(x) := \int_0^1 L_{p_i p_j}(sDu(x + he_k) + (1-s)Du(x)) \, ds \quad (i, j = 1, \dots, n).$$

We substitute (8) into (7) and perform simple calculations, to arrive at the identity:

$$\begin{aligned}
 (10) \quad A_1 + A_2 &:= \sum_{i,j=1}^n \int_U \zeta^2 a^{ij,h} D_k^h u_{x_j} D_k^h u_{x_i} dx \\
 &\quad + \sum_{i,j=1}^n \int_U a^{ij,h} D_k^h u_{x_j} D_k^h u_{x_i} 2\zeta \zeta_{x_i} dx \\
 &= - \int_U f D_k^{-h} (\zeta^2 D_k^h u) dx =: B.
 \end{aligned}$$

Now the uniform convexity condition (6) implies

$$(11) \quad A_1 \geq \theta \int_U \zeta^2 |D_k^h Du|^2 dx.$$

Furthermore we see from (5) that

$$\begin{aligned}
 (12) \quad |A_2| &\leq C \int_W \zeta |D_k^h Du| |D_k^h u| dx \\
 &\leq \epsilon \int_W \zeta^2 |D_k^h Du|^2 dx + \frac{C}{\epsilon} \int_W |D_k^h u|^2 dx.
 \end{aligned}$$

Furthermore, as in the proof of Theorem 1 in §6.3.1, we have

$$|B| \leq \epsilon \int_U \zeta^2 |D_k^h Du|^2 dx + \frac{C}{\epsilon} \int_U f^2 + |Du|^2 dx.$$

We select $\epsilon = \frac{\theta}{4}$, to deduce from the foregoing bounds on A_1, A_2, B the estimate

$$\int_U \zeta^2 |D_k^h Du|^2 dx \leq C \int_W f^2 + |D_k^h u|^2 dx \leq C \int_U f^2 + |Du|^2 dx,$$

the last inequality valid according to Theorem 3(i) in §5.8.2.

2. Since $\zeta \equiv 1$ on V , we find

$$\int_V |D_k^h Du|^2 dx \leq C \int_U f^2 + |Du|^2 dx$$

for $k = 1, \dots, n$ and all sufficiently small $|h| > 0$. Consequently Theorem 3(ii) in §5.8.2 implies $Du \in H^1(V)$, and so $u \in H^2(V)$. This is true for each $V \subset\subset U$; thus $u \in H_{\text{loc}}^2(U)$.

3. If $u \in H_0^1(U)$ is a weak solution of (3) and ∂U is C^2 , we can then mimic the proof of the boundary regularity Theorem 4 in §6.3.2 to prove $u \in H^2(U)$, with estimate

$$\|u\|_{H^2(U)} \leq C(\|f\|_{L^2(U)} + \|u\|_{H^1(U)});$$

details are left to the reader. Now from (6) follows the inequality

$$(DL(p) - DL(0)) \cdot p \geq \theta|p|^2 \quad (p \in \mathbb{R}^n).$$

If we then put $v = u$ in (4), we can employ this estimate to derive the bound

$$\|u\|_{H^1(U)} \leq C\|f\|_{L^2(U)},$$

and so finish the proof. □

8.3.2. Remarks on higher regularity.

We would next like to show that if L is infinitely differentiable, then so is u . By analogy with the regularity theory developed for second-order linear elliptic PDE in §6.3, it may seem natural to try to extend the H_{loc}^2 estimate from the previous section to obtain further estimates in the higher Sobolev spaces $H_{\text{loc}}^k(U)$ for $k = 3, 4, \dots$

This method will *not* work for the nonlinear partial differential equation (3) however. The reason is this. For linear equations we could, roughly speaking, differentiate the equation many times and still obtain a linear PDE of the same general form as that we began with. See for instance the proof of Theorem 2 in §6.3.1. But if we differentiate a nonlinear differential equation many times, the resulting increasingly complicated expressions quickly become impossible to handle. Much deeper ideas are called for, the full development of which is beyond the scope of this book. We will nevertheless at least outline the basic plan.

To start with, choose a test function $w \in C_c^\infty(U)$, select $k \in \{1, \dots, n\}$, and set $v = -w_{x_k}$ in the identity (4), where for simplicity we now take $f \equiv 0$. Since we now know $u \in H_{\text{loc}}^2(U)$, we can integrate by parts to find

$$(13) \quad \int_U \sum_{i,j=1}^n L_{p_i p_j}(Du) u_{x_k x_j} w_{x_i} dx = 0.$$

Next write

$$(14) \quad \tilde{u} := u_{x_k}$$

and

$$(15) \quad a^{ij} := L_{p_i p_j}(Du) \quad (i, j = 1, \dots, n).$$

Fix also any $V \subset\subset U$. Then after an approximation we find from (13)–(15) that

$$(16) \quad \int_U \sum_{i,j=1}^n a^{ij}(x) \tilde{u}_{x_j} w_{x_i} dx = 0$$

for all $w \in H_0^1(V)$. This is to say that $\tilde{u} \in H^1(V)$ is a weak solution of the linear, second-order elliptic PDE

$$(17) \quad - \sum_{i,j=1}^n (a^{ij} \tilde{u}_{x_j})_{x_i} = 0 \quad \text{in } V.$$

But we *cannot* just apply our regularity theory from §6.3 to conclude from (17) that \tilde{u} is smooth, the reason being that we can deduce from (5) and (15) only that

$$a^{ij} \in L^\infty(V) \quad (i, j = 1, \dots, n).$$

However a deep theorem, due independently to DeGiorgi and to Nash, asserts that any weak solution of (17) must in fact be locally Hölder continuous for some exponent $\gamma > 0$. (See Gilbarg–Trudinger [G-T, Chapter 8].) Thus if $W \subset\subset V$, we have $\tilde{u} \in C^{0,\gamma}(W)$, and so

$$u \in C_{\text{loc}}^{1,\gamma}(U).$$

Return to the definition (15). If L is smooth, we now know $a^{ij} \in C_{\text{loc}}^{0,\gamma}(U)$ ($i, j = 1, \dots, n$). Then (3) and an older theorem of Schauder [G-T, Chapters 4 and 6] assert that in fact

$$u \in C_{\text{loc}}^{2,\gamma}(U).$$

But then $a^{ij} \in C_{\text{loc}}^{1,\gamma}(U)$, and so another version of Schauder's estimate implies

$$u \in C_{\text{loc}}^{3,\gamma}(U).$$

We can continue this so-called “bootstrap” argument, eventually to deduce u is $C_{\text{loc}}^{k,\gamma}(U)$ for $k = 1, \dots$, and so $u \in C^\infty(U)$.

See Giaquinta [Gi] for much more about regularity theory in the calculus of variations.

8.4. CONSTRAINTS

In this section we consider applications of the calculus of variations to certain constrained minimization problems and in particular discuss the role of *Lagrange multipliers* in the corresponding Euler–Lagrange PDE.

8.4.1. Nonlinear eigenvalue problems.

We investigate first problems with *integral constraints*. To be specific, let us look at the problem of minimizing the energy functional

$$(1) \quad I[w] := \frac{1}{2} \int_U |Dw|^2 dx$$

over all functions w with, say, $w = 0$ on ∂U but subject now also to the side condition that

$$(2) \quad J[w] := \int_U G(w) dx = 0,$$

where $G : \mathbb{R} \rightarrow \mathbb{R}$ is a given, smooth function.

We will henceforth write $g = G'$. Assume now

$$(3) \quad |g(z)| \leq C(|z| + 1),$$

and so

$$(4) \quad |G(z)| \leq C(|z|^2 + 1) \quad (z \in \mathbb{R})$$

for some constant C .

Let us introduce as well the appropriate admissible class

$$\mathcal{A} := \{w \in H_0^1(U) \mid J[w] = 0\}.$$

We suppose also that the open set U is bounded, connected and has a smooth boundary.

THEOREM 1 (Existence of constrained minimizer). *Assume the admissible set \mathcal{A} is nonempty. Then there exists $u \in \mathcal{A}$ satisfying*

$$I[u] = \min_{w \in \mathcal{A}} I[w].$$

Proof. Choose, as usual, a minimizing sequence $\{u_k\}_{k=1}^\infty \subset \mathcal{A}$ with

$$I[u_k] \rightarrow m = \inf_{w \in \mathcal{A}} I[w].$$

Then as above we can extract a subsequence

$$(5) \quad u_{k_j} \rightharpoonup u \quad \text{weakly in } H_0^1(U),$$

with $I[u] \leq m$. We will be done once we show

$$(6) \quad J[u] = 0,$$

so that $u \in \mathcal{A}$. Utilizing the compactness theory from §5.7, we deduce from (5) that

$$(7) \quad u_{k_j} \rightarrow u \quad \text{in } L^2(U).$$

Consequently

$$(8) \quad \begin{aligned} |J(u)| &= |J(u) - J(u_k)| \leq \int_U |G(u) - G(u_k)| \, dx \\ &\leq C \int_U |u - u_k|(1 + |u| + |u_k|) \, dx \quad \text{by (3)} \\ &\rightarrow 0 \quad \text{as } k \rightarrow \infty. \end{aligned} \quad \square$$

Far more interesting than the mere existence of constrained minimizers is an examination of the corresponding Euler–Lagrange equation.

THEOREM 2 (Lagrange multiplier). *Let $u \in \mathcal{A}$ satisfy*

$$(9) \quad I[u] = \min_{w \in \mathcal{A}} I[w].$$

Then there exists a real number λ such that

$$(10) \quad \int_U Du \cdot Dv \, dx = \lambda \int_U g(u)v \, dx$$

for all $v \in H_0^1(U)$.

Remark. Thus u is a weak solution of the nonlinear boundary-value problem

$$(11) \quad \begin{cases} -\Delta u = \lambda g(u) & \text{in } U \\ u = 0 & \text{on } \partial U, \end{cases}$$

where λ is the *Lagrange multiplier* corresponding to the *integral constraint*

$$(12) \quad J[u] = 0.$$

A problem of the form (11) for the unknowns (u, λ) , with $u \not\equiv 0$, is a *nonlinear eigenvalue problem*.

Proof. 1. Fix any function $v \in H_0^1(U)$. Assume first

$$(13) \quad g(u) \text{ is not equal to zero a.e. within } U.$$

Choose then any function $w \in H_0^1(U)$ with

$$(14) \quad \int_U g(u)w \, dx \neq 0;$$

this is possible because of (13). Now write

$$(15) \quad \begin{aligned} j(\tau, \sigma) &:= J[u + \tau v + \sigma w] \\ &= \int_U G(u + \tau v + \sigma w) \, dx \quad (\tau, \sigma \in \mathbb{R}). \end{aligned}$$

Clearly

$$(16) \quad j(0, 0) = \int_U G(u) \, dx = 0.$$

In addition, j is C^1 and

$$(17) \quad \frac{\partial j}{\partial \tau}(\tau, \sigma) = \int_U g(u + \tau v + \sigma w)v \, dx,$$

$$(18) \quad \frac{\partial j}{\partial \sigma}(\tau, \sigma) = \int_U g(u + \tau v + \sigma w)w \, dx.$$

Consequently (14) implies

$$(19) \quad \frac{\partial j}{\partial \sigma}(0, 0) \neq 0.$$

According to the Implicit Function Theorem (§C.7), there exists a C^1 function $\phi : \mathbb{R} \rightarrow \mathbb{R}$ such that

$$(20) \quad \phi(0) = 0$$

and

$$(21) \quad j(\tau, \phi(\tau)) = 0$$

for all sufficiently small τ , say $|\tau| \leq \tau_0$. Differentiating, we discover

$$\frac{\partial j}{\partial \tau}(\tau, \phi(\tau)) + \frac{\partial j}{\partial \sigma}(\tau, \phi(\tau))\phi'(\tau) = 0,$$

whence (17) and (18) yield

$$(22) \quad \phi'(0) = -\frac{\int_U g(u)v \, dx}{\int_U g(u)w \, dx}.$$

2. Now set

$$w(\tau) := \tau v + \phi(\tau)w \quad (|\tau| \leq \tau_0)$$

and write

$$i(\tau) := I[u + w(\tau)].$$

Since (21) implies $J[u + w(\tau)] = 0$, we see that $u + w(\tau) \in \mathcal{A}$. So the C^1 function $i(\cdot)$ has a minimum at 0. Thus

$$(23) \quad \begin{aligned} 0 = i'(0) &= \int_U (Du + \tau Dv + \phi(\tau)Dw) \cdot (Dv + \phi'(\tau)Dw) \, dx|_{\tau=0} \\ &= \int_U Du \cdot (Dv + \phi'(0)Dw) \, dx. \end{aligned}$$

Recall now (22) and *define*

$$\lambda := \frac{\int_U Du \cdot Dw \, dx}{\int_U g(u)w \, dx},$$

to deduce from (23) the desired equality

$$\int_U Du \cdot Dv \, dx = \lambda \int_U g(u)v \, dx$$

for all $v \in H_0^1(U)$.

3. Suppose now instead of (13) that

$$g(u) = 0 \quad \text{a.e. in } U.$$

Approximating g by bounded functions, we deduce $DG(u) = g(u)Du = 0$ a.e. Hence, since U is connected, $G(u)$ is constant a.e. It follows that $G(u) = 0$ a.e., because $J[u] = \int_U G(u) \, dx = 0$. As $u = 0$ on ∂U in the trace sense, it follows that $G(0) = 0$.

But then $u = 0$ a.e., as otherwise $I[u] > I[0] = 0$. Since $g(u) = 0$ a.e., the identity (10) is trivially valid in this case, for any λ . \square

8.4.2. Unilateral constraints, variational inequalities.

We study now calculus of variation problems with certain pointwise, *one-sided constraints* on the values of $u(x)$ for each $x \in U$. For definiteness let us consider the problem of minimizing, say, the energy functional

$$(24) \quad I[w] := \int_U \frac{1}{2} |Dw|^2 - fw \, dx,$$

among all functions w belonging to the set

$$(25) \quad \mathcal{A} := \{w \in H_0^1(U) \mid w \geq h \text{ a.e. in } U\},$$

where $h : \bar{U} \rightarrow \mathbb{R}$ is a given smooth function, called the *obstacle*. The convex admissible set \mathcal{A} thus comprises those functions $w \in H_0^1(U)$ satisfying the one-sided, or *unilateral*, constraint that $w \geq h$. We suppose as well that f is a given, smooth function.

THEOREM 3 (Existence of minimizer). *Assume the admissible set \mathcal{A} is nonempty. Then there exists a unique function $u \in \mathcal{A}$ satisfying*

$$I[u] = \min_{w \in \mathcal{A}} I[w].$$

Proof. 1. The existence of a minimizer follows very easily from the general ideas discussed before. We need only note explicitly that if $\{u_{k_j}\}_{j=1}^\infty \subset \mathcal{A}$ is a minimizing sequence with $u_{k_j} \rightharpoonup u$ weakly in $H_0^1(U)$, then by compactness we have $u_{k_j} \rightarrow u$ strongly in $L^2(U)$. Since $u_{k_j} \geq h$ a.e., it follows that $u \geq h$ a.e. Therefore $u \in \mathcal{A}$.

2. We now prove uniqueness. Assume u and $\tilde{u} \in \mathcal{A}$ are two minimizers, with $u \neq \tilde{u}$. Then $w := \frac{u+\tilde{u}}{2} \in \mathcal{A}$, and

$$\begin{aligned} I[w] &= \int_U \frac{1}{2} \left| \left(\frac{Du+D\tilde{u}}{2} \right) \right|^2 - f \left(\frac{u+\tilde{u}}{2} \right) dx \\ &= \int_U \frac{1}{8} (|Du|^2 + 2Du \cdot D\tilde{u} + |D\tilde{u}|^2) - f \left(\frac{u+\tilde{u}}{2} \right) dx. \end{aligned}$$

Now $2a \cdot b = |a|^2 + |b|^2 - |a - b|^2$. Thus

$$\begin{aligned} I[w] &= \int_U \frac{1}{8} (2|Du|^2 + 2|D\tilde{u}|^2 - |Du - D\tilde{u}|^2) - f \left(\frac{u+\tilde{u}}{2} \right) dx \\ &< \frac{1}{2} \int_U \frac{1}{2} |Du|^2 - fu \, dx + \frac{1}{2} \int_U \frac{1}{2} |D\tilde{u}|^2 - f\tilde{u} \, dx \\ &= \frac{1}{2} I[u] + \frac{1}{2} I[\tilde{u}], \end{aligned}$$

the strict inequality holding since $u \neq \tilde{u}$. This is a contradiction, since u and \tilde{u} are minimizers. \square

We next compute the analogue of the Euler–Lagrange equation, which for the case at hand turns out to be an inequality.

THEOREM 4 (Variational characterization of minimizer). *Let $u \in \mathcal{A}$ be the unique solution of*

$$I[u] = \min_{w \in \mathcal{A}} I[w].$$

Then

$$(26) \quad \int_U Du \cdot D(w - u) \, dx \geq \int_U f(w - u) \, dx \quad \text{for all } w \in \mathcal{A}.$$

We call (26) a *variational inequality*.

Proof. 1. Fix any element $w \in \mathcal{A}$. Then for each $0 \leq \tau \leq 1$,

$$u + \tau(w - u) = (1 - \tau)u + \tau w \in \mathcal{A},$$

since \mathcal{A} is convex. Thus if we set

$$i(\tau) := I[u + \tau(w - u)],$$

we see that $i(0) \leq i(\tau)$ for all $0 \leq \tau \leq 1$. Hence

$$(27) \quad i'(0) \geq 0.$$

2. Now if $0 < \tau \leq 1$,

$$\begin{aligned} \frac{i(\tau) - i(0)}{\tau} &= \frac{1}{\tau} \int_U \frac{|Du + \tau D(w - u)|^2 - |Du|^2}{2} - f(u + \tau(w - u) - u) \, dx \\ &= \int_U Du \cdot D(w - u) + \frac{\tau |D(w - u)|^2}{2} - f(w - u) \, dx. \end{aligned}$$


Thus (27) implies

$$0 \leq i'(0) = \int_U Du \cdot D(w - u) - f(w - u) \, dx. \quad \square$$

Notice that we obtain the inequality (27), since we can in effect take only “one-sided” variations, away from the constraint.

Interpretation of the variational inequality. To gain some insight into the variational inequality (26), let us quote without proof a regularity assertion (see Kinderlehrer–Stampacchia [K-S]), which states $u \in W^{2,\infty}(U)$, provided ∂U is smooth. Hence the set

$$O := \{x \in U \mid u(x) > h(x)\}$$



The free boundary for the obstacle problem

is open, and

$$C := \{x \in U \mid u(x) = h(x)\}$$

is (relatively) closed.

We claim that in fact $u \in C^\infty(O)$ and

$$(28) \quad -\Delta u = f \quad \text{in } O.$$

To see this, fix any test function $v \in C_c^\infty(O)$. Then if $|\tau|$ is sufficiently small, $w := u + \tau v \geq h$, and so $w \in \mathcal{A}$. Thus (26) implies $\tau \int_O Du \cdot Dv - f v dx \geq 0$. This inequality is valid for all sufficiently small τ , both positive and negative, and so in fact

$$\int_O Du \cdot Dv - f v dx = 0$$


for all $v \in C_c^\infty(O)$. Hence u is a weak solution of (28), whence linear regularity theory (§6.3) shows $u \in C^\infty(O)$.

Now if $v \in C_c^\infty(U)$ satisfies $v \geq 0$ and if $0 < \tau \leq 1$, then $w := u + \tau v \in \mathcal{A}$, whence $\int_U Du \cdot Dv - f v dx \geq 0$. But since $u \in W^{2,\infty}(U)$, we can integrate by parts to deduce $\int_U (-\Delta u - f)v dx \geq 0$ for all nonnegative functions $v \in C_c^\infty(U)$. Thus

$$(29) \quad -\Delta u \geq f \quad \text{a.e. in } U.$$

We summarize our conclusions by observing from (28), (29) that

$$(30) \quad \begin{cases} u \geq h, & -\Delta u \geq f & \text{a.e. in } U \\ & -\Delta u = f & \text{on } U \cap \{u > h\}. \end{cases}$$



A harmonic map into a sphere

Free boundaries. The set

$$F := \partial O \cap U$$

is called the *free boundary*. Many interesting problems in applied mathematics involve partial differential equations with free boundaries. Such of these problems as can be recast as variational inequalities become relatively easy to study, especially since there is no explicit mention of the free boundary in the inequalities (30). Applications arise in stopping time optimal control problems for Brownian motion, in groundwater hydrology, in plasticity theory, etc. See Kinderlehrer–Stampacchia [K-S].

8.4.3. Harmonic maps.

We consider next the case of *pointwise* constraints as exemplified by *harmonic maps into spheres*. We are interested now in the problem of minimizing the energy

$$(31) \quad I[\mathbf{w}] := \frac{1}{2} \int_U |D\mathbf{w}|^2 dx$$

over all functions belonging to the admissible class

$$(32) \quad \mathcal{A} := \{\mathbf{w} \in H^1(U; \mathbb{R}^m) \mid \mathbf{w} = \mathbf{g} \text{ on } \partial U, |\mathbf{w}| = 1 \text{ a.e.}\}.$$

The idea is that we are trying to minimize the energy over all appropriate maps from $U \subset \mathbb{R}^n$ into the unit sphere $S^{m-1} = \partial B(0, 1) \subset \mathbb{R}^m$. This problem and its variants arise for instance as simplified models for the behavior of liquid crystals.

It is straightforward to verify that there exists at least one minimizer in \mathcal{A} , provided $\mathcal{A} \neq \emptyset$.

THEOREM 5 (Euler–Lagrange equation for harmonic maps). *Let $\mathbf{u} \in \mathcal{A}$ satisfy*

$$I[\mathbf{u}] = \min_{\mathbf{w} \in \mathcal{A}} I[\mathbf{w}].$$

Then

$$(33) \quad \int_U D\mathbf{u} : D\mathbf{v} \, dx = \int_U |D\mathbf{u}|^2 \mathbf{u} \cdot \mathbf{v} \, dx$$

for each $\mathbf{v} \in H_0^1(U; \mathbb{R}^m) \cap L^\infty(U; \mathbb{R}^m)$.

We interpret (33) as saying that $\mathbf{u} = (u^1, \dots, u^m)$ is a weak solution of the boundary-value problem

$$(34) \quad \begin{cases} -\Delta \mathbf{u} = |D\mathbf{u}|^2 \mathbf{u} & \text{in } U \\ \mathbf{u} = \mathbf{g} & \text{on } \partial U. \end{cases}$$

The function $\lambda = |D\mathbf{u}|^2$ is the Lagrange multiplier corresponding to the pointwise constraint $|\mathbf{u}| = 1$. Note carefully that for a single, integral constraint (§8.4.1) the Lagrange multiplier is a number, but for a pointwise constraint it is a function.

Proof. 1. Fix $\mathbf{v} \in H_0^1(U; \mathbb{R}^m) \cap L^\infty(U; \mathbb{R}^m)$. Then since $|\mathbf{u}| = 1$ a.e., we have

$$|\mathbf{u} + \tau \mathbf{v}| \neq 0 \text{ a.e.}$$

for each sufficiently small τ . Consequently

$$(35) \quad \mathbf{v}(\tau) := \frac{\mathbf{u} + \tau \mathbf{v}}{|\mathbf{u} + \tau \mathbf{v}|} \in \mathcal{A}.$$

Thus

$$i(\tau) := I[\mathbf{v}(\tau)]$$

has a minimum at $\tau = 0$, and so, as usual,

$$(36) \quad i'(0) = 0.$$

2. Now

$$(37) \quad i'(0) = \int_U D\mathbf{u} : D\mathbf{v}'(0) \, dx \quad \left(' = \frac{d}{d\tau} \right).$$

But we compute directly from (35) that

$$\mathbf{v}'(\tau) = \frac{\mathbf{v}}{|\mathbf{u} + \tau \mathbf{v}|} - \frac{[(\mathbf{u} + \tau \mathbf{v}) \cdot \mathbf{v}](\mathbf{u} + \tau \mathbf{v})}{|\mathbf{u} + \tau \mathbf{v}|^3},$$

whence $\mathbf{v}'(0) = \mathbf{v} - (\mathbf{u} \cdot \mathbf{v})\mathbf{u}$. Inserting this equality into (36), (37), we find

$$(38) \quad 0 = \int_U D\mathbf{u} : D\mathbf{v} - D\mathbf{u} : D((\mathbf{u} \cdot \mathbf{v})\mathbf{u}) \, dx.$$

However since $|\mathbf{u}|^2 \equiv 1$, we have

$$(D\mathbf{u})^T \mathbf{u} = \mathbf{0}.$$

Using this fact, we then verify

$$D\mathbf{u} : D((\mathbf{u} \cdot \mathbf{v})\mathbf{u}) = |D\mathbf{u}|^2(\mathbf{u} \cdot \mathbf{v}) \quad \text{a.e. in } U.$$

This identity employed in (38) gives (33). \square

8.4.4. Incompressibility.

a. Stokes' problem. Suppose $U \subset \mathbb{R}^3$ is open, bounded, simply connected, and set

$$I[\mathbf{w}] := \int_U \frac{1}{2} |D\mathbf{w}|^2 - \mathbf{f} \cdot \mathbf{w} \, dx,$$

for \mathbf{w} belonging to

$$\mathcal{A} := \{\mathbf{w} \in H_0^1(U; \mathbb{R}^3) \mid \operatorname{div} \mathbf{w} = 0 \text{ in } U\}.$$

Here $\mathbf{f} \in L^2(U; \mathbb{R}^3)$ is given.

There is no problem in showing by customary methods that there exists a unique minimizer $\mathbf{u} \in \mathcal{A}$. We interpret \mathbf{u} as representing the velocity field of a steady fluid flow within the region U , subject to the external force \mathbf{f} . The constraint that $\operatorname{div} \mathbf{u} = 0$ ensures that the flow is incompressible: see the Remark at the end of this subsection.

How does the constraint manifest itself in the Euler–Lagrange equation?

THEOREM 6 (Pressure as Lagrange multiplier). *There exists a scalar function $p \in L_{\text{loc}}^2(U)$ such that*

$$(39) \quad \int_U D\mathbf{u} : D\mathbf{v} \, dx = \int_U p \operatorname{div} \mathbf{v} + \mathbf{f} \cdot \mathbf{v} \, dx$$

for all $\mathbf{v} \in H^1(U; \mathbb{R}^3)$ with compact support within U .

We interpret (39) as saying that (\mathbf{u}, p) form a weak solution of *Stokes' problem*

$$(40) \quad \begin{cases} -\Delta \mathbf{u} = \mathbf{f} - Dp & \text{in } U \\ \operatorname{div} \mathbf{u} = 0 & \text{in } U \\ \mathbf{u} = \mathbf{0} & \text{on } \partial U. \end{cases}$$

The function p is the *pressure* and arises as a Lagrange multiplier corresponding to the *incompressibility condition* $\operatorname{div} \mathbf{u} = 0$.

Proof. 1. Assume first $\mathbf{v} \in \mathcal{A}$. Then for each $\tau \in \mathbb{R}$, $\mathbf{u} + \tau\mathbf{v} \in \mathcal{A}$. Thus

$$(41) \quad 0 = i'(0) = \int_U D\mathbf{u} : D\mathbf{v} - \mathbf{f} \cdot \mathbf{v} \, dx.$$

2. Fix now $V \subset\subset U$, V smooth and simply connected, and select $\mathbf{w} \in H_0^1(V; \mathbb{R}^3)$ with $\operatorname{div} \mathbf{w} = 0$. Choose $0 < \epsilon < \operatorname{dist}(V, \partial U)$ and set $\mathbf{v} = \mathbf{v}^\epsilon := \eta_\epsilon * \mathbf{w}$ in (41), η_ϵ denoting the usual mollifier and \mathbf{w} defined to be zero in $U - V$. Then

$$(42) \quad 0 = \int_U D\mathbf{u} : D\mathbf{v}^\epsilon - \mathbf{f} \cdot \mathbf{v}^\epsilon \, dx = \int_U D\mathbf{u}^\epsilon : D\mathbf{w} - \mathbf{f}^\epsilon \cdot \mathbf{w} \, dx$$

for

$$(43) \quad \mathbf{u}^\epsilon := \eta_\epsilon * \mathbf{u}, \quad \mathbf{f}^\epsilon := \eta_\epsilon * \mathbf{f}.$$

As \mathbf{u}^ϵ is smooth, (42) implies

$$(44) \quad \int_V (-\Delta \mathbf{u}^\epsilon - \mathbf{f}^\epsilon) \cdot \mathbf{w} \, dx = 0$$

for each $\mathbf{w} \in H_0^1(V; \mathbb{R}^3)$ with $\operatorname{div} \mathbf{w} = 0$.

3. Fix any smooth vector field $\zeta \in C_c^\infty(V; \mathbb{R}^3)$ and put $\mathbf{w} = \operatorname{curl} \zeta$ in (44). This is legitimate since $\operatorname{div} \mathbf{w} = \operatorname{div}(\operatorname{curl} \zeta) = 0$. Then, temporarily writing $\mathbf{h} := \Delta \mathbf{u}^\epsilon + \mathbf{f}^\epsilon$, we find

$$0 = \int_V \mathbf{h} \cdot \operatorname{curl} \zeta \, dx = \int_V h^1(\zeta_{x_2}^3 - \zeta_{x_3}^2) + h^2(\zeta_{x_3}^1 - \zeta_{x_1}^3) + h^3(\zeta_{x_1}^2 - \zeta_{x_2}^1) \, dx,$$

for $\mathbf{h} = (h^1, h^2, h^3)$, $\zeta = (\zeta^1, \zeta^2, \zeta^3)$. An integration by parts reveals

$$0 = \int_V \zeta^1(h_{x_2}^3 - h_{x_3}^2) + \zeta^2(h_{x_3}^1 - h_{x_1}^3) + \zeta^3(h_{x_1}^2 - h_{x_2}^1) \, dx.$$

As $\zeta^1, \zeta^2, \zeta^3 \in C_c^\infty(V)$ are arbitrary, we deduce $\operatorname{curl} \mathbf{h} = 0$ in V . Since V is simply connected, there consequently exists a smooth function p^ϵ in V such that

$$(45) \quad Dp^\epsilon = \mathbf{h} = \Delta \mathbf{u}^\epsilon + \mathbf{f}^\epsilon \quad \text{in } V.$$

4. If necessary we can add a constant to p^ϵ to ensure $\int_V p^\epsilon \, dx = 0$.

In view of this normalization, there exists a smooth vector field $\mathbf{v}^\epsilon : V \rightarrow \mathbb{R}^3$ solving

$$(46) \quad \begin{cases} \operatorname{div} \mathbf{v}^\epsilon = p^\epsilon & \text{in } V \\ \mathbf{v}^\epsilon = 0 & \text{on } \partial V. \end{cases}$$

In addition we have the estimate

$$(47) \quad \|\mathbf{v}^\epsilon\|_{H^1(V;\mathbb{R}^3)} \leq C\|p^\epsilon\|_{L^2(V)},$$

the constant C depending only on V . (We omit the proof of the existence of the vector field \mathbf{v}^ϵ . The construction both is intricate and requires knowledge of certain estimates for Laplace's equation with Neumann-type boundary conditions beyond the scope of this book. See Dacorogna–Moser (Ann. Inst. H Poincaré 7 (1990), 1–26) for details.)

Now compute

$$\begin{aligned} \int_V (p^\epsilon)^2 dx &= \int_V p^\epsilon \operatorname{div} \mathbf{v}^\epsilon dx \quad \text{by (46)} \\ &= - \int_V Dp^\epsilon \cdot \mathbf{v}^\epsilon dx \\ &= \int_V (-\Delta \mathbf{u}^\epsilon - \mathbf{f}^\epsilon) \cdot \mathbf{v}^\epsilon dx \quad \text{by (45)} \\ &= \int_V D\mathbf{u}^\epsilon : D\mathbf{v}^\epsilon - \mathbf{f}^\epsilon \cdot \mathbf{v}^\epsilon dx \\ &\leq \|\mathbf{v}^\epsilon\|_{H^1(V;\mathbb{R}^3)} (\|\mathbf{u}^\epsilon\|_{H^1(V)} + \|\mathbf{f}^\epsilon\|_{L^2(V)}) \\ &\leq C\|p^\epsilon\|_{L^2(V)} (\|\mathbf{u}\|_{H^1(U)} + \|\mathbf{f}\|_{L^2(U)}) \quad \text{by (47)}. \end{aligned}$$

Thus

$$(48) \quad \|p^\epsilon\|_{L^2(V)} \leq C (\|\mathbf{u}\|_{H^1(U)} + \|\mathbf{f}\|_{L^2(U)}).$$

5. In view of estimate (48) there exists a subsequence $\epsilon_j \rightarrow 0$ so that

$$(49) \quad p^{\epsilon_j} \rightharpoonup p \quad \text{weakly in } L^2(V)$$

for some $p \in L^2(V)$. Now (45) implies

$$\int_V D\mathbf{u}^\epsilon : D\mathbf{v} dx = \int_V p^\epsilon \operatorname{div} \mathbf{v} + \mathbf{f}^\epsilon \cdot \mathbf{v} dx$$

for all $\mathbf{v} \in H_0^1(V;\mathbb{R}^3)$. Sending $\epsilon = \epsilon_j \rightarrow 0$, we find

$$(50) \quad \int_V D\mathbf{u} : D\mathbf{v} dx = \int_V p \operatorname{div} \mathbf{v} + \mathbf{f} \cdot \mathbf{v} dx$$

as well.

6. Finally choose a sequence of sets $V_k \subset\subset U$ ($k = 1, \dots$) as above, with $V_1 \subset V_2 \subset V_3 \subset \dots$ and $U = \bigcup_{k=1}^\infty V_k$. Utilizing steps 2–5, we find $p_k \in L^2(V_k)$ ($k = 1, \dots$) so that

$$(51) \quad \int_{V_k} D\mathbf{u} : D\mathbf{v} dx = \int_{V_k} p_k \operatorname{div} \mathbf{v} + \mathbf{f} \cdot \mathbf{v} dx$$

for each $\mathbf{v} \in H_0^1(V_k;\mathbb{R}^3)$. Adding constants as necessary to each p_k , we deduce from (51) that if $1 \leq l \leq k$, then $p_k = p_l$ on V_l . We finally define $p = p_k$ on V_k ($k = 1, \dots$). \square

b. Incompressible nonlinear elasticity. We return now to the model of nonlinear elasticity discussed before in §8.2.4. Suppose that \mathbf{u} represents the displacement of an elastic body which has the rest configuration U . Let us suppose now that the elastic body is incompressible, which now means

$$\det D\mathbf{u} = 1.$$

We therefore suppose the energy density function $L : \mathbb{M}^{3 \times 3} \times U \rightarrow \mathbb{R}$ is given and consider the problem of minimizing the elastic energy

$$I[\mathbf{w}] := \int_U L(D\mathbf{w}, x) dx$$

over all \mathbf{w} in the admissible set

$$\mathcal{A} := \{\mathbf{w} \in W^{1,q}(U; \mathbb{R}^3) \mid \mathbf{w} = \mathbf{g} \text{ on } \partial U, \det D\mathbf{w} = 1 \text{ a.e.}\}$$

for some $q > 3$.

THEOREM 7 (Minimizers with determinant constraint). *Assume the mapping*

$$P \mapsto L(P, x)$$

is convex and L satisfies the coercivity condition

$$L(P, x) \geq \alpha |P|^q - \beta \quad (P \in \mathbb{M}^{3 \times 3}, x \in U)$$

for some $\alpha > 0$, $\beta \geq 0$. Suppose finally $\mathcal{A} \neq \emptyset$.

Then there exists $\mathbf{u} \in \mathcal{A}$ satisfying

$$I[\mathbf{u}] = \min_{\mathbf{w} \in \mathcal{A}} I[\mathbf{w}].$$

Proof. We as usual select a minimizing sequence, with

$$\mathbf{u}_{k_j} \rightharpoonup \mathbf{u} \quad \text{weakly in } W^{1,q}(U; \mathbb{R}^3).$$

Since

$$I[\mathbf{u}] \leq \liminf_{j \rightarrow \infty} I[\mathbf{u}_{k_j}],$$

we must only show that $\mathbf{u} \in \mathcal{A}$. However, since in view of the lemma in §8.2.4 we have $\det D\mathbf{u}_{k_j} \rightharpoonup \det D\mathbf{u}$ weakly in $L^{q/n}(U)$, we see that $\det D\mathbf{u} = 1$ a.e., as required. \square

Remark on incompressibility. It may seem odd that the incompressibility condition for Stokes' problem is

$$(52) \quad \operatorname{div} \mathbf{u} = 0$$

and for the nonlinear elasticity problem is

$$(53) \quad \det D\mathbf{u} = 1.$$

The explanation is that \mathbf{u} represents a velocity in (52) and a displacement in (53). More generally if \mathbf{b} is a velocity field, say of a fluid, we compute the motion of a particle initially at a point x by solving the ODE

$$\begin{cases} \dot{\mathbf{x}}(t) = \mathbf{b}(\mathbf{x}(t), t) & (t \in \mathbb{R}) \\ \mathbf{x}(0) = x. \end{cases}$$

Write $\mathbf{x}(t) = \mathbf{x}(t, x)$ to display the dependence on the initial position x . Then for each $t > 0$, the mapping $x \mapsto \mathbf{x}(t, x)$ is volume preserving if

$$J(x, t) = \det D_x \mathbf{x}(t, x) = 1 \quad \text{for all } x.$$

Clearly $J(x, 0) = 1$, and a calculation verifies *Euler's formula*:

$$J_t = (\operatorname{div} \mathbf{b})(\mathbf{x}, t)J,$$

the divergence taken with respect to the spatial variables. Hence if $\operatorname{div} \mathbf{b} \equiv 0$, the flow is volume preserving.

8.5. CRITICAL POINTS

Thus far we have studied the problem of locating minimizers of various energy functionals, subject perhaps to constraints, and of discovering the appropriate nonlinear Euler–Lagrange equations they satisfy. For this section we turn our attention to the problem of finding additional solutions of the Euler–Lagrange PDE, by looking for other *critical points*. These critical points will not in general be minimizers, but rather “saddle points” of $I[\cdot]$.

8.5.1. Mountain Pass Theorem.

We develop next some machinery that ensures that an abstract functional $I[\cdot]$ has a critical point.

a. Critical points, deformations. Hereafter H denotes a real Hilbert space, with norm $\| \cdot \|$ and inner product (\cdot , \cdot) . Let $I : H \rightarrow \mathbb{R}$ be a nonlinear functional on H .

DEFINITION. We say I is differentiable at $u \in H$ if there exists $v \in H$ such that

$$(1) \quad I[w] = I[u] + (v, w - u) + o(\|w - u\|) \quad (w \in H).$$

The element v , if it exists, is unique. We then write $I'[u] = v$.

DEFINITION. We say I belongs to $C^1(H; \mathbb{R})$ if $I'[u]$ exists for each $u \in H$ and the mapping $I' : H \rightarrow H$ is continuous.

The theory we will develop below holds if $I \in C^1(H; \mathbb{R})$, but the proofs will be greatly streamlined provided we additionally assume

$$(2) \quad I' : H \rightarrow H \text{ is Lipschitz continuous on bounded subsets of } H.$$

NOTATION. (i) We denote by \mathcal{C} the collection of functions $I \in C^1(H; \mathbb{R})$ satisfying (2).

(ii) If $c \in \mathbb{R}$, we write

$$A_c := \{u \in H \mid I[u] \leq c\}, \quad K_c := \{u \in H \mid I[u] = c, I'[u] = 0\}.$$

DEFINITIONS. (i) We say $u \in H$ is a critical point if $I'[u] = 0$.

(ii) The real number c is a critical value if $K_c \neq \emptyset$.

We now want to prove that if c is *not* a critical value, we can nicely deform the set $A_{c+\epsilon}$ into $A_{c-\epsilon}$ for some $\epsilon > 0$. The idea will be to solve an appropriate ODE in H and to follow the resulting flow “downhill”. As H is generally infinite dimensional, we will need some kind of compactness condition.

DEFINITION. A functional $I \in C^1(H; \mathbb{R})$ satisfies the Palais–Smale compactness condition if each sequence $\{u_k\}_{k=1}^{\infty} \subset H$ such that

$$(i) \quad \{I[u_k]\}_{k=1}^{\infty} \text{ is bounded}$$

and

$$(ii) \quad I'[u_k] \rightarrow 0 \text{ in } H$$

is precompact in H .

THEOREM 1 (Deformation Theorem). *Assume $I \in \mathcal{C}$ satisfies the Palais–Smale condition. Suppose also*

$$(3) \quad K_c = \emptyset.$$

Then for each sufficiently small $\epsilon > 0$, there exists a constant $0 < \delta < \epsilon$ and a function

$$\eta \in C([0, 1] \times H; H)$$

such that the mappings

$$\eta_t(u) = \eta(t, u) \quad (0 \leq t \leq 1, u \in H)$$

satisfy

- (i) $\eta_0(u) = u \quad (u \in H)$,
- (ii) $\eta_1(u) = u \quad (u \notin I^{-1}[c - \epsilon, c + \epsilon])$,
- (iii) $I[\eta_t(u)] \leq I[u] \quad (u \in H, 0 \leq t \leq 1)$,
- (iv) $\eta_1(A_{c+\delta}) \subset A_{c-\delta}$.

Proof. 1. We first claim that there exist constants $0 < \sigma, \epsilon < 1$ such that

$$(4) \quad \|I'[u]\| \geq \sigma \text{ for each } u \in A_{c+\epsilon} - A_{c-\epsilon}.$$

The proof is by contradiction. Were (4) false for all constants $\sigma, \epsilon > 0$, there would exist sequences $\sigma_k \rightarrow 0$, $\epsilon_k \rightarrow 0$ and elements

$$(5) \quad u_k \in A_{c+\epsilon_k} - A_{c-\epsilon_k}$$

with

$$(6) \quad \|I'[u_k]\| \leq \sigma_k.$$

According to the Palais–Smale condition, there is a subsequence $\{u_{k_j}\}_{j=1}^{\infty}$ and an element $u \in H$ with $u_{k_j} \rightarrow u$ in H . But then since $I \in C^1(H; \mathbb{R})$, (5) and (6) imply $I[u] = c$, $I'[u] = 0$. Consequently $K_c \neq \emptyset$, a contradiction to our hypothesis (3).

2. Now fix δ to satisfy

$$(7) \quad 0 < \delta < \epsilon, \quad 0 < \delta < \frac{\sigma^2}{2}.$$

Write

$$A := \{u \in H \mid I[u] \leq c - \epsilon \text{ or } I[u] \geq c + \epsilon\},$$

$$B := \{u \in H \mid c - \delta \leq I[u] \leq c + \delta\}.$$

Since I' is bounded on bounded sets, we verify that the mapping $u \mapsto \text{dist}(u, A) + \text{dist}(u, B)$ is bounded below by a positive constant on each bounded subset of H . Consequently, the function

$$g(u) := \frac{\text{dist}(u, A)}{\text{dist}(u, A) + \text{dist}(u, B)} \quad (u \in H)$$

is Lipschitz continuous on bounded sets and satisfies

$$(8) \quad 0 \leq g \leq 1, \quad g = 0 \text{ on } A, \quad g = 1 \text{ on } B.$$

Set

$$(9) \quad h(t) := \begin{cases} 1, & 0 \leq t \leq 1 \\ 1/t, & t \geq 1. \end{cases}$$

Finally define the mapping $V : H \rightarrow H$ by

$$(10) \quad V(u) := -g(u)h(\|I'[u]\|)I'[u] \quad (u \in H).$$

Observe that V is bounded.

3. Consider now for each $u \in H$ the ODE

$$(11) \quad \begin{cases} \frac{d\eta}{dt}(t) = V(\eta(t)) & (t > 0) \\ \eta(0) = u. \end{cases}$$

As V is bounded and Lipschitz continuous on bounded sets, there is a unique solution, existing for all times $t \geq 0$. We write $\eta = \eta(t, u) = \eta_t(u)$ ($t \geq 0, u \in H$) to display the dependence of the solution on both the time t and the initial position $u \in H$. Restricting ourselves to times $0 \leq t \leq 1$, we see that the mapping $\eta \in C([0, 1] \times H; H)$ so defined satisfies assertions (i) and (ii).

4. We now compute

$$(12) \quad \begin{aligned} \frac{d}{dt}I[\eta_t(u)] &= \left(I'[\eta_t(u)], \frac{d}{dt}\eta_t(u) \right) \\ &= (I'[\eta_t(u)], V(\eta_t(u))) \\ &= -g(\eta_t(u))h(\|I'[\eta_t(u)]\|) \|I'[\eta_t(u)]\|^2. \end{aligned}$$

In particular

$$\frac{d}{dt}I[\eta_t(u)] \leq 0 \quad (u \in H, 0 \leq t \leq 1),$$

and so assertion (iii) is valid.

5. Now fix any point

$$(13) \quad u \in A_{c+\delta}.$$

We want to prove

$$(14) \quad \eta_1(u) \in A_{c-\delta}$$

and thereby verify assertion (iv). If $\eta_t(u) \notin B$ for some $0 \leq t \leq 1$, we are done; and so we may as well suppose instead $\eta_t(u) \in B$ ($0 \leq t \leq 1$). Then $g(\eta_t(u)) = 1$ ($0 \leq t \leq 1$). Consequently, calculation (12) yields

$$(15) \quad \frac{d}{dt}I[\eta_t(u)] = -h(\|I'[\eta_t(u)]\|)\|I'[\eta_t(u)]\|^2.$$

Now if $\|I'[\eta_t(u)]\| \geq 1$, then (9) and (4) imply

$$\frac{d}{dt}I[\eta_t(u)] = -\|I'[\eta_t(u)]\|^2 \leq -\sigma^2.$$

On the other hand, if $\|I'[\eta_t(u)]\| \leq 1$, (9) and (4) yield

$$\frac{d}{dt}I[\eta_t(u)] \leq -\sigma^2.$$

Both these inequalities, and (15), then imply

$$I[\eta_1(u)] \leq I[u] - \sigma^2 \leq c + \delta - \sigma^2 \leq c - \delta \quad \text{by (7).}$$

This estimate establishes (14) and completes the proof. \square

b. Mountain Pass Theorem. Next we employ an interesting “min-max” technique, using the deformation η built above to deduce the existence of a critical point.

THEOREM 2 (Mountain Pass Theorem). *Assume $I \in \mathcal{C}$ satisfies the Palais–Smale condition. Suppose also*

- (i) $I[0] = 0$,
- (ii) *there exist constants $r, a > 0$ such that*

$$I[u] \geq a \quad \text{if} \quad \|u\| = r,$$

and

- (iii) *there exists an element $v \in H$ with*

$$\|v\| > r, \quad I[v] \leq 0.$$

Define

$$\Gamma := \{\mathbf{g} \in C([0, 1]; H) \mid \mathbf{g}(0) = 0, \mathbf{g}(1) = v\}.$$

Then

$$c = \inf_{\mathbf{g} \in \Gamma} \max_{0 \leq t \leq 1} I[\mathbf{g}(t)]$$

is a critical value of I .

Think of the graph of $I[\cdot]$ as a landscape with a low spot at 0, surrounded by a ring of mountains. Beyond these mountains lies another low spot at v . The idea is to look for a path \mathbf{g} connecting 0 to v , which passes through a mountain pass, that is, a saddle point for $I[\cdot]$. But note carefully: we are only asserting the existence of a critical point at the “energy level” c , which may not necessarily correspond to a true saddle point.

Proof. Clearly

$$(16) \quad c \geq a.$$

Assume that c is not a critical value of I , so that

$$(17) \quad K_c = \emptyset.$$

Choose then any sufficiently small number

$$(18) \quad 0 < \epsilon < \frac{a}{2}.$$

According to the Deformation Theorem 1, there exist a constant $0 < \delta < \epsilon$ and a homeomorphism $\eta : H \rightarrow H$ with

$$(19) \quad \eta(A_{c+\delta}) \subset A_{c-\delta}$$

and

$$(20) \quad \eta(u) = u \quad \text{if} \quad u \notin I^{-1}[c - \epsilon, c + \epsilon].$$

Now select $\mathbf{g} \in \Gamma$ satisfying

$$(21) \quad \max_{0 \leq t \leq 1} I[\mathbf{g}(t)] \leq c + \delta.$$

Then $\hat{\mathbf{g}} := \eta \circ \mathbf{g}$ also belongs to Γ , since $\eta(\mathbf{g}(0)) = \eta(0) = 0$ and $\eta(\mathbf{g}(1)) = \eta(v) = v$, according to (20). But then (21) implies $\max_{0 \leq t \leq 1} I[\hat{\mathbf{g}}(t)] \leq c - \delta$, whence $c = \inf_{\mathbf{g} \in \Gamma} \max_{0 \leq t \leq 1} I[\mathbf{g}(t)] \leq c - \delta$, a contradiction. \square

8.5.2. Application to semilinear elliptic PDE.

To illustrate the utility of the Mountain Pass Theorem, let us investigate now the semilinear boundary-value problem:

$$(22) \quad \begin{cases} -\Delta u = f(u) & \text{in } U \\ u = 0 & \text{on } \partial U. \end{cases}$$

We assume f is smooth, and for some

$$1 < p < \frac{n+2}{n-2}$$

we have

$$(23) \quad |f(z)| \leq C(1 + |z|^p), \quad |f'(z)| \leq C(1 + |z|^{p-1}) \quad (z \in \mathbb{R}),$$

where C is a constant. We will suppose also

$$(24) \quad 0 \leq F(z) \leq \gamma f(z)z \quad \text{for some constant } \gamma < \frac{1}{2},$$

where $F(z) := \int_0^z f(s)ds$ and $z \in \mathbb{R}$. We hypothesize finally for constants $0 < a \leq A$ that

$$(25) \quad a|z|^{p+1} \leq |F(z)| \leq A|z|^{p+1} \quad (z \in \mathbb{R}).$$

Now (25) implies $f(0) = 0$, and so obviously $u \equiv 0$ is a trivial solution of (22). We want to find another.

Observe that the PDE

$$-\Delta u = |u|^{p-1}u$$

falls under the hypotheses above. We will return to this particular nonlinearity again in §9.4.2.

THEOREM 3 (Existence). *The boundary-value problem (22) has at least one weak solution $u \neq 0$.*

Proof. 1. Define

$$(26) \quad I[u] := \int_U \frac{1}{2}|Du|^2 - F(u) \, dx$$

for $u \in H_0^1(U)$. We intend to apply the Mountain Pass Theorem to $I[\cdot]$.

We set $H = H_0^1(U)$, with the norm $\|u\| = \left(\int_U |Du|^2 dx\right)^{1/2}$ and inner product $(u, v) = \int_U Du \cdot Dv dx$. Then

$$I[u] = \frac{1}{2}\|u\|^2 - \int_U F(u) dx =: I_1[u] - I_2[u].$$

2. We first claim

$$(27) \quad I \text{ belongs to the class } \mathcal{C}.$$

To see this, note first that for each $u, w \in H$,

$$I_1[w] = \frac{1}{2}\|w\|^2 = \frac{1}{2}\|u + w - u\|^2 = \frac{1}{2}\|u\|^2 + (u, w - u) + \frac{1}{2}\|w - u\|^2.$$

Hence I_1 is differentiable at u , with $I_1'[u] = u$. Consequently, $I_1 \in \mathcal{C}$.

3. We must next examine the term I_2 . Recall from the Lax–Milgram Theorem (§6.2.1) that for each element $v^* \in H^{-1}(U)$, the problem

$$\begin{cases} -\Delta v = v^* & \text{in } U \\ v = 0 & \text{on } \partial U \end{cases}$$

has a unique solution $v \in H_0^1(U)$. We will write $v = Kv^*$, so that

$$(28) \quad K : H^{-1}(U) \rightarrow H_0^1(U) \text{ is an isometry.}$$

Note in particular that if $w \in L^{2n/n+2}(U)$, then the linear functional w^* defined by

$$\langle w^*, u \rangle := \int_U wu dx \quad (u \in H_0^1(U))$$

belongs to $H^{-1}(U)$. (We will misuse notation and say “ $w \in H^{-1}(U)$ ”.) Observe next that $p\left(\frac{2n}{n+2}\right) < \frac{n+2}{n-2} \cdot \frac{2n}{n+2} = 2^*$, and so $f(u) \in L^{2n/n+2}(U) \subset H^{-1}(U)$ if $u \in H_0^1(U)$.

We now demonstrate that if $u \in H_0^1(U)$, then

$$(29) \quad I_2'[u] = K[f(u)].$$

To see this, note first that

$$F(a+b) = F(a) + f(a)b + \int_0^1 (1-s)f'(a+sb) ds b^2.$$

Thus for each $w \in H_0^1(U)$,

$$\begin{aligned} (30) \quad I_2[w] &= \int_U F(w) dx = \int_U F(u + w - u) dx \\ &= \int_U F(u) + f(u)(w - u) dx + R \\ &= I_2(u) + \int_U DK[f(u)] \cdot D(w - u) dx + R, \end{aligned}$$

where the remainder term R satisfies, according to (23),

$$\begin{aligned} |R| &\leq C \int_U (1 + |u|^{p-1} + |w-u|^{p-1}) |w-u|^2 dx \\ &\leq C \left(\int_U |w-u|^2 + |w-u|^{p+1} dx \right) \\ &\quad + C \left(\int_U |u|^{p+1} dx \right)^{\frac{p-1}{p+1}} \left(\int_U |w-u|^{p+1} dx \right)^{\frac{2}{p+1}}. \end{aligned}$$

Since $p+1 < 2^*$, the Sobolev inequalities show $R = o(\|w-u\|)$. Thus we see from (28) that

$$I_2[w] = I_2[u] + (K[f(u)], w-u) + o(\|w-u\|),$$

as required.

Finally we note that if $u, \bar{u} \in H_0^1(U)$ with $\|u\|, \|\bar{u}\| \leq L$, then

$$\begin{aligned} \|I_2'[u] - I_2'[\bar{u}]\| &= \|K[f(u)] - K[f(\bar{u})]\|_{H_0^1(U)} \\ &= \|f(u) - f(\bar{u})\|_{H^{-1}(U)} \\ &\leq \|f(u) - f(\bar{u})\|_{L^{\frac{2n}{n+2}}(U)}. \end{aligned}$$

But

$$\begin{aligned} \|f(u) - f(\bar{u})\|_{L^{\frac{2n}{n+2}}(U)} &\leq C \left(\int_U ((1 + |u|^{p-1} + |\bar{u}|^{p-1}) |u - \bar{u}|)^{\frac{2n}{n+2}} dx \right)^{\frac{n+2}{2n}} \\ &\leq C \left(\int_U (1 + |u|^{p-1} + |\bar{u}|^{p-1})^{\frac{2n}{n+2} \frac{n+2}{4}} dx \right)^{\frac{2}{n}} \|u - \bar{u}\|_{L^{2^*}(U)} \\ &\leq C(L) \|u - \bar{u}\|_{L^{2^*}(U)} \leq C(L) \|u - \bar{u}\|, \end{aligned}$$

where we used (23). Thus $I_2' : H_0^1(U) \rightarrow H_0^1(U)$ is Lipschitz continuous on bounded sets. Consequently $I_2 \in \mathcal{C}$, and we have established assertion (27).

4. Now we verify the Palais–Smale condition. For this suppose $\{u_k\}_{k=1}^\infty \subset H_0^1(U)$, with

$$(31) \quad \{I[u_k]\}_{k=1}^\infty \text{ bounded}$$

and

$$(32) \quad I'[u_k] \rightarrow 0 \quad \text{in } H_0^1(U).$$

According to the foregoing

$$(33) \quad u_k - K(f(u_k)) \rightarrow 0 \quad \text{in } H_0^1(U).$$

Thus for each $\epsilon > 0$ we have

$$|(I'[u_k], v)| = \left| \int_U Du_k \cdot Dv - f(u_k)v \, dx \right| \leq \epsilon \|v\| \quad (v \in H_0^1(U))$$

for k large enough. Let $v = u_k$ above to find

$$\left| \int_U |Du_k|^2 - f(u_k)u_k \, dx \right| \leq \epsilon \|u_k\|$$

for each $\epsilon > 0$ and for all k sufficiently large. For $\epsilon = 1$ in particular, we see that

$$(34) \quad \int_U f(u_k)u_k \, dx \leq \|u_k\|^2 + \|u_k\|$$

for all k sufficiently large. But since (31) says

$$\left(\frac{1}{2} \|u_k\|^2 - \int_U F(u_k) \, dx \right) \leq C < \infty$$

for all k and some constant C , we deduce

$$\begin{aligned} \|u_k\|^2 &\leq C + 2 \int_U F(u_k) \, dx \\ &\leq C + 2\gamma (\|u_k\|^2 + \|u_k\|) \quad \text{by (34), (24)}. \end{aligned}$$

Since $2\gamma < 1$, we discover that $\{u_k\}_{k=1}^\infty$ is bounded in $H_0^1(U)$. Hence there exists a subsequence $\{u_{k_j}\}_{j=1}^\infty$ and $u \in H_0^1(U)$, with $u_{k_j} \rightharpoonup u$ weakly in $H_0^1(U)$ and $u_{k_j} \rightarrow u$ in $L^{p+1}(U)$, the latter assertion holding since $p+1 < 2^*$. But then $f(u_k) \rightarrow f(u)$ in $H^{-1}(U)$, whence $K[f(u_k)] \rightarrow K[f(u)]$ in $H_0^1(U)$. Consequently (33) implies

$$(35) \quad u_{k_j} \rightarrow u \quad \text{in } H_0^1(U).$$

5. We finally verify the remaining hypotheses of the Mountain Pass Theorem. Clearly $I[0] = 0$. Suppose now that $u \in H_0^1(U)$, with $\|u\| = r$, for $r > 0$ to be selected below. Then

$$(36) \quad I[u] = I_1[u] - I_2[u] = \frac{r^2}{2} - I_2[u].$$

Now hypothesis (25) implies, since $p + 1 < 2^*$, that

$$\begin{aligned} |I_2[u]| &\leq C \int_U |u|^{p+1} dx \leq C \left(\int_U |u|^{2^*} dx \right)^{\frac{p+1}{2^*}} \\ &\leq C \|u\|^{p+1} \leq Cr^{p+1}. \end{aligned}$$

In view of (36), then

$$I[u] \geq \frac{r^2}{2} - Cr^{p+1} \geq \frac{r^2}{4} = a > 0,$$

provided $r > 0$ is small enough, since $p + 1 > 2$. Now fix some element $u \in H$, $u \neq 0$. Write $v := tu$ for $t > 0$ to be selected. Then

$$\begin{aligned} I[v] &= I_1[tu] - I_2[tu] \\ &= t^2 I_1[u] - \int_U F(tu) dx \\ &\leq t^2 I_1[u] - at^{p+1} \int_U |u|^{p+1} dx \quad \text{by (25)} \\ &< 0 \end{aligned}$$

for $t > 0$ large enough.

6. We have at last checked all the hypotheses of the Mountain Pass Theorem. There must consequently exist a function $u \in H_0^1(U)$, $u \neq 0$, with

$$I'[u] = u - K[f(u)] = 0.$$

In particular for each $v \in H_0^1(U)$, we have

$$\int_U Du \cdot Dv dx = \int_U f(u)v dx,$$

and so u is a weak solution of (22). □

See §9.4.2 for further discussion about nonlinear Poisson equations and in particular the significance of the *critical exponent* $\frac{n+2}{n-2}$ in hypothesis (23).

8.6. INVARIANCE, NOETHER'S THEOREM

Next we study variational integrands that are invariant under appropriate domain and function variations and show that solutions of the corresponding

Euler–Lagrange equations then automatically solve also certain divergence structure conservation laws.

8.6.1. Invariant variational problems.

We again turn our attention to the functional

$$(1) \quad I[w] := \int_U L(Dw, w, x) dx,$$

where $U \subset \mathbb{R}^n$ and $w : U \rightarrow \mathbb{R}$. We as usual write $L = L(p, z, x)$.

NOTATION. (i) Let $\mathbf{x} : \mathbb{R}^n \times \mathbb{R} \rightarrow \mathbb{R}^n$, $\mathbf{x} = \mathbf{x}(x, \tau)$, be a smooth family of vector fields satisfying $\mathbf{x}(x, 0) = x$ for all $x \in \mathbb{R}^n$. Then for small $|\tau|$, the mapping $x \mapsto \mathbf{x}(x, \tau)$ is a smooth diffeomorphism. We call the mapping $x \mapsto \mathbf{x}(x, \tau)$ a *domain variation*. Define also

$$(2) \quad \mathbf{v}(x) := \mathbf{x}_\tau(x, 0)$$

and

$$(3) \quad U(\tau) := \mathbf{x}(U, \tau).$$

(ii) Next, given a smooth $u : \mathbb{R}^n \rightarrow \mathbb{R}$, we consider a smooth family of *function variations* $w : \mathbb{R}^n \times \mathbb{R} \rightarrow \mathbb{R}$, $w = w(x, \tau)$, such that

$$(4) \quad w(x, 0) = u(x).$$

Write

$$(5) \quad m(x) := w_\tau(x, 0).$$

For reasons that will be clear shortly, we call m a *multiplier*.

Given a functional $I[\cdot]$ of the form (1), we ask if we can find domain and function variations that are compatible with the Lagrangian L , in the sense that $I[\cdot]$ is unchanged under these variations.

DEFINITION. We say that the functional $I[\cdot]$ is invariant under the domain variations \mathbf{x} and the function variations w provided

$$(6) \quad \int_U L(Dw(x, \tau), w(x, \tau), x) dx = \int_{U(\tau)} L(Du, u, x) dx$$

for all small $|\tau|$ and all open sets $U \subset \mathbb{R}^n$.

Here we write $Dw = D_x w$. The idea behind this definition is that given a domain variation \mathbf{x} and a function u , we will look for w as some expression involving $u(\mathbf{x}(x, \tau))$. We will then try to check (6) by changing variables in the integral term on the left, after which the integration will be over the region $U(\tau)$.

8.6.2. Noether's Theorem.

We now show that invariance of the functional implies that the corresponding Euler–Lagrange equation can be transformed into divergence form:

THEOREM 1 (Noether's Theorem). *Suppose that the functional $I[\cdot]$ is invariant under the domain variation \mathbf{x} and the function variation w corresponding to a smooth function u .*

(i) *Then*

$$(7) \quad \sum_{i=1}^n (mL_{p_i}(Du, u, x) - L(Du, u, x)v^i)_{x_i} = m \left(\sum_{i=1}^n (L_{p_i}(Du, u, x))_{x_i} - L_z(Du, u, x) \right),$$

where $\mathbf{v} = (v^1, \dots, v^n)$ is defined by (2) and the multiplier m by (5).

(ii) *In particular, if u a critical point of $I[\cdot]$ and so solves the Euler–Lagrange equation $-\operatorname{div}(DL_p) + L_z = 0$, we have the divergence identity*

$$(8) \quad \sum_{i=1}^n (mL_{p_i}(Du, u, x) - L(Du, u, x)v^i)_{x_i} = 0.$$

So multiplying by m converts the Euler–Lagrange PDE into divergence form.

Proof. Differentiating the invariance identity (6) with respect to τ and then setting $\tau = 0$ yields the identity

$$\int_U D_p L \cdot Dm + L_z m \, dx = \int_{\partial U} L\mathbf{v} \cdot \nu \, dS.$$

The term on the right appears owing to the formula in §C.4 for differentiating integrals over moving regions. Now an integration by parts and the Gauss–Green Theorem imply

$$\begin{aligned} \int_U (-\operatorname{div} D_p L + L_z)m \, dx &= \int_{\partial U} (-mD_p L + L\mathbf{v}) \cdot \nu \, dS \\ &= \int_U \operatorname{div}(-mD_p L + L\mathbf{v}) \, dx. \end{aligned}$$

This identity is valid for all regions U , and so the identity (7) follows. \square

As noted earlier, we can sometimes first guess a domain variation \mathbf{x} and then look for a corresponding function variation w as some formula involving $u(\mathbf{x}(x, \tau))$. Then we will be able to compute the multiplier m in terms of u and its partial derivatives. The following examples illustrate this procedure, in increasingly complicated circumstances.

Example 1 (Lagrangians independent of x). If $L = L(p, z)$ does not depend upon the independent variables x , then the integral (1) is invariant under translations in space. To be specific, select $k \in \{1, \dots, n\}$ and define

$$\mathbf{x}(x, \tau) := x + \tau e_k, \quad w(x, \tau) := u(x + \tau e_k).$$

Then

$$\mathbf{v} = e_k, \quad m = u_{x_k}.$$

Consequently if u is a critical point, (8) becomes the identity

$$(9) \quad \sum_{i=1}^n (L_{p_i} u_{x_k} - L \delta_{ik})_{x_i} = 0 \quad (k = 1, \dots, n).$$

It is a simple exercise to confirm that these formulas follow directly from the Euler–Lagrange equation. The point is that Noether’s Theorem provides a systematic procedure for searching for such identities. \square

Example 2 (Scaling invariance). The functional

$$I[w] = \int_U |Dw|^p dx,$$

smooth minimizers u of which solve the p -Laplacian equation

$$\operatorname{div}(|Du|^{p-2} Du) = 0,$$

is invariant under the scaling transformation

$$x \mapsto \lambda x, \quad u \mapsto \lambda^{\frac{n-p}{p}} u(\lambda x).$$

To be consistent with previous notation, we put $\lambda = e^\tau$ and define

$$\mathbf{x}(x, \tau) := e^\tau x, \quad w(x, \tau) := e^{\tau \frac{n-p}{p}} u(e^\tau x).$$

Then

$$\mathbf{v} = x, \quad m = Du \cdot x + \frac{n-p}{p} u.$$

The corresponding divergence identity (8) for a solution of the p -Laplacian equation is

$$(10) \quad \sum_{i=1}^n \left(\left(Du \cdot x + \frac{n-p}{p} u \right) p |Du|^{p-2} u_{x_i} - |Du|^p x_i \right)_{x_i} = 0.$$

It is again straightforward to check this identity directly.

Application: monotonicity formulas. Assume that u is a smooth solution of the p -Laplacian PDE within some region U and that the ball $B(0, r)$ lies within U . If we integrate the divergence identity (10) over $B(0, r)$ and simplify using the Gauss–Green Theorem, we discover that

$$(n-p) \int_{B(0,r)} |Du|^p dx = r \int_{\partial B(0,r)} |Du|^p - p |Du|^{p-2} u_r^2 dS,$$

where $u_r := Du \cdot \frac{x}{|x|}$. Therefore

$$\begin{aligned} & \frac{d}{dr} \left(\frac{1}{r^{n-p}} \int_{B(0,r)} |Du|^p dx \right) \\ &= \frac{p-n}{r^{n-p+1}} \int_{B(0,r)} |Du|^p dx + \frac{1}{r^{n-p}} \int_{\partial B(0,r)} |Du|^p dS \\ &= \frac{p}{r^{n-p}} \int_{\partial B(0,r)} |Du|^{p-2} u_r^2 dS \geq 0. \end{aligned}$$

This is a *monotonicity formula*, implying

$$r \mapsto \frac{1}{r^{n-p}} \int_{B(0,r)} |Du|^p dx \text{ is nondecreasing.}$$

We discuss in Problem 20 an interesting consequence for $p = 2$. □

In §9.4.2 we will learn the usefulness of the multiplier $Du \cdot x$, suggested by Example 2, in studying certain semilinear elliptic equations.

Time dependent problems. If one of the independent variables is identified with time, then we can interpret (8) as a *conservation law* resulting from the invariance of our variational integral.

Example 3 (Conservation of energy for nonlinear wave equations). Consider the integral expression

$$(11) \quad I[w] = \int_0^T \int_{\mathbb{R}^n} \frac{1}{2} w_t^2 - \left(\frac{1}{2} |Dw|^2 + F(w) \right) dx dt$$

defined for functions $w = w(x, t)$ with, say, compact support. As usual, we write $Dw = D_x w$. We can interpret the Lagrangian as representing the kinetic energy minus the potential energy, by analogy with the ODE example in §3.3.1. The corresponding Euler–Lagrange equation is the *semilinear wave equation*

$$(12) \quad u_{tt} - \Delta u + f(u) = 0$$

for $f := F'$.

The integrand of (11) does not depend on the time variable t and is consequently invariant under shifts in this variable. Noether's Theorem implies that this invariance forces a conservation law, in this case *conservation of energy*. More precisely, we define

$$\mathbf{x}(x, t, \tau) := (x, t + \tau), \quad w(x, t, \tau) := u(x, t + \tau),$$

so that

$$\mathbf{v} = e_{n+1}, \quad m = u_t.$$

Then (8) reads

$$\sum_{i=1}^n (-u_{x_i} u_t)_{x_i} + (u_t^2 - \frac{1}{2}(u_t^2 - |Du|^2) + F(u))_t = 0.$$

This can be rewritten as

$$(13) \quad e_t - \operatorname{div}(u_t Du) = 0$$

for the energy density

$$e := \frac{1}{2}(u_t^2 + |Du|^2) + F(u).$$

The divergence operator in (13) acts in the x variables only. If u has compact support in space at each time, it follows that the total energy is conserved in time:

$$\frac{d}{dt} \int_{\mathbb{R}^n} \frac{1}{2}(u_t^2 + |Du|^2) + F(u) \, dx = 0.$$

In Chapter 12 we will systematically investigate the solvability of semilinear wave equations of the form (12) and will find this conservation law extremely useful. \square

Example 4 (Scaling invariance for the wave equation). The linear wave equation

$$\square u = u_{tt} - \Delta u = 0$$

corresponds to the action functional

$$(14) \quad I[w] = \frac{1}{2} \int_0^T \int_{\mathbb{R}^n} w_t^2 - |Dw|^2 dx dt.$$

Similarly to Example 2 above, the functional (14) is invariant under the scaling transformation

$$(x, t) \mapsto (\lambda x, \lambda t), \quad u \mapsto \lambda^{\frac{n-1}{2}} u(\lambda x, \lambda t).$$

As before, we put $\lambda = e^\tau$ and define

$$\mathbf{x}(x, t, \tau) := (e^\tau x, e^\tau t), \quad w(x, t, \tau) := e^{\tau \frac{n-1}{2}} u(e^\tau x, e^\tau t).$$

Then

$$\mathbf{v} = (x, t), \quad m = tu_t + x \cdot Du + \frac{n-1}{2}u.$$

The conservation law provided by (8) asserts

$$p_t - \operatorname{div} \mathbf{q} = 0$$

for

$$p := \frac{t}{2}(u_t^2 + |Du|^2) + x \cdot Du u_t + \frac{n-1}{2}u u_t,$$

$$\mathbf{q} := \left(tu_t + x \cdot Du + \frac{n-1}{2}u \right) Du + \frac{1}{2}(u_t^2 - |Du|^2)x.$$

We will present in §12.4 an important application of a similar identity for a nonlinear wave equation. \square

Example 5 (Conformal energy for the wave equation). The following much more sophisticated example illustrates how Noether's Theorem, even when not directly applicable, can sometimes help us identify useful multipliers.

The mapping

$$(15) \quad (x, t) \mapsto (\bar{x}, \bar{t}) := \left(\frac{x}{|x|^2 - t^2}, \frac{t}{|x|^2 - t^2} \right),$$

where $|x|^2 \neq t^2$, is called *hyperbolic inversion*. Related is the *hyperbolic Kelvin transform* $\mathcal{K}u = \bar{u}$, defined by

$$(16) \quad \begin{aligned} \bar{u}(x, t) &:= u(\bar{x}, \bar{t}) \left| |\bar{x}|^2 - \bar{t}^2 \right|^{\frac{n-1}{2}} \\ &= u \left(\frac{x}{|x|^2 - t^2}, \frac{t}{|x|^2 - t^2} \right) \frac{1}{\left| |x|^2 - t^2 \right|^{\frac{n-1}{2}}}. \end{aligned}$$

An exercise for Chapter 12 shows that if $\square u = 0$, then $\square \bar{u} = 0$.

Our intention is to use hyperbolic inversion and the hyperbolic Kelvin transform to help us design variations in (x, t) and in u . For the first, let us map (x, t) to (\bar{x}, \bar{t}) , then add τe_{n+1} , and lastly apply hyperbolic inversion again. A calculation shows that the result is the mapping

$$(17) \quad \mathbf{x}(x, t, \tau) := \gamma(x, t + \tau(|x|^2 - t^2)),$$

for

$$(18) \quad \gamma := \frac{|x|^2 - t^2}{|x|^2 - (t + \tau(|x|^2 - t^2))^2}.$$

We next employ a similar procedure to build variations of u : we apply the Kelvin transform to compute $\mathcal{K}u = \bar{u}$, then add τe_{n+1} within the argument of \bar{u} , and lastly reapply the Kelvin transform. Another calculation reveals the resulting function variation to be

$$(19) \quad w(x, t, \tau) := \gamma^{\frac{n-1}{2}} u(\mathbf{x}(x, t, \tau)).$$

We next compute the multiplier corresponding to (17)-(19) by differentiating with respect to τ and then putting $\tau = 0$:

$$(20) \quad \mathbf{v} = (2xt, |x|^2 + t^2), \quad m = (|x|^2 + t^2)u_t + 2tx \cdot Du + (n-1)tu.$$

Now we do not assert that the action functional (14) is invariant under these domain and function variations. Rather, we guess that since the hyperbolic Kelvin transform preserves solutions of the wave equation, then it might be useful to multiply the wave equation $\square u = 0$ by the multiplier m from (20). This turns out to be so, and after a longish calculation we derive *Morawetz's identity*

$$(21) \quad c_t - \operatorname{div} \mathbf{r} = 0,$$

where

$$(22) \quad c := \frac{1}{2}(|x|^2 + t^2)(u_t^2 + |Du|^2) + 2tx \cdot Duu_t + (n-1)tuu_t - \frac{n-1}{2}u^2$$

is the density of the so-called *conformal energy* and

$$(23) \quad \mathbf{r} := ((|x|^2 + t^2)u_t + 2tx \cdot Du + (n-1)tu)Du + t(u_t^2 - |Du|^2)x.$$

Morawetz's identity is important since the conformal energy density c can be written for $n \neq 2$ as a sum of nonnegative terms plus a divergence in the x -variables:

$$(24) \quad c = \frac{(t + |x|)^2}{4} \left(u_t + u_r + \frac{n-1}{2|x|} u \right)^2 + \frac{(t - |x|)^2}{4} \left(u_t - u_r - \frac{n-1}{2|x|} u \right)^2 + \frac{|x|^2 + t^2}{2} \left(|Du|^2 - u_r^2 + \frac{(n-3)(n-1)}{4|x|^2} u^2 \right) - \frac{n-1}{4} \operatorname{div} \left(\frac{|x|^2 + t^2}{|x|^2} ux \right),$$

for $u_r := Du \cdot \frac{x}{|x|}$.

Application: local energy decay. Suppose now that $O \subset \mathbb{R}^n$ denotes a bounded, smooth open set that is star-shaped with respect to the origin: see §9.4.2 for the definition and geometric meaning of this condition. Define the exterior region

$$U := \mathbb{R}^n - \bar{O}.$$

Assume in addition that u is a smooth solution of this initial/boundary-value problem for the wave equation outside of the "obstacle" O :

$$\begin{cases} u_{tt} - \Delta u = 0 & \text{in } U \times (0, \infty) \\ u = 0 & \text{on } \partial U \times \{t = 0\} \\ u = g, u_t = h & \text{on } U \times \{t = 0\}, \end{cases}$$

for which the initial data g, h have compact support.

We assert that if $n \geq 3$ and if $O \subset B(0, R)$, there exists a constant C such that

$$(25) \quad \int_{B(0,R)-O} u_t^2 + |Du|^2 dx \leq \frac{C}{t^2},$$

for times $t \geq 2R$. Consequently *the energy within any bounded region decays to zero as $t \rightarrow \infty$* , although the total energy is conserved.

To prove (25), we first observe from the conservation law (21) that

$$(26) \quad \frac{d}{dt} \int_U c dx = \int_U \operatorname{div} \mathbf{r} dx = \int_{\partial O} \mathbf{r} \cdot \nu dS,$$

where ν denotes the *inward* pointing unit normal to ∂O . Now $u = u_t = 0$ on ∂O , and hence we can compute from (23) that

$$\mathbf{r} \cdot \nu = 2t(x \cdot Du)(Du \cdot \nu) - t|Du|^2(x \cdot \nu)$$

along ∂O . Since $u = 0$ on ∂O , we have $Du = (Du \cdot \nu)\nu$ there. Using this in the formula above, we deduce that

$$\mathbf{r} \cdot \nu = t|Du|^2(x \cdot \nu) \leq 0,$$

since O is star-shaped with respect to the origin and ν is the inward pointing normal.

Then (26) and our formula (24) for c imply for each time $t > 0$ that

$$\begin{aligned} & \int_{B(0,R)-O} \frac{(t+|x|)^2}{4} \left(u_t + u_r + \frac{n-1}{2|x|}u \right)^2 \\ & + \frac{(t-|x|)^2}{4} \left(u_t - u_r - \frac{n-1}{2|x|}u \right)^2 + \frac{|x|^2 + t^2}{2} (|Du|^2 - u_r^2) dx \leq C. \end{aligned}$$

Taking $t \geq 2R$ and making some simple estimates, we derive from this the estimates

$$(27) \quad \int_{B(0,R)-O} |Du|^2 - u_r^2 dx \leq \frac{C}{t^2}$$

and

$$(28) \quad \int_{B(0,R)-O} u_t^2 + u_r^2 + \frac{n-1}{|x|}uu_r + \frac{(n-1)^2}{4|x|^2}u^2 dx \leq \frac{C}{t^2}.$$

Now

$$\frac{u}{|x|}u_r = \operatorname{div} \left(\frac{u^2}{2|x|^2}x \right) - \frac{n-2}{2} \frac{u^2}{|x|^2}.$$

This identity shows that the integral of the last two terms in (28) is non-negative. The energy decay estimate (25) follows. \square

8.7. PROBLEMS

In the exercises U always denotes a bounded, open subset of \mathbb{R}^n , with smooth boundary. All given functions are assumed smooth, unless otherwise stated.

1. This problem illustrates that a weakly convergent sequence can be rather badly behaved.

- (a) Prove $u_k(x) = \sin(kx) \rightharpoonup 0$ as $k \rightarrow \infty$ in $L^2(0, 1)$.
- (b) Fix $a, b \in \mathbb{R}$, $0 < \lambda < 1$. Define

$$u_k(x) := \begin{cases} a & \text{if } j/k \leq x < (j+\lambda)/k \\ b & \text{if } (j+\lambda)/k \leq x < (j+1)/k \end{cases} \quad (j = 0, \dots, k-1).$$

Prove $u_k \rightarrow \lambda a + (1 - \lambda)b$ in $L^2(0, 1)$.

2. Find $L = L(p, z, x)$ so that the PDE

$$-\Delta u + D\phi \cdot Du = f \quad \text{in } U$$

is the Euler–Lagrange equation corresponding to the functional $I[w] := \int_U L(Dw, w, x) dx$.

(Hint: Look for a Lagrangian with an exponential term.)

3. The *elliptic regularization* of the heat equation is the PDE

$$(*) \quad u_t - \Delta u - \epsilon u_{tt} = 0 \quad \text{in } U_T,$$

where $\epsilon > 0$ and $U_T = U \times (0, T]$. Show that $(*)$ is the Euler–Lagrange equation corresponding to an energy functional $I_\epsilon[w] := \int \int_{U_T} L_\epsilon(Dw, w_t, w, x, t) dx dt$.

(Hint: Look for a Lagrangian with an exponential term involving t .)

4. Assume $\eta : \mathbb{R}^n \rightarrow \mathbb{R}$ is C^1 .

(a) Show $L(P, z, x) = \eta(z) \det P$ ($P \in \mathbb{M}^{n \times n}, z \in \mathbb{R}^n$) is a null Lagrangian.

(b) Deduce that if $\mathbf{u} : \mathbb{R}^n \rightarrow \mathbb{R}^n$ is C^2 , then

$$\int_U \eta(\mathbf{u}) \det D\mathbf{u} dx$$

depends only on $\mathbf{u}|_{\partial U}$.

5. (Continuation) Fix $x_0 \notin \mathbf{u}(\partial U)$, and choose a function η as above so that $\int_{\mathbb{R}^n} \eta dz = 1$, $\text{spt } \eta \subset B(x_0, r)$, r taken so small that $B(x_0, r) \cap \mathbf{u}(\partial U) = \emptyset$. Define

$$\text{deg}(\mathbf{u}, x_0) = \int_U \eta(\mathbf{u}) \det D\mathbf{u} dx,$$

the *degree* of \mathbf{u} relative to x_0 . Prove the degree is an integer.

6. Let $\Sigma \subset \mathbb{R}^3$ denote the graph of the smooth function $u : U \rightarrow \mathbb{R}$, $U \subset \mathbb{R}^2$. Then

$$(*) \quad \int_U (1 + |Du|^2)^{-\frac{3}{2}} \det D^2u dx$$

represents the integral of the Gauss curvature over Σ . Prove that this expression depends only upon Du restricted to ∂U . (The Gauss–Bonnet Theorem in differential geometry computes $(*)$ in terms of the geodesic curvature of $\partial\Sigma$.)

7. Let $m = n$. Prove

$$L(P) = \operatorname{tr}(P^2) - \operatorname{tr}(P)^2 \quad (P \in \mathbb{M}^{n \times n})$$

is a null Lagrangian.

8. Explain why the methods in §8.2 will not work to prove the existence of a minimizer of the functional

$$I[w] := \int_U (1 + |Dw|^2)^{1/2} dx$$

over $\mathcal{A} := \{w \in W^{1,q}(U) \mid w = g \text{ on } \partial U\}$, for any $1 \leq q < \infty$.

9. (Second variation for systems) Assume $\mathbf{u} : U \rightarrow \mathbb{R}^m$ is a smooth minimizer of the functional

$$I[\mathbf{w}] := \int_U L(D\mathbf{w}, \mathbf{w}, x) dx.$$

- (a) Show

$$\sum_{i,j=1}^n \sum_{k,l=1}^m \frac{\partial^2 L}{\partial p_i^k \partial p_j^l} (D\mathbf{u}, \mathbf{u}, x) \eta_k \eta_l \xi_i \xi_j \geq 0$$

for all $x \in U$, $\xi \in \mathbb{R}^n$, $\eta \in \mathbb{R}^m$.

- (b) Provide an example of a nonconvex function $L : \mathbb{M}^{m \times n} \rightarrow \mathbb{R}$ satisfying

$$\sum_{i,j=1}^n \sum_{k,l=1}^m \frac{\partial^2 L(P)}{\partial p_i^k \partial p_j^l} \eta_k \eta_l \xi_i \xi_j \geq 0$$

for all $P \in \mathbb{M}^{m \times n}$, $\xi \in \mathbb{R}^n$, $\eta \in \mathbb{R}^m$.

10. Use the methods of §8.4.1 to show the existence of a nontrivial weak solution $u \in H_0^1(U)$, $u \neq 0$, of

$$\begin{cases} -\Delta u = |u|^{q-1}u & \text{in } U \\ u = 0 & \text{on } \partial U \end{cases}$$

for $1 < q < \frac{n+2}{n-2}$, $n > 2$.

11. Assume $\beta : \mathbb{R} \rightarrow \mathbb{R}$ is smooth, with

$$0 < a \leq \beta'(z) \leq b \quad (z \in \mathbb{R})$$

for constants a, b . Let $f \in L^2(U)$. Formulate what it means for $u \in H^1(U)$ to be a weak solution of the nonlinear boundary-value problem

$$\begin{cases} -\Delta u = f & \text{in } U \\ \frac{\partial u}{\partial \nu} + \beta(u) = 0 & \text{on } \partial U. \end{cases}$$

Prove there exists a unique weak solution.

12. Assume u is a smooth minimizer of the area integral

$$I[w] = \int_U (1 + |Dw|^2)^{1/2} dx,$$

subject to given boundary conditions $w = g$ on ∂U and the constraint

$$J[w] = \int_U w dx = 1.$$

Prove the graph of u is a surface of constant mean curvature.

(Hint: Recall Example 4 in §8.1.2.)

13. Assume $f \in L^2(U)$. Prove the *dual variational principle* that

$$\min_{w \in H_0^1(U)} \int_U \frac{1}{2} |Dw|^2 - fw dx = \max_{\substack{\xi \in L^2(U; \mathbb{R}^n) \\ \operatorname{div} \xi = f}} -\frac{1}{2} \int_U |\xi|^2 dx.$$

14. (Multivalued PDE) Show that the variational inequality (26) for the obstacle problem in §8.4.2 can be rewritten as

$$-\Delta u + \beta(u - h) \ni f$$

for the multivalued function

$$\beta(z) := \begin{cases} 0 & \text{if } z > 0 \\ (-\infty, 0] & \text{if } z = 0 \\ \emptyset & \text{if } z < 0. \end{cases}$$

(See also Problem 3 in Chapter 9.)

15. (Pointwise gradient constraint)

(a) Show there exists a unique minimizer $u \in \mathcal{A}$ of

$$I[w] := \int_U \frac{1}{2} |Dw|^2 - fw dx,$$

where $f \in L^2(U)$ and

$$\mathcal{A} := \{w \in H_0^1(U) \mid |Dw| \leq 1 \text{ a.e.}\}.$$

(b) Prove

$$\int_U Du \cdot D(w - u) dx \geq \int_U f(w - u) dx$$

for all $w \in \mathcal{A}$.

16. Assume $n \geq 3$ and U is a bounded open set containing 0. Show that $\mathbf{u} := \frac{x}{|x|}$ belongs to $H^1(U; \mathbb{R}^n)$ and is a harmonic mapping into the sphere S^{n-1} . That is, show \mathbf{u} is a weak solution of

$$\begin{cases} -\Delta \mathbf{u} = |D\mathbf{u}|^2 \mathbf{u} \\ |\mathbf{u}| = 1 \end{cases} \quad \text{in } U.$$

17. Let $u, \hat{u} \in H_0^1(U)$ both be positive minimizers of the Dirichlet energy

$$I[w] := \int_U |Dw|^2.$$

Suppose also that $u, \hat{u} > 0$ within U . Follow the hints to give a new proof that

$$u \equiv \hat{u} \quad \text{in } U.$$

(Hint: Define $w := \left(\frac{u^2 + \hat{u}^2}{2}\right)^{1/2}$, $s := \frac{u^2}{u^2 + \hat{u}^2}$ and $\eta := \frac{u^2 + \hat{u}^2}{2}$; and show that

$$|Dw|^2 = \eta \left| s \frac{Du}{u} + (1-s) \frac{D\hat{u}}{\hat{u}} \right|^2.$$

Deduce

$$|Dw|^2 \leq \eta \left(s \left| \frac{Du}{u} \right|^2 + (1-s) \left| \frac{D\hat{u}}{\hat{u}} \right|^2 \right) = \frac{1}{2} |Du|^2 + \frac{1}{2} |D\hat{u}|^2$$

and therefore $\frac{Du}{u} = \frac{D\hat{u}}{\hat{u}}$ almost everywhere.)

(Belloni–Kawohl, Manuscripta Math. 109 (2002), 229–231)

18. Assume that a_1, a_2 are smooth, positive functions on \bar{U} such that $a_1 \leq a_2$. Let u_1, u_2 be smooth solutions of

$$\operatorname{div}(a_1 Du_1) = 0, \quad \operatorname{div}(a_2 Du_2) = 0 \quad \text{in } U$$

with $Du_2 \neq 0$ a. e. Suppose finally that

$$u_1 = u_2, \quad \frac{\partial u_1}{\partial \nu} = \frac{\partial u_2}{\partial \nu}, \quad a_1 = a_2 \quad \text{on } \partial U.$$

Prove that

$$a_1 \equiv a_2, \quad u_1 \equiv u_2 \quad \text{within } U.$$

(Hint: Observe that $\int_U a_1 |Du_1|^2 dx = \int_U a_2 |Du_2|^2 dx$.)

19. (Momentum conservation) Given a solution u of the nonlinear wave equation $\square u + f(u) = 0$, apply Noether's Theorem to the transformations $\mathbf{x}(x, t, \tau) = (x + \tau e_k, t)$, $w(x, t, \tau) = u(x + \tau e_k, t)$ to calculate the momentum density p_k and the momentum flux \mathbf{j}_k satisfying the conservation laws

$$(p_k)_t - \operatorname{div}(\mathbf{j}_k) = 0 \quad (k = 1, \dots, n).$$

20. Let u be harmonic in some region $U \subseteq \mathbb{R}^n$ and assume $B(0, R) \subset U$, $u(0) = 0$, $u \not\equiv 0$. Define for $0 < r < R$ the functions

$$a(r) := \frac{1}{r^{n-1}} \int_{\partial B(0,r)} u^2 dS, \quad b(r) := \frac{1}{r^{n-2}} \int_{B(0,r)} |Du|^2 dx.$$

We derived in §8.6.2 the monotonicity formula

$$\dot{b} = \frac{2}{r^{n-2}} \int_{\partial B(0,r)} u_r^2 dS.$$

- (a) Prove that

$$\dot{a} = \frac{2}{r^{n-1}} \int_{\partial B(0,r)} uu_r dS = \frac{2}{r} b.$$

- (b) Show

$$b^2 \leq \frac{r}{2} ab.$$

- (c) Define the *frequency function*

$$f := \frac{b}{a}$$

and derive *Almgren's monotonicity formula*: $\dot{f} \geq 0$.

- (d) Demonstrate next that $\frac{\dot{a}}{a} \leq \frac{\beta}{r}$ and consequently

$$a(r) \geq \gamma r^\beta \quad (0 < r < R)$$

for $\beta := \frac{2b(R)}{a(R)}$ and $\gamma := \frac{a(R)}{R^\beta}$. This is an estimate from below on how fast a nonconstant harmonic function must grow near a point where it vanishes.

8.8. REFERENCES

Section 8.1 See Giaquinta [Gi] and Giaquinta–Hildebrandt [G-H] for more about the calculus of variations. Struwe [Str] and Zeidler [Zd, Vol. 3] are good references for variational methods.

- Section 8.2 The proof of Theorem 1 in §8.2.2 is from Ladyzhenskaya–Ural'tseva [L-U]. Section 8.2.5 is based upon Ball–Murat (Pitman Research Notes Math. Series 254, Longman (1991), 25–37).
- Section 8.3 See Giaquinta [Gi] for additional information about regularity (and partial regularity) of minimizers.
- Section 8.4 The books by Kinderlehrer–Stampacchia [K-S] and Friedman [Fr3] explain much more about variational inequalities and free boundary problems. See Hélein [Hi] and Lin–Wang [L-W] for further information about harmonic maps.
- Section 8.5 The Mountain Pass Theorem is due to Ambrosetti and Rabinowitz (J. Functional Analysis 14 (1973), 349–38). Consult Rabinowitz [Ra] (the source of §8.5) for further results on saddle point methods.
- Section 8.6 The discussion of Noether's Theorem and its applications is primarily based upon Reed–Simon [R-S2] and Strauss [St1]. See also Morawetz's appendix to Lax–Phillips [Lx-P] and Lax [Lx3]. The identity (24) is from Costa and Strauss (Quart. Appl. Math. (1981), 351–361).
- Section 8.7 Problem 13: see Ekeland–Temam [E-T] for how to formulate dual variational principles. Problem 18 is due to Alessandrini: see Isakov [I, §4.3].

NONVARIATIONAL TECHNIQUES

- 9.1 Monotonicity methods
- 9.2 Fixed point methods
- 9.3 Method of subsolutions and supersolutions
- 9.4 Nonexistence of solutions
- 9.5 Geometric properties of solutions
- 9.6 Gradient flows
- 9.7 Problems
- 9.8 References

We gather in this chapter various techniques for proving the existence, nonexistence, uniqueness, and various other properties of solutions for non-linear elliptic and parabolic partial differential equations that are not necessarily of variational form.

9.1. MONOTONICITY METHODS

Let us look first at this boundary-value problem for a divergence structure quasilinear PDE:

$$(1) \quad \begin{cases} -\operatorname{div} \mathbf{a}(Du) = f & \text{in } U \\ u = 0 & \text{on } \partial U, \end{cases}$$

where $f \in L^2(U)$ is given, as is the smooth vector field $\mathbf{a} : \mathbb{R}^n \rightarrow \mathbb{R}^n$, $\mathbf{a} = (a^1, \dots, a^n)$. As usual the unknown is $u : \bar{U} \rightarrow \mathbb{R}$, $u = u(x)$, where U is a

bounded, open subset of \mathbb{R}^n with smooth boundary. Now *if* there exists a function $F : \mathbb{R}^n \rightarrow \mathbb{R}$ such that \mathbf{a} is the gradient of F ,

$$(2) \quad \mathbf{a}(p) = DF(p) \quad (p \in \mathbb{R}^n),$$

then (1) is the Euler–Lagrange equation corresponding to the Lagrangian $L(p, z, x) = F(p) - f(x)z$. However if there exists no such potential F , the variational methods from Chapter 8 simply do not apply to the problem (1).

We inquire instead if there is rather some direct method of constructing a solution of (1) and in particular ask what are reasonable conditions to place upon the nonlinearity. For motivation let us note that *if* (2) were valid and *if* F were convex (the natural assumption for the variational theory, as we have seen in §8.2.2), then for each $p, q \in \mathbb{R}^n$

$$\begin{aligned} (\mathbf{a}(p) - \mathbf{a}(q)) \cdot (p - q) &= \sum_{i=1}^n (F_{p_i}(p) - F_{p_i}(q))(p_i - q_i) \\ &= \int_0^1 \sum_{i,j=1}^n F_{p_i p_j}(p + t(q - p))(p_j - q_j)(p_i - q_i) dt \geq 0, \end{aligned}$$

the last inequality following from the convexity of F .

This calculation suggests the following

DEFINITION. A vector field $\mathbf{a} : \mathbb{R}^n \rightarrow \mathbb{R}^n$ is called monotone provided

$$(3) \quad (\mathbf{a}(p) - \mathbf{a}(q)) \cdot (p - q) \geq 0$$

for all $p, q \in \mathbb{R}^n$.

We will show below that the quasilinear PDE does indeed possess a weak solution, under the primary structural assumption that the nonlinearity be monotone. Later we will realize that this condition in effect says that $-\operatorname{div} \mathbf{a}(Du) = f$ is a nonlinear elliptic partial differential equation. So let us henceforth assume that the smooth vector field \mathbf{a} is monotone and that

$$(4) \quad |\mathbf{a}(p)| \leq C(1 + |p|),$$

$$(5) \quad \mathbf{a}(p) \cdot p \geq \alpha|p|^2 - \beta$$

for all $p \in \mathbb{R}^n$ and appropriate constants $C, \alpha > 0, \beta \geq 0$. We will see momentarily that (5) amounts to a coercivity condition on the nonlinearity. We intend now to build a solution of the boundary-value problem (1) as

the limit of certain finite-dimensional approximations, thereby extending Galerkin's method from Chapter 7 to a new class of nonlinear problems.

More precisely, assume that the functions $w_k = w_k(x)$ ($k = 1, \dots$) are smooth and

$$\{w_k\}_{k=1}^{\infty} \text{ is an orthonormal basis of } H_0^1(U),$$

taken with the inner product $(u, v) = \int_U Du \cdot Dv \, dx$. (We could for instance take $\{w_k\}_{k=1}^{\infty}$ to be the set of appropriately normalized eigenfunctions for $-\Delta$ in $H_0^1(U)$.)

We will look for a function $u_m \in H_0^1(U)$ of the form

$$(6) \quad u_m = \sum_{k=1}^m d_m^k w_k,$$

where we hope to select the coefficients d_m^k so that

$$(7) \quad \int_U \mathbf{a}(Du_m) \cdot Dw_k \, dx = \int_U f w_k \, dx \quad (k = 1, \dots, m).$$

This amounts to our requiring that u_m solves the "projection" of the problem (1) onto the finite-dimensional subspace spanned by $\{w_k\}_{k=1}^m$.

We start with a technical assertion.

LEMMA (Zeros of a vector field). *Assume the continuous function $\mathbf{v} : \mathbb{R}^n \rightarrow \mathbb{R}^n$ satisfies*

$$(8) \quad \mathbf{v}(x) \cdot x \geq 0 \quad \text{if } |x| = r,$$

for some $r > 0$. Then there exists a point $x \in B(0, r)$ such that

$$\mathbf{v}(x) = 0.$$

Proof. Suppose the assertion were false; then $\mathbf{v}(x) \neq 0$ for all $x \in B(0, r)$. Define the continuous mapping $\mathbf{w} : B(0, r) \rightarrow \partial B(0, r)$ by setting

$$\mathbf{w}(x) := -\frac{r}{|\mathbf{v}(x)|} \mathbf{v}(x) \quad (x \in B(0, r)).$$

According to Brouwer's fixed point theorem (§8.1.4), there exists a point $z \in B(0, r)$ with

$$(9) \quad \mathbf{w}(z) = z.$$

But then $z \in \partial B(0, r)$, and so (8) and (9) imply the contradiction

$$r^2 = z \cdot z = \mathbf{w}(z) \cdot z = -\frac{r}{|\mathbf{v}(z)|} \mathbf{v}(z) \cdot z \leq 0. \quad \square$$

THEOREM 1 (Construction of approximate solutions). *For each integer $m = 1, \dots$, there exists a function u_m of the form (6) satisfying the identities (7).*

Proof. Define the continuous function $\mathbf{v} : \mathbb{R}^m \rightarrow \mathbb{R}^m$, $\mathbf{v} = (v^1, \dots, v^m)$, by setting

$$(10) \quad v^k(d) := \int_U \mathbf{a} \left(\sum_{j=1}^m d_j Dw_j \right) \cdot Dw_k - f w_k \, dx \quad (k = 1, \dots, m)$$

for each point $d = (d_1, \dots, d_m) \in \mathbb{R}^m$. Now

$$\begin{aligned} \mathbf{v}(d) \cdot d &= \int_U \mathbf{a} \left(\sum_{j=1}^m d_j Dw_j \right) \cdot \left(\sum_{j=1}^m d_j Dw_j \right) - f \left(\sum_{j=1}^m d_j w_j \right) \, dx \\ &\geq \int_U \alpha \left| \sum_{j=1}^m d_j Dw_j \right|^2 - \beta - f \left(\sum_{j=1}^m d_j w_j \right) \, dx \quad \text{by (5)} \\ &= \alpha |d|^2 - \beta |U| - \sum_{j=1}^m d_j \int_U f w_j \, dx \\ &\geq \frac{\alpha}{2} |d|^2 - C. \end{aligned}$$

Hence $\mathbf{v}(d) \cdot d \geq 0$ if $|d| = r$, provided we select $r > 0$ sufficiently large.

We apply the lemma, to conclude that $\mathbf{v}(d) = 0$ for some point $d \in \mathbb{R}^m$. Then (10) implies u_m , defined by (6), satisfies (7). \square

We want to take the limit as $m \rightarrow \infty$, and for this will require some uniform estimates.

THEOREM 2 (Energy estimates). *There exists a constant C , depending only on U and \mathbf{a} , such that*

$$(11) \quad \|u_m\|_{H_0^1(U)} \leq C(1 + \|f\|_{L^2(U)})$$

for $m = 1, 2, \dots$

Proof. Multiply equality (7) by d_m^k and sum for $k = 1, \dots, m$:

$$\int_U \mathbf{a}(Du_m) \cdot Du_m \, dx = \int_U f u_m \, dx.$$

In view of the coercivity inequality (5), we find

$$\alpha \int_U |Du_m|^2 \, dx \leq C + \int_U f u_m \, dx \leq C + \epsilon \int_U u_m^2 \, dx + \frac{1}{4\epsilon} \int_U f^2 \, dx.$$

We recall Poincaré's inequality (Theorem 3 in §5.6.1) and then choose $\epsilon > 0$ small enough to deduce (11). \square

We wish now to employ the L^2 inequalities (11) to pass to limits as $m \rightarrow \infty$, obtaining thereby a weak solution of problem (1), which is to say, a function $u \in H_0^1(U)$ satisfying the identity

$$(12) \quad \int_U \mathbf{a}(Du) \cdot Dv \, dx = \int_U f v \, dx \quad \text{for all } v \in H_0^1(U).$$

Employing estimate (11), we can extract a subsequence $\{u_{m_j}\}_{j=1}^\infty$ that converges weakly in $H_0^1(U)$ to a limit u , which we hope to show verifies (12). However, we encounter a major problem here: we cannot directly conclude that

$$\mathbf{a}(Du_{m_j}) \rightarrow \mathbf{a}(Du)$$

in any sense whatsoever. Take note: *nonlinearities are (usually) not continuous with respect to weak convergence.* (See Problem 2.)

What saves us is the monotonicity assumption on vector field \mathbf{a} .

THEOREM 3 (Existence of weak solution). *There exists a weak solution of the nonlinear boundary-value problem (1).*

Proof. 1. As noted in the foregoing discussion, we can extract a subsequence $\{u_{m_j}\}_{j=1}^\infty \subset \{u_m\}_{m=1}^\infty$ and a function $u \in H_0^1(U)$ such that

$$(13) \quad u_{m_j} \rightharpoonup u \quad \text{weakly in } H_0^1(U)$$

and

$$(14) \quad u_{m_j} \rightarrow u \quad \text{in } L^2(U).$$

We must show u satisfies (12).

2. In view of the growth condition (4), $\{\mathbf{a}(Du_m)\}_{m=1}^\infty$ is bounded in $L^2(U; \mathbb{R}^n)$; and so we may as well suppose—upon passing to a further subsequence if necessary—that

$$(15) \quad \mathbf{a}(Du_{m_j}) \rightharpoonup \boldsymbol{\xi} \quad \text{weakly in } L^2(U; \mathbb{R}^n),$$

for some $\boldsymbol{\xi} \in L^2(U; \mathbb{R}^n)$. Using identity (7), we deduce

$$\int_U \boldsymbol{\xi} \cdot Dw_k \, dx = \int_U f w_k \, dx$$

for each $k = 1, \dots$. And so

$$(16) \quad \int_U \xi \cdot Dv \, dx = \int_U fv \, dx \quad \text{for each } v \in H_0^1(U).$$

3. To proceed further, let us note from the monotonicity condition (3) that

$$(17) \quad \int_U (\mathbf{a}(Du_m) - \mathbf{a}(Dw)) \cdot (Du_m - Dw) \, dx \geq 0$$

for $m = 1, \dots$ and all $w \in H_0^1(U)$. But as observed before, equation (7) yields the identity

$$\int_U \mathbf{a}(Du_m) \cdot Du_m \, dx = \int_U fu_m \, dx.$$

Substitute into (17), to find

$$\int_U fu_m - \mathbf{a}(Du_m) \cdot Dw - \mathbf{a}(Dw) \cdot (Du_m - Dw) \, dx \geq 0.$$

Let $m = m_j \rightarrow \infty$ and recall (13)–(15), to deduce

$$\int_U fu - \xi \cdot Dw - \mathbf{a}(Dw) \cdot (Du - Dw) \, dx \geq 0.$$

We simplify using identity (16) with $v = u$ and discover

$$(18) \quad \int_U (\xi - \mathbf{a}(Dw)) \cdot D(u - w) \, dx \geq 0 \quad \text{for all } w \in H_0^1(U).$$

4. Fix any $v \in H_0^1(U)$ and set $w := u - \lambda v$ ($\lambda > 0$) in (18). We obtain then

$$\int_U (\xi - \mathbf{a}(Du - \lambda Dv)) \cdot Dv \, dx \geq 0.$$

Send $\lambda \rightarrow 0$:

$$(19) \quad \int_U (\xi - \mathbf{a}(Du)) \cdot Dv \, dx \geq 0 \quad \text{for all } v \in H_0^1(U).$$

Replacing v by $-v$, we deduce that in fact equality holds above. Then (16) and (19) taken together yield

$$\int_U \mathbf{a}(Du) \cdot Dv \, dx = \int_U fv \, dx \quad \text{for all } v \in H_0^1(U).$$

Hence u is indeed a weak solution of (1). □

This use of monotonicity is the *method of Browder and Minty*, a remarkable technique which employs the inequality condition (3) to justify passing to weak limits within a nonlinearity.

Let us assume now the vector field \mathbf{a} satisfies the condition of *strict monotonicity*; that is,

$$(20) \quad (\mathbf{a}(p) - \mathbf{a}(q)) \cdot (p - q) \geq \theta |p - q|^2$$

for all $p, q \in \mathbb{R}^n$ and some constant $\theta > 0$.

THEOREM 4 (Uniqueness of weak solution). *Assume the strict monotonicity property (20) holds. Then there exists only one weak solution of (1).*

Proof. Assume that u and \tilde{u} are two weak solutions. Consequently

$$\int_U \mathbf{a}(Du) \cdot Dv \, dx = \int_U \mathbf{a}(D\tilde{u}) \cdot Dv \, dx = \int_U f v \, dx,$$

and so

$$\int_U [\mathbf{a}(Du) - \mathbf{a}(D\tilde{u})] \cdot Dv \, dx = 0$$

for each $v \in H_0^1(U)$. We set $v := u - \tilde{u}$ and use (20) to deduce

$$\int_U |Du - D\tilde{u}|^2 \, dx = 0.$$

Thus $u = \tilde{u}$ a.e. in U . □

H^2 regularity. Under the strengthened monotonicity assumption (20) our weak solution u in fact belongs to $H^2(U)$ and so satisfies

$$-\operatorname{div} \mathbf{a}(Du) = f \quad \text{a.e. in } U.$$

To demonstrate this, we select $q, \xi \in \mathbb{R}^n$ and set $p = q + h\xi$, $h \neq 0$, in (20). We obtain, after dividing by h^2 , the inequality

$$\sum_{i=1}^n \frac{[a^i(q + h\xi) - a^i(q)] \xi_i}{h} \geq \theta |\xi|^2.$$

Now send $h \rightarrow 0$:

$$(21) \quad \sum_{i,j=1}^n a_{p_j}^i(q) \xi_i \xi_j \geq \theta |\xi|^2 \quad (q, \xi \in \mathbb{R}^n).$$

We can thus interpret the nonlinear PDE $-\operatorname{div} \mathbf{a}(Du) = f$ as being uniformly elliptic. The proof of H^2 regularity of the weak solution now follows almost precisely as in the proof of Theorem 1 in §6.3.1.

9.2. FIXED POINT METHODS

We study next the applicability of topological *fixed point theorems* to nonlinear partial differential equations. There are at least three distinct classes of such abstract theorems that are useful. These are:

- (a) fixed point theorems for *strict contractions*,
- (b) fixed point theorems for *compact mappings*,

and

- (c) fixed point theorems for *order-preserving operators*.

We present below applications of types (a) and (b). The utility of order-preserving properties for nonlinear PDE will be explained later, in §9.3.

9.2.1. Banach's Fixed Point Theorem.

Hereafter X denotes a Banach space. The simplest fixed point theorem of all is

THEOREM 1 (Banach's Fixed Point Theorem). *Assume*

$$A : X \rightarrow X$$

is a nonlinear mapping, and suppose that

$$(1) \quad \|A[u] - A[\tilde{u}]\| \leq \gamma \|u - \tilde{u}\| \quad (u, \tilde{u} \in X)$$

for some constant $\gamma < 1$. Then A has a unique fixed point.

DEFINITION. *We say that A is a strict contraction if (1) holds.*

Proof. Fix any point $u_0 \in X$ and thereafter iteratively define $u_{k+1} = A[u_k]$ for $k = 0, 1, \dots$. Then

$$\|A[u_{k+1}] - A[u_k]\| \leq \gamma \|u_{k+1} - u_k\| = \gamma \|A[u_k] - A[u_{k-1}]\|,$$

and so

$$\|A[u_{k+1}] - A[u_k]\| \leq \gamma^k \|A[u_0] - u_0\|$$

for $k = 1, 2, \dots$. Consequently if $k \geq l$,

$$\begin{aligned} \|u_k - u_l\| &= \|A[u_{k-1}] - A[u_{l-1}]\| \leq \sum_{j=l-1}^{k-2} \|A[u_{j+1}] - A[u_j]\| \\ &\leq \|A[u_0] - u_0\| \sum_{j=l-1}^{k-2} \gamma^j. \end{aligned}$$

Hence $\{u_k\}_{k=1}^{\infty}$ is a Cauchy sequence in X , and so there exists a point $u \in X$ with $u_k \rightarrow u$ in X . Clearly then $A[u] = u$. Hence u is a fixed point for A , and hypothesis (1) ensures uniqueness. \square

Applications of Banach's Fixed Point Theorem to nonlinear PDE usually involve perturbation arguments of various sorts: given a well-behaved linear elliptic partial differential equation, it is often straightforward to cast a small nonlinear modification as a contraction mapping. The hallmark of such proofs is the occurrence of a parameter which must be taken small enough to ensure the strict contraction property.

Sometimes however we can eliminate such a smallness hypothesis by an iteration, as now illustrated.

Example 1 (Reaction-diffusion equations). Let us investigate the solvability of the initial/boundary-value problem for the *reaction-diffusion system*

$$(2) \quad \begin{cases} \mathbf{u}_t - \Delta \mathbf{u} = \mathbf{f}(\mathbf{u}) & \text{in } U_T \\ \mathbf{u} = \mathbf{0} & \text{on } \partial U \times [0, T] \\ \mathbf{u} = \mathbf{g} & \text{on } U \times \{t = 0\}. \end{cases}$$

Here $\mathbf{u} = (u^1, \dots, u^m)$, $\mathbf{g} = (g^1, \dots, g^m)$, and as usual $U_T = U \times (0, T]$, where $U \in \mathbb{R}^n$ is open and bounded, with smooth boundary. The time $T > 0$ is fixed. We assume that the initial function \mathbf{g} belongs to $H_0^1(U; \mathbb{R}^m)$. Concerning the nonlinearity, let us suppose

$$(3) \quad \mathbf{f} : \mathbb{R}^m \rightarrow \mathbb{R}^m \text{ is Lipschitz continuous.}$$

This hypothesis in particular implies

$$(4) \quad |\mathbf{f}(z)| \leq C(1 + |z|)$$

for each $z \in \mathbb{R}^m$ and some constant C .

Adapting the terminology from §7.1, we say that a function

$$(5) \quad \mathbf{u} \in L^2(0, T; H_0^1(U; \mathbb{R}^m)), \text{ with } \mathbf{u}' \in L^2(0, T; H^{-1}(U; \mathbb{R}^m)),$$

is a *weak solution* of (2) provided

$$(6) \quad \langle \mathbf{u}', \mathbf{v} \rangle + B[\mathbf{u}, \mathbf{v}] = (\mathbf{f}(\mathbf{u}), \mathbf{v}) \quad \text{a.e. } 0 \leq t \leq T$$

for each $\mathbf{v} \in H_0^1(U; \mathbb{R}^m)$ and

$$(7) \quad \mathbf{u}(0) = \mathbf{g}.$$

In (6) $\langle \cdot, \cdot \rangle$ denotes the pairing of $H^{-1}(U; \mathbb{R}^m)$ and $H_0^1(U; \mathbb{R}^m)$, $B[\cdot, \cdot]$ is the bilinear form associated with $-\Delta$ in $H_0^1(U; \mathbb{R}^m)$, and (\cdot, \cdot) denotes the inner product in $L^2(U; \mathbb{R}^m)$. The norm in $H_0^1(U; \mathbb{R}^m)$ is taken to be

$$\|\mathbf{u}\|_{H_0^1(U; \mathbb{R}^m)} = \left(\int_U |D\mathbf{u}|^2 dx \right)^{\frac{1}{2}}.$$

Recall from §5.9.2 that (5) implies $\mathbf{u} \in C([0, T]; L^2(U; \mathbb{R}^m))$, after possible redefinition of \mathbf{u} on a set of measure zero.

THEOREM 2 (Existence). *There exists a unique weak solution of (2).*

Proof. 1. We will apply Banach's Theorem in the space

$$X = C([0, T]; L^2(U; \mathbb{R}^m)),$$

with the norm

$$\|\mathbf{v}\| := \max_{0 \leq t \leq T} \|\mathbf{v}(t)\|_{L^2(U; \mathbb{R}^m)}.$$

Let the operator A be defined as follows. Given a function $\mathbf{u} \in X$, set $\mathbf{h}(t) := \mathbf{f}(\mathbf{u}(t))$ ($0 \leq t \leq T$). In light of the growth estimate (4), we see $\mathbf{h} \in L^2(0, T; L^2(U; \mathbb{R}^m))$. Consequently the theory set forth in §7.1 ensures that the linear parabolic PDE

$$(8) \quad \begin{cases} \mathbf{w}_t - \Delta \mathbf{w} = \mathbf{h} & \text{in } U_T \\ \mathbf{w} = \mathbf{0} & \text{on } \partial U \times [0, T] \\ \mathbf{w} = \mathbf{g} & \text{on } U \times \{t = 0\} \end{cases}$$

has a unique weak solution

$$(9) \quad \mathbf{w} \in L^2(0, T; H_0^1(U; \mathbb{R}^m)), \text{ with } \mathbf{w}' \in L^2(0, T; H^{-1}(U; \mathbb{R}^m)).$$

Thus $\mathbf{w} \in X$ satisfies

$$(10) \quad \langle \mathbf{w}', \mathbf{v} \rangle + B[\mathbf{w}, \mathbf{v}] = (\mathbf{h}, \mathbf{v}) \quad \text{a.e. } 0 \leq t \leq T$$

for each $\mathbf{v} \in H_0^1(U; \mathbb{R}^m)$ and $\mathbf{w}(0) = \mathbf{g}$.

Define $A : X \rightarrow X$ by setting $A[\mathbf{u}] = \mathbf{w}$.

2. We now claim that

$$(11) \quad \begin{cases} \text{if } T > 0 \text{ is small enough, then} \\ A \text{ is a strict contraction.} \end{cases}$$

To prove this, choose $\mathbf{u}, \tilde{\mathbf{u}} \in X$, and define $\mathbf{w} = A[\mathbf{u}]$, $\tilde{\mathbf{w}} = A[\tilde{\mathbf{u}}]$ as above. Consequently \mathbf{w} verifies (10) for $\mathbf{h} = \mathbf{f}(\mathbf{u})$, and $\tilde{\mathbf{w}}$ satisfies a similar identity for $\tilde{\mathbf{h}} := \mathbf{f}(\tilde{\mathbf{u}})$.

We calculate as in §7.1

$$(12) \quad \begin{aligned} & \frac{d}{dt} \|\mathbf{w} - \tilde{\mathbf{w}}\|_{L^2(U; \mathbb{R}^m)}^2 + 2\|\mathbf{w} - \tilde{\mathbf{w}}\|_{H_0^1(U; \mathbb{R}^m)}^2 \\ &= 2(\mathbf{w} - \tilde{\mathbf{w}}, \mathbf{h} - \tilde{\mathbf{h}}) \\ &\leq \epsilon \|\mathbf{w} - \tilde{\mathbf{w}}\|_{L^2(U; \mathbb{R}^m)}^2 + \frac{1}{\epsilon} \|\mathbf{f}(\mathbf{u}) - \mathbf{f}(\tilde{\mathbf{u}})\|_{L^2(U; \mathbb{R}^m)}^2 \\ &\leq \epsilon C \|\mathbf{w} - \tilde{\mathbf{w}}\|_{H_0^1(U; \mathbb{R}^m)}^2 + \frac{1}{\epsilon} \|\mathbf{f}(\mathbf{u}) - \mathbf{f}(\tilde{\mathbf{u}})\|_{L^2(U; \mathbb{R}^m)}^2, \end{aligned}$$

by Poincaré's inequality. Selecting $\epsilon > 0$ sufficiently small, we deduce

$$\frac{d}{dt} \|\mathbf{w} - \tilde{\mathbf{w}}\|_{L^2(U; \mathbb{R}^m)}^2 \leq C \|\mathbf{f}(\mathbf{u}) - \mathbf{f}(\tilde{\mathbf{u}})\|_{L^2(U; \mathbb{R}^m)}^2 \leq C \|\mathbf{u} - \tilde{\mathbf{u}}\|_{L^2(U; \mathbb{R}^m)}^2,$$

since \mathbf{f} is Lipschitz. Consequently

$$(13) \quad \begin{aligned} \|\mathbf{w}(s) - \tilde{\mathbf{w}}(s)\|_{L^2(U; \mathbb{R}^m)}^2 &\leq C \int_0^s \|\mathbf{u}(t) - \tilde{\mathbf{u}}(t)\|_{L^2(U; \mathbb{R}^m)}^2 dt \\ &\leq CT \|\mathbf{u} - \tilde{\mathbf{u}}\|^2 \end{aligned}$$

for each $0 \leq s \leq T$. Maximizing the left-hand side with respect to s , we discover

$$\|\mathbf{w} - \tilde{\mathbf{w}}\|^2 \leq CT \|\mathbf{u} - \tilde{\mathbf{u}}\|^2.$$

Hence

$$(14) \quad \|A[\mathbf{u}] - A[\tilde{\mathbf{u}}]\| \leq (CT)^{1/2} \|\mathbf{u} - \tilde{\mathbf{u}}\|,$$

and thus A is a strict contraction, provided $T > 0$ is so small that $(CT)^{1/2} = \gamma < 1$.

3. Given any $T > 0$, we select $T_1 > 0$ so small that $(CT_1)^{1/2} < 1$. We can then apply Banach's Fixed Point Theorem to find a weak solution \mathbf{u} of the problem (2) existing on the time interval $[0, T_1]$. Since $\mathbf{u}(t) \in H_0^1(U; \mathbb{R}^m)$ for a.e. $0 \leq t \leq T_1$, we can upon redefining T_1 if necessary assume $\mathbf{u}(T_1) \in H_0^1(U; \mathbb{R}^m)$.

Observe that the time $T_1 > 0$ depends only upon the Lipschitz constant for \mathbf{f} . We can therefore repeat the argument above, to extend our solution to the time interval $[T_1, 2T_1]$. Continuing, after finitely many steps we construct a weak solution existing on the full interval $[0, T]$.

4. To demonstrate uniqueness, suppose both \mathbf{u} and $\tilde{\mathbf{u}}$ are two weak solutions of (2). Then we have $\mathbf{w} = \mathbf{u}$, $\tilde{\mathbf{w}} = \tilde{\mathbf{u}}$ in inequality (13), whence

$$\|\mathbf{u}(s) - \tilde{\mathbf{u}}(s)\|_{L^2(U; \mathbb{R}^m)}^2 \leq C \int_0^s \|\mathbf{u}(t) - \tilde{\mathbf{u}}(t)\|_{L^2(U; \mathbb{R}^m)}^2 dt$$

for $0 \leq s \leq T$. According to Gronwall's inequality, $\mathbf{u} \equiv \tilde{\mathbf{u}}$. □

Interpretation. In common applications problem (2) records the evolution of the densities u^1, \dots, u^m of various chemicals, which both diffuse within a medium and interact with each other. The diffusion term is $\Delta \mathbf{u}$ (or more generally $(a_1 \Delta u^1, \dots, a_m \Delta u^m)$ where the constants $a_k > 0$ characterize the diffusion of the k^{th} chemical). The reaction term $\mathbf{f}(\mathbf{u})$ models the chemistry. In the foregoing example we made the unreasonable assumption that \mathbf{f} is

globally Lipschitz. In more realistic models \mathbf{f} is often a polynomial in \mathbf{u} and there are interesting problems as to the global existence or blow-up of a solution. (A simple such example is treated in §9.4.1.)

We will later employ Banach's Fixed Point Theorem to prove long and short time existence for solutions of certain quasilinear wave equations in §12.2.

9.2.2. Schauder's, Schaefer's Fixed Point Theorems.

Next we extend Brouwer's Fixed Point Theorem (§8.1.4) to Banach spaces. The key assumption is now compactness. Throughout this subsection X continues to denote a real Banach space.

THEOREM 3 (Schauder's Fixed Point Theorem). *Suppose $K \subset X$ is compact and convex, and assume also*

$$A : K \rightarrow K$$

is continuous. Then A has a fixed point in K .

Proof. 1. Fix $\epsilon > 0$ and choose finitely many points $u_1, \dots, u_{N_\epsilon} \in K$, so that the open balls $\{B^0(u_i, \epsilon)\}_{i=1}^{N_\epsilon}$ cover K :

$$(15) \quad K \subset \bigcup_{i=1}^{N_\epsilon} B^0(u_i, \epsilon).$$

This is possible since K is compact. Let K_ϵ denote the closed convex hull of the points $\{u_1, \dots, u_{N_\epsilon}\}$:

$$K_\epsilon := \left\{ \sum_{i=1}^{N_\epsilon} \lambda_i u_i \mid 0 \leq \lambda_i \leq 1, \sum_{i=1}^{N_\epsilon} \lambda_i = 1 \right\}.$$

Then $K_\epsilon \subset K$, since K is convex. Now define $P_\epsilon : K \rightarrow K_\epsilon$ by writing

$$P_\epsilon[u] := \frac{\sum_{i=1}^{N_\epsilon} \text{dist}(u, K - B^0(u_i, \epsilon)) u_i}{\sum_{i=1}^{N_\epsilon} \text{dist}(u, K - B^0(u_i, \epsilon))} \quad (u \in K).$$

The denominator is never zero, because of (15). Now clearly P_ϵ is continuous, and furthermore for each $u \in K$, we have

$$(16) \quad \|P_\epsilon[u] - u\| \leq \frac{\sum_{i=1}^{N_\epsilon} \text{dist}(u, K - B^0(u_i, \epsilon)) \|u_i - u\|}{\sum_{i=1}^{N_\epsilon} \text{dist}(u, K - B^0(u_i, \epsilon))} \leq \epsilon.$$

2. Consider next the operator $A_\epsilon : K_\epsilon \rightarrow K_\epsilon$ defined by

$$A_\epsilon[u] := P_\epsilon[A[u]] \quad (u \in K_\epsilon).$$

Now K_ϵ is homeomorphic to the closed unit ball in \mathbb{R}^{M_ϵ} for some $M_\epsilon \leq N_\epsilon$. Brouwer's Fixed Point Theorem (§8.1.4) therefore ensures the existence of a point $u_\epsilon \in K_\epsilon$ with

$$A_\epsilon[u_\epsilon] = u_\epsilon.$$

3. As K is compact, there exist a subsequence $\epsilon_j \rightarrow 0$ and a point $u \in K$, such that $u_{\epsilon_j} \rightarrow u$ in X . We claim u is a fixed point of A . Indeed, using estimate (16), we deduce

$$\|u_{\epsilon_j} - A[u_{\epsilon_j}]\| = \|A_{\epsilon_j}[u_{\epsilon_j}] - A[u_{\epsilon_j}]\| = \|P_{\epsilon_j}[A[u_{\epsilon_j}]] - A[u_{\epsilon_j}]\| \leq \epsilon_j.$$

Consequently, since A is continuous, we conclude $u = A[u]$. □

We next transform Schauder's Fixed Point Theorem into an alternative form which is often more useful for applications to nonlinear partial differential equations.

DEFINITION. *A nonlinear mapping $A : X \rightarrow X$ is called compact provided for each bounded sequence $\{u_k\}_{k=1}^\infty$ the sequence $\{A[u_k]\}_{k=1}^\infty$ is precompact; that is, there exists a subsequence $\{u_{k_j}\}_{j=1}^\infty$ such that $\{A[u_{k_j}]\}_{j=1}^\infty$ converges in X .*

THEOREM 4 (Schaefer's Fixed Point Theorem). *Suppose*

$$A : X \rightarrow X$$

is a continuous and compact mapping. Assume further that the set

$$\{u \in X \mid u = \lambda A[u] \text{ for some } 0 \leq \lambda \leq 1\}$$

is bounded. Then A has a fixed point.

The assertion is that if we have a bound on any possible fixed points of any of the operators λA for $0 \leq \lambda \leq 1$, then we have the existence of a fixed point for A . This is in accordance with the remarkable informal principle that "if we can prove appropriate estimates for solutions of a nonlinear PDE, under the assumption that such solutions exist, then in fact these solutions do exist". This is the method of *a priori** estimates.

The advantage of Schaefer's Theorem over Schauder's for applications is that we do not have to identify an explicit convex, compact set.

* *a priori* = from before (Latin).

Proof. 1. Choose a constant M so large that

$$(17) \quad \|u\| < M \quad \text{if } u = \lambda A[u] \text{ for some } 0 \leq \lambda \leq 1.$$

Define then

$$(18) \quad \tilde{A}[u] := \begin{cases} A[u] & \text{if } \|A[u]\| \leq M \\ \frac{MA[u]}{\|A[u]\|} & \text{if } \|A[u]\| \geq M. \end{cases}$$

Observe $\tilde{A} : B(0, M) \rightarrow B(0, M)$. Now set $K =$ closed convex hull of $\tilde{A}(B(0, M))$. Then since A and thus \tilde{A} are compact mappings, K is a compact, convex subset of X . Furthermore $\tilde{A} : K \rightarrow K$.

2. Invoking Schauder's Fixed Point Theorem, we infer the existence of a point $u \in K$ with

$$(19) \quad \tilde{A}[u] = u.$$

We now claim additionally that u is a fixed point of A . If not, then according to (18) and (19) we would have

$$\|A[u]\| > M$$

and

$$(20) \quad u = \lambda A[u] \quad \text{for } \lambda = \frac{M}{\|A[u]\|} < 1.$$

But $\|u\| = \|\tilde{A}[u]\| = M$, a contradiction to (17) and (20). \square

A fixed point theorem for convex sets. For certain applications it is convenient to have available a variant of the previous theorem, asserting that if K is a convex subset of a Banach space X , with $0 \in K$, and if $A : K \rightarrow K$ is a continuous and compact mapping for which the set

$$\{u \in K \mid u = \lambda A[u] \text{ for some } 0 \leq \lambda \leq 1\}$$

is bounded, then A has a fixed point in K .

The proof is a slight modification of the argument above.

Application. Employing Schauder's and Schaefer's Fixed Point Theorems for PDE depends upon quite different considerations than applications of Banach's Theorem. The crucial assumption is now not that some parameter be small, but rather that we have some sort of compactness. As the inverses of linear elliptic operators are typically smoothing, compactness is indeed available for certain nonlinear elliptic equations. Following is a quick, albeit fairly crude, example:

Example 2 (A quasilinear elliptic PDE). We present now a simple application of Schaefer's Theorem by solving the semilinear boundary-value problem

$$(21) \quad \begin{cases} -\Delta u + b(Du) + \mu u = 0 & \text{in } U \\ u = 0 & \text{on } \partial U, \end{cases}$$

where U is bounded and ∂U is smooth. We assume $b : \mathbb{R}^n \rightarrow \mathbb{R}$ is smooth, Lipschitz continuous and thus satisfies the growth condition

$$(22) \quad |b(p)| \leq C(|p| + 1)$$

for some constant C and all $p \in \mathbb{R}^n$.

THEOREM 5 (Existence). *If $\mu > 0$ is sufficiently large, there exists a function $u \in H^2(U) \cap H_0^1(U)$ solving the boundary-value problem (21).*

Proof. 1. Given $u \in H_0^1(U)$, set

$$(23) \quad f := -b(Du).$$

Owing to estimate (22), we see that $f \in L^2(U)$. Now let $w \in H_0^1(U)$ be the unique weak solution of the linear problem

$$(24) \quad \begin{cases} -\Delta w + \mu w = f & \text{in } U \\ w = 0 & \text{on } \partial U. \end{cases}$$

By the regularity theory proved in §6.3, we know additionally that $w \in H^2(U)$, with the estimate

$$(25) \quad \|w\|_{H^2(U)} \leq C\|f\|_{L^2(U)}$$

for some constant C .

Let us henceforth write $A[u] = w$ whenever w is derived from u via (23), (24). In light of (22) and (25), we have the estimate

$$(26) \quad \|A[u]\|_{H^2(U)} \leq C(\|u\|_{H_0^1(U)} + 1).$$

2. We now assert that $A : H_0^1(U) \rightarrow H_0^1(U)$ is continuous and compact. Indeed, if

$$(27) \quad u_k \rightarrow u \quad \text{in } H_0^1(U),$$

then in view of estimate (26) we have

$$(28) \quad \sup_k \|w_k\|_{H^2(U)} < \infty,$$

for $w_k = A[u_k]$ ($k = 1, \dots$). Thus there is a subsequence $\{w_{k_j}\}_{j=1}^\infty$ and a function $w \in H_0^1(U)$ with

$$(29) \quad w_{k_j} \rightarrow w \quad \text{in } H_0^1(U).$$

Now

$$\int_U Dw_{k_j} \cdot Dv + \mu w_{k_j} v \, dx = - \int_U b(Du_{k_j})v \, dx$$

for each $v \in H_0^1(U)$. Consequently using (22), (27) and (29), we see

$$\int_U Dw \cdot Dv + \mu wv \, dx = - \int_U b(Du)v \, dx$$

for each $v \in H_0^1(U)$. Thus $w = A[u]$.

Hence (27) implies $A[u_k] \rightarrow A[u]$ in $H_0^1(U)$, and so A is continuous. A similar argument shows that A is compact, since if $\{u_k\}_{k=1}^\infty$ is bounded in $H_0^1(U)$, estimate (22) asserts that $\{A[u_k]\}_{k=1}^\infty$ is bounded in $H^2(U)$ and so possesses a strongly convergent subsequence in $H_0^1(U)$.

3. Finally, we must show that if μ is large enough, the set

$$\{u \in H_0^1(U) \mid u = \lambda A[u] \text{ for some } 0 \leq \lambda \leq 1\}$$

is bounded in $H_0^1(U)$. So assume $u \in H_0^1(U)$,

$$u = \lambda A[u] \quad \text{for some } 0 < \lambda \leq 1.$$

Then $\frac{u}{\lambda} = A[u]$; or, in other words, $u \in H^2(U) \cap H_0^1(U)$ and

$$-\Delta u + \mu u = \lambda b(Du) \quad \text{a.e. in } U.$$

Multiply this identity by u and integrate over U , to compute

$$\begin{aligned} \int_U |Du|^2 + \mu |u|^2 \, dx &= - \int_U \lambda b(Du)u \, dx \leq \int_U C(|Du| + 1)|u| \, dx \\ &\leq \frac{1}{2} \int_U |Du|^2 \, dx + C \int_U |u|^2 + 1 \, dx. \end{aligned}$$

Thus if $\mu > 0$ is sufficiently large, we have $\|u\|_{H_0^1(U)} \leq C$, for some constant C that does not depend on $0 \leq \lambda \leq 1$.

4. Applying Schaefer's Fixed Point Theorem in the space $X = H_0^1(U)$, we conclude that A has a fixed point $u \in H_0^1(U) \cap H^2(U)$, which in turn solves our semilinear PDE (21). \square

Warning. A plausible plan for constructively solving (21) would be to select some u^0 and then iteratively solve the linear boundary-value problems

$$\begin{cases} -\Delta u^{k+1} + \mu u^{k+1} = -b(Du^k) & \text{in } U \\ u^{k+1} = 0 & \text{on } \partial U \end{cases} \quad (k = 0, 1, \dots).$$

However, we *cannot* assert that $\{u^k\}_{k=0}^\infty$ then converges to a solution of (21). Schauder's and Schaefer's Fixed Point Theorems do not say that any sequence converges to a fixed point. (But see the proof in §9.3 following.)

See Gilbarg–Trudinger [G-T] for much more sophisticated applications of fixed point theorems to nonlinear elliptic PDE.

9.3. METHOD OF SUBSOLUTIONS AND SUPERSOLUTIONS

Our application of Schaefer's Theorem above in §9.2.2 depends upon the regularity estimates for solutions of elliptic equations. We turn our attention now to another basic property of elliptic PDE, namely the maximum principle, and demonstrate how various resulting comparison arguments can be used to solve certain semilinear problems. The idea is to exploit *ordering properties* for solutions. More precisely, we will show that if we can find a subsolution \underline{u} and a supersolution \bar{u} of a particular boundary-value problem and if furthermore $\underline{u} \leq \bar{u}$, then there in fact exists a solution satisfying

$$\underline{u} \leq u \leq \bar{u}.$$

We will investigate this boundary-value problem for the nonlinear Poisson equation:

$$(1) \quad \begin{cases} -\Delta u = f(u) & \text{in } U \\ u = 0 & \text{on } \partial U, \end{cases}$$

where $f : \mathbb{R} \rightarrow \mathbb{R}$ is smooth, with

$$(2) \quad |f'| \leq C \quad (z \in \mathbb{R})$$

for some constant C .

DEFINITIONS. (i) We say that $\bar{u} \in H^1(U)$ is a weak supersolution of problem (1) if

$$(3) \quad \int_U D\bar{u} \cdot Dv \, dx \geq \int_U f(\bar{u})v \, dx$$

for each $v \in H_0^1(U)$, $v \geq 0$ a.e.

(ii) Similarly, $\underline{u} \in H^1(U)$ is a weak subsolution provided

$$(4) \quad \int_U D\underline{u} \cdot Dv \, dx \leq \int_U f(\underline{u})v \, dx$$

for each $v \in H_0^1(U)$, $v \geq 0$ a.e.

(iii) We say $u \in H_0^1(U)$ is a weak solution of (1) if

$$\int_U Du \cdot Dv \, dx = \int_U f(u)v \, dx$$

for each $v \in H_0^1(U)$.

Remark. If $\bar{u}, \underline{u} \in C^2(U)$, then from (3) and (4) it follows that

$$-\Delta \bar{u} \geq f(\bar{u}), \quad -\Delta \underline{u} \leq f(\underline{u}) \quad \text{in } U.$$

THEOREM 1 (Existence of a solution between sub- and supersolutions). Assume there exist a weak supersolution \bar{u} and a weak subsolution \underline{u} of (1), satisfying

$$(5) \quad \underline{u} \leq 0, \quad \bar{u} \geq 0 \text{ on } \partial U \text{ in the trace sense, } \underline{u} \leq \bar{u} \text{ a.e. in } U.$$

Then there exists a weak solution u of (1), such that

$$\underline{u} \leq u \leq \bar{u} \quad \text{a.e. in } U.$$

Proof. 1. Fix a number $\lambda > 0$ so large that

$$(6) \quad \text{the mapping } z \mapsto f(z) + \lambda z \text{ is nondecreasing;}$$

this is possible as a consequence of hypothesis (2).

Now write $u_0 = \underline{u}$, and then given u_k ($k = 0, 1, 2, \dots$), inductively define $u_{k+1} \in H_0^1(U)$ to be the unique weak solution of the linear boundary-value problem

$$(7) \quad \begin{cases} -\Delta u_{k+1} + \lambda u_{k+1} = f(u_k) + \lambda u_k & \text{in } U \\ u_{k+1} = 0 & \text{on } \partial U. \end{cases}$$

2. We claim

$$(8) \quad \underline{u} = u_0 \leq u_1 \leq \dots \leq u_k \leq \dots \quad \text{a.e. in } U.$$

To confirm this, first note from (7) for $k = 0$ that

$$(9) \quad \int_U Du_1 \cdot Dv + \lambda u_1 v \, dx = \int_U (f(u_0) + \lambda u_0) v \, dx$$

for each $v \in H_0^1(U)$. Subtract (9) from (4), recall $u_0 = \underline{u}$, and set

$$v := (u_0 - u_1)^+ \in H_0^1(U), \quad v \geq 0 \quad \text{a.e.}$$

We find

$$(10) \quad \int_U D(u_0 - u_1) \cdot D(u_0 - u_1)^+ + \lambda(u_0 - u_1)(u_0 - u_1)^+ \, dx \leq 0.$$

But

$$D(u_0 - u_1)^+ = \begin{cases} D(u_0 - u_1) & \text{a.e. on } \{u_0 \geq u_1\} \\ 0 & \text{a.e. on } \{u_0 \leq u_1\}. \end{cases}$$

(See Problem 18 in Chapter 5.) Consequently,

$$\int_{\{u_0 \geq u_1\}} |D(u_0 - u_1)|^2 + \lambda(u_0 - u_1)^2 \, dx \leq 0,$$

so that $u_0 \leq u_1$ a.e. in U .

Now assume inductively that

$$(11) \quad u_{k-1} \leq u_k \quad \text{a.e. in } U.$$

From (7) we find

$$(12) \quad \int_U Du_{k+1} \cdot Dv + \lambda u_{k+1} v \, dx = \int_U (f(u_k) + \lambda u_k) v \, dx$$

and

$$\int_U Du_k \cdot Dv + \lambda u_k v \, dx = \int_U (f(u_{k-1}) + \lambda u_{k-1}) v \, dx$$

for each $v \in H_0^1(U)$. Subtract and set $v := (u_k - u_{k+1})^+$. We deduce

$$\begin{aligned} & \int_{\{u_k \geq u_{k+1}\}} |D(u_k - u_{k+1})|^2 + \lambda(u_k - u_{k+1})^2 \, dx \\ &= \int_U [(f(u_{k-1}) + \lambda u_{k-1}) - (f(u_k) + \lambda u_k)] (u_k - u_{k+1})^+ \, dx \leq 0, \end{aligned}$$

the last inequality holding in view of (11) and (6). Therefore $u_k \leq u_{k+1}$ a.e. in U , as asserted.

3. Next we show

$$(13) \quad u_k \leq \bar{u} \quad \text{a.e. in } U \quad (k = 0, 1, \dots).$$

Statement (13) is valid for $k = 0$ by hypothesis (5). Assume now for induction that

$$(14) \quad u_k \leq \bar{u} \quad \text{a.e. in } U.$$

Then subtracting (3) from (12) and taking $v := (u_{k+1} - \bar{u})^+$, we find

$$\begin{aligned} & \int_{\{u_{k+1} \geq \bar{u}\}} |D(u_{k+1} - \bar{u})|^2 + \lambda(u_{k+1} - \bar{u})^2 dx \\ & \leq \int_U [(f(u_k) + \lambda u_k) - (f(\bar{u}) + \lambda \bar{u})](u_{k+1} - \bar{u})^+ dx \leq 0, \end{aligned}$$

by (14) and (6). Thus $u_{k+1} \leq \bar{u}$ a.e. in U .

4. In light of (8) and (13), we have

$$(15) \quad \underline{u} \leq \dots \leq u_k \leq u_{k+1} \leq \dots \leq \bar{u} \quad \text{a.e. in } U.$$

Therefore

$$u(x) := \lim_{k \rightarrow \infty} u_k(x)$$

exists for a.e. x . Furthermore we have

$$(16) \quad u_k \rightarrow u \quad \text{in } L^2(U),$$

as guaranteed by the Dominated Convergence Theorem and (15). Finally, since we have $\|f(u_k)\|_{L^2(U)} \leq C(\|u_k\|_{L^2(U)} + 1)$, we deduce from (7) that $\sup_k \|u_k\|_{H_0^1(U)} < \infty$. Hence there is a subsequence $\{u_{k_j}\}_{j=1}^\infty$ which converges weakly in $H_0^1(U)$ to $u \in H_0^1(U)$.

5. We at last verify that u is a weak solution of problem (1). For this, fix $v \in H_0^1(U)$. Then from (7) we find

$$\int_U Du_{k+1} \cdot Dv + \lambda u_{k+1} v dx = \int_U (f(u_k) + \lambda u_k) v dx.$$

Let $k \rightarrow \infty$:

$$\int_U Du \cdot Dv + \lambda uv dx = \int_U (f(u) + \lambda u) v dx.$$

Canceling the term involving λ , we at last confirm that

$$\int_U Du \cdot Dv dx = \int_U f(u) v dx,$$

as desired. □

This proof illustrates the use of integration by parts, rather than the maximum principle, to establish comparisons between sub- and supersolutions.

9.4. NONEXISTENCE OF SOLUTIONS

We now complement the theory in §§9.1–9.3 with some *nonexistence* assertions for solutions of various nonlinear partial differential equations. The overall procedure will be to assume there exists a solution and then to obtain certain inequalities, which in turn force a contradiction.

9.4.1. Blow-up.

Blow-up for large enough initial data. We begin by considering an initial/boundary-value problem for a parabolic equation with a simple quadratic nonlinearity:

$$(1) \quad \begin{cases} u_t - \Delta u = u^2 & \text{in } U_T \\ u = 0 & \text{on } \partial U \times (0, T) \\ u = g & \text{on } U \times \{t = 0\}. \end{cases}$$

We will show that if $T > 0$ and $g \geq 0$ are large enough in an appropriate sense, then there does *not* exist a smooth solution u of (1). We can regard the nonlinear heat equation in (1) as a simple reaction-diffusion equation (cf. Example 1 in §9.2.1). The nonlinear term alone corresponds to the ODE

$$\dot{u} = u^2 \quad \left(\cdot = \frac{d}{dt} \right),$$

which certainly blows up in finite time, provided $u(0) > 0$. The purely diffusive effects on the other hand yield the heat equation, which tends to smooth out irregularities. The following analysis must therefore untangle the competing effects of blow-up from the u^2 term and smoothing from the Δu term.

We proceed by choosing w_1 to be an eigenfunction corresponding to the principal eigenvalue $\lambda_1 > 0$ of $-\Delta$ in $H_0^1(U)$. Then owing to the theory in §6.5.1, w_1 is smooth,

$$\begin{cases} -\Delta w_1 = \lambda_1 w_1 & \text{in } U \\ w_1 = 0 & \text{on } \partial U, \end{cases}$$

and we may furthermore assume

$$(2) \quad w_1 > 0 \text{ in } U, \quad \int_U w_1 \, dx = 1.$$

Suppose u is a smooth solution of (1), with $g \geq 0, g \not\equiv 0$, so that $u > 0$ within U_T by the strong maximum principle.

THEOREM 1 (Blow-up for large data). *Assume that*

$$(3) \quad \int_U g w_1 dx > \lambda_1.$$

Then there cannot exist a smooth solution u of (1) existing for all times $T > 0$.

Proof. Define

$$\eta(t) := \int_U u(x, t) w_1(x) dx \quad (0 \leq t \leq T).$$

Then

$$(4) \quad \begin{aligned} \dot{\eta} &= \int_U u_t w_1 dx = \int_U (\Delta u + u^2) w_1 dx \\ &= \int_U u \Delta w_1 + u^2 w_1 dx = -\lambda_1 \eta + \int_U u^2 w_1 dx. \end{aligned}$$

Furthermore

$$\begin{aligned} \eta &= \int_U u w_1 dx = \int_U u w_1^{1/2} w_1^{1/2} dx \\ &\leq \left(\int_U u^2 w_1 dx \right)^{1/2} \left(\int_U w_1 dx \right)^{1/2} \\ &= \left(\int_U u^2 w_1 dx \right)^{1/2} \quad \text{by (2)}. \end{aligned}$$

Employing this inequality in (4), we find

$$(5) \quad \dot{\eta} \geq -\lambda_1 \eta + \eta^2.$$

Writing $\xi(t) := e^{\lambda_1 t} \eta(t)$ gives

$$\dot{\xi} = e^{\lambda_1 t} \dot{\eta} + \lambda_1 e^{\lambda_1 t} \eta \geq e^{\lambda_1 t} \eta^2 = e^{-\lambda_1 t} \xi^2.$$

Thus

$$\left(\frac{-1}{\xi} \right)' = \frac{\dot{\xi}}{\xi^2} \geq e^{-\lambda_1 t},$$

and therefore

$$\frac{-1}{\xi(t)} \geq \frac{-1}{\xi(0)} + \frac{1 - e^{-\lambda_1 t}}{\lambda_1}.$$

Rearranging, we deduce

$$\xi(t) \geq \frac{\xi(0) \lambda_1}{\lambda_1 - \xi(0)(1 - e^{-\lambda_1 t})},$$

provided the denominator is not zero. But now suppose (3) holds, so that $\eta(0) = \xi(0) > \lambda$. Then $\xi(t) \rightarrow +\infty$ as $t \rightarrow t^*$, for $t^* := \frac{-1}{\lambda_1} \log \left(\frac{\eta(0) - \lambda_1}{\eta(0)} \right)$. \square

Our proof shows that either the solution is not smooth enough to justify the calculation above or else

$$\lim_{t \rightarrow t_*} \int_U u(x, t) w_1(x) dx = \infty$$

for some time $0 < t_* \leq t^*$. In this case we say u “blows up” at time t_* .

Blow-up for small initial data. We discuss next an interesting variant of the foregoing proof, now for this initial-value problem in all of space:

$$(6) \quad \begin{cases} u_t - \Delta u = u^p & \text{in } \mathbb{R}^n \times (0, T) \\ u = g & \text{on } \mathbb{R}^n \times \{t = 0\}. \end{cases}$$

We assume the nonnegative initial function $g \not\equiv 0$ is smooth, with compact support.

THEOREM 2 (Blow-up for small data). *Assume*

$$(7) \quad 1 < p < \frac{n+2}{n}.$$

Then there cannot exist a nonnegative, integrable and smooth solution u of (6) existing for all times $T > 0$.

Proof. Since we are working in all of \mathbb{R}^n , we must replace the eigenfunction used before. Instead, introduce the fundamental solution of the heat equation evaluated at a time $s > 0$ (which we will select later):

$$\Phi(x, s) = \frac{1}{(4\pi s)^{n/2}} e^{-\frac{|x|^2}{4s}}.$$

Recall that

$$\int_{\mathbb{R}^n} \Phi(x, s) dx = 1.$$

Furthermore, a calculation shows that

$$(8) \quad \Delta \Phi(x, s) + \frac{n}{2s} \Phi(x, s) \geq 0.$$

Define

$$\eta(t) := \int_{\mathbb{R}^n} u(x, t) \Phi(x, s) dx.$$

Then (8) implies

$$\begin{aligned}\dot{\eta}(t) &= \int_{\mathbb{R}^n} u_t \Phi \, dx = \int_{\mathbb{R}^n} (\Delta u + u^p) \Phi \, dx \\ &= \int_{\mathbb{R}^n} u \Delta \Phi + u^p \Phi \, dx \geq -\frac{n}{2s} \eta(t) + \int_{\mathbb{R}^n} u^p \Phi \, dx\end{aligned}$$

for $\Phi = \Phi(x, s)$. As in the previous proof,

$$\eta^p \leq \int_{\mathbb{R}^n} u^p \Phi \, dx.$$

Therefore

$$(9) \quad \dot{\eta} \geq -\lambda \eta + \eta^p$$

for

$$\lambda := \frac{n}{2s}.$$

We integrate (9) by setting $\xi(t) := e^{\lambda t} \eta(t)$. Then

$$\dot{\xi} \geq e^{-\lambda(p-1)t} \xi^p.$$

Integrating and rewriting, we discover that

$$\xi^{p-1}(t) \geq \frac{\xi^{p-1}(0) \lambda}{\lambda - \xi^{p-1}(0)(1 - e^{-\lambda(p-1)t})}.$$

This differential inequality shows that $\xi \rightarrow \infty$ in finite time if

$$\eta(0) = \xi(0) > \lambda^{\frac{1}{p-1}}.$$

Recalling the definitions of λ and Φ , we rewrite this condition as saying

$$(10) \quad \frac{1}{(4\pi)^{n/2}} \int_{\mathbb{R}^n} g e^{-\frac{|x|^2}{4s}} \, dx > s^{n/2} \left(\frac{n}{2s} \right)^{\frac{1}{p-1}} = \gamma s^{\frac{n}{2} - \frac{1}{p-1}},$$

where $\gamma > 0$. But our condition (7) on p implies $\frac{n}{2} - \frac{1}{p-1} < 0$, and hence for any initial function $g \not\equiv 0$ we can select s so large that (10) is valid. \square

9.4.2. Derrick–Pohozaev identity.

We investigate next a nonlinear elliptic PDE to which different differential inequality methods apply, namely the nonlinear boundary-value problem

$$(11) \quad \begin{cases} -\Delta u = |u|^{p-1}u & \text{in } U \\ u = 0 & \text{on } \partial U. \end{cases}$$

Now the theory in §8.5.2 applies to (11) provided

$$(12) \quad 1 < p < \frac{n+2}{n-2}$$

and proves the existence of a nontrivial solution $u \neq 0$. Let us now instead suppose


$$(13) \quad \frac{n+2}{n-2} < p < \infty.$$

Our goal is to demonstrate under a certain geometric condition on U that (13) implies $u \equiv 0$ is the only smooth solution of (11). We see therefore that the restriction to condition (12) in §8.5.2 was in some sense natural, and consequently say $p = \frac{n+2}{n-2}$ is a *critical exponent*.

DEFINITION. An open set U is called *star-shaped with respect to 0* provided for each $x \in \bar{U}$, the line segment

$$\{\lambda x \mid 0 \leq \lambda \leq 1\}$$

lies in \bar{U} .



A star-shaped domain

Clearly if U is convex and $0 \in U$, then U is star-shaped with respect to 0. But a general star-shaped region need not be convex.

LEMMA (Normals to a star-shaped region). *Assume ∂U is C^1 and U is star-shaped with respect to 0. Then*

$$x \cdot \nu(x) \geq 0 \quad \text{for all } x \in \partial U,$$

where ν denotes the unit outward normal.

Proof. Since ∂U is C^1 , if $x \in \partial U$, then for each $\epsilon > 0$ there exists $\delta > 0$ such that $|y - x| < \delta$ and $y \in \bar{U}$ imply $\nu(x) \cdot \frac{(y-x)}{|y-x|} \leq \epsilon$. In particular

$$\limsup_{\substack{y \rightarrow x \\ y \in \bar{U}}} \nu(x) \cdot \frac{(y-x)}{|y-x|} \leq 0.$$

Let $y = \lambda x$ for $0 < \lambda < 1$. Then $y \in \bar{U}$, since U is star-shaped. Thus

$$\nu(x) \cdot \frac{x}{|x|} = - \lim_{\lambda \rightarrow 1^-} \nu(x) \cdot \frac{(\lambda x - x)}{|\lambda x - x|} \geq 0. \quad \square$$

We next prove that there can exist no nontrivial solution to problem (11) for supercritical growth, provided U is star-shaped. The proof is a remarkable calculation initiated by multiplying the PDE $-\Delta u = |u|^{p-1}u$ by $x \cdot Du$ and continually integrating by parts. Our selection of the multiplier $x \cdot Du$ is inspired by Example 2 in §8.6.2.

THEOREM 3 (Nonexistence of nontrivial solution). *Assume $u \in C^2(\bar{U})$ is a solution of problem (11) and the exponent p satisfies inequality (13). Suppose further U is star-shaped with respect to 0 and ∂U is C^1 . Then*

$$u \equiv 0 \quad \text{within } U.$$

Proof. 1. We multiply the PDE by $x \cdot Du$ and integrate over U , to find

$$(14) \quad \int_U (-\Delta u)(x \cdot Du) dx = \int_U |u|^{p-1}u(x \cdot Du) dx.$$

We rewrite this expression as

$$A = B.$$

2. The term on the left is

$$\begin{aligned}
 (15) \quad A &:= - \sum_{i,j=1}^n \int_U u_{x_i x_i} x_j u_{x_j} dx \\
 &= \sum_{i,j=1}^n \int_U u_{x_i} (x_j u_{x_j})_{x_i} dx - \sum_{i,j=1}^n \int_{\partial U} u_{x_i} \nu^i x_j u_{x_j} dS \\
 &=: A_1 + A_2.
 \end{aligned}$$

3. Now

$$\begin{aligned}
 (16) \quad A_1 &= \sum_{i,j=1}^n \int_U u_{x_i} \delta_{ij} u_{x_j} + u_{x_i} x_j u_{x_j x_i} dx \\
 &= \int_U |Du|^2 + \sum_{j=1}^n \left(\frac{|Du|^2}{2} \right)_{x_j} x_j dx \\
 &= \left(1 - \frac{n}{2} \right) \int_U |Du|^2 dx + \int_{\partial U} \frac{|Du|^2}{2} (\nu \cdot x) dS.
 \end{aligned}$$

On the other hand, since $u = 0$ on ∂U , $Du(x)$ is parallel to the normal $\nu(x)$ at each point $x \in \partial U$. Thus $Du(x) = \pm |Du(x)| \nu(x)$. Using this equality, we calculate

$$(17) \quad A_2 = - \int_{\partial U} |Du|^2 (\nu \cdot x) dS.$$

Combine (15)–(17), to deduce

$$A = \frac{2-n}{2} \int_U |Du|^2 dx - \frac{1}{2} \int_{\partial U} |Du|^2 (\nu \cdot x) dS.$$

4. Returning to (14), we compute

$$\begin{aligned}
 B &:= \sum_{j=1}^n \int_U |u|^{p-1} u_{x_j} u_{x_j} dx \\
 &= \sum_{j=1}^n \int_U \left(\frac{|u|^{p+1}}{p+1} \right)_{x_j} x_j dx = - \frac{n}{p+1} \int_U |u|^{p+1} dx.
 \end{aligned}$$

5. This calculation and (14) yield

$$(18) \quad \left(\frac{n-2}{2} \right) \int_U |Du|^2 dx + \frac{1}{2} \int_{\partial U} |Du|^2 (\nu \cdot x) dS = \frac{n}{p+1} \int_U |u|^{p+1} dx.$$

In view of the lemma above, we then obtain the inequality

$$(19) \quad \left(\frac{n-2}{2}\right) \int_U |Du|^2 dx \leq \frac{n}{p+1} \int_U |u|^{p+1} dx.$$

But once we multiply the PDE $-\Delta u = |u|^{p-1}u$ by u and integrate by parts, we produce the equality

$$\int_U |Du|^2 dx = \int_U |u|^{p+1} dx.$$

Substituting into (19), we thus conclude

$$\left(\frac{n-2}{2} - \frac{n}{p+1}\right) \int_U |u|^{p+1} dx \leq 0.$$

Hence if $u \not\equiv 0$, it follows that $\frac{n-2}{2} - \frac{n}{p+1} \leq 0$; that is, $p \leq \frac{n+2}{n-2}$. □

The equality (18) is sometimes called the *Derrick-Pohozaev identity*.

9.5. GEOMETRIC PROPERTIES OF SOLUTIONS

9.5.1. Star-shaped level sets.

We explain in this subsection a simple method that is occasionally useful for studying the geometric properties of the level sets of solutions to various PDE. The easiest such case occurs when we look at harmonic functions in an open set U having the form

$$U = W - \bar{V},$$

where $V \subset\subset W$, for open sets V, W , each of which is star-shaped with respect to 0. Write

$$\Gamma_0 = \partial W, \quad \Gamma_1 = \partial V.$$

We consider the problem

$$(1) \quad \begin{cases} \Delta u = 0 & \text{in } U \\ u = 1 & \text{on } \Gamma_1 \\ u = 0 & \text{on } \Gamma_0. \end{cases}$$

Physically u corresponds to the electrostatic potential generated within the region U , once we fix the potential value to be one on Γ_1 and zero on Γ_0 . According to the strong maximum principle $0 < u < 1$ within U .

THEOREM 1 (Star-shaped level sets). *For each $0 < \lambda < 1$ the level set*

$$\Gamma_\lambda := \{x \in U \mid u(x) = \lambda\}$$

is a smooth surface and is the boundary of a set star-shaped with respect to 0.

Proof. 1. For each $\mu > 0$, the function $x \mapsto u(\mu x)$ is harmonic, and thus so is

$$v(x) := \frac{d}{d\mu}(u(\mu x))|_{\mu=1} = Du(x) \cdot x \quad (x \in U).$$

Now since $u = 0$ on Γ_0 , $Du(x)$ points in the direction of $-\nu(x)$ at each point $x \in \partial W$. Additionally, we have $x \cdot \nu(x) \geq 0$ on Γ_0 , since W is star-shaped with respect to 0. Consequently $v = Du \cdot x \leq 0$ on Γ_0 . Similarly $v \leq 0$ on Γ_1 . According then to the strong maximum principle for harmonic functions, $v < 0$ in U . In particular, $Du \neq 0$ within U . Consequently the Implicit Function Theorem (§C.7) implies that Γ_λ is a smooth surface for $0 < \lambda < 1$.

2. Extend u to equal 1 on all of V and write

$$U_\lambda := \{x \in W \mid u > \lambda\}.$$

Then U_λ is an open subset of W and $\partial U_\lambda = \Gamma_\lambda$. By the strong maximum principle, U_λ is connected.

Now let $x \in \Gamma_\lambda$ and let $\nu(x)$ denote the outer unit normal to Γ_λ at x . Then $Du(x)$ points in the direction of $-\nu(x)$. Since $v(x) < 0$, we have $x \cdot \nu(x) > 0$. This inequality holds for each $x \in \Gamma_\lambda$.

3. It follows that Γ_λ is the boundary of a set star-shaped with respect to 0. To see this, return to the proof of the lemma in §9.4.2, and notice that if Γ_λ were not star-shaped with respect to 0, we could then find a point $x \in \Gamma_\lambda$ for which $y = \mu x \notin U_\lambda$ if μ is close to 1, $\mu < 1$. But then we can derive the contradiction

$$\nu(x) \cdot \frac{x}{|x|} = - \lim_{\mu \rightarrow 1^-} \nu(x) \cdot \frac{(\mu x - x)}{|\mu x - x|} \leq 0. \quad \square$$

9.5.2. Radial symmetry.

In this subsection we take $U = B^0(0, 1)$ to be the open unit ball in \mathbb{R}^n and investigate this boundary-value problem for a semilinear Poisson PDE:

$$(2) \quad \begin{cases} -\Delta u = f(u) & \text{in } U \\ u = 0 & \text{on } \partial U. \end{cases}$$

We are interested in *positive solutions*:

$$(3) \quad u > 0 \quad \text{in } U$$

and will assume $f : \mathbb{R} \rightarrow \mathbb{R}$ is Lipschitz continuous but is otherwise arbitrary. Our intention is to prove that u is necessarily radial, that is, $u(x)$ depends only on $r = |x|$. This is quite an unexpectedly strong conclusion, since we are making essentially no assumption on the nonlinearity.

a. Maximum principles. Our proofs will depend upon an extension of the maximum principle for second-order elliptic PDE.

LEMMA 1 (A refinement of Hopf's Lemma). *Suppose $V \subset \mathbb{R}^n$ is open, $v \in C^2(\bar{V})$, and $c \in L^\infty(V)$. Assume*

$$(4) \quad \begin{cases} -\Delta v + cv \geq 0 & \text{in } V \\ v \geq 0 & \text{in } V. \end{cases}$$

Suppose also $v \not\equiv 0$.

(i) *If $x^0 \in \partial V$, $v(x^0) = 0$, and V satisfies the interior ball condition at x^0 , then*

$$(5) \quad \frac{\partial v}{\partial \nu}(x^0) < 0.$$

(ii) *Furthermore,*

$$(6) \quad v > 0 \quad \text{in } V.$$

Observe that we are here making no hypothesis concerning the sign of the zeroth-order coefficient c .

Proof. Let $w := e^{-\lambda x_1} v$, where $\lambda > 0$ will be selected below. Then $v = e^{\lambda x_1} w$, and so

$$cv \geq \Delta v = \Delta(e^{\lambda x_1} w) = \lambda^2 v + 2\lambda e^{\lambda x_1} w_{x_1} + e^{\lambda x_1} \Delta w.$$

Thus

$$-\Delta w - 2\lambda w_{x_1} \geq (\lambda^2 - c)w \geq 0 \quad \text{in } V,$$

if $\lambda = \|c\|_{L^\infty}^{1/2}$.

Consequently w is a supersolution for the elliptic operator $Kw := -\Delta w - 2\lambda w_{x_1}$, which has no zeroth-order term. The strong maximum principle implies $w > 0$ in V . According to Hopf's Lemma (§6.4.2) therefore, $\frac{\partial w}{\partial \nu}(x^0) < 0$. But

$$\frac{\partial w}{\partial \nu}(x^0) = Dw(x^0) \cdot \nu(x^0) = e^{-\lambda x_1^0} \frac{\partial v}{\partial \nu}(x^0)$$

since $v(x^0) = 0$. Assertion (i) therefore holds, and assertion (ii) follows since $w > 0$ in V . \square

LEMMA 2 (Boundary estimates). *Let $u \in C^2(\bar{U})$ satisfy (2), (3). Then for each point $x^0 \in \partial U \cap \{x_n > 0\}$, either*

$$(7) \quad u_{x_n}(x^0) < 0$$

or else

$$(8) \quad u_{x_n}(x^0) = 0, \quad u_{x_n x_n}(x^0) > 0.$$

In either case, u is strictly decreasing as a function of x_n near x^0 .

Proof. 1. Fix any point $x^0 \in \partial U \cap \{x_n > 0\}$ and let $\nu = \nu(x^0) = (\nu_1, \dots, \nu_n)$ denote the outer unit normal to ∂U at x^0 . Note $\nu_n > 0$.

2. We first claim

$$u_{x_n}(x^0) < 0,$$

provided

$$(9) \quad f(0) \geq 0.$$

Indeed

$$\begin{aligned} 0 &= -\Delta u - f(u) = -\Delta u - f(u) + f(0) - f(0) \\ &\leq -\Delta u + cu, \end{aligned}$$

for $c(x) := -\int_0^1 f'(su(x)) ds$. According to Lemma 1, $\frac{\partial u}{\partial \nu}(x^0) < 0$. Since Du is parallel to ν on ∂U and $\nu_n > 0$, we conclude $u_{x_n}(x^0) < 0$.

3. Now suppose

$$(10) \quad f(0) < 0.$$

If $u_{x_n}(x^0) < 0$, we are done. Otherwise, since Du is parallel to ν ,

$$(11) \quad Du(x^0) = 0.$$


As (2) is invariant under a rotation of coordinate axes, we may as well suppose $x^0 = (0, \dots, 1)$, $\nu = (0, \dots, 1)$.

4. We assert

$$(12) \quad u_{x_i x_j}(x^0) = -f(0)\nu_i \nu_j \quad \text{for each } i, j = 1, \dots, n.$$

Since $u = 0$ on ∂U , we have $u(x', \gamma(x')) = 0$ for all $x' \in \mathbb{R}^{n-1}$, $|x'| \leq 1$, where $\gamma(x') = (1 - |x'|^2)^{1/2}$. Differentiating with respect to x_i and x_j ($i, j = 1, \dots, n-1$) and using (11), we conclude

$$(13) \quad u_{x_i x_j}(x^0) = 0 \quad (i, j = 1, \dots, n-1).$$



Reflection through a plane

Since $u_{x_n} \leq 0$ on $\partial U \cap \{x_n > 0\}$ and $u_{x_n}(x^0) = 0$, the mapping $x' \mapsto u_{x_n}(x', \gamma(x'))$ has a maximum at $x' = 0$. Thus

$$(14) \quad u_{x_n x_i}(x^0) = 0 \quad (i = 1, \dots, n-1).$$

Finally, (13), (14) and the PDE (2) force $u_{x_n x_n}(x^0) = -f(0)$. This equality is (12) for $\nu = (0, \dots, 1)$. Returning to the original coordinate axes, we obtain (12).

5. Setting $i = j = n$ in (12), we find using (10) that

$$u_{x_n x_n}(x^0) = -f(0)\nu_n \nu_n > 0. \quad \square$$

b. Moving planes. We introduce next a “moving plane” P_λ , across which we will reflect our partial differential equation.

NOTATION. (i) If $0 \leq \lambda \leq 1$, define the plane

$$P_\lambda := \{x \in \mathbb{R}^n \mid x_n = \lambda\}.$$

(ii) Write $x_\lambda := (x_1, \dots, x_{n-1}, 2\lambda - x_n)$ to denote the *reflection* of x in P_λ .

(iii) $E_\lambda := \{x \in U \mid \lambda < x_n < 1\}$.

THEOREM 2 (Radial symmetry). *Let $u \in C^2(\bar{U})$ solve (2), (3). Then u is radial; that is,*

$$u(x) = v(r) \quad (r = |x|)$$

for some strictly decreasing function $v : [0, 1] \rightarrow [0, \infty)$.

Proof. 1. We consider for each $0 \leq \lambda < 1$ the statement

$$(15_\lambda) \quad u(x) < u(x_\lambda) \text{ for each point } x \in E_\lambda.$$

2. According to Lemma 2, (15_λ) is valid for each $\lambda < 1$, λ sufficiently close to 1. Set

$$(16) \quad \lambda_0 := \inf\{0 \leq \lambda < 1 \mid (15_\mu) \text{ holds for each } \lambda \leq \mu < 1\}.$$

We will prove

$$(17) \quad \lambda_0 = 0.$$

Assume instead $\lambda_0 > 0$. Write $w(x) := u(x_{\lambda_0}) - u(x)$ ($x \in E_{\lambda_0}$). Then

$$-\Delta w = f(u(x_{\lambda_0})) - f(u(x)) = -cw \quad \text{in } E_{\lambda_0},$$

for $c(x) := -\int_0^1 f'(su(x_{\lambda_0}) + (1-s)u(x)) ds$. As $w \geq 0$ in E_{λ_0} , we deduce from Lemma 1 (applied to $V = E_{\lambda_0}$) that $w > 0$ in E_{λ_0} , $w_{x_n} > 0$ on $P_{\lambda_0} \cap U$. Thus

$$(18) \quad u(x) < u(x_{\lambda_0}) \quad \text{in } E_{\lambda_0},$$

and

$$(19) \quad u_{x_n} < 0 \quad \text{on } P_{\lambda_0} \cap U.$$

Using (18), (19) and Lemma 2, we conclude

$$(20) \quad u(x) < u(x_{\lambda_0-\varepsilon}) \quad \text{in } E_{\lambda_0-\varepsilon} \text{ for all } 0 \leq \varepsilon \leq \varepsilon_0,$$

if ε_0 is small enough. Assertion (20) contradicts our choice (16) of λ_0 , if $\lambda_0 > 0$.

3. Since $\lambda_0 = 0$, we see $u(x_1, \dots, x_{n-1}, -x_n) \geq u(x_1, \dots, x_n)$ for all $x \in U \cap \{x_n > 0\}$. A similar argument in $U \cap \{x_n < 0\}$ shows $u(x_1, \dots, x_{n-1}, -x_n) \leq u(x_1, \dots, x_n)$ for all $x \in U \cap \{x_n > 0\}$. Thus u is symmetric in the plane P_0 and $u_{x_n} = 0$ on P_0 .

This argument applies as well after any rotation of coordinate axes, and so the theorem follows. \square

9.6. GRADIENT FLOWS

In this section we augment our discussion of abstract semigroup theory for linear operators (§7.4) by introducing certain nonlinear semigroups, generated by convex functions. Applications include various nonlinear second-order parabolic partial differential equations.

9.6.1. Convex functions on Hilbert spaces.

Convexity has been an essential ingredient in much of our analysis of nonlinear PDE thus far. We now broaden our view by considering convex functions defined on (possibly infinite-dimensional) Hilbert spaces.

Hereafter H will denote a real Hilbert space, with inner product (\cdot, \cdot) and norm $\|\cdot\|$.

DEFINITION. *A function*

$$I : H \rightarrow (-\infty, \infty]$$

is convex provided

$$I[\tau u + (1 - \tau)v] \leq \tau I[u] + (1 - \tau)I[v]$$

for all $u, v \in H$ and each $0 \leq \tau \leq 1$.

Note carefully that we allow I to take on the value $+\infty$ (but not $-\infty$). The function I is called *proper* if I is not identically equal to $+\infty$. The *domain* of I is

$$D(I) := \{u \in H \mid I[u] < +\infty\}.$$

DEFINITION. *We say $I : H \rightarrow (-\infty, +\infty]$ is lower semicontinuous if*

$$\begin{cases} u_k \rightarrow u \text{ in } H \text{ implies} \\ I[u] \leq \liminf_{k \rightarrow \infty} I[u_k]. \end{cases}$$

As in the finite-dimensional case (cf. §B.1), it is important to understand when the graph of I has a supporting hyperplane.

DEFINITIONS. *Let $I : H \rightarrow (-\infty, +\infty]$ be convex and proper.*

(i) *For each $u \in H$, we write*

$$(1) \quad \partial I[u] := \{v \in H \mid I[w] \geq I[u] + (v, w - u) \text{ for all } w \in H\}.$$

The mapping $\partial I : H \rightarrow 2^H$ is the subdifferential of I .

(ii) *We say $u \in D(\partial I)$, the domain of ∂I , provided $\partial I[u] \neq \emptyset$.*

The geometric interpretation of (1) is that $v \in \partial I[u]$ if and only if v is the “slope” of an affine functional touching the graph of I from below at the point u . Since this graph may have a “corner” at u , $\partial I[u]$ could be multivalued.

THEOREM 1 (Properties of subdifferentials). *Let $I : H \rightarrow (-\infty, +\infty]$ be convex, proper and lower semicontinuous. Then*

(i) $D(\partial I) \subseteq D(I)$.

(ii) *If $v \in \partial I[u]$ and $\tilde{v} \in \partial I[\tilde{u}]$, then*

$$(v - \tilde{v}, u - \tilde{u}) \geq 0 \quad (\text{monotonicity}).$$

(iii) $I[u] = \min_{w \in H} I[w]$ *if and only if* $0 \in \partial I[u]$.

(iv) *For each $w \in H$ and $\lambda > 0$, the problem*

$$u + \lambda \partial I[u] \ni w$$

has a unique solution $u \in D(\partial I)$.

Assertion (iv) means that there exist $u \in D(\partial I)$ and $v \in \partial I[u]$ such that

$$u + \lambda v = w.$$

Proof. 1. Let $u \in D(\partial I)$, $v \in \partial I[u]$. Then $I[w] \geq I[u] + (v, w - u)$ for all $w \in H$. Since I is proper, there exists a point u_0 with $I[u_0] < +\infty$. Thus $I[u] \leq I[u_0] + (v, u - u_0) < \infty$ and so $u \in D(I)$. This proves (i).

2. Given $v \in \partial I[u]$, $\tilde{v} \in \partial I[\tilde{u}]$, we know

$$I[\tilde{u}] \geq I[u] + (v, \tilde{u} - u), \quad I[u] \geq I[\tilde{u}] + (\tilde{v}, u - \tilde{u}).$$

As (i) implies $I[u], I[\tilde{u}] < +\infty$, we may add the foregoing inequalities and rearrange to obtain (ii).

3. If $I[u] = \min I$, then

$$(2) \quad I[w] \geq I[u] + (0, w - u) \text{ for all } w \in H.$$

Hence $0 \in \partial I[u]$. If conversely $0 \in \partial I[u]$, then (2) holds, and so $I[u] = \min I$.

4. Given $w \in H$ and $\lambda > 0$, define

$$(3) \quad J[u] := \frac{1}{2} \|u\|^2 + \lambda I[u] - (u, w) \quad (u \in H).$$

We intend to show that J attains its minimum over H .

Let us first claim that

$$(4) \quad \begin{cases} u_k \rightharpoonup u \text{ weakly in } H \text{ implies} \\ I[u] \leq \liminf_{k \rightarrow \infty} I[u_k]. \end{cases}$$

In other words, we are asserting for a convex function I that lower semicontinuity with respect to strong convergence of sequences implies lower semicontinuity with respect to weak convergence. To see this, suppose $u_k \rightharpoonup u$ in H and

$$\liminf_{k \rightarrow \infty} I[u_k] = \lim_{j \rightarrow \infty} I[u_{kj}] = l < \infty$$

for some subsequence $\{u_{kj}\}_{j=1}^{\infty} \subset \{u_k\}_{k=1}^{\infty}$. For each $\varepsilon > 0$ the set $K_\varepsilon = \{w \in H \mid I[w] \leq l + \varepsilon\}$ is closed and convex and is thus weakly closed according to Mazur's Theorem (§D.4). Since all but finitely many of the points $\{u_{kj}\}_{j=1}^{\infty}$ lie in K_ε , u lies in K_ε , and consequently

$$I[u] \leq l + \varepsilon = \liminf_{k \rightarrow \infty} I[u_k] + \varepsilon.$$

This is true for each $\varepsilon > 0$ and thus (4) follows.

5. Next we assert that

$$(5) \quad I[u] \geq -C - C\|u\| \quad (u \in H)$$

for some constant C . To verify this claim, we suppose to the contrary that for each $k = 1, 2, \dots$ there exists a point $u_k \in H$ such that

$$(6) \quad I[u_k] \leq -k - k\|u_k\|.$$

If the sequence $\{u_k\}_{k=1}^{\infty}$ is bounded in H , there exists according to §D.4 a weakly convergent subsequence: $u_{k_j} \rightharpoonup u$. But then (4) and (6) imply the contradiction $I[u] = -\infty$. Thus we may as well assume, passing if necessary to a subsequence, that $\|u_k\| \rightarrow \infty$. Select $u_0 \in H$ so that $I[u_0] < \infty$. Set

$$z_k := \frac{u_k}{\|u_k\|} + \left(1 - \frac{1}{\|u_k\|}\right) u_0 \quad (k = 1, 2, \dots).$$

Then convexity implies

$$I[z_k] \leq \frac{1}{\|u_k\|} I[u_k] + \left(1 - \frac{1}{\|u_k\|}\right) I[u_0] \leq -k + |I[u_0]|.$$

As $\{z_k\}_{k=1}^{\infty}$ is bounded, we can extract a weakly convergent subsequence: $z_{k_j} \rightharpoonup z$ and again derive the contradiction $I[z] = -\infty$. We thereby establish the claim (5).

6. Return now to the function J defined by (3). Choose a minimizing sequence $\{u_k\}_{k=1}^{\infty} \subset H$ so that

$$J[u_k] \rightarrow \inf_{w \in H} J[w] = m.$$

Owing to (3) and (5), m is a finite number. Thus we deduce from (3), (5) that the sequence $\{u_k\}_{k=1}^\infty$ is bounded. We may then extract a weakly convergent subsequence: $u_{k_j} \rightharpoonup u$. As the mapping $u \mapsto \|u\|^2$ is weakly lower semicontinuous, J has a minimum at u . Then assertion (iii) says $0 \in \partial J[u]$. A computation verifies that $\partial J[u] = u - w + \lambda \partial I[u]$, and so

$$u + \lambda \partial I[u] \ni w.$$

7. To confirm uniqueness, suppose as well

$$\tilde{u} + \lambda \partial I[\tilde{u}] \ni w.$$

Then $u + \lambda v = w$, $\tilde{u} + \lambda \tilde{v} = w$ for $v \in \partial I[u]$, $\tilde{v} \in \partial I[\tilde{u}]$. Owing to the monotonicity assertion (ii),

$$0 \leq (u - \tilde{u}, v - \tilde{v}) = \left(u - \tilde{u}, -\frac{u}{\lambda} + \frac{\tilde{u}}{\lambda} \right) = -\frac{1}{\lambda} \|u - \tilde{u}\|^2.$$

Since $\lambda > 0$, $u = \tilde{u}$. □

We introduce next nonlinear analogues of the operators R_λ, A_λ introduced in §7.4.

DEFINITIONS. (1) For each $\lambda > 0$ define the nonlinear resolvent $J_\lambda : H \rightarrow D(\partial I)$ by setting

$$J_\lambda[w] := u,$$

where u is the unique solution of

$$u + \lambda \partial I[u] \ni w.$$

(2) For each $\lambda > 0$ define the Yosida approximation $A_\lambda : H \rightarrow H$ by

$$(7) \quad A_\lambda[w] := \frac{w - J_\lambda[w]}{\lambda} \quad (w \in H).$$

Think of A_λ as a sort of regularization or smoothing of the operator $A = \partial I$.

THEOREM 2 (Properties of J_λ, A_λ). For each $\lambda > 0$ and $w, \hat{w} \in H$, the following statements hold:

$$(i) \quad \|J_\lambda[w] - J_\lambda[\hat{w}]\| \leq \|w - \hat{w}\|,$$

$$(ii) \quad \|A_\lambda[w] - A_\lambda[\hat{w}]\| \leq \frac{2}{\lambda} \|w - \hat{w}\|,$$

$$(iii) \ 0 \leq (w - \hat{w}, A_\lambda[w] - A_\lambda[\hat{w}]),$$

$$(iv) \ A_\lambda[w] \in \partial I[J_\lambda[w]].$$

(v) *If $w \in D(\partial I)$, then*

$$\sup_{\lambda > 0} \|A_\lambda[w]\| \leq |A^0[w]|,$$

where $|A^0[w]| := \min_{z \in \partial I[w]} \|z\|$.

(vi) *For each $w \in \overline{D(\partial I)}$,*

$$\lim_{\lambda \rightarrow 0} J_\lambda[w] = w.$$

Proof. 1. Let $u = J_\lambda[w]$, $\tilde{u} = J_\lambda[\tilde{w}]$. Then $u + \lambda v = w$, $\tilde{u} + \lambda \tilde{v} = \tilde{w}$ for some $v \in \partial I[u]$, $\tilde{v} \in \partial I[\tilde{u}]$. Therefore

$$\begin{aligned} \|w - \tilde{w}\|^2 &= \|u - \tilde{u} + \lambda(v - \tilde{v})\|^2 \\ &= \|u - \tilde{u}\|^2 + 2\lambda(u - \tilde{u}, v - \tilde{v}) + \lambda^2\|v - \tilde{v}\|^2 \\ &\geq \|u - \tilde{u}\|^2, \end{aligned}$$

according to Theorem 1(ii). This proves assertion (i), and assertion (ii) follows at once from the definition (7) of the Yosida approximation A_λ .

2. We verify (iii) by using (7) to compute

$$\begin{aligned} (w - \hat{w}, A_\lambda[w] - A_\lambda[\hat{w}]) &= \frac{1}{\lambda} (\|w - \hat{w}\|^2 - (w - \hat{w}, J_\lambda[w] - J_\lambda[\hat{w}])) \\ &\geq \frac{1}{\lambda} (\|w - \hat{w}\|^2 - \|w - \hat{w}\| \|J_\lambda[w] - J_\lambda[\hat{w}]\|) \geq 0, \end{aligned}$$

according to (i).

3. To prove (iv), note that $u = J_\lambda[w]$ if and only if $u + \lambda v = w$ for some $v \in \partial I[u] = \partial I[J_\lambda[w]]$. But

$$v = \frac{w - u}{\lambda} = \frac{w - J_\lambda[w]}{\lambda} = A_\lambda[w].$$

4. Assume next $w \in D(\partial I)$, $z \in \partial I[w]$. Let $u = J_\lambda[w]$, so that $u + \lambda v = w$, where $v \in \partial I[u]$. By monotonicity

$$0 \leq (w - u, z - v) = \left(w - J_\lambda[w], z - \frac{w - J_\lambda[w]}{\lambda} \right) = (\lambda A_\lambda[w], z - A_\lambda[w]).$$

Consequently

$$\lambda \|A_\lambda[w]\|^2 \leq (\lambda A_\lambda[w], z) \leq \lambda \|A_\lambda[w]\| \|z\|,$$

and so

$$\|A_\lambda[w]\| \leq \|z\|.$$

This estimate is valid for all $\lambda > 0$, $z \in \partial I[w]$. Assertion (v) follows.

5. If $w \in D(\partial I)$, then

$$\|J_\lambda[w] - w\| = \lambda \|A_\lambda[w]\| \leq \lambda |A^0[w]|,$$

and hence $J_\lambda[w] \rightarrow w$ as $\lambda \rightarrow 0$. Now let $w \in \overline{D(\partial I)} - D(\partial I)$. There exists for each $\varepsilon \rightarrow 0$ a point $\hat{w} \in D(\partial I)$ with $\|w - \hat{w}\| \leq \varepsilon$. Then

$$\begin{aligned} \|J_\lambda[w] - w\| &\leq \|J_\lambda[w] - J_\lambda[\hat{w}]\| + \|J_\lambda[\hat{w}] - \hat{w}\| + \|w - \hat{w}\| \\ &\leq 2\|w - \hat{w}\| + \|J_\lambda[\hat{w}] - \hat{w}\| \\ &\leq 2\varepsilon + \|J_\lambda[\hat{w}] - \hat{w}\|. \end{aligned}$$

Since $\hat{w} \in D(\partial I)$, $J_\lambda[\hat{w}] \rightarrow \hat{w}$ as $\lambda \rightarrow 0$. Thus

$$\limsup_{\lambda \rightarrow 0} \|J_\lambda[w] - w\| \leq 2\varepsilon$$

for each $\varepsilon > 0$. □

9.6.2. Subdifferentials and nonlinear semigroups.

As above, let H be a real Hilbert space, and take $I : H \rightarrow (-\infty, +\infty]$ to be convex, proper, lower semicontinuous. Let us for simplicity assume as well

$$(8) \quad \partial I \text{ is densely defined, that is, } \overline{D(\partial I)} = H.$$

By analogy with the theory of linear semigroups set forth in §7.4, we propose now to study the differential equation

$$(9) \quad \begin{cases} \mathbf{u}'(t) + A[\mathbf{u}(t)] \ni 0 & (t \geq 0) \\ \mathbf{u}(0) = u, \end{cases}$$

where $u \in H$ is given and $A = \partial I$ is a nonlinear, discontinuous operator, which is perhaps multivalued. Assuming for the moment (9) has a unique solution for each initial point u , we write

$$(10) \quad \mathbf{u}(t) = S(t)u \quad (t > 0)$$

and regard $S(t)$ so defined as a mapping from H into H for each time $t \geq 0$. We will employ the notation (10) to emphasize similarities with linear semigroup theory, previously introduced in §7.4. But carefully note here and afterwards that the mapping $u \mapsto S(t)u$ is in general nonlinear.

As in §7.4 it is reasonable to expect further that

$$(11) \quad S(0)u = u \quad (u \in H),$$

$$(12) \quad S(t+s)u = S(t)S(s)u \quad (t, s \geq 0, u \in H),$$

and for each $u \in H$

$$(13) \quad \text{the mapping } t \mapsto S(t)u \text{ is continuous from } [0, \infty) \text{ into } H.$$

DEFINITIONS. (i) A family $\{S(t)\}_{t \geq 0}$ of nonlinear operators mapping H into H is called a nonlinear semigroup if conditions (11)–(13) are satisfied.

(ii) We say $\{S(t)\}_{t \geq 0}$ is a contraction semigroup if in addition

$$(14) \quad \|S(t)u - S(t)\hat{u}\| \leq \|u - \hat{u}\| \quad (t \geq 0, u, \hat{u} \in H).$$

Our intention is to show that the operator $A = \partial I$ generates a nonlinear semigroup of contractions on H . In particular we will prove that the ODE

$$(15) \quad \begin{cases} \mathbf{u}'(t) \in -\partial I[\mathbf{u}(t)] & (t \geq 0) \\ \mathbf{u}(0) = u, \end{cases}$$

for a given initial point $u \in H$, is well-posed. This is a kind of infinite-dimensional “gradient flow” governed by ∂I . Later in §9.6.3 we will see that certain quasilinear parabolic PDE can be cast into the abstract form (15).

THEOREM 3 (Solution of gradient flow). *For each $u \in D(\partial I)$ there exists a unique function*

$$(16) \quad \mathbf{u} \in C([0, \infty); H), \text{ with } \mathbf{u}' \in L^\infty(0, \infty; H),$$

such that

$$(i) \quad \mathbf{u}(0) = u,$$

$$(ii) \quad \mathbf{u}(t) \in D(\partial I) \quad \text{for each } t > 0,$$

and

$$(iii) \quad \mathbf{u}'(t) \in -\partial I[\mathbf{u}(t)] \quad \text{for a.e. } t \geq 0.$$

Proof. 1. We first build approximate solutions by solving for each $\lambda > 0$ the ODE

$$(17) \quad \begin{cases} \mathbf{u}'_\lambda(t) + A_\lambda[\mathbf{u}_\lambda(t)] = 0 & (t \geq 0) \\ \mathbf{u}_\lambda(0) = u. \end{cases}$$

According to Theorem 2(ii) the Yosida approximation $A_\lambda : H \rightarrow H$ is an everywhere defined, Lipschitz continuous mapping, and thus (17) has a unique solution $\mathbf{u}_\lambda \in C^1([0, \infty); H)$.

Our plan is to show that as $\lambda \rightarrow 0^+$, the functions \mathbf{u}_λ converge to a solution of (15). This is subtle, however, as the operator $A = \partial I$ is in general nonlinear, multivalued, and not everywhere defined.

2. First, let us take another point $v \in H$ and consider as well the ODE

$$(18) \quad \begin{cases} \mathbf{v}'_\lambda(t) + A_\lambda[\mathbf{v}_\lambda(t)] = 0 & (t \geq 0) \\ \mathbf{v}_\lambda(0) = v. \end{cases}$$

Then

$$\begin{aligned} \frac{1}{2} \frac{d}{dt} \|\mathbf{u}_\lambda - \mathbf{v}_\lambda\|^2 &= (\mathbf{u}'_\lambda - \mathbf{v}'_\lambda, \mathbf{u}_\lambda - \mathbf{v}_\lambda) \\ &= (-A_\lambda[\mathbf{u}_\lambda] + A_\lambda[\mathbf{v}_\lambda], \mathbf{u}_\lambda - \mathbf{v}_\lambda) \leq 0, \end{aligned}$$

owing to Theorem 2(iii). Hence

$$(19) \quad \|\mathbf{u}_\lambda(t) - \mathbf{v}_\lambda(t)\| \leq \|u - v\| \quad (t \geq 0).$$

In particular, if $h > 0$ and $v = \mathbf{u}_\lambda(h)$, then by uniqueness $\mathbf{v}_\lambda(t) = \mathbf{u}_\lambda(t+h)$. Consequently (19) implies

$$\|\mathbf{u}_\lambda(t+h) - \mathbf{u}_\lambda(t)\| \leq \|\mathbf{u}(h) - u\|.$$

Divide by h and send $h \rightarrow 0$:

$$(20) \quad \|\mathbf{u}'_\lambda(t)\| \leq \|\mathbf{u}'_\lambda(0)\| = \|A_\lambda[u]\| \leq |A^0[u]|,$$

the last inequality resulting from Theorem 2(v).

3. We next take $\lambda, \mu > 0$ and compute

$$(21) \quad \begin{aligned} \frac{1}{2} \frac{d}{dt} \|\mathbf{u}_\lambda - \mathbf{u}_\mu\|^2 &= (\mathbf{u}'_\lambda - \mathbf{u}'_\mu, \mathbf{u}_\lambda - \mathbf{u}_\mu) \\ &= (-A_\lambda[\mathbf{u}_\lambda] + A_\mu[\mathbf{u}_\mu], \mathbf{u}_\lambda - \mathbf{u}_\mu). \end{aligned}$$

Now

$$\begin{aligned} \mathbf{u}_\lambda - \mathbf{u}_\mu &= (\mathbf{u}_\lambda - J_\lambda[\mathbf{u}_\lambda]) + (J_\lambda[\mathbf{u}_\lambda] - J_\mu[\mathbf{u}_\mu]) + (J_\mu[\mathbf{u}_\mu] - \mathbf{u}_\mu) \\ &= \lambda A_\lambda[\mathbf{u}_\lambda] + J_\lambda[\mathbf{u}_\lambda] - J_\mu[\mathbf{u}_\mu] - \mu A_\mu[\mathbf{u}_\mu]. \end{aligned}$$

Consequently

$$(22) \quad (A_\lambda[\mathbf{u}_\lambda] - A_\mu[\mathbf{u}_\mu], \mathbf{u}_\lambda - \mathbf{u}_\mu) = (A_\lambda[\mathbf{u}_\lambda] - A_\mu[\mathbf{u}_\mu], J_\lambda[\mathbf{u}_\lambda] - J_\mu[\mathbf{u}_\mu]) \\ + (A_\lambda[\mathbf{u}_\lambda] - A_\mu[\mathbf{u}_\mu], \lambda A_\lambda[\mathbf{u}_\lambda] - \mu A_\mu[\mathbf{u}_\mu]).$$

Since $A_\lambda[\mathbf{u}_\lambda] \in \partial I[J_\lambda[\mathbf{u}_\lambda]]$ and $A_\mu[\mathbf{u}_\mu] \in \partial I[J_\mu[\mathbf{u}_\mu]]$, the monotonicity property implies that the first term on the right-hand side of (22) is nonnegative. Thus

$$(A_\lambda[\mathbf{u}_\lambda] - A_\mu[\mathbf{u}_\mu], \mathbf{u}_\lambda - \mathbf{u}_\mu) \geq \lambda \|A_\lambda[\mathbf{u}_\lambda]\|^2 + \mu \|A_\mu[\mathbf{u}_\mu]\|^2 \\ - (\lambda + \mu) \|A_\lambda[\mathbf{u}_\lambda]\| \|A_\mu[\mathbf{u}_\mu]\|.$$

Since

$$(\lambda + \mu) \|A_\lambda[\mathbf{u}_\lambda]\| \|A_\mu[\mathbf{u}_\mu]\| \leq \lambda (\|A_\lambda[\mathbf{u}_\lambda]\|^2 + \frac{1}{4} \|A_\mu[\mathbf{u}_\mu]\|^2) \\ + \mu (\|A_\mu[\mathbf{u}_\mu]\|^2 + \frac{1}{4} \|A_\lambda[\mathbf{u}_\lambda]\|^2),$$

we deduce

$$(A_\lambda[\mathbf{u}_\lambda] - A_\mu[\mathbf{u}_\mu], \mathbf{u}_\lambda - \mathbf{u}_\mu) \geq -\frac{\lambda}{4} \|A_\mu[\mathbf{u}_\mu]\|^2 - \frac{\mu}{4} \|A_\lambda[\mathbf{u}_\lambda]\|^2.$$

But $\|A_\lambda[\mathbf{u}_\lambda]\| = \|\mathbf{u}'_\lambda\| \leq |A^0[u]|$ according to (20), whence

$$(A_\lambda[\mathbf{u}_\lambda] - A_\mu[\mathbf{u}_\mu], \mathbf{u}_\lambda - \mathbf{u}_\mu) \geq -\frac{\lambda + \mu}{4} |A^0[u]|^2.$$

Recalling (21), (22), we obtain the inequality

$$\frac{d}{dt} \|\mathbf{u}_\lambda - \mathbf{u}_\mu\|^2 \leq \frac{(\lambda + \mu)}{2} |A^0[u]|^2 \quad (t \geq 0);$$

and hence

$$(23) \quad \|\mathbf{u}_\lambda(t) - \mathbf{u}_\mu(t)\|^2 \leq \frac{(\lambda + \mu)}{2} t |A^0[u]|^2 \quad (t \geq 0).$$

In view of estimate (23) there exists a function $\mathbf{u} \in C([0, \infty); H)$ such that

$$\mathbf{u}_\lambda \rightarrow \mathbf{u} \quad \text{uniformly in } C([0, T], H)$$

as $\lambda \rightarrow 0$, for each time $T > 0$. Furthermore estimate (20) implies

$$(24) \quad \mathbf{u}'_\lambda \rightharpoonup \mathbf{u}' \quad \text{weakly in } L^2(0, T; H)$$

for each $T > 0$ and

$$(25) \quad \|\mathbf{u}'(t)\| \leq |A^0[u]| \quad \text{for a.e. } t.$$

4. We must show $\mathbf{u}(t) \in D(\partial I)$ for each $t \geq 0$ and

$$\mathbf{u}'(t) + \partial I[\mathbf{u}(t)] \ni 0 \quad \text{for a.e. } t \geq 0.$$

Now

$$\|J_\lambda[\mathbf{u}_\lambda](t) - \mathbf{u}_\lambda(t)\| = \lambda \|A_\lambda[\mathbf{u}_\lambda](t)\| = \lambda \|\mathbf{u}'_\lambda(t)\| \leq \lambda |A^0[u]|$$

by (20). Hence

$$(26) \quad J_\lambda[\mathbf{u}_\lambda] \rightarrow \mathbf{u} \quad \text{uniformly in } C([0, T]; H)$$

for each $T > 0$.

For each time $t \geq 0$,

$$-\mathbf{u}'_\lambda(t) = A_\lambda[\mathbf{u}_\lambda(t)] \in \partial I[J_\lambda[\mathbf{u}_\lambda(t)]].$$

Thus given $w \in H$, we have

$$I[w] \geq I[J_\lambda[\mathbf{u}_\lambda(t)]] - (\mathbf{u}'_\lambda(t), w - J_\lambda[\mathbf{u}_\lambda(t)]).$$

Consequently if $0 \leq s \leq t$,

$$(t - s)I[w] \geq \int_s^t I[J_\lambda[\mathbf{u}_\lambda(r)]] dr - \int_s^t (\mathbf{u}'_\lambda(r), w - J_\lambda[\mathbf{u}_\lambda(r)]) dr.$$

In view of (26), the lower semicontinuity of I , and Fatou's Lemma (§E.3), we conclude upon sending $\lambda \rightarrow 0$ that

$$(t - s)I[w] \geq \int_s^t I[\mathbf{u}(r)] dr - \int_s^t (\mathbf{u}'(r), w - \mathbf{u}(r)) dr$$

for each $0 \leq s \leq t$. Therefore

$$I[w] \geq I[\mathbf{u}(t)] + (-\mathbf{u}'(t), w - \mathbf{u}(t))$$

if t is a Lebesgue point of \mathbf{u}' , $I[\mathbf{u}]$. Hence for a.e. $t \geq 0$,

$$I[w] \geq I[\mathbf{u}(t)] + (-\mathbf{u}'(t), w - \mathbf{u}(t))$$

for all $w \in H$. Thus $\mathbf{u}(t) \in D(\partial I)$, with

$$-\mathbf{u}'(t) \in \partial I[\mathbf{u}(t)]$$

for a.e. $t \geq 0$.

5. Finally we prove $\mathbf{u}(t) \in D(\partial I)$ for each $t \geq 0$. To see this, fix $t \geq 0$ and choose $t_k \rightarrow t$ such that $\mathbf{u}(t_k) \in D(\partial I)$, $-\mathbf{u}'(t_k) \in \partial I[\mathbf{u}(t_k)]$. In view of (25) we may assume, upon passing if necessary to a subsequence, that

$$\mathbf{u}'(t_k) \rightharpoonup \mathbf{v} \quad \text{weakly in } H.$$

Fix $w \in H$. Then

$$I[w] \geq I[\mathbf{u}(t_k)] + (-\mathbf{u}'(t_k), w - \mathbf{u}(t_k)).$$

Let $t_k \rightarrow t$ and recall that $\mathbf{u} \in C([0, \infty]; H)$ and I is lower semicontinuous. We obtain the inequality

$$I[w] \geq I[\mathbf{u}(t)] + (-\mathbf{v}, w - \mathbf{u}(t)).$$

Hence $\mathbf{u}(t) \in D(\partial I)$ and $-\mathbf{v} \in \partial I[\mathbf{u}(t)]$.

6. We have shown \mathbf{u} satisfies assertions (i)–(iii). To prove uniqueness, assume $\tilde{\mathbf{u}}$ is another solution and compute

$$\frac{1}{2} \frac{d}{dt} \|\mathbf{u} - \tilde{\mathbf{u}}\|^2 = (\mathbf{u}' - \tilde{\mathbf{u}}', \mathbf{u} - \tilde{\mathbf{u}}) \leq 0 \quad \text{for a.e. } t \geq 0,$$

since $-\mathbf{u}' \in \partial I[\mathbf{u}]$, $-\tilde{\mathbf{u}}' \in \partial I[\tilde{\mathbf{u}}]$. □

Remarks. (i) The operator $A = \partial I$ in fact generates a nonlinear contraction semigroup on all of H . If $u, v \in D(\partial I)$, we write as above

$$\lim_{\lambda \rightarrow 0} \mathbf{u}_\lambda(t) = \mathbf{u}(t) = S(t)u$$

and

$$\lim_{\lambda \rightarrow 0} \mathbf{v}_\lambda(t) = \mathbf{v}(t) = S(t)v.$$

Owing to (19), we see

$$\|S(t)u - S(t)v\| \leq \|u - v\| \quad (t \geq 0)$$

if $u, v \in D(\partial I)$. Using this inequality, we uniquely extend the semigroup of nonlinear operators $\{S(t)\}_{t \geq 0}$ to $H = \overline{D(\partial I)}$.

(ii) We have assumed that $D(\partial I)$ is dense in H solely to streamline the exposition. In general $\{S(t)\}_{t \geq 0}$ is a semigroup of contractions on $\overline{D(\partial I)} \subseteq H$.

9.6.3. Applications.

We now illustrate how some of the abstract theory set forth in §9.6.1 and §9.6.2 applies to certain nonlinear parabolic partial differential equations.

Let us hereafter suppose U is a bounded, open subset of \mathbb{R}^n , with smooth boundary ∂U . We choose $H = L^2(U)$, and set

$$(27) \quad I[u] := \begin{cases} \int_U L(Du) \, dx & \text{if } u \in H_0^1(U) \\ +\infty & \text{otherwise,} \end{cases}$$

where $L : \mathbb{R}^n \rightarrow \mathbb{R}$ is smooth, convex and satisfies

$$(28) \quad |D^2L(p)| \leq C \quad (p \in \mathbb{R}^n),$$

$$(29) \quad \sum_{i,j=1}^n L_{p_i p_j}(p) \xi_i \xi_j \geq \theta |\xi|^2 \quad (p, \xi \in \mathbb{R}^n)$$

for constants $C, \theta > 0$.

THEOREM 4 (Characterization of ∂I).

- (i) $I : H \rightarrow (-\infty, +\infty]$ is convex, proper and lower semicontinuous.
- (ii) $D(\partial I) = H^2(U) \cap H_0^1(U)$.
- (iii) If $u \in D(\partial I)$, then ∂I is single-valued and

$$\partial I[u] = - \sum_{i=1}^n (L_{p_i}(Du))_{x_i} \quad a.e.$$

Proof. 1. I is clearly proper and convex. Furthermore, since I is weakly sequentially lower semicontinuous (cf. Theorem 1 in §8.2.2), I is lower semicontinuous.

2. Define the nonlinear operator A by setting

$$\begin{cases} D(A) := H^2(U) \cap H_0^1(U), \\ A[u] := - \sum_{i=1}^n (L_{p_i}(Du))_{x_i} \quad (u \in D(A)). \end{cases}$$

We must prove $A = \partial I$.

First let $u \in D(A)$, $v = A[u]$, $w \in L^2(U)$. If $w \notin H_0^1(U)$, then $I[w] = +\infty$ and so clearly

$$(30) \quad I[w] \geq I[u] + (v, w - u).$$

Assume next $w \in H_0^1(U)$. Thus

$$\begin{aligned} (v, w - u) &= - \int_U \sum_{i=1}^n (L_{p_i}(Du))_{x_i} (w - u) dx \\ &= \int_U \sum_{i=1}^n L_{p_i}(Du) \cdot (Dw - Du) dx. \end{aligned}$$

Since L is convex,

$$L(Dw) \geq L(Du) + D_p L(Du) \cdot (Dw - Du) \quad \text{a.e. in } U.$$

Integrating over U gives (30).

3. We have thus far shown that $A \subseteq \partial I$; that is, $D(A) \subseteq D(\partial I)$ and $Au \in \partial I[u]$ for $u \in D(A)$. To conclude, we must prove that $A \supseteq \partial I$.

Select any function $f \in L^2(U)$. If we minimize the functional

$$J[w] := \int_U L(Dw) + \frac{w^2}{2} - fw dx$$

over the admissible class $\mathcal{A} = H_0^1(U)$, we will find $u \in H_0^1(U)$, which is a weak solution of

$$(31) \quad u - \sum_{i=1}^n (L_{p_i}(Du))_{x_i} = f \quad \text{in } U.$$

According to calculations similar to those for the proof of Theorem 1(ii) in §8.3, we see that in fact $u \in H^2(U)$, with the estimate

$$(32) \quad \|u\|_{H^2(U)} \leq C \|f\|_{L^2(U)}.$$

Thus $u \in D(A)$, and $u + A[u] = f$. Consequently the range of $I + A$ is all of H . But this implies $A = \partial I$. For if $v \in D(\partial I)$, $w \in \partial I[v]$, then there exists $u \in D(A)$ such that $u + A[u] = v + w$. Since $A[u] \in \partial I[u]$, $w \in \partial I[v]$, the uniqueness assertion of Theorem 1(iv) implies $u = v$, $w = A[u]$. Thus $A = \partial I$. \square

We can now look at the initial/boundary-value problem:

$$(33) \quad \begin{cases} u_t - \sum_{i=1}^n (L_{p_i}(Du))_{x_i} = 0 & \text{in } U \times (0, \infty) \\ u = 0 & \text{on } \partial U \times (0, \infty) \\ u = g & \text{on } U \times \{t = 0\}, \end{cases}$$

where $g \in L^2(U)$. In accordance with Theorem 4, we can recast this problem into the abstract form

$$(34) \quad \begin{cases} \mathbf{u}'(t) = -\partial I[\mathbf{u}(t)] & (t \geq 0) \\ \mathbf{u}(0) = g. \end{cases}$$

We apply Theorem 3. If $g \in H^2(U) \cap H_0^1(U)$, there exists a unique function

$$\mathbf{u} \in C([0, \infty); L^2(U)), \text{ with } \mathbf{u}' \in L^\infty((0, \infty); L^2(U)),$$

that is, a weak solution of (33). In view of the estimate

$$\|\mathbf{u}(t)\|_{H^2(U)} \leq C\|\mathbf{u}'(t)\|_{L^2(U)},$$

we see $\mathbf{u} \in L^\infty((0, \infty), H^2(U) \cap H_0^1(U))$ as well.

9.7. PROBLEMS

In these problems U always denotes a bounded, open subset of \mathbb{R}^n , with smooth boundary.

1. Assume the vector field \mathbf{v} is smooth. Give another proof of the lemma in §9.1 for this case by solving the ODE

$$\begin{cases} \dot{\mathbf{x}}(t) = -\mathbf{v}(\mathbf{x}(t)) & (t \geq 0) \\ \mathbf{x}(0) = y. \end{cases}$$

Let us write the solution as $\mathbf{x}(t, y)$ to display the dependence on the initial point y . For each fixed time $t > 0$, the map $y \mapsto \mathbf{x}(t, y)$ is continuous and so has a fixed point. Conclude that \mathbf{v} has a zero in the closed ball $B(0, r)$.

2. Assume $a : \mathbb{R} \rightarrow \mathbb{R}$ is continuous and $a(f_n) \rightharpoonup a(f)$ weakly in $L^2(0, 1)$ whenever $f_n \rightharpoonup f$ weakly in $L^2(0, 1)$. Show a is an affine function; that is, a has the form

$$a(z) = \alpha z + \beta \quad (z \in \mathbb{R})$$

for constants α, β .

3. (Penalty method) Let $\epsilon > 0$. Define

$$\beta_\epsilon(z) := \begin{cases} 0 & \text{if } z \geq 0 \\ \frac{z}{\epsilon} & \text{if } z \leq 0, \end{cases}$$

and suppose $u_\epsilon \in H_0^1(U)$ is the weak solution of

$$(*) \quad \begin{cases} -\Delta u_\epsilon + \beta_\epsilon(u_\epsilon) = f & \text{in } U \\ u_\epsilon = 0 & \text{on } \partial U, \end{cases}$$

where $f \in L^2(U)$. Prove that as $\epsilon \rightarrow 0$, $u^\epsilon \rightharpoonup u$ weakly in $H_0^1(U)$, u being the unique nonnegative solution of the variational inequality

$$\int_U Du \cdot D(w - u) dx \geq \int_U f(w - u) dx$$

for all $w \in H_0^1(U)$ with $w \geq 0$ a.e.

Approximating the variational inequality by (*) is the *penalty method*.

4. (Solutions periodic in time) Assume

$$\begin{cases} u_t - \Delta u = f & \text{in } U \times (0, \infty) \\ u = 0 & \text{on } \partial U \times (0, \infty) \\ u = g & \text{on } U \times \{t = 0\}, \end{cases}$$

where $g \in L^2(U)$, $f \in L^\infty(U_T)$ for each $T > 0$. Suppose $\tau > 0$ and f is τ -periodic in t ; that is, $f(x, t) = f(x, t + \tau)$ ($x \in U, t \geq 0$). Prove there exists a unique function $g \in L^2(U)$ for which the corresponding solution u is τ -periodic.

5. Consider the nonlinear boundary-value problem

$$\begin{cases} -\Delta u + b(Du) = f & \text{in } U \\ u = 0 & \text{on } \partial U. \end{cases}$$

Use Banach's Fixed Point Theorem to show there exists a unique weak solution $u \in H^2(U) \cap H_0^1(U)$ provided $b : \mathbb{R}^n \rightarrow \mathbb{R}$ is Lipschitz continuous, with $\text{Lip}(b)$ small enough.

6. Assume $f : \mathbb{R} \rightarrow \mathbb{R}$ is Lipschitz continuous, bounded, with $f(0) = 0$ and $f'(0) > \lambda_1$, λ_1 denoting the principal eigenvalue for $-\Delta$ on $H_0^1(U)$. Use the method of sub- and supersolutions to show there exists a weak solution u of

$$\begin{cases} -\Delta u = f(u) & \text{in } U \\ u = 0 & \text{on } \partial U \\ u > 0 & \text{in } U. \end{cases}$$

7. Assume that \underline{u}, \bar{u} are smooth sub- and supersolutions of the boundary-value problem (1) in §9.3. Use the maximum principle to verify directly

$$\underline{u} = u_0 \leq u_1 \leq \cdots \leq u_k \leq \cdots \leq \bar{u},$$

where the $\{u_k\}_{k=0}^\infty$ are defined as in §9.3.

8. (Noncompact families of solutions)

(a) Assume $n \geq 3$. Find a constant c such that

$$u(x) := (1 + |x|^2)^{\frac{2-n}{2}}$$

solves *Yamabe's equation*

$$-\Delta u = c u^{\frac{n+2}{n-2}} \quad \text{in } \mathbb{R}^n.$$

Note the appearance of the critical exponent $\frac{n+2}{n-2}$. (Compare with Problem 6 in Chapter 4.)

- (b) Check that for each $\lambda > 0$,

$$u_\lambda(x) := \left(\frac{\lambda}{\lambda^2 + |x|^2} \right)^{\frac{n-2}{2}}$$

is also a solution.

- (c) Show that

$$\|u_\lambda\|_{L^{2^*}(\mathbb{R}^n)} = \|u\|_{L^{2^*}(\mathbb{R}^n)}, \quad \|Du_\lambda\|_{L^2(\mathbb{R}^n)} = \|Du\|_{L^2(\mathbb{R}^n)}$$

for each λ and thus that $\{u_\lambda\}_{\lambda>0}$ is not precompact in $L^{2^*}(\mathbb{R}^n)$.

9. Let u solve the semilinear heat equation

$$u_t - \Delta u = f(u) \quad \text{in } \mathbb{R}^n \times (0, \infty).$$

Assume that u and its derivatives go to zero rapidly as $|x| \rightarrow \infty$.

- (a) Show that

$$\frac{d}{dt} \int_{\mathbb{R}^n} \frac{1}{2} |Du|^2 - F(u) \, dx = - \int_{\mathbb{R}^n} u_t^2 \, dx,$$

where $F' = f$, $F(0) = 0$.

- (b) Now show that

$$\begin{aligned} \frac{d}{dt} \int_{\mathbb{R}^n} |x|^2 \left(\frac{1}{2} |Du|^2 - F(u) \right) dx \\ = - \int_{\mathbb{R}^n} u_t^2 |x|^2 - 2nF(u) + (n-2)|Du|^2 \, dx. \end{aligned}$$

This is a parabolic analogue of the Derrick–Pohozaev identity.

(Bauman, Chen, Phillips, Euro. J. Applied Math. 6 (1995), 115–126)

10. Let $K \subset \mathbb{R}^n$ be a closed, convex, nonempty set. Define

$$I[x] := \begin{cases} 0 & \text{if } x \in K \\ \infty & \text{if } x \notin K. \end{cases}$$

Explicitly determine $A = \partial I$, $J_\lambda = (I + \lambda A)^{-1}$, $A_\lambda = \frac{I - J_\lambda}{\lambda}$ ($\lambda > 0$) in terms of the geometry of K .

11. Give a simple example showing that the flow

$$(*) \quad \mathbf{u}' \in -\partial I[\mathbf{u}] \quad (t > 0)$$

may be irreversible. (That is, find a Hilbert space H and a convex, proper, lower semicontinuous function $I : H \rightarrow (-\infty, +\infty]$ such that the semigroup solution of $(*)$ satisfies

$$S(t)u = S(t)\hat{u}$$

for some $t > 0$ and $u \neq \hat{u}$.)

12. Let $u = u(x, t)$ denote the height at $x \in \mathbb{R}^2$ of a sandpile that grows as sand is added at rate $f = f(x, t) \geq 0$. We assume the stability condition

$$|Du| \leq 1,$$

meaning that nowhere can the sandpile have slope greater than 1. As usual, $D_x u = Du$. We propose the dynamics

$$u_t - \operatorname{div}(aDu) = f \quad \text{in } \mathbb{R}^2 \times (0, \infty),$$

where $a = a(x, t) \geq 0$ describes the flow rate of sand rolling downhill, that is, in the direction $-Du$. Suppose lastly that

$$\operatorname{spt} a \subseteq \{|Du| = 1\},$$

so that the sand flows downhill only if the slope is one.

Show that the foregoing implies

$$f - u_t \in \partial I[u],$$

for the convex function

$$I[u] := \begin{cases} 0 & \text{if } u \in L^2(\mathbb{R}^2), |Du| \leq 1 \text{ a.e.} \\ \infty & \text{otherwise.} \end{cases}$$

13. Assume u is a smooth solution of the gradient flow system (33) in §9.6.3, where L satisfies the uniform convexity condition (29). Show there exist constants $C, \gamma > 0$ such that

$$\int_U u_t^2(x, t) dx \leq Ce^{-\gamma t} \quad (t > 0).$$

14. Let u be a smooth solution of the nonlinear heat equation

$$u_t - \Delta\phi(u) = 0 \quad \text{in } \mathbb{R}^n \times (0, \infty),$$

where $\phi' > 0$. Assuming $|Du| > 0$, derive the formula

$$\frac{d}{dt} \int_{\mathbb{R}^n} |Du| dx = - \int_{\mathbb{R}^n} \frac{|D^2v|^2}{|Dv|} - \frac{|D^2vDv|^2}{|Dv|^3} dx \leq 0,$$

for $v := \phi(u)$.

9.8. REFERENCES

- Section 9.1 See J.-L. Lions [**L2**] and Zeidler [**Zd**, Vol. 2]. The booklet [**E**] is a survey of methods for confronting weak convergence problems for nonlinear PDE.
- Section 9.2 Zeidler [**Zd**, Vol. 1] has more on fixed point techniques. Consult also Gilbarg–Trudinger [**G-T**, Chapter 11]. S. Armstrong helped me with Schaefer’s Theorem.
- Section 9.3 See Smoller [**S**, Chapter 10].
- Section 9.4 Section 9.4.1 is based upon Payne [**Pa**, §9]. Theorem 2 in §9.4.1 is due to H. Fujita (J. Fac. Sci. U. Tokyo Sect. I 13 (1966), 109–124).
- Section 9.5 Section 9.5.2 is due to Gidas, Ni, and Nirenberg (Comm. Math. Phys. 68 (1979), 209–243).
- Section 9.6 This material is from Brezis [**Br2**], which contains much more on nonlinear semigroups on Hilbert spaces. See Barbu [**Ba**] or Zeidler [**Zd**, Vol. 2] for nonlinear semigroup theory in Banach spaces.

This page is intentionally left blank.

HAMILTON–JACOBI EQUATIONS

- 10.1 Introduction, viscosity solutions
- 10.2 Uniqueness
- 10.3 Control theory, dynamic programming
- 10.4 Problems
- 10.5 References

10.1. INTRODUCTION, VISCOSITY SOLUTIONS

This chapter investigates the existence, uniqueness and other properties of appropriately defined weak solutions of the initial-value problem for the *Hamilton–Jacobi equation*:

$$(1) \quad \begin{cases} u_t + H(Du, x) = 0 & \text{in } \mathbb{R}^n \times (0, \infty) \\ u = g & \text{on } \mathbb{R}^n \times \{t = 0\}. \end{cases}$$

Here the Hamiltonian $H : \mathbb{R}^n \times \mathbb{R}^n \rightarrow \mathbb{R}$ is given, as is the initial function $g : \mathbb{R}^n \rightarrow \mathbb{R}$. The unknown is $u : \mathbb{R}^n \times [0, \infty) \rightarrow \mathbb{R}$, $u = u(x, t)$, and $Du = D_x u = (u_{x_1}, \dots, u_{x_n})$. We will write $H = H(p, x)$, so that “ p ” is the name of the variable for which we substitute the gradient Du in the PDE.

We recall from our study of characteristics in §3.2 that in general there can be no smooth solution of (1) lasting for all times $t \geq 0$. We recall further that if H depends only on p and is convex, then the Hopf–Lax formula (expression (21) in §3.3.2) provides us with a type of generalized solution.

In this chapter we consider the general case that H depends also on x and, more importantly, is no longer necessarily convex in the variable p . We

will discover in these new circumstances a different way to define a weak solution of (1).

An approximation. Our approach is to consider first this problem:

$$(2) \quad \begin{cases} u_t^\epsilon + H(Du^\epsilon, x) - \epsilon \Delta u^\epsilon = 0 & \text{in } \mathbb{R}^n \times (0, \infty) \\ u^\epsilon = g & \text{on } \mathbb{R}^n \times \{t = 0\}, \end{cases}$$

for $\epsilon > 0$. The idea is that whereas (1) involves a fully nonlinear first-order PDE, (2) is an initial-value problem for a quasilinear parabolic PDE, which turns out to have a smooth solution. The term $\epsilon \Delta$ in (2) in effect regularizes the Hamilton–Jacobi equation. Then of course we hope that as $\epsilon \rightarrow 0$ the solutions u^ϵ of (2) will converge to some sort of weak solution of (1). This technique is the method of *vanishing viscosity*.

However, as $\epsilon \rightarrow 0$ we can expect to lose control over the various estimates of the function u^ϵ and its derivatives: these estimates depend strongly on the regularizing effect of $\epsilon \Delta$ and blow up as $\epsilon \rightarrow 0$. However, it turns out that we can often in practice at least be sure that the family $\{u^\epsilon\}_{\epsilon > 0}$ is bounded and equicontinuous on compact subsets of $\mathbb{R}^n \times [0, \infty)$. Consequently the Arzela–Ascoli compactness criterion, §C.7, ensures that

$$(3) \quad u^{\epsilon_j} \rightarrow u \quad \text{locally uniformly in } \mathbb{R}^n \times [0, \infty),$$

for some subsequence $\{u^{\epsilon_j}\}_{j=1}^\infty$ and some limit function

$$(4) \quad u \in C(\mathbb{R}^n \times [0, \infty)).$$

Now we can surely expect that u is some kind of solution of our initial-value problem (1), but as we only know that u is continuous and have absolutely no information as to whether Du and u_t exist in any sense, such an interpretation is difficult.

Similar problems have arisen before in Chapters 8 and 9, where we had to deal with the weak convergence of various would-be approximate solutions to other nonlinear partial differential equations. Remember in particular that in §9.1 we solved a divergence structure quasilinear elliptic PDE by passing to limits using the method of Browder and Minty. Roughly speaking, we there integrated by parts to throw “hard-to-control” derivatives onto a fixed test function and only then tried to go to limits to discover a solution. We will for the Hamilton–Jacobi equation (1) attempt something similar. We will fix a smooth test function v and will pass from (2) to (1) as $\epsilon \rightarrow 0$ by first “putting the derivatives onto v ”.

But since (1) is fully nonlinear and in particular is not of divergence structure, we cannot just integrate by parts to switch to differentiations

on v , as we did in §9.1. Instead we will exploit the maximum principle to accomplish this transition, at least at certain points.

We will call the solution we build a *viscosity solution*, in honor of the vanishing viscosity technique. Our main goal will then be to discover an intrinsic characterization of such generalized solutions of (1).

10.1.1. Definitions.

Motivation for definition of viscosity solution. We henceforth assume that H, g are continuous and will as necessary later add further hypotheses.

The technique alluded to above works as follows. Fix any smooth test function $v \in C^\infty(\mathbb{R}^n \times (0, \infty))$ and suppose

$$(5) \quad \begin{cases} u - v \text{ has a strict local maximum at some point} \\ (x_0, t_0) \in \mathbb{R}^n \times (0, \infty). \end{cases}$$

This means

$$(u - v)(x_0, t_0) > (u - v)(x, t)$$

for all points (x, t) sufficiently close to (x_0, t_0) , with $(x, t) \neq (x_0, t_0)$.

Now recall (3). We claim for each sufficiently small $\epsilon_j > 0$, there exists a point $(x_{\epsilon_j}, t_{\epsilon_j})$ such that

$$(6) \quad u^{\epsilon_j} - v \text{ has a local maximum at } (x_{\epsilon_j}, t_{\epsilon_j})$$

and

$$(7) \quad (x_{\epsilon_j}, t_{\epsilon_j}) \rightarrow (x_0, t_0) \quad \text{as } j \rightarrow \infty.$$

To confirm this, note that for each sufficiently small $r > 0$, (5) implies $\max_{\partial B}(u - v) < (u - v)(x_0, t_0)$, B denoting the closed ball in \mathbb{R}^{n+1} with center (x_0, t_0) and radius r . In view of (3), $u^{\epsilon_j} \rightarrow u$ uniformly on B , and so $\max_{\partial B}(u^{\epsilon_j} - v) < (u^{\epsilon_j} - v)(x_0, t_0)$ provided ϵ_j is small enough. Consequently $u^{\epsilon_j} - v$ attains a local maximum at some point in the interior of B . We can next replace r by a sequence of radii tending to zero to obtain (6), (7).

Now owing to (6), we see that the equations

$$(8) \quad Du^{\epsilon_j}(x_{\epsilon_j}, t_{\epsilon_j}) = Dv(x_{\epsilon_j}, t_{\epsilon_j}),$$

$$(9) \quad u_t^{\epsilon_j}(x_{\epsilon_j}, t_{\epsilon_j}) = v_t(x_{\epsilon_j}, t_{\epsilon_j})$$

and the inequality

$$(10) \quad -\Delta u^{\epsilon_j}(x_{\epsilon_j}, t_{\epsilon_j}) \geq -\Delta v(x_{\epsilon_j}, t_{\epsilon_j})$$

hold. We consequently can calculate

$$\begin{aligned}
 (11) \quad & v_t(x_{\epsilon_j}, t_{\epsilon_j}) + H(Dv(x_{\epsilon_j}, t_{\epsilon_j}), x_{\epsilon_j}) \\
 & = u_t^{\epsilon_j}(x_{\epsilon_j}, t_{\epsilon_j}) + H(Du^{\epsilon_j}(x_{\epsilon_j}, t_{\epsilon_j}), x_{\epsilon_j}) \quad \text{by (8),(9)} \\
 & = \epsilon_j \Delta u^{\epsilon_j}(x_{\epsilon_j}, t_{\epsilon_j}) \quad \text{by (2)} \\
 & \leq \epsilon_j \Delta v(x_{\epsilon_j}, t_{\epsilon_j}) \quad \text{by (10)}.
 \end{aligned}$$

Now let $\epsilon_j \rightarrow 0$ and remember (7). Since v is smooth and H is continuous, we deduce

$$(12) \quad v_t(x_0, t_0) + H(Dv(x_0, t_0), x_0) \leq 0.$$

We have established this inequality assuming (5). Suppose now instead that

$$(13) \quad u - v \text{ has a local maximum at } (x_0, t_0)$$

but that this maximum is not necessarily strict. Then $u - \tilde{v}$ has a strict local maximum at (x_0, t_0) , for $\tilde{v}(x, t) := v(x, t) + \delta(|x - x_0|^2 + (t - t_0)^2)$ ($\delta > 0$). We thus conclude as above that $\tilde{v}_t(x_0, t_0) + H(D\tilde{v}(x_0, t_0), x_0) \leq 0$, whereupon (12) again follows.

Consequently (13) implies inequality (12). Similarly, we deduce the reverse inequality

$$(14) \quad v_t(x_0, t_0) + H(Dv(x_0, t_0), x_0) \geq 0,$$

provided

$$(15) \quad u - v \text{ has a local minimum at } (x_0, t_0).$$

The proof is exactly like that above, except that the inequalities in (10), and thus in (11), are reversed.

In summary, we have discovered for any smooth function v that inequality (12) follows from (13), and (14) from (15). We have in effect put the derivatives onto v , at the expense of certain inequalities holding. \square

Our intention now is to *define* a weak solution of (1) in terms of (12), (13) and (14), (15).

DEFINITION. *Assume that u is bounded and uniformly continuous on $\mathbb{R}^n \times [0, T]$, for each $T > 0$. We say that u is a viscosity solution of the initial-value problem (1) for the Hamilton–Jacobi equation provided*

$$(i) \quad u = g \text{ on } \mathbb{R}^n \times \{t = 0\},$$

and

(ii) for each $v \in C^\infty(\mathbb{R}^n \times (0, \infty))$,

$$(16) \quad \begin{cases} \text{if } u - v \text{ has a local maximum at a point } (x_0, t_0) \in \mathbb{R}^n \times (0, \infty), \\ \text{then} \\ v_t(x_0, t_0) + H(Dv(x_0, t_0), x_0) \leq 0, \end{cases}$$

and

$$(17) \quad \begin{cases} \text{if } u - v \text{ has a local minimum at a point } (x_0, t_0) \in \mathbb{R}^n \times (0, \infty), \\ \text{then} \\ v_t(x_0, t_0) + H(Dv(x_0, t_0), x_0) \geq 0. \end{cases}$$

Remark. Note carefully that by definition a viscosity solution satisfies (16), (17), and so all subsequent deductions must be based on these inequalities. The previous discussion was purely motivational.

For emphasis, we repeat the same point, which has caused some confusion among students. To verify that a given function u is a viscosity solution of the Hamilton–Jacobi equation $u_t + H(Du, x) = 0$, we must confirm that (16), (17) hold for all smooth functions v . Now the argument above shows that *if* u is constructed using the vanishing viscosity method, it is indeed a viscosity solution. But we will also see later in §10.3 that viscosity solutions can be built in entirely different ways, which have nothing whatsoever to do with vanishing viscosity.

The point is that the inequalities (16), (17) provide an intrinsic characterization, and indeed the very definition, of our generalized solutions.

We devote the rest of this chapter to demonstrating that viscosity solutions provide an appropriate and useful notion of weak solutions for our Hamilton–Jacobi PDE.

10.1.2. Consistency.

Let us begin by checking that the notion of viscosity solution is consistent with that of a classical solution. First of all, note that if $u \in C^1(\mathbb{R}^n \times [0, \infty))$ solves (1) and if u is bounded and uniformly continuous, then u is a viscosity solution. That is, we assert that any *classical solution* of $u_t + H(Du, x) = 0$ is also a viscosity solution. The proof is easy. If v is smooth and $u - v$ obtains a local maximum at (x_0, t_0) , then

$$\begin{cases} Du(x_0, t_0) = Dv(x_0, t_0) \\ u_t(x_0, t_0) = v_t(x_0, t_0). \end{cases}$$

Consequently

$$\begin{aligned} v_t(x_0, t_0) + H(Dv(x_0, t_0), x_0) \\ = u_t(x_0, t_0) + H(Du(x_0, t_0), x_0) = 0, \end{aligned}$$

since u solves (1). A similar equality holds at any point (x_0, t_0) where $u - v$ has a local minimum.

Next we assert that *any sufficiently smooth viscosity solution is a classical solution* and, even more, that if a viscosity solution is differentiable at some point, then it solves the Hamilton–Jacobi PDE there. We will need the following calculus fact:

LEMMA (Touching by a C^1 function). *Assume $u : \mathbb{R}^n \rightarrow \mathbb{R}$ is continuous and is also differentiable at the point x_0 . Then there exists a function $v \in C^1(\mathbb{R}^n)$ such that*

$$(18) \quad u(x_0) = v(x_0)$$

and

$$(19) \quad u - v \text{ has a strict local maximum at } x_0.$$

Proof. 1. We may as well assume

$$(20) \quad x_0 = 0, \quad u(0) = Du(0) = 0,$$

for otherwise we could consider $\tilde{u}(x) := u(x + x_0) - u(x_0) - Du(x_0) \cdot x$ in place of u .

2. In view of (20) and our hypothesis, we have

$$(21) \quad u(x) = |x|\rho_1(x),$$

where

$$(22) \quad \rho_1 : \mathbb{R}^n \rightarrow \mathbb{R} \text{ is continuous, } \rho_1(0) = 0.$$

Set

$$(23) \quad \rho_2(r) := \max_{x \in B(0,r)} \{|\rho_1(x)|\} \quad (r \geq 0).$$

Then

$$(24) \quad \rho_2 : [0, \infty) \rightarrow [0, \infty) \text{ is continuous, } \rho_2(0) = 0,$$

and

$$(25) \quad \rho_2 \text{ is nondecreasing.}$$

3. Now write

$$v(x) := \int_{|x|}^{2|x|} \rho_2(r) dr + |x|^2 \quad (x \in \mathbb{R}^n).$$

Since $|v(x)| \leq |x|\rho_2(2|x|) + |x|^2$, we observe

$$(26) \quad v(0) = Dv(0) = 0.$$

Furthermore if $x \neq 0$, we have

$$Dv(x) = \frac{2x}{|x|} \rho_2(2|x|) - \frac{x}{|x|} \rho_2(|x|) + 2x,$$

and so $v \in C^1(\mathbb{R}^n)$.

4. Finally note that if $x \neq 0$,

$$\begin{aligned} u(x) - v(x) &= |x|\rho_1(x) - \int_{|x|}^{2|x|} \rho_2(r) dr - |x|^2 \\ &\leq |x|\rho_2(|x|) - \int_{|x|}^{2|x|} \rho_2(r) dr - |x|^2 \\ &\leq -|x|^2 \quad \text{by (25)} \\ &< 0 = u(0) - v(0). \end{aligned}$$

Thus $u - v$ has a strict local maximum at 0, as required. \square

THEOREM 1 (Consistency of viscosity solutions). *Let u be a viscosity solution of (1), and suppose u is differentiable at some point $(x_0, t_0) \in \mathbb{R}^n \times (0, \infty)$. Then*

$$u_t(x_0, t_0) + H(Du(x_0, t_0), x_0) = 0.$$

Proof. 1. Applying the lemma above to u , with \mathbb{R}^{n+1} replacing \mathbb{R}^n and (x_0, t_0) replacing x_0 , we deduce there exists a C^1 function v such that

$$(27) \quad u - v \text{ has a strict maximum at } (x_0, t_0).$$

2. Now set $v^\epsilon := \eta_\epsilon * v$, η_ϵ denoting the usual mollifier in the $n + 1$ variables (x, t) . Then

$$(28) \quad \begin{cases} v^\epsilon \rightarrow v \\ Dv^\epsilon \rightarrow Dv \quad \text{uniformly near } (x_0, t_0) \\ v_t^\epsilon \rightarrow v_t; \end{cases}$$

and so (27) implies

$$(29) \quad u - v^\epsilon \text{ has a maximum at some point } (x_\epsilon, t_\epsilon),$$

with

$$(30) \quad (x_\epsilon, t_\epsilon) \rightarrow (x_0, t_0) \quad \text{as } \epsilon \rightarrow 0.$$

Applying then the definition of viscosity solution, we see

$$v_t^\epsilon(x_\epsilon, t_\epsilon) + H(Dv^\epsilon(x_\epsilon, t_\epsilon), x_\epsilon) \leq 0.$$

Let $\epsilon \rightarrow 0$ and use (28), (30) to deduce

$$(31) \quad v_t(x_0, t_0) + H(Dv(x_0, t_0), x_0) \leq 0.$$

But in view of (27), we see that since u is differentiable at (x_0, t_0) ,

$$Du(x_0, t_0) = Dv(x_0, t_0), \quad u_t(x_0, t_0) = v_t(x_0, t_0).$$

Substitute above, to conclude from (31) that

$$(32) \quad u_t(x_0, t_0) + H(Du(x_0, t_0), x_0) \leq 0.$$

3. Now apply the lemma above to $-u$ in \mathbb{R}^{n+1} , to find a C^1 function v such that $u - v$ has a strict minimum at (x_0, t_0) . Then, arguing as above, we likewise deduce

$$u_t(x_0, t_0) + H(Du(x_0, t_0), x_0) \geq 0.$$

This inequality and (32) complete the proof. □

10.2. UNIQUENESS

Our goal now is to establish the uniqueness of a viscosity solution of our initial-value problem for Hamilton–Jacobi PDE. To be slightly more general, let us fix a time $T > 0$ and consider the problem

$$(1) \quad \begin{cases} u_t + H(Du, x) = 0 & \text{in } \mathbb{R}^n \times (0, T] \\ u = g & \text{on } \mathbb{R}^n \times \{t = 0\}. \end{cases}$$

We say that a bounded, uniformly continuous function u is a viscosity solution of (1) provided $u = g$ on $\mathbb{R}^n \times \{t = 0\}$, and the inequalities in (16) (or (17)) from §10.1.1 hold if $u - v$ has a local maximum (or minimum) at a point $(x_0, t_0) \in \mathbb{R}^n \times (0, T)$.

LEMMA (Extrema at a terminal time). *Assume u is a viscosity solution of (1) and $u - v$ has a local maximum (minimum) at a point $(x_0, t_0) \in \mathbb{R}^n \times (0, T]$. Then*

$$(2) \quad v_t(x_0, t_0) + H(Dv(x_0, t_0), x_0) \leq 0 \quad (\geq 0).$$

The point is that we are now allowing for $t_0 = T$.

Proof. Assume $u - v$ has a local maximum at the point (x_0, T) ; as before we may assume that this is a strict local maximum. Write

$$\tilde{v}(x, t) := v(x, t) + \frac{\epsilon}{T - t} \quad (x \in \mathbb{R}^n, 0 < t < T).$$

Then for $\epsilon > 0$ small enough, $u - \tilde{v}$ has a local maximum at a point (x_ϵ, t_ϵ) , where $0 < t_\epsilon < T$ and $(x_\epsilon, t_\epsilon) \rightarrow (x_0, T)$. Consequently

$$\tilde{v}_t(x_\epsilon, t_\epsilon) + H(D\tilde{v}(x_\epsilon, t_\epsilon), x_\epsilon) \leq 0,$$

and so

$$v_t(x_\epsilon, t_\epsilon) + \frac{\epsilon}{(T - t_\epsilon)^2} + H(Dv(x_\epsilon, t_\epsilon), x_\epsilon) \leq 0.$$

Letting $\epsilon \rightarrow 0$, we find

$$v_t(x_0, T) + H(Dv(x_0, T), x_0) \leq 0.$$

This proves (2) if $u - v$ has a maximum at (x_0, T) . A similar proof gives the reverse inequality should $u - v$ have a minimum at (x_0, T) . \square

To go further, let us hereafter suppose the Hamiltonian H to satisfy these conditions of Lipschitz continuity:

$$(3) \quad \begin{cases} |H(p, x) - H(q, x)| \leq C|p - q| \\ |H(p, x) - H(p, y)| \leq C|x - y|(1 + |p|) \end{cases}$$

for $x, y, p, q \in \mathbb{R}^n$ and some constant $C \geq 0$.

We come next to the central fact concerning viscosity solutions of the initial-value problem (1), namely uniqueness. This important assertion justifies our taking the inequalities (16) and (17) from §10.1.1 as the foundation of our theory.

THEOREM 1 (Uniqueness of viscosity solution). *Under assumption (3) there exists at most one viscosity solution of (1).*

The following proof is based upon an unusual idea of “doubling the number of variables”. See the proof of Theorem 3 in §11.4.3 for a related technique.

Proof*. 1. Assume u and \tilde{u} are both viscosity solutions with the same initial conditions, but

$$(4) \quad \sup_{\mathbb{R}^n \times [0, T]} (u - \tilde{u}) =: \sigma > 0.$$

*Omit on first reading.

Choose $0 < \epsilon, \lambda < 1$ and set

$$(5) \quad \begin{aligned} \Phi(x, y, t, s) := & u(x, t) - \tilde{u}(y, s) - \lambda(t + s) \\ & - \frac{1}{\epsilon^2}(|x - y|^2 + (t - s)^2) - \epsilon(|x|^2 + |y|^2), \end{aligned}$$

for $x, y \in \mathbb{R}^n$, $t, s \geq 0$. Then there exists a point $(x_0, y_0, t_0, s_0) \in \mathbb{R}^{2n} \times [0, T]^2$ such that

$$(6) \quad \Phi(x_0, y_0, t_0, s_0) = \max_{\mathbb{R}^{2n} \times [0, T]^2} \Phi(x, y, t, s).$$

2. We may fix $0 < \epsilon, \lambda < 1$ so small that (4) implies

$$(7) \quad \Phi(x_0, y_0, t_0, s_0) \geq \sup_{\mathbb{R}^n \times [0, T]} \Phi(x, x, t, t) \geq \frac{\sigma}{2}.$$

In addition, $\Phi(x_0, y_0, t_0, s_0) \geq \Phi(0, 0, 0, 0)$; and therefore

$$(8) \quad \begin{aligned} \lambda(t_0 + s_0) + \frac{1}{\epsilon^2}(|x_0 - y_0|^2 + (t_0 - s_0)^2) + \epsilon(|x_0|^2 + |y_0|^2) \\ \leq u(x_0, t_0) - \tilde{u}(y_0, s_0) - u(0, 0) + \tilde{u}(0, 0). \end{aligned}$$

Since u and \tilde{u} are bounded, we deduce

$$(9) \quad |x_0 - y_0|, |t_0 - s_0| = O(\epsilon) \quad \text{as } \epsilon \rightarrow 0.$$

Furthermore (8) implies $\epsilon(|x_0|^2 + |y_0|^2) = O(1)$, and consequently

$$\begin{aligned} \epsilon(|x_0| + |y_0|) &= \epsilon^{1/4} \epsilon^{3/4} (|x_0| + |y_0|) \\ &\leq \epsilon^{1/2} + C \epsilon^{3/2} (|x_0|^2 + |y_0|^2) \\ &\leq C \epsilon^{1/2}. \end{aligned}$$

Thus

$$(10) \quad \epsilon(|x_0| + |y_0|) = O(\epsilon^{1/2}).$$

3. Since $\Phi(x_0, y_0, t_0, s_0) \geq \Phi(x_0, x_0, t_0, t_0)$, we also have

$$\begin{aligned} u(x_0, t_0) - \tilde{u}(y_0, s_0) - \lambda(t_0 + s_0) - \frac{1}{\epsilon^2}(|x_0 - y_0|^2 + (t_0 - s_0)^2) \\ - \epsilon(|x_0|^2 + |y_0|^2) \geq u(x_0, t_0) - \tilde{u}(x_0, t_0) - 2\lambda t_0 - 2\epsilon|x_0|^2. \end{aligned}$$

Hence

$$\begin{aligned} \frac{1}{\epsilon^2}(|x_0 - y_0|^2 + (t_0 - s_0)^2) &\leq \tilde{u}(x_0, t_0) - \tilde{u}(y_0, s_0) + \lambda(t_0 - s_0) \\ &\quad + \epsilon(x_0 + y_0) \cdot (x_0 - y_0). \end{aligned}$$

In view of (9), (10) and the uniform continuity of \tilde{u} , we deduce

$$(11) \quad |x_0 - y_0|, |t_0 - s_0| = o(\epsilon).$$

4. Now write $\omega(\cdot)$ to denote the modulus of continuity of u ; that is,

$$|u(x, t) - u(y, s)| \leq \omega(|x - y| + |t - s|)$$

for all $x, y \in \mathbb{R}^n$, $0 \leq t, s \leq T$, and $\omega(r) \rightarrow 0$ as $r \rightarrow 0$. Similarly, $\tilde{\omega}(\cdot)$ will denote the modulus of continuity of \tilde{u} .

Then (7) implies

$$\begin{aligned} \frac{\sigma}{2} &\leq u(x_0, t_0) - \tilde{u}(y_0, s_0) = u(x_0, t_0) - u(x_0, 0) + u(x_0, 0) - \tilde{u}(x_0, 0) \\ &\quad + \tilde{u}(x_0, 0) - \tilde{u}(x_0, t_0) + \tilde{u}(x_0, t_0) - \tilde{u}(y_0, s_0) \\ &\leq \omega(t_0) + \tilde{\omega}(t_0) + \tilde{\omega}(o(\epsilon)), \end{aligned}$$

by (9), (11) and the initial condition. We can now take $\epsilon > 0$ to be so small that the foregoing implies $\frac{\sigma}{4} \leq \omega(t_0) + \tilde{\omega}(t_0)$; and this in turn implies $t_0 \geq \mu > 0$ for some constant $\mu > 0$. Similarly we have $s_0 \geq \mu > 0$.

5. Now observe in light of (6) that the mapping $(x, t) \mapsto \Phi(x, y_0, t, s_0)$ has a maximum at the point (x_0, t_0) . In view of (5) then,

$$u - v \text{ has a maximum at } (x_0, t_0)$$

for

$$v(x, t) := \tilde{u}(y_0, s_0) + \lambda(t + s_0) + \frac{1}{\epsilon^2}(|x - y_0|^2 + (t - s_0)^2) + \epsilon(|x|^2 + |y_0|^2).$$

Since u is a viscosity solution of (1), we conclude, using the lemma if necessary, that

$$v_t(x_0, t_0) + H(D_x v(x_0, t_0), x_0) \leq 0.$$

Therefore

$$(12) \quad \lambda + \frac{2(t_0 - s_0)}{\epsilon^2} + H\left(\frac{2}{\epsilon^2}(x_0 - y_0) + 2\epsilon x_0, x_0\right) \leq 0.$$

We further observe that since the mapping $(y, s) \mapsto -\Phi(x_0, y, t_0, s)$ has a minimum at the point (y_0, s_0) ,

$$\tilde{u} - \tilde{v} \text{ has a minimum at } (y_0, s_0)$$

for

$$\tilde{v}(y, s) := u(x_0, t_0) - \lambda(t_0 + s) - \frac{1}{\epsilon^2}(|x_0 - y|^2 + (t_0 - s)^2) - \epsilon(|x_0|^2 + |y|^2).$$

As \tilde{u} is a viscosity solution of (1), we know then that

$$\tilde{v}_s(y_0, s_0) + H(D_y \tilde{v}(y_0, s_0), y_0) \geq 0.$$

Consequently

$$(13) \quad -\lambda + \frac{2(t_0 - s_0)}{\epsilon^2} + H\left(\frac{2}{\epsilon^2}(x_0 - y_0) - 2\epsilon y_0, y_0\right) \geq 0.$$

6. Next, subtract (13) from (12):

$$(14) \quad 2\lambda \leq H\left(\frac{2}{\epsilon^2}(x_0 - y_0) - 2\epsilon y_0, y_0\right) - H\left(\frac{2}{\epsilon^2}(x_0 - y_0) + 2\epsilon x_0, x_0\right).$$

In view of hypothesis (3) therefore,

$$(15) \quad \lambda \leq C\epsilon(|x_0| + |y_0|) + C|x_0 - y_0| \left(1 + \frac{|x_0 - y_0|}{\epsilon^2} + \epsilon(|x_0| + |y_0|)\right).$$


We employ estimates (10), (11) in (15) and then let $\epsilon \rightarrow 0$, to discover $0 < \lambda \leq 0$. This contradiction completes the proof. \square

10.3. CONTROL THEORY, DYNAMIC PROGRAMMING

It remains for us to establish the existence of a viscosity solution to our initial-value problem for the Hamilton–Jacobi partial differential equation. One method would be now to prove the existence of a smooth solution u^ϵ of the regularized equation (2) in §10.1 and then to make good enough uniform estimates. This technique in fact works but requires knowledge of certain bounds for the heat equation beyond the scope of this book.

In this section we provide an alternative approach of independent interest, which is suitable for Hamiltonians which are convex in p .

We will first of all introduce some of the basic issues concerning *control theory* for ordinary differential equations and the connection with Hamilton–Jacobi PDE afforded by the method of *dynamic programming*. This discussion will make clearer the connections of the theory developed above in §§10.1–10.2 with that set forth earlier in §3.3.1. The remarkable fact is that *the defining viscosity solution inequalities* (16), (17) in §10.1.1 *are a consequence of the optimality conditions of control theory*.



Response of system to the control $\alpha(\cdot)$

10.3.1. Introduction to optimal control theory.

We will now study the possibility of optimally controlling the solution $\mathbf{x}(\cdot)$ of the ordinary differential equation

$$(1) \quad \begin{cases} \dot{\mathbf{x}}(s) = \mathbf{f}(\mathbf{x}(s), \alpha(s)) & (t < s < T) \\ \mathbf{x}(t) = x. \end{cases}$$

Here $\dot{\cdot} = \frac{d}{ds}$, $T > 0$ is a fixed terminal time, and $x \in \mathbb{R}^n$ is a given initial point, taken on by our solution $\mathbf{x}(\cdot)$ at the starting time $t \geq 0$. At later times $t < s < T$, $\mathbf{x}(\cdot)$ evolves according to the ODE, where

$$\mathbf{f} : \mathbb{R}^n \times A \rightarrow \mathbb{R}^n$$

is a given bounded, Lipschitz continuous function and A is some given compact subset of, say, \mathbb{R}^m . The function $\alpha(\cdot)$ appearing in (1) is a *control*, that is, some appropriate scheme for adjusting parameters from the set A as time evolves, thereby affecting the dynamics of the system modeled by (1).

Let us write

$$(2) \quad \mathcal{A} := \{\alpha : [0, T] \rightarrow A \mid \alpha(\cdot) \text{ is measurable}\}$$

to denote the set of *admissible controls*. Then since

$$(3) \quad |\mathbf{f}(x, a)| \leq C, \quad |\mathbf{f}(x, a) - \mathbf{f}(y, a)| \leq C|x - y| \quad (x, y \in \mathbb{R}^n, a \in A)$$

for some constant C , we see that for each control $\alpha(\cdot) \in \mathcal{A}$, the ODE (1) has a unique, Lipschitz continuous solution $\mathbf{x}(\cdot) = \mathbf{x}^{\alpha(\cdot)}(\cdot)$, existing on the time interval $[t, T]$ and solving the ODE for a.e. time $t < s < T$. We call $\mathbf{x}(\cdot)$ the *response* of the system to the control $\alpha(\cdot)$, and $\mathbf{x}(s)$ the *state* of the system at time s .

Our goal is to find a control $\alpha^*(\cdot)$ which optimally steers the system. However in order to define what “optimal” means, we must first introduce a *cost criterion*. Given $x \in \mathbb{R}^n$ and $0 \leq t \leq T$, let us define for each admissible control $\alpha(\cdot) \in \mathcal{A}$ the corresponding *cost functional*

$$(4) \quad C_{x,t}[\alpha(\cdot)] := \int_t^T r(\mathbf{x}(s), \alpha(s)) ds + g(\mathbf{x}(T)),$$

where $\mathbf{x}(\cdot) = \mathbf{x}^{\alpha(\cdot)}(\cdot)$ solves the ODE (1) and

$$r : \mathbb{R}^n \times A \rightarrow \mathbb{R}, \quad g : \mathbb{R}^n \rightarrow \mathbb{R}$$

are given functions. We call r the *running cost per unit time* and g the *terminal cost*, and will henceforth assume

$$(5) \quad \begin{cases} |r(x, a)|, |g(x)| \leq C \\ |r(x, a) - r(y, a)|, |g(x) - g(y)| \leq C|x - y| \end{cases} \quad (x, y \in \mathbb{R}^n, a \in A)$$

for some constant C .

Given now $x \in \mathbb{R}^n$ and $0 \leq t \leq T$, we would like to find if possible a control $\alpha^*(\cdot)$ which minimizes the cost functional (4) among all other admissible controls. This is a *finite horizon* optimal control problem. (See Problems 10 and 11 for infinite horizon problems.)

10.3.2. Dynamic programming.

The method of *dynamic programming* investigates the above problem by turning attention to the *value function*

$$(6) \quad u(x, t) := \inf_{\alpha(\cdot) \in \mathcal{A}} C_{x,t}[\alpha(\cdot)] \quad (x \in \mathbb{R}^n, 0 \leq t \leq T).$$

The plan is this: having defined $u(x, t)$ as the least cost given that we start at the position x at time t , we want to study u as a function of x and t . We are therefore embedding our given control problem (1), (4) into the larger class of all such problems, as x and t vary. The idea then is to show that u solves a certain Hamilton–Jacobi type PDE and to show conversely that a solution of this PDE helps us to synthesize an optimal feedback control.

Hereafter, we fix $x \in \mathbb{R}^n$, $0 \leq t < T$.

THEOREM 1 (Optimality conditions). *For each $h > 0$ so small that $t + h \leq T$, we have*

$$(7) \quad u(x, t) = \inf_{\alpha(\cdot) \in \mathcal{A}} \left\{ \int_t^{t+h} r(\mathbf{x}(s), \alpha(s)) ds + u(\mathbf{x}(t+h), t+h) \right\},$$

where $\mathbf{x}(\cdot) = \mathbf{x}^{\alpha(\cdot)}(\cdot)$ solves the ODE (1) for the control $\alpha(\cdot)$.

Proof. 1. Choose any control $\alpha_1(\cdot) \in \mathcal{A}$ and solve the ODE

$$(8) \quad \begin{cases} \dot{\mathbf{x}}_1(s) = \mathbf{f}(\mathbf{x}_1(s), \alpha_1(s)) & (t < s < t+h) \\ \mathbf{x}_1(t) = x. \end{cases}$$

Fix $\epsilon > 0$ and choose then $\alpha_2(\cdot) \in \mathcal{A}$ so that

$$(9) \quad u(\mathbf{x}_1(t+h), t+h) + \epsilon \geq \int_{t+h}^T r(\mathbf{x}_2(s), \alpha_2(s)) ds + g(\mathbf{x}_2(T)),$$

where

$$(10) \quad \begin{cases} \dot{\mathbf{x}}_2(s) = \mathbf{f}(\mathbf{x}_2(s), \alpha_2(s)) & (t+h < s < T) \\ \mathbf{x}_2(t+h) = \mathbf{x}_1(t+h). \end{cases}$$

Now define the control

$$(11) \quad \alpha_3(s) := \begin{cases} \alpha_1(s) & \text{if } t \leq s < t+h \\ \alpha_2(s) & \text{if } t+h \leq s \leq T, \end{cases}$$

and let

$$(12) \quad \begin{cases} \dot{\mathbf{x}}_3(s) = \mathbf{f}(\mathbf{x}_3(s), \alpha_3(s)) & (t < s < T) \\ \mathbf{x}_3(t) = x. \end{cases}$$

By uniqueness of solutions to the differential equation (1), we have

$$(13) \quad \mathbf{x}_3(s) = \begin{cases} \mathbf{x}_1(s) & \text{if } t \leq s \leq t+h \\ \mathbf{x}_2(s) & \text{if } t+h \leq s \leq T. \end{cases}$$

Thus the definition (6) implies

$$\begin{aligned} u(x, t) &\leq C_{x,t}[\alpha_3(\cdot)] \\ &= \int_t^T r(\mathbf{x}_3(s), \alpha_3(s)) ds + g(\mathbf{x}_3(T)) \\ &= \int_t^{t+h} r(\mathbf{x}_1(s), \alpha_1(s)) ds + \int_{t+h}^T r(\mathbf{x}_2(s), \alpha_2(s)) ds + g(\mathbf{x}_2(T)) \\ &\leq \int_t^{t+h} r(\mathbf{x}_1(s), \alpha_1(s)) ds + u(\mathbf{x}_1(t+h), t+h) + \epsilon, \end{aligned}$$

the last inequality resulting from (9). As $\alpha_1(\cdot) \in \mathcal{A}$ was arbitrary, we conclude

$$(14) \quad u(x, t) \leq \inf_{\alpha(\cdot) \in \mathcal{A}} \left\{ \int_t^{t+h} r(\mathbf{x}(s), \alpha(s)) ds + u(\mathbf{x}(t+h), t+h) \right\} + \epsilon,$$

$\mathbf{x}(\cdot) = \mathbf{x}^{\alpha(\cdot)}(\cdot)$ solving (1).

2. Fixing again $\epsilon > 0$, select now $\alpha_4(\cdot) \in \mathcal{A}$ so that

$$(15) \quad u(x, t) + \epsilon \geq \int_t^T r(\mathbf{x}_4(s), \alpha_4(s)) ds + g(\mathbf{x}_4(T)),$$

where

$$\begin{cases} \dot{\mathbf{x}}_4(s) = \mathbf{f}(\mathbf{x}_4(s), \alpha_4(s)) & (t < s < T) \\ \mathbf{x}_4(t) = x. \end{cases}$$

Observe then from (6) that

$$(16) \quad u(\mathbf{x}_4(t+h), t+h) \leq \int_{t+h}^T r(\mathbf{x}_4(s), \alpha_4(s)) ds + g(\mathbf{x}_4(T)).$$

Therefore

$$u(x, t) + \epsilon \geq \inf_{\alpha(\cdot) \in \mathcal{A}} \left\{ \int_t^{t+h} r(\mathbf{x}(s), \alpha(s)) ds + u(\mathbf{x}(t+h), t+h) \right\},$$

$\mathbf{x}(\cdot) = \mathbf{x}^{\alpha(\cdot)}(\cdot)$ solving (1). This inequality and (14) complete the proof of (7). \square

10.3.3. Hamilton–Jacobi–Bellman equation.

Our eventual goal is writing down as a PDE an “infinitesimal version” of the optimality conditions (7). But first we must check that the value function u is bounded and Lipschitz continuous.

LEMMA (Estimates for value function). *There exists a constant C such that*

$$|u(x, t)| \leq C,$$

$$|u(x, t) - u(\hat{x}, \hat{t})| \leq C(|x - \hat{x}| + |t - \hat{t}|)$$

for all $x, \hat{x} \in \mathbb{R}^n$, $0 \leq t, \hat{t} \leq T$.

Proof. 1. Clearly hypothesis (5) implies u is bounded on $\mathbb{R}^n \times [0, T]$.

2. Fix $x, \hat{x} \in \mathbb{R}^n$, $0 \leq t < T$. Let $\epsilon > 0$ and then choose $\hat{\alpha}(\cdot) \in \mathcal{A}$ so that

$$(17) \quad u(\hat{x}, t) + \epsilon \geq \int_t^T r(\hat{\mathbf{x}}(s), \hat{\alpha}(s)) ds + g(\hat{\mathbf{x}}(T)),$$

where $\hat{\mathbf{x}}(\cdot)$ solves the ODE

$$(18) \quad \begin{cases} \dot{\hat{\mathbf{x}}}(s) = \mathbf{f}(\hat{\mathbf{x}}(s), \hat{\alpha}(s)) & (t < s < T) \\ \hat{\mathbf{x}}(t) = \hat{x}. \end{cases}$$

Then

$$(19) \quad \begin{aligned} u(x, t) - u(\hat{x}, t) &\leq \int_t^T r(\mathbf{x}(s), \hat{\alpha}(s)) ds + g(\mathbf{x}(T)) \\ &\quad - \int_t^T r(\hat{\mathbf{x}}(s), \hat{\alpha}(s)) ds - g(\hat{\mathbf{x}}(T)) + \epsilon, \end{aligned}$$

where $\mathbf{x}(\cdot)$ solves

$$(20) \quad \begin{cases} \dot{\mathbf{x}}(s) = \mathbf{f}(\mathbf{x}(s), \hat{\alpha}(s)) & (t < s < T) \\ \mathbf{x}(t) = x. \end{cases}$$

Since \mathbf{f} is Lipschitz continuous, (18), (20) and Gronwall's inequality (§B.2) imply $|\mathbf{x}(s) - \hat{\mathbf{x}}(s)| \leq C|x - \hat{x}|$ ($t \leq s \leq T$). Hence we deduce from (5) and (19) that $u(x, t) - u(\hat{x}, t) \leq C|x - \hat{x}| + \epsilon$. The same argument with the roles of x and \hat{x} reversed implies

$$|u(x, t) - u(\hat{x}, t)| \leq C|x - \hat{x}| \quad (x, \hat{x} \in \mathbb{R}^n, 0 \leq t \leq T).$$

3. Now let $x \in \mathbb{R}^n$, $0 \leq t < \hat{t} \leq T$. Take $\epsilon > 0$ and choose $\alpha(\cdot) \in \mathcal{A}$ so that

$$u(x, t) + \epsilon \geq \int_t^T r(\mathbf{x}(s), \alpha(s)) ds + g(\mathbf{x}(T)),$$

$\mathbf{x}(\cdot)$ solving the ODE (1). Define

$$\hat{\alpha}(s) := \alpha(s + t - \hat{t}) \quad \text{for } \hat{t} \leq s \leq T$$

and let $\hat{\mathbf{x}}(\cdot)$ solve

$$\begin{cases} \dot{\hat{\mathbf{x}}}(s) = \mathbf{f}(\hat{\mathbf{x}}(s), \hat{\alpha}(s)) & (\hat{t} < s < T) \\ \hat{\mathbf{x}}(\hat{t}) = x. \end{cases}$$

Then $\hat{\mathbf{x}}(s) = \mathbf{x}(s + t - \hat{t})$. Hence

$$(21) \quad \begin{aligned} u(x, \hat{t}) - u(x, t) &\leq \int_{\hat{t}}^T r(\hat{\mathbf{x}}(s), \hat{\alpha}(s)) ds + g(\hat{\mathbf{x}}(T)) \\ &\quad - \int_t^T r(\mathbf{x}(s), \alpha(s)) ds - g(\mathbf{x}(T)) + \epsilon \\ &= - \int_{T+t-\hat{t}}^T r(\mathbf{x}(s), \alpha(s)) ds + g(\mathbf{x}(T + t - \hat{t})) - g(\mathbf{x}(T)) + \epsilon \\ &\leq C|t - \hat{t}| + \epsilon. \end{aligned}$$

Next pick $\hat{\alpha}(\cdot)$ so that

$$u(x, \hat{t}) + \epsilon \geq \int_{\hat{t}}^T r(\hat{\mathbf{x}}(s), \hat{\alpha}(s)) ds + g(\hat{\mathbf{x}}(T)),$$

where

$$\begin{cases} \dot{\hat{\mathbf{x}}}(s) = \mathbf{f}(\hat{\mathbf{x}}(s), \hat{\alpha}(s)) & (\hat{t} < s < T) \\ \hat{\mathbf{x}}(\hat{t}) = x. \end{cases}$$

Define

$$\alpha(s) := \begin{cases} \hat{\alpha}(s + \hat{t} - t) & \text{if } t \leq s \leq T + t - \hat{t} \\ \hat{\alpha}(T) & \text{if } T + t - \hat{t} \leq s \leq T, \end{cases}$$

and let $\mathbf{x}(\cdot)$ solve (1). Then $\alpha(s) = \hat{\alpha}(s + \hat{t} - t)$, $\mathbf{x}(s) = \hat{\mathbf{x}}(s + \hat{t} - t)$ for $t \leq s \leq T + t - \hat{t}$. Consequently

$$\begin{aligned} u(x, t) - u(x, \hat{t}) &\leq \int_t^T r(\mathbf{x}(s), \alpha(s)) ds + g(\mathbf{x}(T)) \\ &\quad - \int_{\hat{t}}^T r(\hat{\mathbf{x}}(s), \alpha(s)) ds - g(\hat{\mathbf{x}}(T)) + \epsilon \\ &= \int_{T+t-\hat{t}}^T r(\mathbf{x}(s), \alpha(s)) ds + g(\mathbf{x}(T)) - g(\mathbf{x}(T + t - \hat{t})) + \epsilon \\ &\leq C|t - \hat{t}| + \epsilon. \end{aligned}$$

This inequality and (21) prove

$$|u(x, t) - u(x, \hat{t})| \leq C|t - \hat{t}| \quad (0 \leq t \leq \hat{t} \leq T, x \in \mathbb{R}^n). \quad \square$$

We prove next that the value function solves a Hamilton–Jacobi type partial differential equation.

THEOREM 2 (A PDE for the value function). *The value function u is the unique viscosity solution of this terminal-value problem for the Hamilton–Jacobi–Bellman equation:*

$$(22) \quad \begin{cases} u_t + \min_{a \in A} \{ \mathbf{f}(x, a) \cdot Du + r(x, a) \} = 0 & \text{in } \mathbb{R}^n \times (0, T) \\ u = g & \text{on } \mathbb{R}^n \times \{t = T\}. \end{cases}$$

Remarks. (i) The Hamilton–Jacobi–Bellman PDE has the form

$$u_t + H(Du, x) = 0 \quad \text{in } \mathbb{R}^n \times (0, T),$$

for the Hamiltonian

$$(23) \quad H(p, x) := \min_{a \in A} \{ \mathbf{f}(x, a) \cdot p + r(x, a) \} \quad (p, x \in \mathbb{R}^n).$$

From the inequalities (5), we deduce that H satisfies the estimates (3) in §10.2.

(ii) Since (22) is a *terminal-value problem*, we must specify what we mean by a solution. Let us say that a bounded, uniformly continuous function u is a *viscosity solution* of (22) provided

$$(a) \quad u = g \text{ on } \mathbb{R}^n \times \{t = T\},$$

and

$$(b) \text{ for each } v \in C^\infty(\mathbb{R}^n \times (0, T))$$

$$(24) \quad \begin{cases} \text{if } u - v \text{ has a local maximum at a point } (x_0, t_0) \in \mathbb{R}^n \times (0, T), \\ \text{then} \\ v_t(x_0, t_0) + H(Dv(x_0, t_0), x_0) \geq 0, \end{cases}$$

and

$$(25) \quad \begin{cases} \text{if } u - v \text{ has a local minimum at a point } (x_0, t_0) \in \mathbb{R}^n \times (0, T), \\ \text{then} \\ v_t(x_0, t_0) + H(Dv(x_0, t_0), x_0) \leq 0. \end{cases}$$

Observe that for our terminal-value problem (22) we *reverse* the sense of the inequalities from those for the initial-value problem.

(iii) The reader should check that if u is the viscosity solution of (22), then $w(x, t) := u(x, T - t)$ ($x \in \mathbb{R}^n, 0 \leq t \leq T$) is the viscosity solution of the initial-value problem

$$\begin{cases} w_t - H(Dw, x) = 0 & \text{in } \mathbb{R}^n \times (0, T) \\ w = g & \text{on } \mathbb{R}^n \times \{t = 0\}. \end{cases}$$

Proof. 1. In view of the lemma, u is bounded and Lipschitz continuous. In addition, we see directly from (4) and (6) that

$$u(x, T) = \inf_{\alpha(\cdot) \in \mathcal{A}} C_{x, T}[\alpha(\cdot)] = g(x) \quad (x \in \mathbb{R}^n).$$

2. Now let $v \in C^\infty(\mathbb{R}^n \times (0, T))$, and assume

$$u - v \text{ has a local maximum at a point } (x_0, t_0) \in \mathbb{R}^n \times (0, T).$$

We must prove

$$(26) \quad v_t(x_0, t_0) + \min_{a \in A} \{f(x_0, a) \cdot Dv(x_0, t_0) + r(x_0, a)\} \geq 0.$$

Suppose not. Then there exist $a \in A$ and $\theta > 0$ such that

$$(27) \quad v_t(x, t) + \mathbf{f}(x, a) \cdot Dv(x, t) + r(x, a) \leq -\theta < 0$$

for all points (x, t) sufficiently close to (x_0, t_0) , say

$$(28) \quad |x - x_0| + |t - t_0| < \delta.$$

Since $u - v$ has a local maximum at (x_0, t_0) , we may as well also suppose

$$(29) \quad \begin{cases} (u - v)(x, t) \leq (u - v)(x_0, t_0) \\ \text{for all } (x, t) \text{ satisfying (28).} \end{cases}$$

Consider now the constant control $\alpha(s) \equiv a$ ($t_0 \leq s \leq T$) and the corresponding dynamics

$$(30) \quad \begin{cases} \dot{\mathbf{x}}(s) = \mathbf{f}(\mathbf{x}(s), a) & (t_0 < s < T) \\ \mathbf{x}(t_0) = x_0. \end{cases}$$

Choose $0 < h < \delta$ so small that $|\mathbf{x}(s) - x_0| < \delta$ for $t_0 \leq s \leq t_0 + h$. Then

$$(31) \quad v_t(\mathbf{x}(s), s) + \mathbf{f}(\mathbf{x}(s), a) \cdot Dv(\mathbf{x}(s), s) + r(\mathbf{x}(s), a) \leq -\theta \quad (t_0 \leq s \leq t_0 + h),$$

according to (27), (28). But utilizing (29), we find

$$(32) \quad \begin{aligned} & u(\mathbf{x}(t_0 + h), t_0 + h) - u(x_0, t_0) \leq v(\mathbf{x}(t_0 + h), t_0 + h) - v(x_0, t_0) \\ & = \int_{t_0}^{t_0+h} \frac{d}{ds} v(\mathbf{x}(s), s) ds = \int_{t_0}^{t_0+h} v_t(\mathbf{x}(s), s) + Dv(\mathbf{x}(s), s) \cdot \dot{\mathbf{x}}(s) ds \\ & = \int_{t_0}^{t_0+h} v_t(\mathbf{x}(s), s) + \mathbf{f}(\mathbf{x}(s), a) \cdot Dv(\mathbf{x}(s), s) ds. \end{aligned}$$

In addition, the optimality condition (7) provides us with the inequality

$$(33) \quad u(x_0, t_0) \leq \int_{t_0}^{t_0+h} r(\mathbf{x}(s), a) ds + u(\mathbf{x}(t_0 + h), t_0 + h).$$

Combining (32) and (33), we discover

$$0 \leq \int_{t_0}^{t_0+h} v_t(\mathbf{x}(s), s) + \mathbf{f}(\mathbf{x}(s), a) \cdot Dv(\mathbf{x}(s), s) + r(\mathbf{x}(s), a) ds \leq -\theta h,$$

according to (31). This contradiction establishes (26).

3. Now suppose

$u - v$ has a local minimum at a point $(x_0, t_0) \in \mathbb{R}^n \times (0, T)$;

we must prove

$$(34) \quad v_t(x_0, t_0) + \min_{a \in A} \{ \mathbf{f}(x_0, a) \cdot Dv(x_0, t_0) + r(x_0, a) \} \leq 0.$$

Suppose not. Then there exists $\theta > 0$ such that

$$(35) \quad v_t(x, t) + \mathbf{f}(x, a) \cdot Dv(x, t) + r(x, a) \geq \theta > 0$$

for all $a \in A$ and all (x, t) sufficiently close to (x_0, t_0) , say

$$(36) \quad |x - x_0| + |t - t_0| < \delta.$$

Since $u - v$ has a local minimum at (x_0, t_0) , we may as well also suppose

$$(37) \quad \begin{cases} (u - v)(x, t) \geq (u - v)(x_0, t_0) \\ \text{for all } (x, t) \text{ satisfying (36).} \end{cases}$$

Choose $0 < h < \delta$ so small that $|\mathbf{x}(s) - x_0| < \delta$ for $t_0 \leq s \leq t_0 + h$, where $\mathbf{x}(\cdot)$ solves

$$(38) \quad \begin{cases} \dot{\mathbf{x}}(s) = \mathbf{f}(\mathbf{x}(s), \alpha(s)) & (t_0 < s < T) \\ \mathbf{x}(t_0) = x_0 \end{cases}$$

for some control $\alpha(\cdot) \in \mathcal{A}$. This is possible owing to hypothesis (3).

Then utilizing (37), we find for any control $\alpha(\cdot)$ that

$$(39) \quad \begin{aligned} u(\mathbf{x}(t_0 + h), t_0 + h) - u(x_0, t_0) & \\ & \geq v(\mathbf{x}(t_0 + h), t_0 + h) - v(x_0, t_0) \\ & = \int_{t_0}^{t_0+h} \frac{d}{ds} v(\mathbf{x}(s), s) ds \\ & = \int_{t_0}^{t_0+h} v_t(\mathbf{x}(s), s) + \mathbf{f}(\mathbf{x}(s), \alpha(s)) \cdot Dv(\mathbf{x}(s), s) ds, \end{aligned}$$

by (38). On the other hand, according to the optimality condition (7) we can select a control $\alpha(\cdot) \in \mathcal{A}$ so that

$$(40) \quad u(x_0, t_0) \geq \int_{t_0}^{t_0+h} r(\mathbf{x}(s), \alpha(s)) ds + u(\mathbf{x}(t_0 + h), t_0 + h) - \frac{\theta h}{2}.$$

Combining (39) and (40), we discover

$$\begin{aligned} \frac{\theta h}{2} & \geq \int_{t_0}^{t_0+h} v_t(\mathbf{x}(s), s) + \mathbf{f}(\mathbf{x}(s), \alpha(s)) \cdot Dv(\mathbf{x}(s), s) \\ & \quad + r(\mathbf{x}(s), \alpha(s)) ds \geq \theta h, \end{aligned}$$

according to (35). This contradiction proves (34). \square

Design of optimal controls. We have now shown that the value function u , defined by (6), is the unique viscosity solution of the terminal-value problem (22) for the Hamilton–Jacobi–Bellman equation. How does this PDE help us solve the problem of synthesizing an optimal control? In informal terms, the method is this. Given an initial time $0 < t \leq T$ and an initial state $x \in \mathbb{R}^n$, we consider the optimal ODE

$$(41) \quad \begin{cases} \dot{\mathbf{x}}^*(s) = \mathbf{f}(\mathbf{x}^*(s), \boldsymbol{\alpha}^*(s)) & (t < s < T) \\ \mathbf{x}^*(t) = x, \end{cases}$$

where at each time s , $\boldsymbol{\alpha}^*(s) \in A$ is selected so that

$$(42) \quad \begin{aligned} \mathbf{f}(\mathbf{x}^*(s), \boldsymbol{\alpha}^*(s)) \cdot Du(\mathbf{x}^*(s), s) + r(\mathbf{x}^*(s), \boldsymbol{\alpha}^*(s)) \\ = H(Du(\mathbf{x}^*(s), s), \mathbf{x}^*(s)). \end{aligned}$$

In other words, given that the system is at the point $\mathbf{x}^*(s)$ at time s , we adjust the optimal control value $\boldsymbol{\alpha}^*(s)$ so as to attain the minimum in the definition (23) of the Hamiltonian H . We call $\boldsymbol{\alpha}^*(\cdot)$ so defined a *feedback control*.

It is fairly easy to check that this prescription does in fact generate a minimum cost trajectory, at least in regions where u and $\boldsymbol{\alpha}^*(\cdot)$ are smooth (so that (42) makes sense). There are however problems in interpreting (42) at points where the gradient Du does not exist.

10.3.4. Hopf–Lax formula revisited.

Remember that earlier in §3.3 we investigated this initial-value problem for the Hamilton–Jacobi equation:

$$(43) \quad \begin{cases} u_t + H(Du) = 0 & \text{in } \mathbb{R}^n \times (0, T] \\ u = g & \text{on } \mathbb{R}^n \times \{t = 0\}, \end{cases}$$

under the assumptions that

$$p \mapsto H(p) \text{ is convex, } \lim_{|p| \rightarrow \infty} \frac{H(p)}{|p|} = +\infty,$$

and

$$g : \mathbb{R}^n \rightarrow \mathbb{R} \text{ is Lipschitz continuous.}$$

Notice that we are now taking $0 \leq t \leq T$, to be consistent with §10.2. We introduced as well the Hopf–Lax formula for a solution:

$$(44) \quad u(x, t) = \min_{y \in \mathbb{R}^n} \left\{ tL \left(\frac{x - y}{t} \right) + g(y) \right\} \quad (x \in \mathbb{R}^n, t > 0),$$

where L is the Legendre transform of H :

$$(45) \quad L(v) = \sup_{p \in \mathbb{R}^n} \{p \cdot v - H(p)\} \quad (q \in \mathbb{R}^n).$$

In order to tie together the theory set forth here and in §3.3, let us now check that the Hopf–Lax formula gives the correct viscosity solution, as defined in §10.1.1. (The proof is really just a special case of that for Theorem 2.)

THEOREM 3 (Hopf–Lax formula as viscosity solution). *Assume in addition that g is bounded. Then the unique viscosity solution of the initial-value problem (43) is given by the formula (44).*

Proof. 1. As shown in §3.3 the function u defined by (44) is Lipschitz continuous and takes on the initial function g at time $t = 0$. It is easy to verify as well that u is also bounded on $\mathbb{R}^n \times (0, T]$, since g is bounded.

2. Now let $v \in C^\infty(\mathbb{R}^n \times (0, \infty))$ and assume $u - v$ has a local maximum at $(x_0, t_0) \in \mathbb{R}^n \times (0, \infty)$. According to Lemma 1 in §3.3.2,

$$(46) \quad u(x_0, t_0) = \min_{x \in \mathbb{R}^n} \left\{ (t_0 - t)L \left(\frac{x_0 - x}{t_0 - t} \right) + u(x, t) \right\}$$

for each $0 \leq t < t_0$. Thus for each $0 \leq t < t_0$, $x \in \mathbb{R}^n$

$$(47) \quad u(x_0, t_0) \leq (t_0 - t)L \left(\frac{x_0 - x}{t_0 - t} \right) + u(x, t).$$

But since $u - v$ has a local maximum at (x_0, t_0) ,

$$u(x_0, t_0) - v(x_0, t_0) \geq u(x, t) - v(x, t)$$

for (x, t) close to (x_0, t_0) . Combining this estimate with (47), we find

$$(48) \quad v(x_0, t_0) - v(x, t) \leq (t_0 - t)L \left(\frac{x_0 - x}{t_0 - t} \right)$$

for $t < t_0$, (x, t) close to (x_0, t_0) . Now write $h = t_0 - t$ and set $x = x_0 - hv$, where $v \in \mathbb{R}^n$ is given. Inequality (48) becomes

$$v(x_0, t_0) - v(x_0 - hv, t_0 - h) \leq hL(v).$$

Divide by $h > 0$ and send $h \rightarrow 0$:

$$v_t(x_0, t_0) + Dv(x_0, t_0) \cdot v - L(v) \leq 0.$$

This is true for all $v \in \mathbb{R}^n$ and so

$$(49) \quad v_t(x_0, t_0) + H(Dv(x_0, t_0)) \leq 0,$$

since

$$(50) \quad H(p) = \sup_{v \in \mathbb{R}^n} \{p \cdot v - L(v)\},$$

by the convex duality of H and L . We have, as desired, established the inequality (49) whenever $u - v$ has a local maximum at (x_0, t_0) .

3. Now suppose instead $u - v$ has a local minimum at a point $(x_0, t_0) \in \mathbb{R}^n \times (0, T)$. We must prove

$$(51) \quad v_t(x_0, t_0) + H(Dv(x_0, t_0)) \geq 0.$$

Suppose to the contrary that estimate (51) fails, in which case

$$v_t(x, t) + H(Dv(x, t)) \leq -\theta < 0$$

for some $\theta > 0$ and all points (x, t) close enough to (x_0, t_0) . In view of (50)

$$(52) \quad v_t(x, t) + Dv(x, t) \cdot v - L(v) \leq -\theta$$

for all (x, t) near (x_0, t_0) and all $v \in \mathbb{R}^n$.

Now from (46) we see that if $h > 0$ is small enough,

$$(53) \quad u(x_0, t_0) = hL\left(\frac{x_0 - x_1}{h}\right) + u(x_1, t_0 - h)$$

for some point x_1 close to x_0 . We then compute

$$\begin{aligned} v(x_0, t_0) - v(x_1, t_0 - h) &= \int_0^1 \frac{d}{ds} v(sx_0 + (1-s)x_1, t_0 + (s-1)h) ds \\ &= \int_0^1 Dv(sx_0 + (1-s)x_1, t_0 + (s-1)h) \cdot (x_0 - x_1) \\ &\quad + v_t(sx_0 + (1-s)x_1, t_0 + (s-1)h) h ds \\ &= h \int_0^1 Dv(\dots) \cdot \left(\frac{x_0 - x_1}{h}\right) + v_t(\dots) ds. \end{aligned}$$

Now if $h > 0$ is sufficiently small, we may apply (52), to find

$$v(x_0, t_0) - v(x_1, t_0 - h) \leq hL\left(\frac{x_0 - x_1}{h}\right) - \theta h.$$

But then (53) forces

$$v(x_0, t_0) - v(x_1, t_0 - h) \leq u(x_0, t_0) - u(x_1, t_0 - h) - \theta h,$$

a contradiction, since $u - v$ has a local minimum at (x_0, t_0) . Consequently the desired inequality (51) is indeed valid. \square

10.4. PROBLEMS

1. Assume
- u
- is a viscosity solution of

$$u_t + H(Du, x) = 0 \quad \text{in } \mathbb{R}^n \times (0, \infty).$$

Show that $\hat{u} := -u$ is a viscosity solution of

$$\hat{u}_t + \hat{H}(D\hat{u}, x) = 0 \quad \text{in } \mathbb{R}^n \times (0, \infty),$$

for $\hat{H}(p, x) := -H(-p, x)$.

2. Let
- $\{u^k\}_{k=1}^\infty$
- be viscosity solutions of the Hamilton–Jacobi equations

$$u_t^k + H(Du^k, x) = 0 \quad \text{in } \mathbb{R}^n \times (0, \infty)$$

($k = 1, \dots$), and suppose $u^k \rightarrow u$ uniformly. Assume as well that H is continuous. Show u is a viscosity solution of

$$u_t + H(Du, x) = 0 \quad \text{in } \mathbb{R}^n \times (0, \infty).$$

Hence the uniform limits of viscosity solutions are viscosity solutions.

3. Suppose for each
- $\epsilon > 0$
- that
- u^ϵ
- is a smooth solution of the parabolic equation

$$u_t^\epsilon + H(Du^\epsilon, x) - \epsilon \sum_{i,j=1}^n a^{ij} u_{x_i x_j}^\epsilon = 0$$

in $\mathbb{R}^n \times (0, \infty)$, where the smooth coefficients a^{ij} ($i, j = 1, \dots, n$) satisfy the uniform ellipticity condition from Chapter 6. Suppose also that H is continuous and that $u^\epsilon \rightarrow u$ uniformly as $\epsilon \rightarrow 0$.

Prove that u is a viscosity solution of $u_t + H(Du, x) = 0$. (This exercise shows that viscosity solutions do not depend upon the precise structure of the parabolic smoothing.)

4. Let
- u^i
- (
- $i = 1, 2$
-) be viscosity solutions of

$$\begin{cases} u_t^i + H(Du^i, x) = 0 & \text{in } \mathbb{R}^n \times (0, \infty) \\ u^i = g^i & \text{on } \mathbb{R}^n \times \{t = 0\}. \end{cases}$$

Assume H satisfies condition (3) in §10.2. Prove the *contraction* property

$$\sup_{\mathbb{R}^n} |u^1(\cdot, t) - u^2(\cdot, t)| \leq \sup_{\mathbb{R}^n} |g^1 - g^2| \quad (t \geq 0).$$

5. (a) Show that
- $u(x) := 1 - |x|$
- is a viscosity solution of

$$(*) \quad \begin{cases} |u'| = 1 & \text{in } (-1, 1) \\ u(-1) = u(1) = 0. \end{cases}$$

This means that for each $v \in C^\infty(-1, 1)$, if $u - v$ has a maximum (minimum) at a point $x_0 \in (-1, 1)$, then $|v'(x_0)| \leq 1$ (≥ 1).

(b) Show that $\tilde{u}(x) := |x| - 1$ is *not* a viscosity solution of (*).

(c) Show that \tilde{u} is a viscosity solution of

$$(**) \quad \begin{cases} -|\tilde{u}'| = -1 & \text{in } (-1, 1) \\ \tilde{u}(-1) = \tilde{u}(1) = 0. \end{cases}$$

(Hint: What is the meaning of a viscosity solution of (**)?)

(d) Why do problems (*), (**) have different viscosity solutions?

6. Let $U \subset \mathbb{R}^n$ be open, bounded. Set $u(x) := \text{dist}(x, \partial U)$ ($x \in U$). Prove that u is Lipschitz continuous and that it is a viscosity solution of the eikonal equation

$$|Du| = 1 \quad \text{in } U.$$

This means that for each $v \in C^\infty(U)$, if $u - v$ has a maximum (minimum) at a point $x_0 \in U$, then $|Dv(x_0)| \leq 1$ (≥ 1).

7. Suppose an open set $U \subset \mathbb{R}^n$ is subdivided by a smooth hypersurface Γ into the subregions V^+ and V^- . Let ν denote the unit normal to Γ , pointing into V^+ . Assume that u is a viscosity solution of

$$H(Du) = 0 \quad \text{in } U$$

and that u is smooth in \bar{V}^+ and \bar{V}^- . Write u_ν^+ for the limit of $Du \cdot \nu$ along Γ from within V^+ , and write u_ν^- for the limit from within V^- .

Prove that along Γ we have the inequalities

$$H(\lambda u_\nu^- + (1 - \lambda)u_\nu^+) \geq 0 \quad \text{if } u_\nu^- \leq u_\nu^+$$

and

$$H(\lambda u_\nu^- + (1 - \lambda)u_\nu^+) \leq 0 \quad \text{if } u_\nu^+ \leq u_\nu^-,$$

for each $0 \leq \lambda \leq 1$.

8. A surface described by the graph of $u : \mathbb{R}^2 \rightarrow \mathbb{R}$ is illuminated by parallel light rays from the vertical e_3 direction. We assume the surface has constant albedo and in addition is *Lambertian*, meaning that incoming right rays are scattered equally in all directions. Then the intensity $i = i(x)$ of the reflected light above the point $x \in \mathbb{R}^2$ is given by the formula $i = e_3 \cdot \nu$, where ν is the upward pointing unit normal to the surface.

Show that u solves a PDE of the form

$$|Du| = n$$

for a given function $n = n(x)$, computed in terms of the intensity i . (Finding the surface by solving this PDE for u is the *shape from shading* problem.)

9. A yacht starts at the point $(x_1, 0)$ on the positive x_1 -axis and sails to the right at speed $b_1 > 0$. Another yacht is initially at the point $(0, x_2)$ along the positive x_2 -axis and starts in pursuit, sailing always towards the first yacht at speed $b_2 > b_1$.

Find the PDE solved by

$$u(x_1, x_2) := \text{time it takes the second yacht to intercept the first.}$$

(Think of this as a dynamic programming problem, but with no controls.)

10. (Infinite horizon control problem) Assume \mathbf{f} and r satisfy the conditions given in §10.3. Given a point $x \in \mathbb{R}^n$ and a control belonging to $\mathcal{A} := \{\alpha : [0, \infty) \rightarrow A \mid \alpha(\cdot) \text{ is measurable}\}$, let $\mathbf{x}(\cdot)$ be the unique solution of the ODE

$$\begin{cases} \dot{\mathbf{x}}(s) = \mathbf{f}(\mathbf{x}(s), \alpha(s)) & (s > 0) \\ \mathbf{x}(0) = x. \end{cases}$$

Fix $\lambda > 0$ and define the discounted cost

$$C_x[\alpha(\cdot)] := \int_0^\infty e^{-\lambda s} r(\mathbf{x}(s), \alpha(s)) ds.$$

Define the value function

$$u(x) := \inf_{\alpha(\cdot) \in \mathcal{A}} C_x[\alpha(\cdot)].$$

- (a) Show that u is bounded and that if $\lambda > \text{Lip}[\mathbf{f}]$, then u is Lipschitz continuous.
- (b) Show that if $0 < \lambda \leq \text{Lip}[\mathbf{f}]$, then u is Hölder continuous for some exponent $0 < \alpha < 1$.
11. (Continuation) Prove that the value function u is a viscosity solution of the PDE

$$\lambda u - \min_{a \in A} \{\mathbf{f}(x, a) \cdot Du + r(x, a)\} = 0 \quad \text{in } \mathbb{R}^n.$$

(This means that if v is smooth and $u - v$ has a local maximum at a point x_0 , then

$$\lambda u - \min_{a \in A} \{f(x, a) \cdot Dv + r(x, a)\} \leq 0$$

at x_0 , and that the opposite inequality holds if $u - v$ has a local minimum at x_0 .)

The next sequence of exercises develops some of the theory of viscosity solutions for fully nonlinear elliptic PDE of second order.

12. Remember from §A.1 that if $R, S \in \mathbb{S}^n$, we write $R \geq S$ if $R - S$ is nonnegative definite. A function $F : \mathbb{S}^n \times \mathbb{R}^n \times \mathbb{R}^n \rightarrow \mathbb{R}$, $F = F(R, p, x)$, is *elliptic* provided

$$R \geq S \text{ implies } F(R, p, x) \leq F(S, p, x).$$

Here \mathbb{S}^n denotes the space of real, $n \times n$ symmetric matrices.

- (a) Show that $F(R) = -\operatorname{tr} R$ is elliptic.
 (b) More generally, show that if $A \in \mathbb{S}^n$ and $A \geq 0$, then $F(R) = -A : R = -\operatorname{tr}(AR^T)$ is elliptic.
 (c) Show that if for each $k = 1, \dots, m$, F_k is elliptic, then so are $\max_k F_k$ and $\min_k F_k$.
13. Let F be continuous and elliptic. We say that a function $u \in C(U)$ is a *viscosity solution* of the fully nonlinear elliptic PDE

$$(*) \quad F(D^2u, Du, x) = 0 \quad \text{in } U,$$

provided for each $v \in C^\infty(U)$, (i) if $u - v$ has a local maximum at a point $x_0 \in U$, then $F(D^2u(x_0), Du(x_0), x_0) \leq 0$ and (ii) if $u - v$ has a local minimum at a point $x_0 \in U$, then $F(D^2u(x_0), Du(x_0), x_0) \geq 0$. Show that if u is a C^2 solution of $(*)$, then u is a viscosity solution.

14. Assume that u_k is a viscosity solution of

$$F(D^2u_k, Du_k, x) = 0 \quad \text{in } U$$

for $k = 1, \dots$. Suppose $u_k \rightarrow u$ uniformly and show u is a viscosity solution of

$$F(D^2u, Du, x) = 0 \quad \text{in } U.$$

10.5. REFERENCES

Section 10.1 The definition of viscosity solutions presented here is due to Crandall, Evans and Lions (Trans. AMS 282 (1984), 487–502), who recast an earlier definition set forth in the basic

paper of Crandall and Lions (Trans. AMS 277 (1983), 1–42).

- Section 10.2 The uniqueness proof follows the first paper cited above, with improvements I learned from M. Crandall.
- Section 10.3 P.-L. Lions [Li] observed the connection between the definition of viscosity solution and the optimality conditions of control theory. The books of Fleming–Soner [F-S] and Bardi–Capuzzo Dolcetta [B-CD] provide much more information about viscosity solutions and the connections with deterministic and stochastic optimal control.
- Section 10.4 D. Ostrov contributed Problem 8. Consult Cabre–Caffarelli [C-C] and Gilbarg–Trudinger [G-T] for more about fully nonlinear elliptic equations.

This page is intentionally left blank.

SYSTEMS OF CONSERVATION LAWS

- 11.1 Introduction
- 11.2 Riemann's problem
- 11.3 Systems of two conservation laws
- 11.4 Entropy criteria
- 11.5 Problems
- 11.6 References

11.1. INTRODUCTION

In this chapter we study *systems* of nonlinear, divergence structure first-order hyperbolic PDE, which arise as models of *conservation laws*.

Physical interpretation. In the most general circumstance we would like to investigate a vector function

$$\mathbf{u} = \mathbf{u}(x, t) = (u^1(x, t), \dots, u^m(x, t)) \quad (x \in \mathbb{R}^n, t \geq 0),$$

the components of which are the densities of various conserved quantities in some physical system under investigation. Given then any smooth, bounded region $U \subset \mathbb{R}^n$, we note that the integral

$$(1) \quad \int_U \mathbf{u}(x, t) dx$$

represents the total amount of these quantities within U at time t . Now conservation laws typically assert that the rate of change within U is governed by a *flux* function $\mathbf{F} : \mathbb{R}^m \rightarrow \mathbb{M}^{m \times n}$, which controls the rate of loss or increase of \mathbf{u} through ∂U . Otherwise stated, it is appropriate to assume for each time t

$$(2) \quad \frac{d}{dt} \int_U \mathbf{u} \, dx = - \int_{\partial U} \mathbf{F}(\mathbf{u}) \boldsymbol{\nu} \, dS,$$

$\boldsymbol{\nu}$ denoting as usual the outward unit normal along U . Rewriting (2), we deduce

$$(3) \quad \int_U \mathbf{u}_t \, dx = - \int_{\partial U} \mathbf{F}(\mathbf{u}) \boldsymbol{\nu} \, dS = - \int_U \operatorname{div} \mathbf{F}(\mathbf{u}) \, dx.$$

As the region $U \subset \mathbb{R}^n$ was arbitrary, we derive from (3) this initial-value problem for a *general system of conservation laws*:

$$(4) \quad \begin{cases} \mathbf{u}_t + \operatorname{div} \mathbf{F}(\mathbf{u}) = 0 & \text{in } \mathbb{R}^n \times (0, \infty) \\ \mathbf{u} = \mathbf{g} & \text{on } \mathbb{R}^n \times \{t = 0\}, \end{cases}$$

the given function $\mathbf{g} = (g^1, \dots, g^m)$ describing the initial distribution of $\mathbf{u} = (u^1, \dots, u^m)$.

At present a good mathematical understanding of problem (4) is largely unavailable (but see Zheng [Zh]). For this reason we shall henceforth consider instead the initial-value problem for a *system of conservation laws in one space dimension*:

$$(5) \quad \begin{cases} \mathbf{u}_t + \mathbf{F}(\mathbf{u})_x = 0 & \text{in } \mathbb{R} \times (0, \infty) \\ \mathbf{u} = \mathbf{g} & \text{on } \mathbb{R} \times \{t = 0\}, \end{cases}$$

where $\mathbf{F} : \mathbb{R}^m \rightarrow \mathbb{R}^m$ and $\mathbf{g} : \mathbb{R} \rightarrow \mathbb{R}^m$ are given and $\mathbf{u} : \mathbb{R} \times [0, \infty) \rightarrow \mathbb{R}^m$ is the unknown, $\mathbf{u} = \mathbf{u}(x, t)$. We call \mathbb{R}^m the *state space* and write

$$\mathbf{F} = \mathbf{F}(z) = (F^1(z), \dots, F^m(z)) \quad (z \in \mathbb{R}^m)$$

for the smooth flux function.

We intend to study the solvability of problem (5), properties of its solutions, etc.

Example 1. The *p-system* is this collection of two conservation laws:

$$(6) \quad \begin{cases} u_t^1 - u_x^2 = 0 & \text{(compatibility condition)} \\ u_t^2 - p(u^1)_x = 0 & \text{(Newton's law)} \end{cases}$$

in $\mathbb{R} \times (0, \infty)$, where $p : \mathbb{R} \rightarrow \mathbb{R}$ is given. Here

$$(7) \quad \mathbf{F}(z) = (-z_2, -p(z_1))$$

for $z = (z_1, z_2)$. The p -system arises as a rewritten form of the scalar *quasilinear wave equation*

$$(8) \quad u_{tt} - (p(u_x))_x = 0 \quad \text{in } \mathbb{R} \times (0, \infty).$$

Taking $u^1 := u_x$, $u^2 := u_t$, we obtain the system (6), with the stated interpretations. \square

Example 2. *Euler's equations* for compressible gas flow in one dimension are

$$(9) \quad \begin{cases} \rho_t + (\rho v)_x = 0 & \text{(conservation of mass)} \\ (\rho v)_t + (\rho v^2 + p)_x = 0 & \text{(conservation of momentum)} \\ (\rho E)_t + (\rho E v + p v)_x = 0 & \text{(conservation of energy)} \end{cases}$$

in $\mathbb{R} \times (0, \infty)$. Here ρ is the *mass density*, v the *velocity*, and E the *energy density* per unit mass. We assume

$$E = e + \frac{v^2}{2},$$

where e is the *internal energy* per unit mass and the term $\frac{v^2}{2}$ corresponds to the *kinetic energy* per unit mass. The letter p in (9) denotes the *pressure*. We assume p is a known function

$$(10) \quad p = p(\rho, e)$$

of ρ and e ; formula (10) is a *constitutive relation*. Writing $\mathbf{u} = (u^1, u^2, u^3) = (\rho, \rho v, \rho E)$, we check that (9) is a system of conservation laws of the requisite form

$$\mathbf{u}_t + \mathbf{F}(\mathbf{u})_x = 0 \quad \text{in } \mathbb{R} \times (0, \infty)$$

for $\mathbf{F} = (F^1, F^2, F^3)$,

$$(11) \quad \begin{cases} F^1(z) = z_2 \\ F^2(z) = \frac{(z_2)^2}{z_1} + p(z_1, \frac{z_3}{z_1} - \frac{1}{2}(\frac{z_2}{z_1})^2) \\ F^3(z) = \frac{z_2 z_3}{z_1} + p(z_1, \frac{z_3}{z_1} - \frac{1}{2}(\frac{z_2}{z_1})^2) \frac{z_2}{z_1}, \end{cases}$$

where $z = (z_1, z_2, z_3)$, $z_1 > 0$. \square

Example 3. The one-dimensional *shallow water equations* are

$$\begin{cases} h_t + (vh)_x = 0 & \text{(conservation of mass)} \\ (vh)_t + \left(v^2h + \frac{h^2}{2}\right)_x = 0 & \text{(conservation of momentum)} \end{cases}$$

in $\mathbb{R} \times [0, \infty)$, where v is the horizontal velocity and $h > 0$ is the height. Putting $q := vh$, we can rewrite the equations into standard conservation law form:

$$\begin{cases} h_t + q_x = 0 \\ q_t + \left(\frac{q^2}{h} + \frac{h^2}{2}\right)_x = 0. \end{cases}$$

Here

$$\mathbf{F}(z) = \left(z_2, \frac{z_2^2}{z_1} + \frac{z_1^2}{2}\right)$$

for $z = (z_1, z_2)$, $z_1 > 0$. □

11.1.1. Integral solutions.

The great difficulty in this subject is discovering a proper notion of weak solution for the initial-value problem (5). We have already encountered similar issues in §3.4 in our study of the much simpler case of a single or *scalar* conservation law (i.e., $m = 1$ above).

Following then the development in §3.4.1, let us suppose

$$(12) \quad \begin{cases} \mathbf{v} : \mathbb{R} \times [0, \infty) \rightarrow \mathbb{R}^m \text{ is smooth,} \\ \text{with compact support, } \mathbf{v} = (v^1, \dots, v^m). \end{cases}$$


We temporarily assume \mathbf{u} is a smooth solution of our problem (5), take the dot product of the PDE $\mathbf{u}_t + \mathbf{F}(\mathbf{u})_x = 0$ with the test function \mathbf{v} , and integrate by parts, to obtain the equality

$$(13) \quad \int_0^\infty \int_{-\infty}^\infty \mathbf{u} \cdot \mathbf{v}_t + \mathbf{F}(\mathbf{u}) \cdot \mathbf{v}_x \, dxdt + \int_{-\infty}^\infty \mathbf{g} \cdot \mathbf{v} \, dx|_{t=0} = 0.$$

This identity, which we derived supposing \mathbf{u} to be a smooth solution, makes sense if \mathbf{u} is merely bounded.

DEFINITION. We say that $\mathbf{u} \in L^\infty(\mathbb{R} \times (0, \infty); \mathbb{R}^m)$ is an integral solution of the initial-value problem (5) provided the equality (13) holds for all test functions \mathbf{v} satisfying (12).

Continuing now to parallel the development in §3.4.1 for a single conservation law, let us now consider the situation that we have an integral solution \mathbf{u} of (5) which is smooth on either side of a curve C , along which \mathbf{u} has simple jump discontinuities. More precisely, let us assume that $V \subset \mathbb{R} \times (0, \infty)$



Rankine-Hugoniot condition

is some region cut by a smooth curve C into a left-hand part V_l and a right-hand part V_r .

Assuming that \mathbf{u} is smooth in V_l , we select the test function \mathbf{v} with compact support in V_l and deduce from (13) that

$$(14) \quad \mathbf{u}_t + \mathbf{F}(\mathbf{u})_x = 0 \quad \text{in } V_l.$$

Similarly, we have

$$(15) \quad \mathbf{u}_t + \mathbf{F}(\mathbf{u})_x = 0 \quad \text{in } V_r,$$

provided \mathbf{u} is smooth in V_r . Now choose a test function \mathbf{v} with compact support in V but which does not necessarily vanish along the curve C . Then utilizing the identity (13), we find

$$(16) \quad \begin{aligned} 0 &= \int_0^\infty \int_{-\infty}^\infty \mathbf{u} \cdot \mathbf{v}_t + \mathbf{F}(\mathbf{u}) \cdot \mathbf{v}_x \, dx dt \\ &= \iint_{V_l} \mathbf{u} \cdot \mathbf{v}_t + \mathbf{F}(\mathbf{u}) \cdot \mathbf{v}_x \, dx dt + \iint_{V_r} \mathbf{u} \cdot \mathbf{v}_t + \mathbf{F}(\mathbf{u}) \cdot \mathbf{v}_x \, dx dt. \end{aligned}$$

As \mathbf{v} has compact support within V , we deduce

$$\begin{aligned}
 \iint_{V_l} \mathbf{u} \cdot \mathbf{v}_t + \mathbf{F}(\mathbf{u}) \cdot \mathbf{v}_x \, dxdt &= - \iint_{V_l} [\mathbf{u}_t + \mathbf{F}(\mathbf{u})_x] \cdot \mathbf{v} \, dxdt \\
 (17) \qquad \qquad \qquad &+ \int_C (\mathbf{u}_l \nu^2 + \mathbf{F}(\mathbf{u}_l) \nu^1) \cdot \mathbf{v} \, dl \\
 &= \int_C (\mathbf{u}_l \nu^2 + \mathbf{F}(\mathbf{u}_l) \nu^1) \cdot \mathbf{v} \, dl,
 \end{aligned}$$

owing to (14). Here $\boldsymbol{\nu} = (\nu^1, \nu^2)$ is the unit normal to the curve C pointing from V_l into V_r , and the subscript “ l ” denotes the limit from the left. Likewise (16) implies

$$\iint_{V_r} \mathbf{u} \cdot \mathbf{v}_t + \mathbf{F}(\mathbf{u}) \cdot \mathbf{v}_x \, dxdt = - \int_C (\mathbf{u}_r \nu^2 + \mathbf{F}(\mathbf{u}_r) \nu^1) \cdot \mathbf{v} \, dl,$$

“ r ” denoting the limit from the right. Adding this identity to (17) and remembering (16), we deduce

$$\int_C [(\mathbf{F}(\mathbf{u}_l) - \mathbf{F}(\mathbf{u}_r)) \nu^1 + (\mathbf{u}_l - \mathbf{u}_r) \nu^2] \cdot \mathbf{v} \, dl = 0.$$

This identity obtains for all smooth functions \mathbf{v} as above, whence

$$(18) \qquad (\mathbf{F}(\mathbf{u}_l) - \mathbf{F}(\mathbf{u}_r)) \nu^1 + (\mathbf{u}_l - \mathbf{u}_r) \nu^2 = 0 \quad \text{along } C.$$

Suppose now the curve C is represented parametrically as $\{(x, t) \mid x = s(t)\}$ for some smooth function $s(\cdot) : [0, \infty) \rightarrow \mathbb{R}$. Then $\boldsymbol{\nu} = (\nu^1, \nu^2) = (1 + \dot{s}^2)^{-1/2}(1, -\dot{s})$. Consequently (18) reads

$$(19) \qquad \mathbf{F}(\mathbf{u}_l) - \mathbf{F}(\mathbf{u}_r) = \dot{s}(\mathbf{u}_l - \mathbf{u}_r)$$

in V along the curve C .

NOTATION.

$$\begin{cases}
 [[\mathbf{u}]] = \mathbf{u}_l - \mathbf{u}_r = \text{jump in } \mathbf{u} \text{ across the curve } C \\
 [[\mathbf{F}(\mathbf{u})]] = \mathbf{F}(\mathbf{u}_l) - \mathbf{F}(\mathbf{u}_r) = \text{jump in } \mathbf{F}(\mathbf{u}) \\
 \sigma = \dot{s} = \text{speed of the curve } C.
 \end{cases}$$

Let us then rewrite (19) as the identity

$$(20) \qquad [[\mathbf{F}(\mathbf{u})]] = \sigma [[\mathbf{u}]]$$

along the discontinuity curve, the *Rankine–Hugoniot jump condition*. Note carefully this is a vector equality.

11.1.2. Traveling waves, hyperbolic systems.

We have seen in §3.4 that the notion of an integral solution for conservation laws is not adequate: such solutions need not be unique. We are therefore intent upon discovering some additional requirements for a good definition of a generalized solution. This will presumably entail as in §3.4 an entropy criterion based upon an analysis of shock waves. This expectation, now as carried over to systems, is largely correct, but first of all we must study more carefully the nonlinearity \mathbf{F} in the hopes of discovering mathematically appropriate and physically correct structural conditions to impose.

Let us start by first considering the wider class of *semilinear systems* having the nondivergence form:

$$(21) \quad \mathbf{u}_t + \mathbf{B}(\mathbf{u})\mathbf{u}_x = 0 \quad \text{in } \mathbb{R} \times (0, \infty),$$

where $\mathbf{B} : \mathbb{R}^m \rightarrow \mathbb{M}^{m \times m}$. This system is for smooth functions equivalent to the conservation law in (5), provided

$$\mathbf{B} = D\mathbf{F} = \begin{pmatrix} F_{z_1}^1 & \cdots & F_{z_m}^1 \\ \vdots & \ddots & \vdots \\ F_{z_1}^m & \cdots & F_{z_m}^m \end{pmatrix}_{m \times m}.$$

We consider now the possibility of finding particular solutions which have the form of a traveling wave:

$$(22) \quad \mathbf{u}(x, t) = \mathbf{v}(x - \sigma t) \quad (x \in \mathbb{R}, t > 0),$$

where the profile $\mathbf{v} : \mathbb{R} \rightarrow \mathbb{R}^m$ and the velocity $\sigma \in \mathbb{R}$ are to be found. We substitute expression (22) into the PDE (21) and thereby obtain the equality

$$(23) \quad -\sigma \mathbf{v}'(x - \sigma t) + \mathbf{B}(\mathbf{v}(x - \sigma t))\mathbf{v}'(x - \sigma t) = 0.$$

Observe that (23) says σ is an eigenvalue of the matrix $\mathbf{B}(\mathbf{v})$ corresponding to the eigenvector \mathbf{v}' .

This conclusion suggests (exactly as for the linear theory in §7.3) that if we wish to find traveling waves or, more generally, wavelike solutions of our system of PDE, we should make some sort of hyperbolicity hypothesis concerning the eigenvalues of \mathbf{B} .

DEFINITION. *If for each $z \in \mathbb{R}^m$ the eigenvalues of $\mathbf{B}(z)$ are real and distinct, we call the system (21) strictly hyperbolic.*

We henceforth assume the system of partial differential equations (21) (and the special case $\mathbf{B} = D\mathbf{F}$ of conservation laws) to be strictly hyperbolic.

NOTATION. (i) We will write

$$(24) \quad \lambda_1(z) < \lambda_2(z) < \cdots < \lambda_m(z) \quad (z \in \mathbb{R}^m)$$

to denote the real and distinct eigenvalues of $\mathbf{B}(z)$, in increasing order.

(ii) Then for each $k = 1, \dots, m$, we let

$$\mathbf{r}_k(z)$$

denote a corresponding nonzero eigenvector, so that

$$(25) \quad \mathbf{B}(z)\mathbf{r}_k(z) = \lambda_k(z)\mathbf{r}_k(z) \quad (k = 1, \dots, m, z \in \mathbb{R}^m).$$

Since we are always assuming the strict hyperbolicity condition, the vectors $\{\mathbf{r}_k(z)\}_{k=1}^m$ span \mathbb{R}^m for each $z \in \mathbb{R}^m$.

(iii) Next, since a matrix and its transpose have the same spectrum, we can introduce for each $k = 1, \dots, m$ a nonzero eigenvector

$$\mathbf{l}_k(z)$$

for the matrix $\mathbf{B}(z)^T$, corresponding to the eigenvalue $\lambda_k(z)$. Thus

$$(26) \quad \mathbf{B}(z)^T\mathbf{l}_k(z) = \lambda_k(z)\mathbf{l}_k(z) \quad (k = 1, \dots, m, z \in \mathbb{R}^m).$$

This equality is usually written

$$(27) \quad \mathbf{l}_k(z)\mathbf{B}(z) = \lambda_k(z)\mathbf{l}_k(z) \quad (k = 1, \dots, m, z \in \mathbb{R}^m).$$

Thus $\{\mathbf{l}_k(z)\}_{k=1}^m$ can be regarded as *left eigenvectors* of $\mathbf{B}(z)$, and $\{\mathbf{r}_k(z)\}_{k=1}^m$ are *right eigenvectors*.

Remark. Additionally, we observe

$$(28) \quad \mathbf{l}_l(z) \cdot \mathbf{r}_k(z) = 0 \quad \text{if } k \neq l \quad (z \in \mathbb{R}^m).$$

To confirm this, we compute using (25) and (26) that

$$\begin{aligned} \lambda_k(z)(\mathbf{l}_l(z) \cdot \mathbf{r}_k(z)) &= \mathbf{l}_l(z) \cdot (\mathbf{B}(z)\mathbf{r}_k(z)) = (\mathbf{B}(z)^T\mathbf{l}_l(z)) \cdot \mathbf{r}_k(z) \\ &= \lambda_l(z)(\mathbf{l}_l(z) \cdot \mathbf{r}_k(z)), \end{aligned}$$

whence (28) follows since $\lambda_k(z) \neq \lambda_l(z)$ if $k \neq l$.

Let us first show that the notion of strict hyperbolicity is independent of coordinates.

THEOREM 1 (Invariance of hyperbolicity under change of coordinates).
 Let \mathbf{u} be a smooth solution of the strictly hyperbolic system (21). Assume also $\Phi : \mathbb{R}^m \rightarrow \mathbb{R}^m$ is a smooth diffeomorphism, with inverse Ψ . Then

$$(29) \quad \tilde{\mathbf{u}} := \Phi(\mathbf{u})$$

solves the strictly hyperbolic system

$$(30) \quad \tilde{\mathbf{u}}_t + \tilde{\mathbf{B}}(\tilde{\mathbf{u}})\tilde{\mathbf{u}}_x = 0 \quad \text{in } \mathbb{R} \times (0, \infty),$$

for

$$(31) \quad \tilde{\mathbf{B}}(\tilde{z}) := D\Phi(\Psi(\tilde{z}))\mathbf{B}(\Psi(\tilde{z}))D\Psi(\tilde{z}) \quad (\tilde{z} \in \mathbb{R}^m).$$

Proof. 1. We compute $\tilde{\mathbf{u}}_t = D\Phi(\mathbf{u})\mathbf{u}_t$, $\tilde{\mathbf{u}}_x = D\Phi(\mathbf{u})\mathbf{u}_x$, and so equation (30) is valid for $\mathbf{B}(\tilde{z}) = D\Phi(z)\mathbf{B}(z)D\Phi^{-1}(z)$, where $\tilde{z} = \Phi(z)$. Substituting $z = \Psi(\tilde{z})$, we obtain (31).

2. We must prove that the system (30) is strictly hyperbolic. If $\lambda_k(z)$ is an eigenvalue of $\mathbf{B}(z)$, with corresponding right eigenvector $\mathbf{r}_k(z)$, we have

$$\mathbf{B}(z)\mathbf{r}_k(z) = \lambda_k(z)\mathbf{r}_k(z).$$

Setting

$$(32) \quad \tilde{\mathbf{r}}_k(\tilde{z}) := D\Phi(\Psi(\tilde{z}))\mathbf{r}_k(\Psi(\tilde{z})),$$

$$(33) \quad \tilde{\lambda}_k(\tilde{z}) := \lambda_k(\Psi(\tilde{z})),$$

we compute

$$(34) \quad \tilde{\mathbf{B}}(\tilde{z})\tilde{\mathbf{r}}_k(\tilde{z}) = \tilde{\lambda}_k(\tilde{z})\tilde{\mathbf{r}}_k(\tilde{z}).$$

Similarly if $\mathbf{l}_k(z)$ is a left eigenvector, we write

$$(35) \quad \tilde{\mathbf{l}}_k(\tilde{z}) := \mathbf{l}_k(\Psi(\tilde{z}))D\Psi(\tilde{z})$$

and calculate

$$(36) \quad \tilde{\mathbf{l}}_k(\tilde{z})\tilde{\mathbf{B}}(\tilde{z}) = \tilde{\lambda}_k(\tilde{z})\tilde{\mathbf{l}}_k(\tilde{z}).$$

In view of (32)–(36), we conclude that the system (30) is strictly hyperbolic. \square

Next we study how $\lambda_k(z)$, $\mathbf{r}_k(z)$ and $\mathbf{l}_k(z)$ change as z varies:

THEOREM 2 (Dependence of eigenvalues and eigenvectors on parameters). *Assume the matrix function \mathbf{B} is smooth, strictly hyperbolic.*

(i) *Then the eigenvalues $\lambda_k(z)$ depend smoothly on $z \in \mathbb{R}^m$ ($k = 1, \dots, m$).*

(ii) *Furthermore, we can select the right eigenvectors $\mathbf{r}_k(z)$ and left eigenvectors $\mathbf{l}_k(z)$ to depend smoothly on $z \in \mathbb{R}^m$ and satisfy the normalization*

$$|\mathbf{r}_k(z)|, |\mathbf{l}_k(z)| = 1 \quad (k = 1, \dots, m).$$

Proof. 1. Since $\mathbf{B}(z)$ is strictly hyperbolic, for each $z_0 \in \mathbb{R}^m$ we have

$$(37) \quad \lambda_1(z_0) < \lambda_2(z_0) < \dots < \lambda_m(z_0).$$

Fix $k \in \{1, \dots, m\}$ and any point $z_0 \in \mathbb{R}^m$, and let $\mathbf{r}_k(z_0)$ satisfy

$$\begin{cases} \mathbf{B}(z_0)\mathbf{r}_k(z_0) = \lambda_k(z_0)\mathbf{r}_k(z_0) \\ |\mathbf{r}_k(z_0)| = 1. \end{cases}$$

Upon rotating coordinates if necessary, we may assume

$$(38) \quad \mathbf{r}_k(z_0) = e_m = (0, \dots, 1).$$

We first show that near z_0 , there exist smooth functions $\lambda_k(z), \mathbf{r}_k(z)$ such that

$$\begin{cases} \mathbf{B}(z)\mathbf{r}_k(z) = \lambda_k(z)\mathbf{r}_k(z) \\ |\mathbf{r}_k(z)| = 1. \end{cases}$$

2. We will apply the Implicit Function Theorem (§C.7) to the smooth function $\Phi : \mathbb{R}^m \times \mathbb{R} \times \mathbb{R}^m \rightarrow \mathbb{R}^{m+1}$ defined by

$$\Phi(r, \lambda, z) = (\mathbf{B}(z)r - \lambda r, |r|^2) \quad (r, z \in \mathbb{R}^m, \lambda \in \mathbb{R}).$$

Now

$$\frac{\partial \Phi(r, \lambda, z)}{\partial (r, \lambda)} = \begin{pmatrix} & -r_1 & \\ \mathbf{B}(z) - \lambda I & \vdots & \\ & -r_m & \\ 2r_1 \dots 2r_m & 0 & \end{pmatrix}_{(m+1) \times (m+1)},$$

and so, according to (38), it suffices to check that

$$(39) \quad \det \begin{pmatrix} & 0 & \\ \mathbf{B}(z_0) - \lambda_k(z_0)I & \vdots & \\ & -1 & \\ 0 \dots \dots 2 & 0 & \end{pmatrix} \neq 0.$$

3. Note that for $\epsilon > 0$ sufficiently small, the matrix

$$(40) \quad \mathbf{B}_\epsilon = \mathbf{B}(z_0) - (\lambda_k(z_0) + \epsilon)I$$

is invertible. In light of (38),

$$\mathbf{B}_\epsilon e_m = -\epsilon e_m.$$

Therefore

$$\begin{pmatrix} 0 \\ \vdots \\ \mathbf{B}_\epsilon \\ -1 \\ 0 \dots 2 \end{pmatrix} \begin{pmatrix} 0 \\ \vdots \\ I \\ (-\epsilon)^{-1} \\ 0 \dots 0 \end{pmatrix} = \begin{pmatrix} 0 \\ \vdots \\ \mathbf{B}_\epsilon \\ 0 \\ 0 \dots 2 \end{pmatrix} \begin{pmatrix} 0 \\ \vdots \\ 0 \\ 2(-\epsilon)^{-1} \\ 0 \dots 0 \end{pmatrix}.$$

Consequently, since the determinant of the second matrix before the equals sign is one, we have

$$\begin{aligned} \det \begin{pmatrix} 0 \\ \vdots \\ \mathbf{B}_\epsilon \\ -1 \\ 0 \dots 2 \end{pmatrix} &= 2(\det \mathbf{B}_\epsilon)(-\epsilon)^{-1} \\ &= 2 \prod_{j \neq k} (\lambda_j(z_0) - (\lambda_k(z_0) + \epsilon))(-\epsilon)(-\epsilon)^{-1} \\ &\rightarrow 2 \prod_{j \neq k} (\lambda_j(z_0) - \lambda_k(z_0)) \quad \text{as } \epsilon \rightarrow 0. \end{aligned}$$

As $B(z_0)$ is strictly hyperbolic, the last expression is nonzero. Condition (39) is verified. We may thus invoke the Implicit Function Theorem (§C.7) to find near z_0 smooth functions $\lambda_k(z)$ and $\mathbf{r}_k(z)$, satisfying the conclusion of the theorem.

4. It remains to show that we can define $\lambda_k(z)$ and $\mathbf{r}_k(z)$ for all $z \in \mathbb{R}^m$ and not just near any particular point z_0 . To do so, let us write

$$R := \sup\{r > 0 \mid \lambda_k(z), \mathbf{r}_k(z) \text{ as above exist and are smooth on } B(0, r)\}.$$

If $R = \infty$, we are done. Otherwise, we cover $\partial B(0, R)$ with finitely many open balls into which we can smoothly extend $\lambda_k(\cdot)$ and $\mathbf{r}_k(\cdot)$, using steps 1–3 above. This yields a contradiction to the definition of R .

A similar proof works for the left eigenvectors. □

Observe that we are not only globally and smoothly defining the eigenvalues and eigenspaces of \mathbf{B} , but are also globally providing the eigenspaces with an orientation.

This proof depends fundamentally upon the one-dimensionality of the eigenspaces. See Problem 3 for an example of what could go wrong in the hyperbolic, but not strictly hyperbolic, setting.

Example 1 (continued). For the p -system (6), we have

$$\mathbf{u}_t + \mathbf{B}(\mathbf{u})\mathbf{u}_x = 0,$$

for

$$\mathbf{B}(z) = D\mathbf{F}(z) = \begin{pmatrix} 0 & -1 \\ -p'(z_1) & 0 \end{pmatrix}.$$

The eigenvalues are $\lambda_1 = -\sigma$, $\lambda_2 = \sigma$, for $\sigma := p'(z_1)^{1/2}$. These are real and distinct provided we hereafter suppose the strict hyperbolicity condition

$$(41) \quad p' > 0.$$

For the nonlinear wave equation (8) this is the physical assumption that the *stress* $p(u_x)$ is a strictly increasing function of the *strain* u_x . \square

Example 2 (continued). Euler's equations (9) comprise a strictly hyperbolic system provided we assume $p > 0$ and

$$(42) \quad \frac{\partial p}{\partial \rho} > 0, \quad \frac{\partial p}{\partial e} > 0,$$

where $p = p(\rho, e)$ is the constitutive relation between the mass density, the internal energy density and the pressure. This assertion is however difficult to verify directly, as the flux function \mathbf{F} defined by (11) is complicated.

Let us rather change variables and regard the density ρ , velocity v and internal energy e as the unknowns. We can then rewrite Euler's equations (9) in terms of these quantities and, in so doing, obtain after some calculations the system

$$(43) \quad \begin{cases} \rho_t + v\rho_x + \rho v_x = 0 \\ v_t + vv_x + \frac{1}{\rho}p_x = 0 \\ e_t + ve_x + \frac{p}{\rho}v_x = 0, \end{cases}$$

provided $\rho > 0$. These equations are not in conservation form. Setting now $\mathbf{u} = (u^1, u^2, u^3) = (\rho, v, e)$, we rewrite (43) as

$$(44) \quad \mathbf{u}_t + \mathbf{B}(\mathbf{u})\mathbf{u}_x = 0 \quad \text{in } \mathbb{R} \times (0, \infty),$$

for

$$(45) \quad \mathbf{B}(z) = z_2 I + \hat{\mathbf{B}}(z),$$

where

$$\hat{\mathbf{B}}(z) := \begin{pmatrix} 0 & z_1 & 0 \\ \frac{1}{z_1} \frac{\partial p}{\partial \rho}(z_1, z_3) & 0 & \frac{1}{z_1} \frac{\partial p}{\partial e}(z_1, z_3) \\ 0 & \frac{1}{z_1} p(z_1, z_3) & 0 \end{pmatrix}.$$

The characteristic polynomial of $\hat{\mathbf{B}}$ is $-\lambda(\lambda^2 - \sigma^2)$, for $\sigma^2 = \frac{p}{z_1^2} \frac{\partial p}{\partial e} + \frac{\partial p}{\partial \rho}$. Recalling (45) and reverting to physical notation, we see that the eigenvalues of \mathbf{B} are

$$(46) \quad \lambda_1 = v - \sigma, \quad \lambda_2 = v, \quad \lambda_3 = v + \sigma,$$

where

$$\sigma := \left(\frac{p}{\rho^2} \frac{\partial p}{\partial e} + \frac{\partial p}{\partial \rho} \right)^{1/2} > 0$$

is the local *sound speed*. We therefore see that the system (44) is strictly hyperbolic, provided assumption (42) is valid. Remembering now Theorem 1, we deduce that Euler's equations (9) are also strictly hyperbolic, with eigenvalues given by (46). \square

11.2. RIEMANN'S PROBLEM

In this section we investigate in detail the system of conservation laws

$$(1) \quad \mathbf{u}_t + \mathbf{F}(\mathbf{u})_x = 0 \quad \text{in } \mathbb{R} \times (0, \infty),$$

with the piecewise-constant initial data

$$(2) \quad \mathbf{g} = \begin{cases} u_l & \text{if } x < 0 \\ u_r & \text{if } x > 0. \end{cases}$$

This is *Riemann's problem*. We call the given vectors u_l and u_r the left and right *initial states*.

11.2.1. Simple waves.

We commence our study of (1), (2) in very much the same spirit as in §11.1.2, in that we look for solutions of (1) having a special form. Before we searched for traveling waves, that is, solutions of the type $\mathbf{u}(x, t) = \mathbf{v}(x - \sigma t)$. We now seek *simple waves*. These are solutions of (1) having the structure

$$(3) \quad \mathbf{u}(x, t) = \mathbf{v}(w(x, t)) \quad (x \in \mathbb{R}, t > 0),$$

where $\mathbf{v} : \mathbb{R} \rightarrow \mathbb{R}^m$, $\mathbf{v} = (v^1, \dots, v^m)$, and $w : \mathbb{R} \times [0, \infty) \rightarrow \mathbb{R}$ are to be found. To discover the requisite properties of \mathbf{v} and w , let us substitute (3) into (1) and obtain the equality

$$(4) \quad \dot{\mathbf{v}}(w)w_t + D\mathbf{F}(\mathbf{v}(w))\dot{\mathbf{v}}(w)w_x = 0.$$

Now in view of equation (25) from §11.1.2, with $\mathbf{B} = D\mathbf{F}$, we see (4) will be valid if for some $k \in \{1, \dots, m\}$, w solves the PDE

$$(5) \quad w_t + \lambda_k(\mathbf{v}(w))w_x = 0$$

and \mathbf{v} solves the ODE

$$(6) \quad \dot{\mathbf{v}}(s) = \mathbf{r}_k(\mathbf{v}(s)) \quad \left(\cdot = \frac{d}{ds} \right).$$


If (5) and (6) hold, we call the function \mathbf{u} defined by (3) a *k-simple wave*. The point of all this is that we can regard (6) as an ODE for the vector function \mathbf{v} , and then—once \mathbf{v} has been found by solving (6)—we can interpret equation (5) as a *scalar* conservation law for w .

Let us next identify circumstances under which we can employ the construction (3)–(6) to build a continuous solution \mathbf{u} of (1). We must examine first the ODE (6).

DEFINITION. Given a fixed state $z_0 \in \mathbb{R}^m$, we define the k^{th} -rarefaction curve

$$R_k(z_0)$$

to be the path in \mathbb{R}^m of the solution of the ODE (6) which passes through z_0 .



Rarefaction curve

Given then the solution \mathbf{v} of (6), we turn to the PDE (5), which we rewrite as the scalar conservation law

$$(7) \quad w_t + F_k(w)_x = 0$$

for

$$(8) \quad F_k(s) := \int_0^s \lambda_k(\mathbf{v}(t)) dt \quad (s \in \mathbb{R}).$$

The PDE (7) will fall under the general theory developed in §3.4 provided F_k is strictly convex (or else strictly concave). Let us therefore compute

$$(9) \quad F'_k(s) = \lambda_k(\mathbf{v}(s)),$$

$$(10) \quad F_k''(s) = D\lambda_k(\mathbf{v}(s)) \cdot \dot{\mathbf{v}}(s) = D\lambda_k(\mathbf{v}(s)) \cdot \mathbf{r}_k(\mathbf{v}(s)).$$

Owing to (10), the function F_k will be convex if

$$D\lambda_k(z) \cdot \mathbf{r}_k(z) > 0 \quad (z \in \mathbb{R}^m)$$

and concave if

$$D\lambda_k(z) \cdot \mathbf{r}_k(z) < 0 \quad (z \in \mathbb{R}^m).$$

The function F_k is linear provided

$$D\lambda_k(z) \cdot \mathbf{r}_k(z) \equiv 0 \quad (z \in \mathbb{R}^m).$$

These possibilities motivate the following

DEFINITIONS. (i) *The pair $(\lambda_k(z), \mathbf{r}_k(z))$ is called genuinely nonlinear provided*

$$(11) \quad D\lambda_k(z) \cdot \mathbf{r}_k(z) \neq 0 \quad \text{for all } z \in \mathbb{R}^m.$$

(ii) *We say $(\lambda_k(z), \mathbf{r}_k(z))$ is linearly degenerate if*

$$(12) \quad D\lambda_k(z) \cdot \mathbf{r}_k(z) = 0 \quad \text{for all } z \in \mathbb{R}^m.$$

NOTATION. If the pair $(\lambda_k, \mathbf{r}_k)$ is genuinely nonlinear, write


$$R_k^+(z_0) := \{z \in R_k(z_0) \mid \lambda_k(z) > \lambda_k(z_0)\}$$

and

$$R_k^-(z_0) := \{z \in R_k(z_0) \mid \lambda_k(z) < \lambda_k(z_0)\}.$$

Then

$$R_k(z_0) = R_k^+(z_0) \cup \{z_0\} \cup R_k^-(z_0).$$



Two parts of the rarefaction curve

11.2.2. Rarefaction waves.


We turn our attention again to Riemann's problem (1), (2).

THEOREM 1 (Existence of k -rarefaction waves). *Suppose that for some $k \in \{1, \dots, m\}$,*

- (i) *the pair $(\lambda_k, \mathbf{r}_k)$ is genuinely nonlinear, and*
- (ii) *$u_r \in R_k^+(u_l)$.*

Then there exists a continuous integral solution \mathbf{u} of Riemann's problem (1), (2), which is a k -simple wave constant along lines through the origin.

We call \mathbf{u} a (centered) k -rarefaction wave.



k -rarefaction wave

Proof. We first choose w_l and $w_r \in \mathbb{R}$ so that $u_l = \mathbf{v}(w_l)$, $u_r = \mathbf{v}(w_r)$. Suppose for the moment

$$(13) \quad w_l < w_r.$$

Consider then the scalar Riemann problem consisting of the PDE (7) together with the initial condition

$$(14) \quad g = \begin{cases} w_l & \text{if } x < 0 \\ w_r & \text{if } x > 0. \end{cases}$$

Now in view of hypothesis (ii) we have $\lambda_k(u_r) > \lambda_k(u_l)$; that is, according to (9), $F'_k(w_r) > F'_k(w_l)$. But then it follows from (i) that the function F_k defined by (8) is strictly convex. Accordingly we can apply Theorem 4 in §3.4.4 to the scalar Riemann problem (7), (14), whose unique weak solution is a continuous rarefaction wave connecting the states w_l and w_r . More specifically,


$$w(x, t) = \begin{cases} w_l & \text{if } \frac{x}{t} < F'_k(w_l) \\ G_k\left(\frac{x}{t}\right) & \text{if } F'_k(w_l) < \frac{x}{t} < F'_k(w_r) \\ w_r & \text{if } \frac{x}{t} > F'_k(w_r), \end{cases} \quad (x \in \mathbb{R}, t > 0)$$

where $G_k = (F'_k)^{-1}$. Thus $\mathbf{u}(x, t) = \mathbf{v}(w(x, t))$, where \mathbf{v} solves the ODE (6) and passes through u_l , is a continuous integral solution of (1), (2).

The case $w_l > w_r$ is treated similarly, since F_k is then concave. □

11.2.3. Shock waves, contact discontinuities.

We consider next the possibility that the states u_l and u_r may be joined not by a rarefaction wave as above, but rather by a shock.



k -shock wave

a. The shock set. Recalling the Rankine–Hugoniot condition from §11.1.1, we see that necessarily we must have the equality $\mathbf{F}(u_l) - \mathbf{F}(u_r) = \sigma(u_l - u_r)$, where $\sigma \in \mathbb{R}$, for such a shock wave to exist. This observation motivates the following

DEFINITION. Given a fixed state $z_0 \in \mathbb{R}^m$, we define the shock set


$$S(z_0) := \{z \in \mathbb{R}^m \mid \mathbf{F}(z) - \mathbf{F}(z_0) = \sigma(z - z_0) \text{ for a constant } \sigma = \sigma(z, z_0)\}.$$

THEOREM 2 (Structure of the shock set). Fix $z_0 \in \mathbb{R}^m$. In some neighborhood of z_0 , $S(z_0)$ consists of the union of m smooth curves $S_k(z_0)$ ($k = 1, \dots, m$), with the following properties:

- (i) The curve $S_k(z_0)$ passes through z_0 , with tangent $\mathbf{r}_k(z_0)$.
- (ii) $\lim_{\substack{z \rightarrow z_0 \\ z \in S_k(z_0)}} \sigma(z, z_0) = \lambda_k(z_0)$.
- (iii) $\sigma(z, z_0) = \frac{\lambda_k(z) + \lambda_k(z_0)}{2} + O(|z - z_0|^2)$, as $z \rightarrow z_0$ with $z \in S_k(z_0)$.

Proof. 1. Define

$$\mathbf{B}(z) := \int_0^1 D\mathbf{F}(z_0 + t(z - z_0)) dt \quad (z \in \mathbb{R}^m).$$



Contact between R_k and S_k

Then

$$(15) \quad \mathbf{B}(z)(z - z_0) = \mathbf{F}(z) - \mathbf{F}(z_0).$$

In particular $z \in S(z_0)$ if and only if

$$(16) \quad (\mathbf{B}(z) - \sigma I)(z - z_0) = 0$$

for some scalar $\sigma = \sigma(z, z_0)$.

2. We study equation (16) by first of all noting

$$(17) \quad \mathbf{B}(z_0) = D\mathbf{F}(z_0).$$

Now in view of the strict hyperbolicity, the characteristic polynomial $\lambda \mapsto \det(\lambda I - \mathbf{B}(z_0))$ has m distinct, real roots, and hence the polynomial $\lambda \mapsto \det(\lambda I - \mathbf{B}(z))$ likewise has m distinct roots if z is close to z_0 . Recalling Theorem 2 in §11.1.2, we see that near z_0 there exist smooth functions $\hat{\lambda}_1(z) < \dots < \hat{\lambda}_m(z)$ and unit vectors $\{\hat{\mathbf{r}}_k(z), \hat{\mathbf{l}}_k(z)\}_{k=1}^m$ satisfying

$$\hat{\lambda}_k(z_0) = \lambda_k(z_0), \quad \hat{\mathbf{r}}_k(z_0) = \mathbf{r}_k(z_0), \quad \hat{\mathbf{l}}_k(z_0) = \mathbf{l}_k(z_0) \quad (k = 1, \dots, m)$$

and

$$(18) \quad \begin{cases} \mathbf{B}(z)\hat{\mathbf{r}}_k(z) = \hat{\lambda}_k(z)\hat{\mathbf{r}}_k(z) \\ \hat{\mathbf{l}}_k(z)\mathbf{B}(z) = \hat{\lambda}_k(z)\hat{\mathbf{l}}_k(z) \end{cases} \quad (k = 1, \dots, m).$$

Note that $\{\hat{\mathbf{r}}_k(z)\}, \{\hat{\mathbf{l}}_k(z)\}_{k=1}^m$ are bases of \mathbb{R}^m , and also

$$(19) \quad \hat{\mathbf{l}}_l(z) \cdot \hat{\mathbf{r}}_k(z) = 0 \quad (k \neq l).$$

3. Equation (16) will hold provided $\sigma = \hat{\lambda}_k(z)$ for some $k \in \{1, \dots, m\}$ and $(z - z_0)$ is parallel to $\hat{\mathbf{r}}_k(z)$. In light of (19), these conditions are equivalent to asking that

$$(20) \quad \hat{\mathbf{l}}_l(z) \cdot (z - z_0) = 0 \quad (l \neq k).$$

These equalities amount to $m - 1$ equations for the m unknown components of z , which we intend to solve using the Implicit Function Theorem (§C.7). So define $\Phi_k : \mathbb{R}^m \rightarrow \mathbb{R}^{m-1}$ by setting

$$\Phi_k(z) := (\dots, \hat{\mathbf{l}}_{k-1}(z) \cdot (z - z_0), \hat{\mathbf{l}}_{k+1}(z) \cdot (z - z_0), \dots).$$

Now $\Phi_k(z_0) = 0$ and

$$D\Phi_k(z_0) = \begin{pmatrix} \mathbf{l}_1(z_0) \\ \vdots \\ \mathbf{l}_{k-1}(z_0) \\ \mathbf{l}_{k+1}(z_0) \\ \vdots \\ \mathbf{l}_m(z_0) \end{pmatrix}_{(m-1) \times m},$$

the entries of this matrix being regarded as row vectors. Since the vectors $\{\mathbf{l}_k(z_0)\}_{k=1}^m$ form a basis of \mathbb{R}^m , we see

$$\text{rank } D\Phi_k(z_0) = m - 1.$$

Accordingly, there exists a smooth curve $\phi_k : \mathbb{R} \rightarrow \mathbb{R}^m$ such that

$$(21) \quad \phi_k(0) = z_0$$

and

$$(22) \quad \Phi_k(\phi_k(t)) = 0 \quad \text{for all } t \text{ close to } 0.$$

The path of the curve $\phi_k(\cdot)$ for t near zero defines $S_k(z_0)$. We may reparameterize as necessary to ensure

$$(23) \quad |\dot{\phi}_k(t)| = 1 \quad \left(\cdot = \frac{d}{dt} \right).$$

4. Now (20)–(22) imply

$$(24) \quad \phi_k(t) = z_0 + \mu(t)\hat{\mathbf{r}}_k(\phi_k(t))$$

for all t near zero, where $\mu : \mathbb{R} \rightarrow \mathbb{R}$ is a smooth function satisfying $\mu(0) = 0$, $\dot{\mu}(0) = 1$. Differentiating (24) with respect to t and setting $t = 0$, we thus find

$$(25) \quad \dot{\phi}_k(0) = \mathbf{r}_k(z_0).$$

Hence the curve $S_k(z_0)$ has tangent $\mathbf{r}_k(z_0)$ at z_0 . Assertion (i) is proved.

5. In light of the foregoing analysis, there exists a smooth function $\sigma : \mathbb{R}^m \times \mathbb{R}^m \rightarrow \mathbb{R}$ such that

$$(26) \quad \mathbf{F}(\phi_k(t)) - \mathbf{F}(z_0) = \sigma(\phi_k(t), z_0)(\phi_k(t) - z_0)$$

for all t close to zero. Differentiating with respect to t and setting $t = 0$, we deduce from (21) that

$$D\mathbf{F}(z_0)\dot{\phi}_k(0) = \sigma(z_0, z_0)\dot{\phi}_k(0).$$

In light of (25), we see that $\sigma(z_0, z_0) = \lambda_k(z_0)$. This establishes assertion (ii).

6. Now write $\sigma(t) := \sigma(\phi_k(t), z_0)$, so that (26) reads

$$\mathbf{F}(\phi_k(t)) - \mathbf{F}(z_0) = \sigma(t)(\phi_k(t) - z_0).$$

Differentiate twice with respect to t :

$$\begin{aligned} & (D^2\mathbf{F}(\phi_k(t))\dot{\phi}_k(t))\dot{\phi}_k(t) + D\mathbf{F}(\phi_k(t))\ddot{\phi}_k(t) \\ & = \ddot{\sigma}(t)(\phi_k(t) - z_0) + 2\dot{\sigma}(t)\dot{\phi}_k(t) + \sigma(t)\ddot{\phi}_k(t). \end{aligned}$$

Evaluate this expression at $t = 0$ and recall $\sigma(0) = \lambda_k(z_0)$, $\phi_k(0) = z_0$, $\dot{\phi}_k(0) = \mathbf{r}_k(z_0)$:

$$(27) \quad (2\dot{\sigma}(0)I - D^2\mathbf{F}(z_0)\mathbf{r}_k(z_0))\mathbf{r}_k(z_0) = (D\mathbf{F}(z_0) - \lambda_k(z_0)I)\ddot{\phi}_k(0).$$

7. Let $\psi_k(t) = \mathbf{v}(t)$ be a unit speed parameterization of the rarefaction curve $R_k(z_0)$ near z_0 (as in (6) above). Then

$$(28) \quad \psi_k(0) = z_0, \quad \dot{\psi}_k(t) = \mathbf{r}_k(\psi_k(t)).$$

Thus

$$D\mathbf{F}(\psi_k(t))\mathbf{r}_k(t) = \lambda_k(t)\mathbf{r}_k(t),$$

for

$$\lambda_k(t) := \lambda_k(\psi_k(t)), \quad \mathbf{r}_k(t) := \mathbf{r}_k(\psi_k(t)).$$

Next differentiate with respect to t and set $t = 0$:

$$(29) \quad (D^2\mathbf{F}(z_0)\mathbf{r}_k(z_0) - \dot{\lambda}_k(0)I)\mathbf{r}_k(z_0) = -(D\mathbf{F}(z_0) - \lambda_k(z_0)I)\dot{\mathbf{r}}_k(0).$$

Add (27) and (29), to obtain

$$(30) \quad (2\dot{\sigma}(0) - \dot{\lambda}_k(0))\mathbf{r}_k(z_0) = (D\mathbf{F}(z_0) - \lambda_k(z_0)I)(\ddot{\phi}_k(0) - \dot{\mathbf{r}}_k(0)).$$

Take the dot product with $\mathbf{l}_k(z_0)$ and observe $\mathbf{l}_k \cdot \mathbf{r}_k \neq 0$, to conclude

$$(31) \quad 2\dot{\sigma}(0) = \dot{\lambda}_k(0).$$

We deduce from (31) that

$$2\sigma(t) - \lambda_k(z_0) - \lambda_k(t) = O(t^2) \quad \text{as } t \rightarrow 0.$$

Assertion (iii) follows. □

We see from Theorem 2(iii) that the curves $R_k(z_0)$ and $S_k(z_0)$ agree at least to first order at z_0 . Next is the assertion that in the linearly degenerate case these curves in fact coincide.

THEOREM 3 (Linear degeneracy). *Suppose for some $k \in \{1, \dots, m\}$ that the pair $(\lambda_k, \mathbf{r}_k)$ is linearly degenerate. Then for each $z_0 \in \mathbb{R}^m$,*

$$(i) \quad R_k(z_0) = S_k(z_0)$$

and

$$(ii) \quad \sigma(z, z_0) = \lambda_k(z) = \lambda_k(z_0) \text{ for all } z \in S_k(z_0).$$

Proof. Let $\mathbf{v} = \mathbf{v}(s)$ solve the ODE

$$\begin{cases} \dot{\mathbf{v}}(s) = \mathbf{r}_k(\mathbf{v}(s)) & (s \in \mathbb{R}) \\ \mathbf{v}(0) = z_0. \end{cases}$$


Then the mapping $s \mapsto \lambda_k(\mathbf{v}(s))$ is constant, and so

$$\begin{aligned} \mathbf{F}(\mathbf{v}(s)) - \mathbf{F}(z_0) &= \int_0^s D\mathbf{F}(\mathbf{v}(t))\dot{\mathbf{v}}(t) dt = \int_0^s D\mathbf{F}(\mathbf{v}(t))\mathbf{r}_k(\mathbf{v}(t)) dt \\ &= \int_0^s \lambda_k(\mathbf{v}(t))\mathbf{r}_k(\mathbf{v}(t)) dt = \lambda_k(z_0) \int_0^s \dot{\mathbf{v}}(t) dt \\ &= \lambda_k(z_0)(\mathbf{v}(s) - z_0). \end{aligned} \quad \square$$

b. Contact discontinuities, shock waves. We next undertake to analyze in light of Theorems 2 and 3 the possibility of solving Riemann's problem by joining two given states u_l and u_r by some kind of shock wave.

Contact discontinuities. Suppose first that $(\lambda_k, \mathbf{r}_k)$ is *linearly degenerate* and

$$(32) \quad u_r \in S_k(u_l).$$



k -contact discontinuity

We then define an integral solution of our system of conservation laws by setting

$$(33) \quad \mathbf{u}(x, t) = \begin{cases} u_l & \text{if } x < \sigma t \\ u_r & \text{if } x > \sigma t, \end{cases}$$

for

$$(34) \quad \sigma = \sigma(u_r, u_l) = \lambda_k(u_l) = \lambda_k(u_r).$$

Now observe from our analysis in §3.2.1 that since $\lambda_k(u_l) = \lambda_k(u_r) = \sigma$, the projected characteristics to the left and right are *parallel* to the line of discontinuity. We interpret this situation physically by saying that fluid particles do not cross the discontinuity. The line $x = \sigma t$ is called a *k -contact discontinuity*.

Shock waves. We next turn our attention to the case that $(\lambda_k, \mathbf{r}_k)$ is *genuinely nonlinear* and

$$(35) \quad u_r \in S_k(u_l),$$

as before. If we consider the integral solution

$$(36) \quad \mathbf{u}(x, t) = \begin{cases} u_l & \text{if } x < \sigma t \\ u_r & \text{if } x > \sigma t, \end{cases}$$

for

$$(37) \quad \sigma = \sigma(u_r, u_l),$$

we see that there are two essentially different cases according as to whether

$$(38) \quad \lambda_k(u_r) < \lambda_k(u_l)$$

or else

$$(39) \quad \lambda_k(u_l) < \lambda_k(u_r).$$

Now in view of assertion (iii) from Theorem 2, we have then either

$$(40) \quad \lambda_k(u_r) < \sigma(u_r, u_l) < \lambda_k(u_l)$$

or

$$(41) \quad \lambda_k(u_l) < \sigma(u_r, u_l) < \lambda_k(u_r),$$

provided u_r is close enough to u_l .

This dichotomy is reminiscent of a corresponding situation in §3.4, for a scalar conservation law. By analogy with the entropy conditions introduced there, let us hereafter agree to reject the inequalities (41) as allowing for “nonphysical shocks” from which characteristics emanate as we move forward in time. We rather take (40) as being physically correct. The informal viewpoint is that then the characteristics from the left and right run into the line of discontinuity, whereupon “information is lost” and so “entropy increases”. This interpretation was largely justified mathematically in §3.4 with our uniqueness theorem for weak solutions that satisfied this sort of entropy condition.

Refocusing our attention again to systems, we therefore agree to regard (40) as the correct inequalities to be satisfied:

DEFINITION. Assume the pair $(\lambda_k, \mathbf{r}_k)$ is genuinely nonlinear at u_l . We say that the pair (u_l, u_r) is admissible provided

$$u_r \in S_k(u_l)$$

and


$$(42) \quad \lambda_k(u_r) < \sigma(u_r, u_l) < \lambda_k(u_l).$$

We refer to (42) as the *Lax entropy condition*. If (u_l, u_r) is admissible, we call our solution \mathbf{u} defined by (36), (37) a *k-shock wave*.

By analogy with our decomposition of $R_k(z_0)$ into $R_k^\pm(z_0)$, let us introduce this

DEFINITION. If the pair $(\lambda_k, \mathbf{r}_k)$ is genuinely nonlinear, we write

$$S_k^+(z_0) := \{z \in S_k(z_0) \mid \lambda_k(z_0) < \sigma(z, z_0) < \lambda_k(z)\}$$



Shock curve

and

$$S_k^-(z_0) := \{z \in S_k(z_0) \mid \lambda_k(z) < \sigma(z, z_0) < \lambda_k(z_0)\}.$$

Then

$$S_k(z_0) = S_k^+(z_0) \cup \{z_0\} \cup S_k^-(z_0)$$

near z_0 . Note then that the pair (u_l, u_r) is admissible if and only if

$$u_r \in S_k^-(u_l).$$

11.2.4. Local solution of Riemann's problem.

Next we glue together the physically relevant parts of the rarefaction and shock curves.

DEFINITIONS. (i) If the pair $(\lambda_k, \mathbf{r}_k)$ is genuinely nonlinear, write

$$T_k(z_0) := R_k^+(z_0) \cup \{z_0\} \cup S_k^-(z_0).$$


(ii) If the pair $(\lambda_k, \mathbf{r}_k)$ is linearly degenerate, we set

$$T_k(z_0) := R_k(z_0) = S_k(z_0).$$

Owing to Theorem 2(ii), the curve $T_k(z_0)$ is C^1 . Employing the notation of §11.2.3, we see that nearby states u_l and u_r can be joined by a k -rarefaction wave, a shock wave or a contact discontinuity provided

$$(43) \quad u_l \in T_k(u_r).$$

We now at last ask if we can find a solution to Riemann's problem provided only that u_r is close to u_l (but (43) may fail for each $k = 1, \dots, m$). The hope is that by moving along various paths T_k for different values of k , we may be able to connect u_l to u_r , utilizing a sequence of rarefaction waves, shock waves, and/or contact discontinuities.



Structure of the T -curve

THEOREM 4 (Local solution of Riemann's problem). *Assume for each $k = 1, \dots, m$ that the pair $(\lambda_k, \mathbf{r}_k)$ is either genuinely nonlinear or else linearly degenerate. Suppose further the left state u_l is given. Then for each right state u_r sufficiently close to u_l there exists an integral solution \mathbf{u} of Riemann's problem, which is constant on lines through the origin.*

Proof. 1. We intend to apply the Inverse Function Theorem (§C.6) to a mapping $\Phi : \mathbb{R}^m \rightarrow \mathbb{R}^m$, defined near 0 as follows.

First, for each family of curves T_k ($k = 1, \dots, m$) choose the nonsingular parameter τ_k to measure arc length; that is, if $z, \tilde{z} \in \mathbb{R}^m$ with $\tilde{z} \in T_k(z)$, then

$$\tau_k(\tilde{z}) - \tau_k(z) = (\text{signed}) \text{ distance from } z \text{ to } \tilde{z} \text{ along the curve } T_k(z).$$

We take the plus sign for $\tau_k(\tilde{z})$ if $\tilde{z} \in R_k^+(z)$, the minus sign if $\tilde{z} \in S_k^-(z)$.

2. Given then $t = (t_1, \dots, t_m) \in \mathbb{R}^m$, with $|t|$ small, we define $\Phi(t) = z$ as follows. First, temporarily write

$$(44) \quad u_l = z_0.$$

Then choose states z_1, \dots, z_m to satisfy

$$(45) \quad \begin{cases} z_1 \in T_1(z_0), \tau_1(z_1) - \tau_1(z_0) = t_1, \\ z_2 \in T_2(z_1), \tau_2(z_2) - \tau_2(z_1) = t_2, \\ \vdots \\ z_m \in T_m(z_{m-1}), \tau_m(z_m) - \tau_m(z_{m-1}) = t_m. \end{cases}$$

Now write

$$(46) \quad z = z_m,$$

and define

$$(47) \quad \Phi(t) = z.$$

Note Φ is C^1 and

$$(48) \quad \Phi(0) = z_0.$$

3. We claim

$$(49) \quad D\Phi(0) \text{ is nonsingular.}$$

To see this, observe

$$\Phi(0, \dots, t_k, \dots, 0) - \Phi(0, \dots, 0) = t_k \mathbf{r}_k(z_0) + o(t_k) \quad \text{as } t_k \rightarrow 0.$$

Thus

$$\frac{\partial \Phi}{\partial t_k}(0) = \mathbf{r}_k(z_0) \quad (k = 1, \dots, m),$$

and so

$$D\Phi(0) = (\mathbf{r}_1(z_0), \dots, \mathbf{r}_m(z_0))_{m \times m},$$

the entries regarded as column vectors. This matrix is nonsingular, since $\{\mathbf{r}_k(z_0)\}_{k=1}^m$ is a basis.

4. In light of (49), the Inverse Function Theorem applies: for each state u_τ sufficiently close to u_l there exists a unique parameter $t = (t_1, \dots, t_m)$ close to zero such that $\Phi(t) = u_\tau$.

Recall next that if z_{k-1} and z_k are joined by a k -rarefaction wave, this wave is

$$\begin{cases} z_{k-1} & \text{if } \frac{x}{t} < \lambda_k(z_{k-1}) \\ G_k\left(\frac{x}{t}\right) & \text{if } \lambda_k(z_{k-1}) < \frac{x}{t} < \lambda_k(z_k), \quad \text{for } G_k = (F'_k)^{-1} \\ z_k & \text{if } \lambda_k(z_k) < \frac{x}{t}. \end{cases}$$

Moreover if z_{k-1}, z_k are joined by a k -shock, it has the form

$$\begin{cases} z_{k-1} & \text{if } \frac{x}{t} < \sigma(z_k, z_{k-1}) \\ z_k & \text{if } \sigma(z_k, z_{k-1}) < \frac{x}{t}, \end{cases}$$

where $\lambda_k(z_k) < \sigma(z_k, z_{k-1}) < \lambda_k(z_{k-1})$. In both cases the waves are constant outside the regions $\lambda_k(z_0) - \epsilon < \frac{x}{t} < \lambda_k(z_0) + \epsilon$, for small $\epsilon > 0$, provided z_k, z_{k-1} are close enough to z_0 . This is true for $k = 1, \dots, m$.

Since $\lambda_1(z_0) < \dots < \lambda_m(z_0)$, we see then that the rarefactions, shock waves and/or contact discontinuities connecting $u_l = z_0$ to z_1 , z_1 to z_2 , z_2 to z_3 , \dots , z_{m-1} to $z_m = u_\tau$ do not intersect. \square

11.3. SYSTEMS OF TWO CONSERVATION LAWS

In this section we more deeply analyze the initial-value problem for $m = 2$, which is to say, for a pair of conservation laws:

$$(1) \quad \begin{cases} u_t^1 + F^1(u^1, u^2)_x = 0 \\ u_t^2 + F^2(u^1, u^2)_x = 0 \\ u^1 = g^1, u^2 = g^2 \end{cases} \begin{array}{l} \text{in } \mathbb{R} \times (0, \infty) \\ \\ \text{on } \mathbb{R} \times \{t = 0\}. \end{array}$$

Here $\mathbf{F} = (F^1, F^2)$, $\mathbf{g} = (g^1, g^2)$, $\mathbf{u} = (u^1, u^2)$.

11.3.1. Riemann invariants.

Our intention is first to demonstrate that we can transform (1) into a much simpler form by performing an appropriate nonlinear change of dependent variables. The idea is to find two functions $w^1, w^2 : \mathbb{R}^2 \rightarrow \mathbb{R}$ with nice properties along the rarefaction curves R_1, R_2 :

DEFINITION. We say

$$w^i : \mathbb{R}^2 \rightarrow \mathbb{R}$$

is an i^{th} -Riemann invariant provided

$$(2) \quad Dw^i(z) \text{ is parallel to } \mathbf{l}_j(z) \quad (z \in \mathbb{R}^2, i \neq j).$$

We will see momentarily how useful condition (2) is, but let us first pause to ask whether Riemann invariants exist. It turns out that since we are now taking $m = 2$, this is easy. Indeed, because $\mathbf{l}_j(z) \cdot \mathbf{r}_i(z) = 0$ ($i \neq j$), we see (2) is equivalent in \mathbb{R}^2 to the statement

$$(2') \quad Dw^i(z) \cdot \mathbf{r}_i(z) = 0 \quad (i = 1, 2, z \in \mathbb{R}^2),$$

which is to say

$$(3) \quad w^i \text{ is constant along the rarefaction curve } R_i \quad (i = 1, 2).$$

In particular, any smooth function w^i satisfying (3) satisfies also (2'), (2) and so is an i^{th} -Riemann invariant.

In the case that $m > 2$, Riemann invariants do not in general exist.

Now we can regard $w = (w_1, w_2) = (w^1(z_1, z_2), w^2(z_1, z_2))$ as being new coordinates on the state space \mathbb{R}^2 , replacing $z = (z_1, z_2)$. More precisely, we define $\mathbf{w} : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ by setting

$$(4) \quad \mathbf{w}(z) = \mathbf{w}(z_1, z_2) = (w^1(z_1, z_2), w^2(z_1, z_2)).$$

The inverse mapping is $\mathbf{z}(w) = \mathbf{z}(w_1, w_2) = (z^1(w_1, w_2), z^2(w_1, w_2))$.

Let us now utilize the transformation (4) to simplify our system of two conservation laws (1). For this, let us suppose henceforth $\mathbf{u} = (u^1, u^2)$ is a smooth solution of (1). We now change dependent variables by setting

$$(5) \quad \mathbf{v}(x, t) := \mathbf{w}(\mathbf{u}(x, t)) \quad (x \in \mathbb{R}, t > 0).$$

What system of PDE does $\mathbf{v} = (v^1, v^2)$ satisfy?

THEOREM 1 (Conservation laws and Riemann invariants). *The functions v^1, v^2 solve the system*

$$(6) \quad \begin{cases} v_t^1 + \lambda_2(\mathbf{u})v_x^1 = 0 \\ v_t^2 + \lambda_1(\mathbf{u})v_x^2 = 0 \end{cases} \quad \text{in } \mathbb{R} \times (0, \infty).$$

The point is that the system (6), although not in conservation law form, is in many ways rather simpler than (1). Note in particular that whereas the PDE for u^1 involves the term u_x^2 , the PDE for v^1 does *not* entail v_x^2 . Similarly, the PDE for v^2 does not involve v_x^1 .

Proof. According to (5), we see that for $i = 1, 2, i \neq j$,

$$\begin{aligned} v_t^i + \lambda_j(\mathbf{u})v_x^i &= Dw^i(\mathbf{u}) \cdot \mathbf{u}_t + \lambda_j(\mathbf{u})Dw^i(\mathbf{u}) \cdot \mathbf{u}_x \\ &= Dw^i(\mathbf{u}) \cdot (-\mathbf{F}(\mathbf{u})_x + \lambda_j(\mathbf{u})\mathbf{u}_x) \\ &= Dw^i(\mathbf{u}) \cdot (-D\mathbf{F}(\mathbf{u}) + \lambda_j(\mathbf{u})I)\mathbf{u}_x = 0, \end{aligned}$$

since, by definition, Dw^i is parallel to \mathbf{l}_j . □

Remarks. (i) We can interpret the system of PDE (6) by introducing the ODE

$$(7) \quad \dot{x}_i(s) = \lambda_j(\mathbf{u}(x_i(s), s)) \quad (s \geq 0)$$

for $i = 1, 2, j \neq i$. Then we see from (6) that

$$(8) \quad v^i \text{ is constant along the curve } (x_i(s), s) \quad (s \geq 0)$$

for $i = 1, 2$.

(ii) Recall from §11.2 that our condition of genuine nonlinearity reads

$$(9) \quad D\lambda_i(z) \cdot \mathbf{r}_i(z) \neq 0 \quad (z \in \mathbb{R}^2, i = 1, 2).$$

Since we can also regard λ_i as a function of $w = (w_1, w_2)$, we can rewrite (9) to read

$$(10) \quad \frac{\partial \lambda_i}{\partial w_j} \neq 0 \quad (w \in \mathbb{R}^2, i \neq j).$$

To see (9) is equivalent to (10), observe that if (10) fails, then

$$(11) \quad 0 = \frac{\partial \lambda_i}{\partial w_j} = \sum_{k=1}^2 \frac{\partial \lambda_i}{\partial z_k} \frac{\partial z^k}{\partial w_j}.$$

But since $\sum_{k=1}^2 \frac{\partial w^i}{\partial z_k} \frac{\partial z^k}{\partial w_j} = \delta_{ij} = 0$ for $i \neq j$, we see that (11) asserts that $D\lambda_i$ is parallel to Dw^i . However, Dw^i is perpendicular to \mathbf{r}_i , and so we obtain a contradiction to (9). Hence (9) implies (10), and the converse implication is established in the same way. \square

Example (Barotropic compressible gas dynamics). We illustrate the foregoing ideas by examining in detail Euler's equations for compressible gas dynamics (Example 2 in §11.1) in the special case that the internal energy e is constant. The relevant PDE are

$$(12) \quad \begin{cases} \rho_t + (\rho v)_x = 0 & \text{(conservation of mass)} \\ (\rho v)_t + (\rho v^2 + p)_x = 0 & \text{(conservation of momentum),} \end{cases}$$

where we now assume

$$(13) \quad p = p(\rho)$$

for some smooth function $p : \mathbb{R} \rightarrow \mathbb{R}$. Formula (13) is called a *barotropic equation of state*. We assume the strict hyperbolicity condition

$$(14) \quad p' > 0.$$

Setting $\mathbf{u} = (u^1, u^2) = (\rho, \rho v)$, we can rewrite (12), (13) to read

$$\mathbf{u}_t + \mathbf{F}(\mathbf{u})_x = 0,$$

for

$$\mathbf{F} = (F^1, F^2) = (z_2, (z_2)^2/z_1 + p(z_1))$$

and $z = (z_1, z_2)$, provided $z_1 > 0$. Then

$$D\mathbf{F} = \begin{pmatrix} 0 & 1 \\ -(\frac{z_2}{z_1})^2 + p'(z_1) & \frac{2z_2}{z_1} \end{pmatrix}.$$

Consequently

$$(15) \quad \lambda_1 = \frac{z_2}{z_1} - p'(z_1)^{1/2}, \quad \lambda_2 = \frac{z_2}{z_1} + p'(z_1)^{1/2}.$$

In physical notation,

$$(16) \quad \lambda_1 = v - \sigma, \quad \lambda_2 = v + \sigma,$$

for the sound speed

$$(17) \quad \sigma := p'(\rho)^{1/2}.$$

Remembering (7), we consider next the ODE

$$(18) \quad \dot{x}_1(t) = v(x_1(t), t) + \sigma(x_1(t), t),$$

$$(19) \quad \dot{x}_2(t) = v(x_2(t), t) - \sigma(x_2(t), t),$$

where $\sigma(x, t) := p'(\rho(x, t))^{\frac{1}{2}}$, $t \geq 0$. We know from (8) that the Riemann invariant $v^1 = w^1(\mathbf{u})$ is constant along the trajectories of (18) and $v^2 = w^2(\mathbf{u})$ is constant along trajectories of (19).

To compute w^1 and w^2 directly, let us carry out some computations on the system (12). First we transform (12) in nondivergence form:

$$(20) \quad \rho_t + \rho v_x + \rho_x v = 0,$$

$$(21) \quad \rho_t v + \rho v_t + \rho_x v^2 + 2\rho v v_x + p_x = 0.$$

Multiplying (20) by $\sigma^2 = p'(\rho)$ and recalling (13) gives us

$$(22) \quad p_t + v p_x + \sigma^2 \rho v_x = 0.$$

In addition (20), (21) combine to yield

$$(23) \quad \rho v_t + \rho v v_x + p_x = 0.$$

We now manipulate (22), (23) so that the directions $\lambda_1, \lambda_2 = v \mp \sigma$ appear explicitly. To accomplish this, we multiply (23) by σ and then add to and subtract from (22):

$$(24) \quad \begin{cases} p_t + (v + \sigma)p_x + \rho\sigma(v_t + (v + \sigma)v_x) = 0 \\ p_t + (v - \sigma)p_x - \rho\sigma(v_t + (v - \sigma)v_x) = 0. \end{cases}$$

We then deduce from (24) that

$$(25) \quad \begin{cases} \frac{d}{dt}[p(x_1(t), t)] + \rho(x_1(t), t)\sigma(x_1(t), t)\frac{d}{dt}[v(x_1(t), t)] = 0 \\ \frac{d}{dt}[p(x_2(t), t)] - \rho(x_2(t), t)\sigma(x_2(t), t)\frac{d}{dt}[v(x_2(t), t)] = 0. \end{cases}$$

As $\frac{dp}{dt} = \sigma^2 \frac{d\rho}{dt}$, we see

$$(26) \quad \frac{\sigma}{\rho} \frac{d\rho}{dt} \pm \frac{dv}{dt} = 0 \text{ along the trajectories of (18), (19),}$$

provided $\rho > 0$.

Think now of the Riemann invariants as functions of ρ and v . Then since $v^1 = w^1(\rho, v)$ is constant along the curve determined by $x_1(\cdot)$, we have

$$\begin{aligned} 0 &= \frac{d}{dt}[w^1(\rho(x_1(t), t), v(x_1(t), t))] \\ &= \frac{\partial w^1}{\partial \rho} \frac{d}{dt}[\rho(x_1(t), t)] + \frac{\partial w^1}{\partial v} \frac{d}{dt}[v(x_1(t), t)]. \end{aligned}$$

This is consistent with (26) if

$$\frac{\partial w^1}{\partial \rho} = \frac{\sigma(\rho)}{\rho}, \quad \frac{\partial w^1}{\partial v} = 1.$$

We similarly deduce

$$\frac{\partial w^2}{\partial \rho} = \frac{\sigma(\rho)}{\rho}, \quad \frac{\partial w^2}{\partial v} = -1.$$

Integrating, we conclude that the Riemann invariants are, up to additive constants,

$$w^1 = \int_1^\rho \frac{\sigma(s)}{s} ds + v, \quad w^2 = \int_1^\rho \frac{\sigma(s)}{s} ds - v.$$

We leave it as an exercise to check that w^1, w^2 , taken now as functions of $z = (z_1, z_2)$, satisfy the definition of Riemann invariants. \square

11.3.2. Nonexistence of smooth solutions.

Illustrating now the usefulness of Riemann invariants, we establish the following criterion for the nonexistence of a smooth solution:

THEOREM 2 (Riemann invariants and blow-up). *Assume \mathbf{g} is smooth, with compact support. Suppose also the genuine nonlinearity condition*

$$(27) \quad \frac{\partial \lambda_i}{\partial w_j} > 0 \text{ in } \mathbb{R}^2 \quad (i = 1, 2, i \neq j)$$

holds. Then the initial-value problem (1) cannot have a smooth solution \mathbf{u} existing for all times $t \geq 0$ if

$$(28) \quad \text{either } v_x^1 < 0 \text{ or } v_x^2 < 0 \text{ somewhere on } \mathbb{R} \times \{t = 0\}.$$

Proof. 1. Assume for the time being that \mathbf{u} is a smooth solution of (1). Write

$$(29) \quad a := v_x^1, \quad b := v_x^2,$$

where $\mathbf{v} = \mathbf{w}(\mathbf{u})$, $\mathbf{v} = (v^1, v^2)$, solves the system of PDE (6). We differentiate the first equation of (6) with respect to x , to compute

$$(30) \quad a_t + \lambda_2 a_x + \frac{\partial \lambda_2}{\partial w_1} a^2 + \frac{\partial \lambda_2}{\partial w_2} ab = 0.$$

We employ then the second equation of (6), which we rewrite as

$$v_t^2 + \lambda_2 v_x^2 = (\lambda_2 - \lambda_1)b.$$

Substituting this expression into (30) gives

$$(31) \quad a_t + \lambda_2 a_x + \frac{\partial \lambda_2}{\partial w_1} a^2 + \left[\frac{1}{\lambda_2 - \lambda_1} \frac{\partial \lambda_2}{\partial w_2} (v_t^2 + \lambda_2 v_x^2) \right] a = 0.$$

2. To integrate (31), fix $x_0 \in \mathbb{R}$ and set

$$(32) \quad \xi(t) := \exp \left(\int_0^t \frac{1}{\lambda_2 - \lambda_1} \frac{\partial \lambda_2}{\partial w_2} (v_t^2 + \lambda_2 v_x^2)(x_1(s), s) ds \right),$$

where

$$(33) \quad \begin{cases} \dot{x}_1(s) = \lambda_2(\mathbf{u}(x_1(s), s)) & (s \geq 0) \\ x_1(0) = x_0. \end{cases}$$

Next is the key observation from (8) that v^1 is constant along the curve $(x_1(s), s)$. So write

$$v^1(x_1(s), s) = v_0^1 = v^1(x_0, 0) \quad (s \geq 0).$$

Thus we see that the expression $\left(\frac{1}{\lambda_2 - \lambda_1} \frac{\partial \lambda_2}{\partial w_2} \right)$, considered as a function of $\mathbf{v} = \mathbf{w}(\mathbf{u})$, depends only on v^2 . Let us set

$$\gamma(\mu) := \int_0^\mu \left(\frac{1}{\lambda_2 - \lambda_1} \frac{\partial \lambda_2}{\partial w_2} \right) (v_0^1, v) dv.$$

Then (32), (33) imply

$$(34) \quad \begin{aligned} \xi(t) &= \exp \left(\int_0^t \frac{d}{ds} [\gamma(v^2(x_1(s), s))] ds \right) \\ &= \exp(\gamma(v^2(x_1(t), t)) - \gamma(v^2(x_0, 0))). \end{aligned}$$

3. We now transform (31) to read

$$\frac{d}{dt}((\xi\alpha)^{-1}) = \frac{-1}{(\xi\alpha)^2} \frac{d}{dt}(\xi\alpha) = \frac{\partial\lambda_2}{\partial w_1} \xi^{-1}$$

where $\alpha(t) := a(x_1(t), t)$ and we assume $\alpha \neq 0$. Consequently

$$(\xi(t)\alpha(t))^{-1} = (\alpha(0))^{-1} + \int_0^t \frac{\partial\lambda_2}{\partial w_1} \xi^{-1}(s) ds.$$

This equality in turn rearranges to become

$$(35) \quad \alpha(t) = \alpha(0)\xi^{-1}(t) \left(1 + \alpha(0) \int_0^t \frac{\partial\lambda_2}{\partial w_1} \xi^{-1}(s) ds \right)^{-1}.$$

4. Now in view of the system of PDE (6), \mathbf{v} is bounded. Thus we deduce from (34) that $0 < \theta \leq \xi(t) \leq \Theta$ for all times $t > 0$, for appropriate constants θ, Θ . Therefore it follows from (27) and (35) that α is bounded for all $t > 0$ if and only if $\alpha(0) \geq 0$, that is, if

$$v_x^1(x_0, 0) \geq 0.$$

A similar calculation holds with v^2 replacing v^1 . We conclude that if either $v_x^1 < 0$ or else $v_x^2 < 0$ somewhere on $\mathbb{R} \times \{t = 0\}$, there cannot then exist a smooth solution of (1), lasting for all times $t \geq 0$. \square

11.4. ENTROPY CRITERIA

In our study of Riemann's problem in §11.2 we have taken Lax's entropy condition

$$(1) \quad \lambda_k(u_r) < \sigma(u_r, u_l) < \lambda_k(u_l)$$

for some $k \in \{1, \dots, m\}$ as the selection criteria for admissible shock waves.

There is great ongoing interest in discovering other mathematically correct and physically appropriate entropy conditions of various sorts, with the aim of applying these to more complicated integral solutions of our system of conservation laws, so as to obtain uniqueness criteria, more information concerning allowable discontinuities, etc.

One general principle, instances of which we have already seen for scalar conservation laws in §4.5.1 and for Hamilton–Jacobi equations in §10.1, is

that physically and mathematically correct solutions should arise as the limit of solutions to the regularized system

$$(2) \quad \mathbf{u}_t^\varepsilon + \mathbf{F}(\mathbf{u}^\varepsilon)_x - \varepsilon \mathbf{u}_{xx}^\varepsilon = 0 \quad \text{in } \mathbb{R} \times (0, \infty)$$

as $\varepsilon \rightarrow 0$. The idea is to interpret the term “ $\varepsilon \mathbf{u}_{xx}^\varepsilon$ ” as providing a small viscosity effect, which will presumably “smear out” sharp shocks. The hope is to study various aspects of the problem (2) in the limit $\varepsilon \rightarrow 0$, and thereby to discover more general entropy criteria, to augment Lax’s condition (1).

The next subsections discuss aspects of this general program.

11.4.1. Vanishing viscosity, traveling waves.

We begin our investigation of the parabolic system (2) by first seeking a traveling wave solution, having the form

$$(3) \quad \mathbf{u}^\varepsilon(x, t) = \mathbf{v}\left(\frac{x - \sigma t}{\varepsilon}\right) \quad (x \in \mathbb{R}, t \geq 0),$$

where as usual the speed σ and profile \mathbf{v} must be found. Substituting (3) into (2), we find $\mathbf{v} : \mathbb{R} \rightarrow \mathbb{R}^m$, $\mathbf{v} = \mathbf{v}(s)$, must solve the ODE

$$(4) \quad \ddot{\mathbf{v}} = -\sigma \dot{\mathbf{v}} + D\mathbf{F}(\mathbf{v})\dot{\mathbf{v}} \quad \left(\dot{\cdot} = \frac{d}{ds}\right).$$

Assume now $u_l, u_r \in \mathbb{R}^m$ are given and furthermore

$$(5) \quad \lim_{s \rightarrow -\infty} \mathbf{v} = u_l, \quad \lim_{s \rightarrow +\infty} \mathbf{v} = u_r, \quad \lim_{s \rightarrow \pm\infty} \dot{\mathbf{v}} = 0.$$

Then from (3) we deduce

$$(6) \quad \lim_{\varepsilon \rightarrow 0} \mathbf{u}^\varepsilon(x, t) = \begin{cases} u_l & \text{if } x < \sigma t \\ u_r & \text{if } x > \sigma t. \end{cases}$$

Hence the limit as $\varepsilon \rightarrow 0$ of our solution to (2) gives us a shock wave connecting the states u_l, u_r . The plan now is to study carefully the form of σ and \mathbf{v} and thereby glean more detailed information about the structure of the shock determined by (6).

The first and primary question is whether there in fact exist σ and \mathbf{v} solving (4), (5). Integrating (4), we deduce

$$(7) \quad \dot{\mathbf{v}} = \mathbf{F}(\mathbf{v}) - \sigma \mathbf{v} + c$$

for some constant $c \in \mathbb{R}^m$. We conclude from (5) that

$$(8) \quad \mathbf{F}(u_l) - \sigma u_l + c = \mathbf{F}(u_r) - \sigma u_r + c.$$

Hence

$$(9) \quad \mathbf{F}(u_l) - \mathbf{F}(u_r) = \sigma(u_l - u_r).$$

In view of (5), (8) and (9), our ODE (7) becomes

$$(10) \quad \dot{\mathbf{v}} = \mathbf{F}(\mathbf{v}) - \mathbf{F}(u_l) - \sigma(\mathbf{v} - u_l).$$

Now think of the left state u_l as being given, and suppose we are trying to build a traveling wave connecting u_l to a nearby state u_r . From (9) we see that necessarily $u_r \in S_k(u_l)$ for some $k \in \{1, \dots, m\}$ and

$$(11) \quad \sigma = \sigma(u_r, u_l).$$

We refine this observation as follows:

THEOREM 1 (Existence of traveling waves for genuinely nonlinear systems). *Assume the pair $(\lambda_k, \mathbf{r}_k)$ is genuinely nonlinear for $k = 1, \dots, m$. Let u_r be selected sufficiently close to u_l . Then there exists a traveling wave solution of (2) connecting u_l to u_r if and only if*

$$(12) \quad u_r \in S_k^-(u_l)$$

for some $k \in \{1, \dots, m\}$.

Proof. 1. Assume first σ and \mathbf{v} solve (4), (5). Then, as noted above, necessarily $u_r \in S_k(u_l)$ for some $k \in \{1, \dots, m\}$, $\sigma = \sigma(u_r, u_l)$. Now set

$$(13) \quad \mathbf{G}(z) := \mathbf{F}(z) - \mathbf{F}(u_l) - \sigma(z - u_l).$$

Our ODE (10) then reads

$$(14) \quad \dot{\mathbf{v}} = \mathbf{G}(\mathbf{v}),$$

and we have

$$(15) \quad \mathbf{G}(u_l) = \mathbf{G}(u_r) = 0,$$

according to (9). We compute

$$D\mathbf{G}(u_l) = D\mathbf{F}(u_l) - \sigma I;$$

and so the eigenvalues of $D\mathbf{G}$ at u_l are $\{\lambda_k(u_l) - \sigma\}_{k=1}^m$, with corresponding right and left eigenvectors $\{\mathbf{r}_k, \mathbf{l}_k\}_{k=1}^m$, $\mathbf{r}_k = \mathbf{r}_k(u_l)$, $\mathbf{l}_k = \mathbf{l}_k(u_l)$.

2. Now since $u_r \in S_k(u_l)$ and $|u_r - u_l|$ is small, we know from Theorem 2(iii) in §11.2.3 that

$$\sigma = \frac{\lambda_k(u_r) + \lambda_k(u_l)}{2} + o(|u_r - u_l|).$$

Thus

$$\lambda_k(u_l) - \sigma = \frac{\lambda_k(u_l) - \lambda_k(u_r)}{2} + o(|u_r - u_l|).$$

In order that there be an orbit of the ODE (14) connecting u_l at $s = -\infty$ to $u_r \in S_k(u_r)$ at $s = +\infty$, it must be that $\lambda_k(u_l) - \sigma > 0$, for otherwise the trajectory would not converge to u_l as $s \rightarrow -\infty$. Thus if $|u_r - u_l|$ is small enough, $\lambda_k(u_r) < \lambda_k(u_l)$, which is to say $u_r \in S_k^-(u_l)$.

3. We omit proof of the sufficiency of condition (12): see Majda–Pego (J. Diff. Eq. 56 (1985), 229–262). \square

The preceding result employs the genuine nonlinearity assumption, but the assertion holds in general, provided we introduce an appropriate variant of Lax's entropy condition (1). So let us suppose now $u_r \in S_k(u_l)$ for some $k \in \{1, \dots, m\}$ and furthermore

$$(16) \quad \begin{cases} \sigma(z, u_l) > \sigma(u_r, u_l) \text{ for each } z \text{ lying} \\ \text{on the curve } S_k(u_l) \text{ between } u_r \text{ and } u_l. \end{cases}$$

Condition (16) is *Liu's entropy criterion*. (Observe that this condition is automatic provided $(\lambda_k, \mathbf{r}_k)$ is genuinely nonlinear, $u_r \in S_k^-(u_l)$, and u_r is sufficiently close to u_l .)

We can motivate (16) by again seeking traveling wave solutions of system (2). So assume u_l is given. Then provided $|u_r - u_l|$ is small enough, it turns out that there exists a traveling wave solution $\mathbf{u}^\varepsilon(x, t) = \mathbf{v}\left(\frac{x - \sigma t}{\varepsilon}\right)$, \mathbf{v} solving (4), (5), if and only if the entropy condition (16) is satisfied. See Conlon (Adv. in Math. 35 (1980), 1–18) for a proof.

To make this all a bit clearer, we next present in detail a specific application.

Example (Traveling waves for the p -system). Let us consider again the p -system

$$(17) \quad \begin{cases} u_t^1 - u_x^2 = 0 & \text{(compatibility condition)} \\ u_t^2 - p(u^1)_x = 0 & \text{(Newton's law),} \end{cases}$$

under the usual strict hyperbolicity condition

$$(18) \quad p' > 0.$$

We investigate the existence of traveling wave solutions to the regularized system

$$(19) \quad \begin{cases} u_t^{\varepsilon,1} - u_x^{\varepsilon,2} = 0 \\ u_t^{\varepsilon,2} - p(u^{\varepsilon,1})_x = \varepsilon u_{xx}^{\varepsilon,2}. \end{cases}$$

Notice we have added the viscosity term only to the second equation. This makes sense physically, as the first line of (17) is only a mathematical compatibility condition.

Assume now $u^\varepsilon = v\left(\frac{x-\sigma t}{\varepsilon}\right)$ is a traveling wave solution of (19), with

$$(20) \quad \lim_{s \rightarrow -\infty} v = u_l, \quad \lim_{s \rightarrow \infty} v = u_r, \quad \lim_{s \rightarrow \pm\infty} \dot{v} = 0.$$

Writing $v = (v^1, v^2)$, we compute from (19) that

$$(21) \quad \begin{cases} -\sigma \dot{v}^1 - \dot{v}^2 = 0 & (\dot{\cdot} = \frac{d}{ds}) \\ -\sigma \dot{v}^2 - p(v^1) \dot{\cdot} = \ddot{v}^2. \end{cases}$$

An integration using (20) gives

$$(22) \quad \begin{cases} \sigma v^1 + v^2 = \sigma v_l^1 + v_l^2 = \sigma v_r^1 + v_r^2 \\ \dot{v}^2 = \sigma(v_l^2 - v^2) + p(v_l^1) - p(v^1) = \sigma(v_r^2 - v^2) + p(v_r^1) - p(v^1), \end{cases}$$

for $u_l = (v_l^1, v_l^2)$, $u_r = (v_r^1, v_r^2)$. In particular,

$$\begin{cases} \sigma v_l^1 + v_l^2 = \sigma v_r^1 + v_r^2 \\ \sigma v_l^2 + p(v_l^1) = \sigma v_r^2 + p(v_r^1). \end{cases}$$

Solving these equations for σ , we obtain

$$(23) \quad \sigma^2 = \frac{p(v_r^1) - p(v_l^1)}{v_r^1 - v_l^1}.$$

Suppose hereafter $v_r^1 > v_l^1$. In view of (18) we can take $\sigma > 0$. In this situation the Liu entropy criterion reads

$$(24) \quad \frac{p(z_1) - p(v_l^1)}{z_1 - v_l^1} > \frac{p(v_r^1) - p(v_l^1)}{v_r^1 - v_l^1}$$

for all z on the curve $S_k(u_l)$ between u_l and u_r , $z = (z_1, z_2)$.

We now claim the system of ODE (22), with asymptotic boundary conditions (20), has a solution if and only if the entropy condition (24) holds. To confirm this, combine the two equations in (22) to eliminate v^2 :

$$\dot{v}^1 = \frac{p(v^1) - p(v_l^1)}{\sigma} - \sigma(v^1 - v_l^1) =: g(v^1).$$

Now $g(v_l^1) = 0$ and $g(v_r^1) = 0$, according to (23). Thus in order that the ODE (24) have a solution, with $\lim_{s \rightarrow -\infty} v^1 = v_l^1$, $\lim_{s \rightarrow \infty} v^1 = v_r^1$, we require

$$g(z_1) > 0 \quad \text{for } v_l^1 < z_1 < v_r^1.$$

But this is precisely the entropy criterion (24). A similar calculation works if $v_r^1 < v_l^1$. \square

11.4.2. Entropy/entropy-flux pairs.

Both Lax's and Liu's entropy criteria provide restrictions on possible left- and right-hand states joined by a shock wave (or a traveling wave for the viscous approximation). It is however of considerable interest to widen still further the entropy criteria, so as to apply to more general integral solutions of our conservation laws.

One idea is to require that an integral solution satisfy certain "entropy-type" inequalities.

DEFINITION. *Two smooth functions $\Phi, \Psi : \mathbb{R}^m \rightarrow \mathbb{R}$ comprise an entropy/entropy-flux pair for the conservation law $\mathbf{u}_t + \mathbf{F}(\mathbf{u})_x = 0$ provided*

$$(25) \quad \Phi \text{ is convex}$$

and

$$(26) \quad D\Phi(z)DF(z) = D\Psi(z) \quad (z \in \mathbb{R}^m).$$

To motivate condition (26), suppose for the moment \mathbf{u} is a smooth solution of the system of PDE $\mathbf{u}_t + \mathbf{F}(\mathbf{u})_x = 0$. We then compute

$$(27) \quad \begin{aligned} \Phi(\mathbf{u})_t + \Psi(\mathbf{u})_x &= D\Phi(\mathbf{u}) \cdot \mathbf{u}_t + D\Psi(\mathbf{u}) \cdot \mathbf{u}_x \\ &= (-D\Phi(\mathbf{u})DF(\mathbf{u}) + D\Psi(\mathbf{u})) \cdot \mathbf{u}_x = 0 \end{aligned}$$

by (26). This computation says the quantity $\Phi(\mathbf{u})$ satisfies a scalar conservation law, with flux $\Psi(\mathbf{u})$.

Now in general integral solutions of (1) will not be smooth enough, owing to shocks and other irregularities, to justify the foregoing computation. The new idea is instead to replace (27) with an *inequality*:

$$(28) \quad \Phi(\mathbf{u})_t + \Psi(\mathbf{u})_x \leq 0 \quad \text{in } \mathbb{R} \times (0, \infty).$$

In applications $\Phi(\mathbf{u})$ will sometimes be the negative of physical entropy and $\Psi(\mathbf{u})$ the entropy flux. The inequality (28) therefore asserts that entropy evolves according to its flux but may also undergo sharp increases, for instance along shocks.

Let us hereafter rigorously understand (28) to mean

$$(29) \quad \begin{cases} \int_0^\infty \int_{-\infty}^\infty \Phi(\mathbf{u})v_t + \Psi(\mathbf{u})v_x \, dxdt \geq 0 \\ \text{for each } v \in C_c^\infty(\mathbb{R} \times (0, \infty)), v \geq 0. \end{cases}$$

We consider once more the initial-value problem

$$(30) \quad \begin{cases} \mathbf{u}_t + \mathbf{F}(\mathbf{u})_x = 0 & \text{in } \mathbb{R} \times (0, \infty) \\ \mathbf{u} = \mathbf{g} & \text{on } \mathbb{R} \times \{t = 0\}. \end{cases}$$

DEFINITION. We call \mathbf{u} an entropy solution of (30) provided \mathbf{u} is an integral solution and \mathbf{u} satisfies the inequalities (29) for each entropy/entropy-flux pair (Φ, Ψ) .

Let us now attempt to build for general initial data \mathbf{g} an entropy solution. As in §11.4.1 we expect such a “physically correct” solution \mathbf{u} to be a limit of solutions \mathbf{u}^ε of approximating viscous problems

$$(31) \quad \begin{cases} \mathbf{u}_t^\varepsilon + \mathbf{F}(\mathbf{u}^\varepsilon)_x - \varepsilon \mathbf{u}_{xx}^\varepsilon = 0 & \text{in } \mathbb{R} \times (0, \infty) \\ \mathbf{u}^\varepsilon = \mathbf{g} & \text{on } \mathbb{R} \times \{t = 0\}. \end{cases}$$

We assume \mathbf{u}^ε is a smooth solution of (31), converging to 0 as $|x| \rightarrow \infty$ sufficiently rapidly to justify the calculations below. Let us further suppose $\{\mathbf{u}^\varepsilon\}_{0 < \varepsilon \leq 1}$ is uniformly bounded in L^∞ and furthermore

$$(32) \quad \mathbf{u}^\varepsilon \rightarrow \mathbf{u} \quad \text{a.e. as } \varepsilon \rightarrow 0$$

for some limit function \mathbf{u} . (In practice it is extremely difficult to verify this a.e. convergence.)

THEOREM 2 (Entropy and vanishing viscosity). *The function \mathbf{u} is an entropy solution of the conservation law (30).*

Proof. 1. Choose any smooth entropy/entropy-flux pair (Φ, Ψ) . Left multiplying (30) by $D\Phi(\mathbf{u}^\varepsilon)$ and recalling (26), we compute

$$(33) \quad \begin{aligned} \Phi(\mathbf{u}^\varepsilon)_t + \Psi(\mathbf{u}^\varepsilon)_x &= \varepsilon D\Phi(\mathbf{u}^\varepsilon) \mathbf{u}_{xx}^\varepsilon \\ &= \varepsilon \Phi(\mathbf{u}^\varepsilon)_{xx} - \varepsilon (D^2\Phi(\mathbf{u}^\varepsilon) \mathbf{u}_x^\varepsilon) \cdot \mathbf{u}_x^\varepsilon. \end{aligned}$$

As Φ is convex,

$$(34) \quad (D^2\Phi(\mathbf{u}^\varepsilon) \mathbf{u}_x^\varepsilon) \cdot \mathbf{u}_x^\varepsilon \geq 0.$$

2. Multiply (33) by $v \in C_c^\infty(\mathbb{R} \times (0, \infty))$, $v \geq 0$. We integrate by parts and discover that

$$\begin{aligned} \int_0^\infty \int_{-\infty}^\infty \Phi(\mathbf{u}^\varepsilon) v_t + \Psi(\mathbf{u}^\varepsilon) v_x \, dx dt \\ &= \int_0^\infty \int_{-\infty}^\infty \varepsilon (D^2\Phi(\mathbf{u}^\varepsilon) \mathbf{u}_x^\varepsilon) \cdot \mathbf{u}_x^\varepsilon v - \varepsilon \Phi(\mathbf{u}^\varepsilon) v_{xx} \, dx dt \\ &\geq - \int_0^\infty \int_{-\infty}^\infty \varepsilon \Phi(\mathbf{u}^\varepsilon) v_{xx} \, dx dt, \end{aligned}$$

the last inequality holding in view of (34) and the nonnegativity of v .

Now let $\varepsilon \rightarrow 0$. Recalling (32) and the Dominated Convergence Theorem, we obtain

$$\int_0^\infty \int_{-\infty}^\infty \Phi(\mathbf{u}) v_t + \Psi(\mathbf{u}) v_x \, dx dt \geq 0.$$

Thus \mathbf{u} verifies the entropy/entropy-flux inequalities (29). If Φ and Ψ are not smooth, we obtain the same conclusion after an approximation.

3. Finally fix $\mathbf{v} \in C_c^\infty(\mathbb{R} \times [0, \infty); \mathbb{R}^m)$ and take the dot product of the PDE in (31) with \mathbf{v} . After integrating by parts, we obtain

$$\int_0^\infty \int_{-\infty}^\infty \mathbf{u}^\varepsilon \cdot \mathbf{v}_t + \mathbf{F}(\mathbf{u}^\varepsilon) \mathbf{v}_x + \varepsilon \mathbf{u}^\varepsilon \cdot \mathbf{v}_{xx} \, dx dt + \int_{-\infty}^\infty \mathbf{g} \cdot \mathbf{v} \, dx|_{t=0} = 0.$$

We send $\varepsilon \rightarrow 0$, to deduce \mathbf{u} is an integral solution of (30). □

Example 1. In the case of a scalar conservation law (i.e. $m = 1$), for any convex Φ we can find a corresponding flux function Ψ , namely

$$\Psi(z) := \int_{z_0}^z \Phi'(w) F'(w) \, dw \quad (z \in \mathbb{R}).$$

See §11.4.3 following for an application. □

Example 2. For the p -system we have $m = 2$. To verify (25), (26) we must find Φ, Ψ , with Φ convex and

$$(\Phi_{z_1}, \Phi_{z_2}) \begin{pmatrix} 0 & -1 \\ -p'(z_1) & 0 \end{pmatrix} = \begin{pmatrix} \Psi_{z_1} \\ \Psi_{z_2} \end{pmatrix}.$$

A solution is

$$\Phi(z) = \frac{z_2^2}{2} + P(z_1), \quad \Psi(z) = -p(z_1)z_2 \quad (z \in \mathbb{R}^2),$$

where $P' = p$. Note Φ is convex, since $p' > 0$. □

See the exercises for further examples.

11.4.3. Uniqueness for scalar conservation laws.

As a further illustration of the ideas in §11.4.2, let us now consider again the initial-value problem for a *scalar* conservation law

$$(35) \quad \begin{cases} u_t + F(u)_x = 0 & \text{in } \mathbb{R} \times (0, \infty) \\ u = g & \text{on } \mathbb{R} \times \{t = 0\}. \end{cases}$$

Hence the unknown $u = u(x, t)$ is real-valued and $F : \mathbb{R} \rightarrow \mathbb{R}$ is a given smooth flux function.

In §3.4 we carefully studied the problem (35), making use of the primary assumption that F be strictly convex, to derive the Lax–Oleinik formula (see §3.4.2). Let us now drop the assumption that F be convex and devise an appropriate notion of weak solution. As above, we introduce entropies:

DEFINITION. Two smooth functions $\Phi, \Psi : \mathbb{R} \rightarrow \mathbb{R}$ comprise an entropy/entropy-flux pair for the conservation law $u_t + F(u)_x = 0$ provided

$$(36) \quad \Phi \text{ is convex}$$

and

$$(37) \quad \Phi'(z)F'(z) = \Psi'(z) \quad (z \in \mathbb{R}).$$

As noted in Example 1 above, for each convex Φ there exists a corresponding flux Ψ .

The entropy condition for u reads

$$\Phi(u)_t + \Psi(u)_x \leq 0 \quad \text{on } \mathbb{R} \times (0, \infty)$$

for each entropy/entropy-flux pair Φ, Ψ . This means

$$(38) \quad \begin{cases} \int_0^\infty \int_{-\infty}^\infty \Phi(u)v_t + \Psi(u)v_x \, dxdt \geq 0 \\ \text{for each } v \in C_c^\infty(\mathbb{R} \times (0, \infty)), \, v \geq 0. \end{cases}$$

DEFINITION. We call $u \in C([0, \infty), L^1(\mathbb{R})) \cap L^\infty(\mathbb{R} \times (0, \infty))$ an entropy solution of (35) provided u satisfies the inequalities (38) for each entropy/entropy-flux pair (Φ, Ψ) and $u(\cdot, t) \rightarrow g$ in L^1 as $t \rightarrow 0$.

Remarks. (i) This definition supersedes our earlier definition of “entropy solution” in §3.4.3.

(ii) Taking $\Phi(z) = \pm z$, $\Psi(z) = \pm F(z)$ in (38), we deduce

$$\int_0^\infty \int_{-\infty}^\infty uv_t + F(u)v_x \, dxdt = 0$$

for all $v \geq 0$ and thus for all $v \in C_c^1(\mathbb{R} \times (0, \infty))$. It is an exercise to prove then that

$$\int_0^\infty \int_{-\infty}^\infty uv_t + F(u)v_x \, dxdt + \int_{-\infty}^\infty gv \, dx|_{t=0} = 0$$

for all $v \in C_c^1(\mathbb{R} \times [0, \infty))$, since $u(\cdot, t) \rightarrow g$ in L^1 . Thus an entropy solution is an integral solution.

We discussed in §11.4.2 the construction of an entropy solution, and we now prove uniqueness.

THEOREM 3 (Uniqueness of entropy solutions for a single conservation law). *There exists—up to a set of measure zero—at most one entropy solution of (35).*

As in the proof of Theorem 1 in §10.2, the basic idea will be to “double the variables” in the problem.

Proof*. 1. Let u be an entropy solution of (35). Then

$$(39) \quad \int_0^\infty \int_{-\infty}^\infty \Phi(u)v_t + \Psi(u)v_x \, dxdt \geq 0$$

for all $v \in C_c^\infty(\mathbb{R} \times (0, \infty))$, $v \geq 0$, where Φ is smooth, convex and

$$\Psi(z) = \int_{z_0}^z \Phi'(w)F'(w) \, dw$$

for any z_0 . Fix $\alpha \in \mathbb{R}$ and take

$$(40) \quad \Phi_k(z) := \beta_k(z - \alpha) \quad (z \in \mathbb{R}),$$

*Omit on first reading.

where for each $k = 1, \dots$, the function $\beta_k : \mathbb{R} \rightarrow \mathbb{R}$ is smooth, convex and

$$\begin{cases} \beta_k(z) \rightarrow |z| & \text{uniformly} \\ \beta'_k(z) \rightarrow \text{sgn}(z) & \text{boundedly, a.e.} \end{cases}$$

Thus $\Phi_k(z) \rightarrow |z - \alpha|$ uniformly for $z \in \mathbb{R}$. A flux corresponding to (40) is

$$\Psi_k(z) = \int_{\alpha}^z \beta'_k(w - \alpha)F'(w) dw.$$

Consequently for each z

$$\Psi_k(z) \rightarrow \int_{\alpha}^z \text{sgn}(w - \alpha)F'(w) dw = \text{sgn}(z - \alpha)(F(z) - F(\alpha)).$$

Putting Φ_k, Ψ_k into (39) and sending $k \rightarrow \infty$, we deduce

$$(41) \quad \int_0^{\infty} \int_{-\infty}^{\infty} |u - \alpha|v_t + \text{sgn}(u - \alpha)(F(u) - F(\alpha))v_x dxdt \geq 0$$

for each $\alpha \in \mathbb{R}$ and v as above.

2. Next let \tilde{u} be another entropy solution. Then

$$(42) \quad \int_0^{\infty} \int_{-\infty}^{\infty} |\tilde{u} - \tilde{\alpha}|\tilde{v}_s + \text{sgn}(\tilde{u} - \tilde{\alpha})(F(\tilde{u}) - F(\tilde{\alpha}))\tilde{v}_y dyds \geq 0$$

where $\tilde{\alpha} \in \mathbb{R}$ and $\tilde{v} \in C_c^{\infty}(\mathbb{R} \times (0, \infty))$, $\tilde{v} \geq 0$.

Now let $w \in C_c^{\infty}(\mathbb{R} \times \mathbb{R} \times (0, \infty) \times (0, \infty))$, $w \geq 0$, $w = w(x, y, t, s)$. Fixing $(y, s) \in \mathbb{R} \times (0, \infty)$, we take $\alpha = \tilde{u}(y, s)$, $v(x, t) = w(x, y, t, s)$ in (41). Integrating with respect to y, s , we produce the inequality

$$(43) \quad \int_0^{\infty} \int_0^{\infty} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} |u(x, t) - \tilde{u}(y, s)|w_t + \text{sgn}(u(x, t) - \tilde{u}(y, s))(F(u(x, t)) - F(\tilde{u}(y, s)))w_x dx dy dt ds \geq 0.$$

Likewise, for each fixed $(x, t) \in \mathbb{R} \times (0, \infty)$ we take $\tilde{\alpha} = u(x, t)$, $\tilde{v}(y, s) = w(x, y, t, s)$ in (42). Integrating with respect to x, t gives

$$(44) \quad \int_0^{\infty} \int_0^{\infty} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} |\tilde{u}(y, s) - u(x, t)|w_s + \text{sgn}(\tilde{u}(y, s) - u(x, t))(F(\tilde{u}(y, s)) - F(u(x, t)))w_y dx dy dt ds \geq 0.$$

Add (43), (44):

$$(45) \quad \int_0^{\infty} \int_0^{\infty} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} |u(x, t) - \tilde{u}(y, s)|(w_t + w_s) + \text{sgn}(u(x, t) - \tilde{u}(y, s))(F(u(x, t)) - F(\tilde{u}(y, s)))(w_x + w_y) dx dy dt ds \geq 0.$$

3. We design as follows a clever choice for w in (45). Select η to be a standard mollifier as in §C.4 (with $n = 1$) and, as usual, write $\eta_\varepsilon(x) = \frac{1}{\varepsilon}\eta\left(\frac{x}{\varepsilon}\right)$. Take

$$w(x, y, t, s) := \eta_\varepsilon\left(\frac{x-y}{2}\right) \eta_\varepsilon\left(\frac{t-s}{2}\right) \phi\left(\frac{x+y}{2}, \frac{t+s}{2}\right),$$

where $\phi \in C_c^\infty(\mathbb{R} \times (0, \infty))$, $\phi \geq 0$. We insert this choice of w into (45) and thereby obtain

$$(46) \quad \int_0^\infty \int_0^\infty \int_{-\infty}^\infty \int_{-\infty}^\infty \left\{ |u(x, t) - \tilde{u}(y, s)| \phi_t\left(\frac{x+y}{2}, \frac{t+s}{2}\right) \right. \\ \left. + \operatorname{sgn}(u(x, t) - \tilde{u}(y, s)) (F(u(x, t)) - F(\tilde{u}(y, s))) \phi_x\left(\frac{x+y}{2}, \frac{t+s}{2}\right) \right\} \\ \eta_\varepsilon\left(\frac{x-y}{2}\right) \eta_\varepsilon\left(\frac{t-s}{2}\right) dx dy dt ds \geq 0.$$

Change variables by writing

$$\begin{cases} \bar{x} = \frac{x+y}{2}, \bar{t} = \frac{t+s}{2} \\ \bar{y} = \frac{x-y}{2}, \bar{s} = \frac{t-s}{2}. \end{cases}$$

Then (46) implies

$$(47) \quad \int_{-\infty}^\infty \int_{-\infty}^\infty G(\bar{y}, \bar{s}) \eta_\varepsilon(\bar{y}) \eta_\varepsilon(\bar{s}) d\bar{y} d\bar{s} \geq 0,$$

where

$$(48) \quad G(\bar{y}, \bar{s}) := \int_0^\infty \int_{-\infty}^\infty |u(\bar{x} + \bar{y}, \bar{t} + \bar{s}) - \tilde{u}(\bar{x} - \bar{y}, \bar{t} - \bar{s})| \phi_t(\bar{x}, \bar{t}) \\ + \operatorname{sgn}(u(\bar{x} + \bar{y}, \bar{t} + \bar{s}) - \tilde{u}(\bar{x} - \bar{y}, \bar{t} - \bar{s})) \\ (F(u(\bar{x} + \bar{y}, \bar{t} + \bar{s})) - F(\tilde{u}(\bar{x} - \bar{y}, \bar{t} - \bar{s}))) \phi_x(\bar{x}, \bar{t}) d\bar{x} d\bar{t}.$$

Now $u(\bar{x} + \bar{y}, \bar{t} + \bar{s}) \rightarrow u(\bar{x}, \bar{t})$, $\tilde{u}(\bar{x} - \bar{y}, \bar{t} - \bar{s}) \rightarrow \tilde{u}(\bar{x}, \bar{t})$ in L^1_{loc} as $\bar{y}, \bar{s} \rightarrow 0$. Since the mappings $(a, b) \mapsto |a - b|$, $\operatorname{sgn}(a - b)(F(a) - F(b))$ are Lipschitz continuous, we deduce upon letting $\varepsilon \rightarrow 0$ in (47) that

$$\int_0^\infty \int_{-\infty}^\infty |u(\bar{x}, \bar{t}) - \tilde{u}(\bar{x}, \bar{t})| \phi_t(\bar{x}, \bar{t}) \\ + \operatorname{sgn}(u(\bar{x}, \bar{t}) - \tilde{u}(\bar{x}, \bar{t})) (F(u(\bar{x}, \bar{t})) - F(\tilde{u}(\bar{x}, \bar{t}))) \phi_x(\bar{x}, \bar{t}) d\bar{x} d\bar{t} \geq 0.$$

Rewriting $x = \bar{x}$, $t = \bar{t}$, we have therefore

$$(49) \quad \int_0^\infty \int_{-\infty}^\infty a(x, t) \phi_t(x, t) + b(x, t) \phi_x(x, t) dx dt \geq 0,$$

for

$$\begin{cases} a(x, t) := |u(x, t) - \tilde{u}(x, t)| \\ b(x, t) := \operatorname{sgn}(u(x, t) - \tilde{u}(x, t))(F(u(x, t)) - F(\tilde{u}(x, t))). \end{cases}$$

4. We now employ the inequality (49) to establish the L^1 -contraction inequalities

$$(50) \quad \begin{cases} \int_{-\infty}^{\infty} |u(x, t) - \tilde{u}(x, t)| \, dx \leq \int_{-\infty}^{\infty} |u(x, s) - \tilde{u}(x, s)| \, dx \\ \text{for a.e. } 0 \leq s \leq t. \end{cases}$$

To prove this assertion, we take $0 < s < t$, $r > 0$ and let $\phi(x, t) = \alpha(x)\beta(t)$ in (49), where

$$\begin{cases} \alpha : \mathbb{R} \rightarrow \mathbb{R} \text{ is smooth,} \\ \alpha(x) = 1 \text{ if } |x| \leq r, \alpha(x) = 0 \text{ if } |x| \geq r + 1, \\ |\alpha'(x)| \leq 2 \end{cases}$$

and

$$\begin{cases} \beta : \mathbb{R} \rightarrow \mathbb{R} \text{ is Lipschitz,} \\ \beta(\tau) = 0 \text{ if } 0 \leq \tau \leq s \text{ or } \tau \geq t + \delta, \\ \beta(\tau) = 1 \text{ if } s + \delta \leq \tau \leq t, \\ \beta \text{ is linear on } [s, s + \delta] \text{ and } [t, t + \delta], \end{cases}$$

for $0 < \delta < t - s$. We deduce

$$\begin{aligned} \frac{1}{\delta} \int_s^{s+\delta} \int_{-\infty}^{\infty} a(x, \tau) \alpha(x) \, dx d\tau &\geq \frac{1}{\delta} \int_t^{t+\delta} \int_{-\infty}^{\infty} a(x, \tau) \alpha(x) \, dx d\tau \\ &\quad - \int_s^{t+\delta} \int_{\{r \leq |x| \leq r+1\}} b(x, \tau) \alpha'(x) b(\tau) \, dx d\tau. \end{aligned}$$

Let $r \rightarrow \infty$:

$$\frac{1}{\delta} \int_t^{t+\delta} \int_{-\infty}^{\infty} a(x, \tau) \, dx d\tau \leq \frac{1}{\delta} \int_s^{s+\delta} \int_{-\infty}^{\infty} a(x, \tau) \, dx d\tau.$$

Next let $\delta \rightarrow 0$ to deduce (50) for a.e. $0 \leq s \leq t$.

5. In light of (50) and the fact that $u(\cdot, t), \tilde{u}(\cdot, t) \rightarrow g$ in L^1 as $t \rightarrow 0$, we at last conclude $u = \tilde{u}$ a.e. □

11.5. PROBLEMS

1. Show that this generalization of the p -system is a strictly hyperbolic system if $p', q' > 0$:

$$\begin{cases} u_t^1 - q(u^2)_x = 0 \\ u_t^2 - p(u^1)_x = 0. \end{cases}$$

2. (a) Verify that the shallow water equations (Example 3 in §11.1) form a strictly hyperbolic system, provided $h > 0$.
 (b) Show that for a smooth solution $(h, q) = (h, vh)$, with $h > 0$, the shallow water equations can be recast into this alternate conservation law form:

$$(*) \quad \begin{cases} h_t + (vh)_x = 0 \\ v_t + \left(\frac{v^2}{2} + h\right)_x = 0. \end{cases}$$

Check that this is a strictly hyperbolic system.

3. Define for $z \in \mathbb{R}$, $z \neq 0$, the matrix function

$$\mathbf{B}(z) := e^{-\frac{1}{z^2}} \begin{pmatrix} \cos(\frac{2}{z}) & \sin(\frac{2}{z}) \\ \sin(\frac{2}{z}) & -\cos(\frac{2}{z}) \end{pmatrix},$$

and set $\mathbf{B}(0) = 0$. Show that \mathbf{B} is C^∞ and has real eigenvalues, but we cannot find unit-length right eigenvectors $\{\mathbf{r}_1(z), \mathbf{r}_2(z)\}$ depending continuously on z near 0. What happens to the eigenspaces as $z \rightarrow 0$?

4. (Rarefaction curves) Show that for the shallow water wave equations in the form (*) from Problem 2 the rarefaction curves R_1, R_2 in the (h, v) -plane are given by the formulas

$$2\sqrt{h} \pm v = c,$$

where c is a constant.

5. (Shock curves) For the shallow water wave equations in the variables (h, q) , find a formula describing the shock set $S(z_0)$ in the (h, q) -plane, where $z_0 = (h_0, q_0)$ and $h_0 > 0$. For the particular case that $q_0 = 0$, show that the shock set is given by the expression

$$q = \pm \frac{(h - h_0)h}{\sqrt{2}} \left(\frac{1}{h} + \frac{1}{h_0} \right)^{\frac{1}{2}}.$$

6. (Continuation) Construct an entropy solution $\mathbf{u} = (h, q)$ of the shallow water equations, corresponding to the Riemann initial data $u_l = (h_l, 0)$ for $x < 0$ and $u_r = (h_r, 0)$ for $x > 0$, where $h_l > h_r > 0$.

Your answer should consist of a rarefaction wave connecting u_l to an intermediate state $u_i = (h_i, q_i) = (h_i, v_i h_i)$ and a shock wave connecting u_i to u_r . To do this, transfer the formula from Problem 5 to the (h, v) -plane, and show that there exists a point of intersection (h_i, v_i) of a rarefaction curve through $(h_l, 0)$ and a shock curve through $(h_r, 0)$.

7. Confirm that the functions w^1, w^2 computed for the barotropic gas dynamics in §11.3.1 are indeed Riemann invariants.
8. Suppose that Φ is an entropy for the shallow water equations in the form (*) from Problem 2. Prove

$$\frac{\partial^2 \Phi}{\partial v^2} = h \frac{\partial^2 \Phi}{\partial h^2}.$$

9. Show that $\Phi = \rho v^2/2 + P(\rho)$ is an entropy for the barotropic Euler equations (from §11.3.1), provided $P''(\rho) = p'(\rho)/\rho$, $\rho > 0$. Confirm that Φ is convex in the proper variables. What is the corresponding entropy flux Ψ ?
10. (Gradient flux function) Suppose that $\mathbf{F} = D\phi$, where $\phi : \mathbb{R}^m \rightarrow \mathbb{R}$. Show that

$$\Phi := \frac{1}{2}|z|^2, \quad \Psi := z \cdot D\phi(z) - \phi(z)$$

form an entropy/entropy-flux pair for the system of conservation laws $\mathbf{u}_t + \mathbf{F}(\mathbf{u})_x = 0$.

11. (Antigradient flux function)
 - (a) Assume that $m = 2$ and \mathbf{F} has the “antigradient” form $\mathbf{F} = (\phi_{z_2}, \phi_{z_1})$ for a convex function $\phi : \mathbb{R}^2 \rightarrow \mathbb{R}$. Confirm that

$$\Phi := \phi, \quad \Psi := \phi_{z_1} \phi_{z_2}$$

are an entropy/entropy-flux pair for the system $\mathbf{u}_t + \mathbf{F}(\mathbf{u})_x = 0$.

- (b) Find an entropy/entropy-flux pair for the generalization of the p -system given in Problem 1.
12. Formulate what it means for Φ, Ψ to be an entropy/entropy-flux pair for the general system of conservation laws

$$\mathbf{u}_t + \operatorname{div} \mathbf{F}(\mathbf{u}) = 0 \quad \text{in } \mathbb{R}^n \times (0, \infty).$$

13. *Maxwell's equations for nonlinear dielectrics* read in part

$$\begin{cases} \mathbf{D}_t = \operatorname{curl} \mathbf{H} \\ \mathbf{B}_t = -\operatorname{curl} \mathbf{E} \end{cases}$$

in $\mathbb{R}^3 \times (0, \infty)$, where \mathbf{E} denotes the electric field; \mathbf{H} the magnetic field; \mathbf{D} the electric displacement; and \mathbf{B} the magnetic induction. (We suppose the current \mathbf{J} vanishes.) We introduce the constitutive relations

$$\mathbf{H} = D_B \eta(\mathbf{B}, \mathbf{D}), \quad \mathbf{E} = D_D \eta(\mathbf{B}, \mathbf{D}),$$

determined by a given convex electromagnetic field energy $\eta = \eta(B, D)$.

Show that

$$\Phi := \eta, \quad \Psi := D_D \eta \times D_B \eta$$

are an entropy/entropy-flux pair.

14. (Nonconvex flux function) Assume that u is an entropy solution of the scalar conservation law $u_t + F(u)_x = 0$ and that, as in §3.4.1, u is smooth on either side of a curve $\{x = s(t)\}$.

(a) Prove that along this curve the left- and right-hand limits of u satisfy

$$F(\lambda u_l + (1 - \lambda)u_r) \geq \lambda F(u_l) + (1 - \lambda)F(u_r) \quad \text{if } u_l \leq u_r$$

and

$$F(\lambda u_l + (1 - \lambda)u_r) \leq \lambda F(u_l) + (1 - \lambda)F(u_r) \quad \text{if } u_r \leq u_l$$

for each $0 \leq \lambda \leq 1$. These inequalities are called *condition E*. Draw pictures illustrating the geometric meaning of these inequalities.

(Compare with Problem 7 in Chapter 10.)

(b) What does condition *E* imply if F is uniformly convex?

15. (Explicit solutions for nonconvex flux) Compute the unique entropy solution of the Riemann problem

$$(*) \quad \begin{cases} u_t + (u^3 - u)_x = 0 & \text{in } \mathbb{R} \times (0, \infty) \\ u = g & \text{on } \mathbb{R} \times \{t = 0\}, \end{cases}$$

for

$$g(x) = \begin{cases} -1 & \text{if } x < 0 \\ 1 & \text{if } x > 0. \end{cases}$$

(Hint: The solution involves a rarefaction wave with a shock along its left edge. Make sure your solution verifies condition *E*, introduced in the previous problem.)

16. (Continuation) Discuss the structure of the entropy solution of the conservation law (*) from Problem 15, for

$$g(x) = \begin{cases} -1 + \lambda & \text{if } x < -1 \\ -1 - \lambda x & \text{if } -1 < x < 0 \\ 1 & \text{if } x > 0, \end{cases}$$

when $\lambda > 0$ is small. How does this perturbation affect the behavior of the shock in the solution found in the previous problem?

17. Assume u is a smooth solution with compact support in space of the scalar conservation law in several variables

$$\begin{cases} u_t + \operatorname{div} \mathbf{F}(u) = 0 & \text{in } \mathbb{R}^n \times (0, \infty) \\ u = g & \text{on } \mathbb{R}^n \times \{t = 0\}, \end{cases}$$

where $\mathbf{F} = (F^1, \dots, F^n)$. Show that if \hat{u} is another smooth solution with initial data \hat{g} , we have the L^1 -contraction estimate

$$\|u(\cdot, t) - \hat{u}(\cdot, t)\|_{L^1(\mathbb{R}^n)} \leq \|g - \hat{g}\|_{L^1(\mathbb{R}^n)}$$

for each time $t \geq 0$.

(Hints: Show that $w := u - \hat{u}$ satisfies $w_t + \operatorname{div}(\mathbf{b}w) = 0$ for an appropriate vector field $\mathbf{b} = \mathbf{b}(x, t)$. Let $T > 0$ and introduce the solution v of the adjoint problem $v_t + \mathbf{b} \cdot Dv = 0$ in $\mathbb{R}^n \times [0, T]$, with the terminal condition $v(\cdot, T) = \operatorname{sgn}(w(\cdot, T))$.)

11.6. REFERENCES

T.-P. Liu wrote the initial draft of this entire chapter.

- Section 11.1 The shallow water equations are discussed in Holden–Risebro [H-R] and LeVeque [LV]. P. Colella helped me with the calculations in §11.1.2. The proof of Theorem 2 follows suggestions of W. Han. See Bresson [Bs], Courant–Friedrichs [C-F], Dafermos [D], Majda [Mj], Xiao–Zhang [X-Z], and Zheng [Zh] for more.
- Section 11.3 I followed Logan [Lo] for the example. Consult Neu [Ne] for more on using Riemann invariants.
- Section 11.4 See Smoller [S] for more about viscous traveling waves. The uniqueness theorem in §11.4.3 is due to S. Kružkov (Mat. Sbornik 123 (1970), 228–255).
- Section 11.5 Problem 3 is taken from Kato [K]. Problems 4–6 are based upon Holden–Risebro [H-R]. Problems 10 and 11 are from Smoller [S], and Problem 13 is from Dafermos [D].

This page is intentionally left blank.

NONLINEAR WAVE EQUATIONS

- 12.1 Introduction
- 12.2 Existence of solutions
- 12.3 Semilinear wave equations
- 12.4 Critical power nonlinearity
- 12.5 Nonexistence of solutions
- 12.6 Problems
- 12.7 References

12.1. INTRODUCTION

This final chapter studies the existence (or sometimes nonexistence) of solutions to the initial-value problem for the semilinear wave equation

$$(1) \quad u_{tt} - \Delta u + f(u) = 0.$$

The linear case that $f(u) = m^2u$ is the *Klein–Gordon* equation. We will also discuss certain mildly quasilinear wave equations having the form

$$(2) \quad u_{tt} - \Delta u + f(Du, u_t, u) = 0,$$

where as usual we write $Du = D_x u$ for the gradient in the x -variables.

We follow the custom of putting the nonlinearity on the left of the equals sign in (1) and (2): this simplifies some later formulas a bit. More complicated quasilinear wave equations, in which the coefficients of the second-order derivatives depend on Du, u_t, u , are beyond the scope of this book.

12.1.1. Conservation of energy.

Considering first the semilinear PDE (1), we hereafter set

$$F(z) := \int_0^z f(w) dw \quad (z \in \mathbb{R}).$$

Then $F(0) = 0$ and $F' = f$. We recall from §8.6.2 that the *energy* of a solution u of (1) at time $t \geq 0$ is

$$E(t) := \int_{\mathbb{R}^n} \frac{1}{2}(u_t^2 + |Du|^2) + F(u) dx$$

and that this energy is conserved:

THEOREM 1 (Conservation of energy). *Assume that u is a smooth solution of the semilinear wave equation (1) and that $u(\cdot, t)$ has compact support in space for each time t . Then*

$$t \mapsto E(t) \quad \text{is constant.}$$

Proof. We calculate

$$\dot{E}(t) = \int_{\mathbb{R}^n} u_t u_{tt} + Du \cdot Du_t + f(u)u_t dx = \int_{\mathbb{R}^n} u_t(u_{tt} - \Delta u + f(u)) dx = 0,$$

where $\dot{\cdot} = \frac{d}{dt}$. □

The integration by parts in this proof is valid, since u has compact support in space for each time. In many subsequent proofs we will similarly integrate by parts, implicitly relying upon our solution's vanishing for large $|x|$ to justify the computation.

12.1.2. Finite propagation speed.

Recalling the domain of dependence calculation for the linear wave equation in §2.4.3, we reintroduce the backwards wave cone:

DEFINITION. *Fix $x_0 \in \mathbb{R}^n$, $t_0 > 0$ and define the backwards wave cone with apex (x_0, t_0) :*

$$K(x_0, t_0) := \{(x, t) \mid 0 \leq t \leq t_0, |x - x_0| \leq t_0 - t\}.$$

The curved part of the boundary of $K(x_0, t_0)$ is

$$\Gamma(x_0, t_0) := \{(x, t) \mid 0 \leq t \leq t_0, |x - x_0| = t_0 - t\}.$$

THEOREM 2 (Flux estimate for semilinear wave equation). *Assume that u is a smooth solution of the semilinear wave equation (1).*

(i) *For each point $(x_0, t_0) \in \mathbb{R}^n \times (0, \infty)$ we have the identity*

$$(3) \quad \frac{1}{\sqrt{2}} \int_{\Gamma(x_0, t_0)} \frac{1}{2} |u_t \nu - Du|^2 + F(u) dS = e(0),$$

where $\nu := \frac{x-x_0}{|x-x_0|}$ and

$$e(t) := \int_{B(x_0, t_0-t)} \frac{1}{2} (u_t^2 + |Du|^2) + F(u) dx \quad (0 \leq t \leq t_0).$$

(ii) *If*

$$F \geq 0$$

and

$$u(\cdot, 0), u_t(\cdot, 0) \equiv 0 \quad \text{within } B(x_0, t_0),$$

then $u \equiv 0$ within the cone $K(x_0, t_0)$.

The expression on the left-hand side of (3) is the *energy flux* through the curved surface $\Gamma(x_0, t_0)$.

Proof. 1. We compute that

$$(4) \quad \begin{aligned} \dot{e}(t) &= \int_{B(x_0, t_0-t)} u_t u_{tt} + Du \cdot Du_t + f(u) u_t dx \\ &\quad - \int_{\partial B(x_0, t_0-t)} \frac{1}{2} (u_t^2 + |Du|^2) + F(u) dS \\ &= \int_{\partial B(x_0, t_0-t)} \frac{\partial u}{\partial \nu} u_t - \frac{1}{2} (u_t^2 + |Du|^2) - F(u) dS \\ &= - \int_{\partial B(x_0, t_0-t)} \frac{1}{2} |u_t \nu - Du|^2 + F(u) dS, \end{aligned}$$

since

$$|u_t \nu - Du|^2 = u_t^2 - 2u_t \frac{\partial u}{\partial \nu} + |Du|^2.$$

Now integrate in time between 0 and t_0 to derive (3). Notice that the factor $\frac{1}{\sqrt{2}}$ appears when we switch to integration over $\Gamma(x_0, t_0)$, since this surface is tilted at constant angle $\frac{\pi}{4}$ above $B(x_0, t_0) \times \{t = 0\}$.

2. If $u(\cdot, 0), u_t(\cdot, 0) \equiv 0$ on $B(x_0, t_0)$, then $e(0) = 0$ since $F(0) = 0$. As $F \geq 0$, it follows from (4) that $e \equiv 0$. We deduce that $u_t, Du \equiv 0$, and therefore $u \equiv 0$, within the cone $K(x_0, t_0)$. \square

Although the mildly quasilinear wave equation (2) does not in general have a conserved energy, we can nevertheless adapt the previous proof to show finite propagation speed.

THEOREM 3 (Domain of dependence). *Assume that*

$$f(0, 0, 0) = 0$$

and that u is a smooth solution of the quasilinear wave equation (2). If

$$u(\cdot, 0), u_t(\cdot, 0) \equiv 0 \quad \text{within } B(x_0, t_0),$$

then $u \equiv 0$ within the cone $K(x_0, t_0)$.

So any disturbance originating outside $B(x_0, t_0)$ does not affect the solution within $K(x_0, t_0)$. Consequently the effects of nonzero initial data propagate with speed at most one.

Proof. Define

$$e(t) := \frac{1}{2} \int_{B(x_0, t_0-t)} u_t^2 + |Du|^2 + u^2 dx \quad (0 \leq t \leq t_0).$$

Then

$$\begin{aligned} \dot{e}(t) &= \int_{B(x_0, t_0-t)} u_t u_{tt} + Du \cdot Du_t + u u_t dx \\ &\quad - \frac{1}{2} \int_{\partial B(x_0, t_0-t)} u_t^2 + |Du|^2 + u^2 dS \\ &= \int_{B(x_0, t_0-t)} u_t (u_{tt} - \Delta u + u) dx \\ &\quad + \int_{\partial B(x_0, t_0-t)} \frac{\partial u}{\partial \nu} u_t dS - \frac{1}{2} \int_{\partial B(x_0, t_0-t)} u_t^2 + |Du|^2 + u^2 dS \\ &\leq \int_{B(x_0, t_0-t)} u_t (-f(Du, u_t, u) + u) dx. \end{aligned}$$

Since $f(0, 0, 0) = 0$ and u is smooth,

$$|f(Du, u_t, u)| \leq C(|Du| + |u_t| + |u|)$$

for some constant C depending upon $\|Du, u_t, u\|_{L^\infty}$. We conclude that

$$\dot{e}(t) \leq C \int_{B(x_0, t_0-t)} u_t^2 + |Du|^2 + u^2 dx = Ce(t).$$

As $e(0) = 0$, Gronwall's inequality (§B.2) implies $e \equiv 0$. Therefore $u \equiv 0$ within the cone $K(x_0, t_0)$. \square

12.2. EXISTENCE OF SOLUTIONS

We devote this section to proving existence theorems for solutions of the mildly quasilinear initial-value problem

$$(1) \quad \begin{cases} u_{tt} - \Delta u + f(Du, u_t, u) = 0 & \text{in } \mathbb{R}^n \times (0, T] \\ u = g, \quad u_t = h & \text{on } \mathbb{R}^n \times \{t = 0\}. \end{cases}$$

Similarly to §7.2, we say that a function $\mathbf{u} \in L^2(0, T; H_{\text{loc}}^1(\mathbb{R}^n))$, with $\mathbf{u}' \in L^2(0, T; L_{\text{loc}}^2(\mathbb{R}^n))$ and $\mathbf{u}'' \in L^2(0, T; H_{\text{loc}}^{-1}(\mathbb{R}^n))$, is a weak solution of the initial-value problem (1) provided $\mathbf{u}(0) = g$, $\mathbf{u}'(0) = h$,

$$\mathbf{f} := -f(D\mathbf{u}, \mathbf{u}', \mathbf{u}) \in L^2(0, T; L_{\text{loc}}^2(\mathbb{R}^n)),$$

and

$$\langle \mathbf{u}'', v \rangle + B[\mathbf{u}, v] = (\mathbf{f}, v)$$

for each $v \in H^1(\mathbb{R}^n)$ with compact support and a.e. time $0 \leq t \leq T$. Here $B[u, v] := \int_{\mathbb{R}^n} Du \cdot Dv \, dx$.

We always assume

$$f(0, 0, 0) = 0.$$

12.2.1. Lipschitz nonlinearities.

We start with a strong assumption on the nonlinearity, namely that

$$(2) \quad f : \mathbb{R}^n \times \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R} \text{ is Lipschitz continuous.}$$

THEOREM 1 (Existence and uniqueness).

(i) Assume $g \in H_{\text{loc}}^1(\mathbb{R}^n)$, $h \in L_{\text{loc}}^2(\mathbb{R}^n)$. Then for each $T > 0$ there exists a unique weak solution \mathbf{u} of the initial-value problem (1).

(ii) If in addition $g \in H_{\text{loc}}^2(\mathbb{R}^n)$, $h \in H_{\text{loc}}^1(\mathbb{R}^n)$, then

$$\begin{cases} \mathbf{u} \in L^\infty((0, T); H_{\text{loc}}^2(\mathbb{R}^n)), \\ \mathbf{u}' \in L^\infty((0, T); H_{\text{loc}}^1(\mathbb{R}^n)), \\ \mathbf{u}'' \in L^\infty((0, T); L_{\text{loc}}^2(\mathbb{R}^n)). \end{cases}$$

Proof. 1. We will first suppose that $T > 0$ is sufficiently small, as determined below. Given $R \gg 1$, let us consider first the initial/boundary-value problem

$$(3) \quad \begin{cases} u_{tt} - \Delta u + f(Du, u_t, u) = 0 & \text{in } B(0, R) \times (0, T] \\ u = 0 & \text{on } \partial B(0, R) \times [0, T] \\ u = g, \quad u_t = h & \text{on } B(0, R) \times \{t = 0\}. \end{cases}$$

We temporarily also assume $g \in H_0^1(B(0, R))$.

Introduce the space of functions

$$X := \{\mathbf{u} \in L^\infty(0, T; H_0^1(B(0, R))) \mid \mathbf{u}' \in L^\infty(0, T; L^2(B(0, R)))\},$$

with the norm

$$\|\mathbf{u}\| := \operatorname{ess\,sup}_{0 \leq t \leq T} (\|\mathbf{u}(t)\|_{H_0^1(B(0, R))} + \|\mathbf{u}'(t)\|_{L^2(B(0, R))}).$$

Given $\mathbf{v} \in X$, we hereafter write $\mathbf{u} = A[\mathbf{v}]$ to mean that $\mathbf{u} \in X$ is the unique weak solution of the linear problem

$$(4) \quad \begin{cases} u_{tt} - \Delta u = -f(Dv, v_t, v) & \text{in } B(0, R) \times (0, T) \\ u = 0 & \text{on } \partial B(0, R) \times (0, T) \\ u = g, \quad u_t = h & \text{on } B(0, R) \times \{t = 0\}. \end{cases}$$

This weak solution exists according to Theorems 3–5 in §7.2.

Suppose that we are given also $\hat{\mathbf{v}} \in X$, and likewise write $\hat{\mathbf{u}} = A[\hat{\mathbf{v}}]$. Put $\mathbf{w} := \mathbf{u} - \hat{\mathbf{u}}$. Then \mathbf{w} is the unique weak solution of

$$\begin{cases} w_{tt} - \Delta w = f(D\hat{v}, \hat{v}_t, \hat{v}) - f(Dv, v_t, v) & \text{in } B(0, R) \times (0, T) \\ w = 0 & \text{on } \partial B(0, R) \times (0, T) \\ w = 0, \quad w_t = 0 & \text{on } B(0, R) \times \{t = 0\}. \end{cases}$$

Consequently estimate (50) from §7.2 provides us with the bound

$$\|\mathbf{w}\| \leq C \|f(D\hat{\mathbf{v}}, \hat{\mathbf{v}}', \hat{\mathbf{v}}) - f(D\mathbf{v}, \mathbf{v}', \mathbf{v})\|_{L^2(0, T; L^2(B(0, R)))}.$$

In view of the Lipschitz continuity of f , it follows that

$$\|\mathbf{w}\|^2 \leq C \int_0^T \int_{B(0, R)} |D\mathbf{v} - D\hat{\mathbf{v}}|^2 + |\mathbf{v}' - \hat{\mathbf{v}}'|^2 + |\mathbf{v} - \hat{\mathbf{v}}|^2 \, dxdt \leq CT \|\mathbf{v} - \hat{\mathbf{v}}\|^2.$$

Since $\mathbf{w} = \mathbf{u} - \hat{\mathbf{u}} = A[\mathbf{v}] - A[\hat{\mathbf{v}}]$, we deduce that

$$\|A[\mathbf{v}] - A[\hat{\mathbf{v}}]\| \leq \frac{1}{2} \|\mathbf{v} - \hat{\mathbf{v}}\|$$

provided $T > 0$ is small enough, depending only upon the Lipschitz constant of f . Banach's Theorem (§9.2.1) now implies the existence of a unique fixed point $\mathbf{u} \in X$, which is the unique weak solution of (3).

2. In particular we may assume that $T < 1$. Let $S = R - 1$. Then the finite propagation speed (Theorem 3 in §12.1) implies that the solution

12.2.2. Short time existence.

Consider again the problem

$$(5) \quad \begin{cases} u_{tt} - \Delta u + f(Du, u_t, u) = 0 & \text{in } \mathbb{R}^n \times (0, T] \\ u = g, u_t = h & \text{on } \mathbb{R}^n \times \{t = 0\}. \end{cases}$$

We hereafter drop the restrictive assumption (2) that the nonlinearity f be Lipschitz continuous and instead just suppose that f is a given smooth function. Our goal is proving there is a unique solution, existing for at least some short time interval $[0, T]$. We will need more smoothness on the initial data, requiring $g \in H^k(\mathbb{R}^n)$, $h \in H^{k-1}(\mathbb{R}^n)$ for a possibly large integer k (depending on n).

We first introduce some new estimates for the Sobolev space $H^k(\mathbb{R}^n)$:

THEOREM 2 (Sobolev inequalities for H^k). *Suppose that the functions u_1, \dots, u_m belong to $H^k(\mathbb{R}^n)$, where $k > \frac{n}{2}$.*

(i) *If $|\beta_1| + \dots + |\beta_m| \leq k$, then*

$$(6) \quad \|D^{\beta_1} u_1 \cdots D^{\beta_m} u_m\|_{L^2(\mathbb{R}^n)} \leq C \prod_{j=1}^m \|u_j\|_{H^k(\mathbb{R}^n)}$$

for a constant $C = C(n, m, k)$.

(ii) *Let $f : \mathbb{R}^m \rightarrow \mathbb{R}$ be a smooth function satisfying $f(0) = 0$. Then $f(u_1, \dots, u_m) \in H^k(\mathbb{R}^n)$ and*

$$(7) \quad \|f(u_1, \dots, u_m)\|_{H^k(\mathbb{R}^n)} \leq \Phi(\|u_1\|_{H^k(\mathbb{R}^n)}, \dots, \|u_m\|_{H^k(\mathbb{R}^n)}),$$

where Φ is a continuous function, nondecreasing in each argument and depending only upon f, k, n, m .

Proof. 1. We leave the proof of (6) to the reader: see Problem 12.

2. Let $0 < |\alpha| \leq k$. Then $D^\alpha f(u_1, \dots, u_m)$ can be written as a finite sum of terms of the form

$$AD^{\beta_1} u_{j_1} \cdots D^{\beta_l} u_{j_l},$$

where A depends on the partial derivatives of f of order at most k evaluated at (u_1, \dots, u_m) , $l \leq k$, $0 \leq |\beta_j| \leq |\alpha|$, and $\beta_1 + \dots + \beta_l = \alpha$.

Recall from Theorem 6 in §5.6.3 that $k > \frac{n}{2}$ implies the estimate

$$\|u\|_{L^\infty(\mathbb{R}^n)} \leq C\|u\|_{H^k(\mathbb{R}^n)}.$$

Therefore $\|A\|_{L^\infty}$ is bounded by a term depending only upon f and $\|u_j\|_{H^k}$ ($j = 1, \dots, m$). According then to estimate (6),

$$\|AD^{\beta_1}u_{j_1} \cdots D^{\beta_l}u_{j_l}\|_{L^2}$$

is bounded by an expression involving only $\|u_j\|_{H^k}$, for $j = 1, \dots, m$. Since $f(0) = 0$, we can similarly estimate $\|f(u_1, \dots, u_m)\|_{L^2}$. This establishes (7). \square

We next demonstrate that if we select the time $T > 0$ sufficiently small, depending upon the initial data g, h , we can find a solution existing on $\mathbb{R}^n \times (0, T]$:

THEOREM 3 (Short time existence). *Assume $f : \mathbb{R}^n \times \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$ is smooth, $f(0, 0, 0) = 0$. Suppose also $g \in H^k(\mathbb{R}^n)$, $h \in H^{k-1}(\mathbb{R}^n)$ for $k > \frac{n}{2} + 1$.*

There exists a time $T > 0$ such that the initial-value problem

$$(8) \quad \begin{cases} u_{tt} - \Delta u + f(Du, u_t, u) = 0 & \text{in } \mathbb{R}^n \times (0, T] \\ u = g, u_t = h & \text{on } \mathbb{R}^n \times \{t = 0\} \end{cases}$$

has a unique weak solution \mathbf{u} , with

$$\mathbf{u} \in L^\infty(0, T; H^k(\mathbb{R}^n)), \mathbf{u}' \in L^\infty(0, T; H^{k-1}(\mathbb{R}^n)).$$

The time T of existence provided by the proof depends in a complicated way upon both f and $\|g\|_{H^k(\mathbb{R}^n)}, \|h\|_{H^{k-1}(\mathbb{R}^n)}$ and can be very short if $\|g\|_{H^k(\mathbb{R}^n)}$ and $\|h\|_{H^{k-1}(\mathbb{R}^n)}$ are large. We will see in §12.5 that solutions of even the simpler semilinear wave equation need not exist for all time.

Proof. 1. Let

$$X := \{\mathbf{u} \in L^\infty(0, T; H^1(\mathbb{R}^n)) \mid \mathbf{u}' \in L^\infty(0, T; L^2(\mathbb{R}^n))\}$$

with the norm

$$\|\mathbf{u}\| := \operatorname{ess\,sup}_{0 \leq t \leq T} (\|\mathbf{u}(t)\|_{H^1(\mathbb{R}^n)} + \|\mathbf{u}'(t)\|_{L^2(\mathbb{R}^n)}).$$

We introduce also the stronger norm

$$\|\|\mathbf{u}\|\| := \operatorname{ess\,sup}_{0 \leq t \leq T} (\|\mathbf{u}(t)\|_{H^k(\mathbb{R}^n)} + \|\mathbf{u}'(t)\|_{H^{k-1}(\mathbb{R}^n)}).$$

For $\lambda > 0$ define

$$X_\lambda := \{\mathbf{u} \in X \mid \|\mathbf{u}\| \leq \lambda, \mathbf{u}(0) = g, \mathbf{u}'(0) = h\}.$$

If $\mathbf{v} \in X_\lambda$, we write $\mathbf{u} = A[\mathbf{v}]$ to mean that \mathbf{u} solves the linear initial-value problem

$$(9) \quad \begin{cases} u_{tt} - \Delta u = -f(Dv, v_t, v) & \text{in } \mathbb{R}^n \times (0, T] \\ u = g, u_t = h & \text{on } \mathbb{R}^n \times \{t = 0\}. \end{cases}$$

Define

$$E_k(t) := \|\mathbf{u}(t)\|_{H^k(\mathbb{R}^n)}^2 + \|\mathbf{u}'(t)\|_{H^{k-1}(\mathbb{R}^n)}^2$$

and

$$F_k(t) := \|\mathbf{v}(t)\|_{H^k(\mathbb{R}^n)}^2 + \|\mathbf{v}'(t)\|_{H^{k-1}(\mathbb{R}^n)}^2.$$

2. We claim now that we have the estimate

$$(10) \quad E_k(t) \leq E_k(0) + C \int_0^t \Psi(F_k(s)) ds \quad (0 \leq t \leq T)$$

for some continuous and monotone function Ψ depending only upon n , k and f .

To prove (10), let $|\alpha| \leq k - 1$ and apply $D^\alpha = D_x^\alpha$ to the PDE (9), to discover

$$w_{tt} - \Delta w = -D^\alpha(f(Dv, v_t, v))$$

for $w := D^\alpha u$. Therefore we can use estimate (7), with $k - 1$ in place of k and with $m = n + 2$, to compute

$$\begin{aligned} \frac{d}{dt} \int_{\mathbb{R}^n} (w_t^2 + |Dw|^2) dx &= 2 \int_{\mathbb{R}^n} w_t w_{tt} + Dw \cdot Dw_t dx \\ &= 2 \int_{\mathbb{R}^n} w_t (w_{tt} - \Delta w) dx \\ &= -2 \int_{\mathbb{R}^n} w_t D^\alpha f(Dv, v_t, v) dx \\ &\leq \int_{\mathbb{R}^n} w_t^2 + |D^\alpha f(Dv, v_t, v)|^2 dx \\ &\leq \int_{\mathbb{R}^n} w_t^2 dx + C \Phi^2(\|v_{x_1}\|_{H^{k-1}(\mathbb{R}^n)}, \dots, \|v_{x_n}\|_{H^{k-1}(\mathbb{R}^n)}, \\ &\quad \|v_t\|_{H^{k-1}(\mathbb{R}^n)}, \|v\|_{H^{k-1}(\mathbb{R}^n)}) \\ &\leq \int_{\mathbb{R}^n} w_t^2 dx + \Psi(F_k(t)), \end{aligned}$$

for some appropriate function Ψ . Apply Gronwall's inequality and then sum the above over all $|\alpha| \leq k - 1$ to deduce (10).

3. We assert next that if $\lambda > 0$ is large enough and $T > 0$ is small enough, then

$$A : X_\lambda \rightarrow X_\lambda.$$

To see this, observe that (10) implies

$$(11) \quad |||\mathbf{u}|||^2 \leq \|g\|_{H^k(\mathbb{R}^n)}^2 + \|h\|_{H^{k-1}(\mathbb{R}^n)}^2 + CT\Psi(|||\mathbf{v}|||^2)$$

for some function Ψ . Let

$$\lambda^2 := 2(\|g\|_{H^k(\mathbb{R}^n)}^2 + \|h\|_{H^{k-1}(\mathbb{R}^n)}^2)$$

and then fix $T > 0$ so small that

$$CT\Psi(\lambda^2) \leq \frac{\lambda^2}{2}.$$

Then (11) forces $|||A[\mathbf{v}]|||^2 \leq \lambda^2$, and consequently $A[\mathbf{v}] = \mathbf{u} \in X_\lambda$.

4. Next we claim that if λ is large enough and T is small enough, then

$$(12) \quad \|A[\mathbf{v}] - A[\hat{\mathbf{v}}]\| \leq \frac{1}{2}\|\mathbf{v} - \hat{\mathbf{v}}\|$$

for all $\mathbf{v}, \hat{\mathbf{v}} \in X_\lambda$.

To confirm this, let us write $\mathbf{u} = A[\mathbf{v}]$, $\hat{\mathbf{u}} = A[\hat{\mathbf{v}}]$. Put $\mathbf{w} := \mathbf{u} - \hat{\mathbf{u}}$. Then

$$\begin{aligned} & \frac{d}{dt} \int_{\mathbb{R}^n} |Dw|^2 + w_t^2 + w^2 \, dx \\ &= 2 \int_{\mathbb{R}^n} w_t (f(D\hat{v}, \hat{v}_t, \hat{v}) - f(Dv, v_t, v) + w) \, dx \\ &\leq \int_{\mathbb{R}^n} w_t^2 + w^2 \, dx + C \int_{\mathbb{R}^n} |D\hat{v} - Dv|^2 + |\hat{v}_t - v_t|^2 + |\hat{v} - v|^2 \, dx, \end{aligned}$$

the constant C depending on $\|D\hat{v}, \hat{v}_t, \hat{v}, Dv, v_t, v\|_{L^\infty}$. This quantity is bounded since the functions $\mathbf{v}, \hat{\mathbf{v}}$ belong to X_λ and $k > \frac{n}{2} + 1$. Invoking Gronwall's inequality, we deduce that

$$\begin{aligned} & \max_{0 \leq t \leq T} \int |Dw|^2 + |w_t|^2 + w^2 \, dx \\ & \leq C \int_0^T \int_{\mathbb{R}^n} |Dv - D\hat{v}|^2 + |v_t - \hat{v}_t|^2 + |v - \hat{v}|^2 \, dxdt \\ & \leq CT\|v - \hat{v}\|^2. \end{aligned}$$

We can now select T small enough to ensure (12).

5. Select any $\mathbf{u}_0 \in X_\lambda$. According to the proof of Banach's Theorem from §9.2.1, if we inductively define $\mathbf{u}_{k+1} := A[\mathbf{u}_k]$ for $k = 0, \dots$, then $\mathbf{u}_k \rightarrow \mathbf{u}$ in X and

$$A[\mathbf{u}] = \mathbf{u}.$$

Furthermore, since $\|\|\mathbf{u}_k\|\| \leq \lambda$, we have $\mathbf{u} \in X_\lambda$. Uniqueness follows from (12). \square

Note carefully the strategy of this proof. We showed that for small $T > 0$ the operator A is a strict contraction in the weaker norm $\|\cdot\|$ and also preserves certain estimates in the stronger norm $\|\|\cdot\|\|$. Consequently the iteration scheme from Banach's Theorem provides a sequence that converges in X , to a fixed point that actually lies in the better space X_λ . We did not have to show that A is a strict contraction in the stronger norm.

12.3. SEMILINEAR WAVE EQUATIONS

This section and the next section discuss the initial-value problem for semilinear wave equations:

$$(1) \quad \begin{cases} u_{tt} - \Delta u + f(u) = 0 & \text{in } \mathbb{R}^n \times (0, \infty) \\ u = g, \quad u_t = h & \text{on } \mathbb{R}^n \times \{t = 0\}. \end{cases}$$

We will prove much stronger existence theorems than those in §12.2 not only because the nonlinear term $f(u)$ is simpler than $f(u, Du, u_t)$ but also because we have for (1) the conserved energy functional

$$E(t) := \int_{\mathbb{R}^n} \frac{1}{2}(u_t^2 + |Du|^2) + F(u) \, dx,$$

unavailable for the general quasilinear wave equations. Our main goal is discovering when solutions exist for all times $t \geq 0$.

12.3.1. Sign conditions.

Our first existence theorem holds for nonlinearities such that $f(z)$ and z have the same sign.

THEOREM 1 (Sign condition on f). *Suppose f is smooth and*

$$(2) \quad z f(z) \geq 0 \quad (z \in \mathbb{R}).$$

Assume $g \in H^1(\mathbb{R}^n)$, $h \in L^2(\mathbb{R}^n)$, $F(g) \in L^1(\mathbb{R}^n)$.

Then the initial-value problem (1) has a global weak solution \mathbf{u} existing for all times, with

$$\begin{cases} \mathbf{u} \in L^2_{\text{loc}}((0, \infty); L^2(\mathbb{R}^n)) \cap L^\infty((0, \infty); H^1(\mathbb{R}^n)) \\ \mathbf{u}' \in L^\infty((0, \infty); L^2(\mathbb{R}^n)), F(\mathbf{u}) \in L^\infty((0, \infty); L^1(\mathbb{R}^n)). \end{cases}$$

Furthermore, we have the energy inequality

$$(3) \quad E(t) \leq E(0) \quad \text{for all times } t \geq 0.$$

Note that we do not assert equality in (3): our solution is not known to be smooth enough for us to calculate rigorously that $\dot{E} \equiv 0$.

Change of notation. Starting with the following proof, we transition away from boldface notation, indicating a mapping of time t into a space of functions of x , and will instead hereafter regard our solution as a function $u = u(x, t)$ of both variables x and t together.

Proof. 1. According to the sign condition (2), $F(z) = \int_0^z f(w) dw$ is non-decreasing for $z \geq 0$ and is nonincreasing for $z \leq 0$. Select a sequence of smooth function $F_k : \mathbb{R} \rightarrow \mathbb{R}$ so that

$$F_k \geq 0, F_k \rightarrow F \text{ pointwise, } F_k \leq F, F_k \equiv F \text{ on } [-k, k]$$

and $f_k := F'_k$ is Lipschitz continuous, with $z f_k(z) \geq 0$ for all $z \in \mathbb{R}$.

We solve the problems

$$(4) \quad \begin{cases} u^k_{tt} - \Delta u^k + f_k(u^k) = 0 & \text{in } \mathbb{R}^n \times (0, \infty) \\ u^k = g, u^k_t = h & \text{on } \mathbb{R}^n \times \{t = 0\}. \end{cases}$$

Since f_k is Lipschitz continuous, there exists according to Theorem 1 in §12.2 a solution u^k satisfying

$$\begin{cases} u^k, Du^k, u^k_t \in C([0, \infty); L^2(\mathbb{R}^n)), \\ D_x^2 u^k, D_x u^k_t, u^k_{tt} \in L^\infty_{\text{loc}}((0, \infty); L^2(\mathbb{R}^n)). \end{cases}$$

2. This is enough regularity to allow us to calculate for almost every time t that

$$\dot{E}_k(t) = \frac{d}{dt} \int_{\mathbb{R}^n} \frac{1}{2} ((u^k_t)^2 + |Du^k|^2) + F_k(u^k) dx = 0.$$

Therefore

$$(5) \quad E_k(t) = E_k(0) = \int_{\mathbb{R}^n} \frac{1}{2} (h^2 + |Dg|^2) + F_k(g) dx.$$

Now $F_k(g) \rightarrow F(g)$ pointwise as $k \rightarrow \infty$, and $0 \leq F_k(g) \leq F(g)$. Since $F(g) \in L^1$, we can apply the Dominated Convergence Theorem to deduce

$$E^k(0) \rightarrow \int_{\mathbb{R}^n} \frac{1}{2} (|Dg|^2 + |h|^2) + F(g) dx = E(0).$$

Since $F_k \geq 0$, we also have the bound

$$(6) \quad \max_{0 \leq t \leq T} \|u_t^k, Du^k\|_{L^2(\mathbb{R}^n)} \leq C$$

and consequently for each time $T > 0$

$$(7) \quad \max_{0 \leq t \leq T} \|u^k\|_{L^2(\mathbb{R}^n)} \leq C.$$

We extract a subsequence (which we reindex and still denote “ u^k ”) such that

$$\begin{cases} u^k \rightarrow u & \text{strongly in } L^2_{\text{loc}}(\mathbb{R}^n \times (0, \infty)) \text{ and a.e.,} \\ Du^k, u_t^k \rightharpoonup Du, u_t & \text{weakly in } L^2_{\text{loc}}(\mathbb{R}^n \times (0, \infty)). \end{cases}$$

3. Next, multiply the PDE (4) by u^k and integrate over $\mathbb{R}^n \times (0, T)$:

$$\int_0^T \int_{\mathbb{R}^n} |Du^k|^2 - (u_t^k)^2 + u^k f_k(u^k) dxdt = \int_{\mathbb{R}^n} u_t^k u^k dx \Big|_{t=0}^{t=T}.$$

Since $u^k f_k(u^k) \geq 0$, (6) and (7) imply the estimate

$$(8) \quad \int_0^T \int_{\mathbb{R}^n} |u^k f_k(u^k)| dxdt \leq C.$$

4. We next assert that the functions $\{g^k := f_k(u^k)\}_{k=1}^\infty$ are *uniformly integrable*. This means that for each $\varepsilon > 0$, there exists $\delta > 0$ such that if E is a measurable subset of $\mathbb{R}^n \times (0, T)$ and $|E| \leq \delta$, then $\iint_E |g^k| dxdt \leq \varepsilon$ for all k .

To confirm this, we calculate using (8) that

$$\begin{aligned} \iint_E |g^k| dxdt &= \iint_{E \cap \{|u^k| \geq \lambda\}} |f_k(u^k)| dxdt + \iint_{E \cap \{|u^k| \leq \lambda\}} |f_k(u^k)| dxdt \\ &\leq \frac{1}{\lambda} \iint_E |u^k f_k(u^k)| dxdt + |E| \max_{|y| \leq \lambda} |f_k(y)| \\ &\leq \frac{1}{\lambda} C(T) + |E| \max_{|y| \leq \lambda} |f_k(y)|. \end{aligned}$$

Take λ so large that $\frac{C(T)}{\lambda} \leq \frac{\varepsilon}{2}$. Then for all but finitely many k

$$\iint_E |g^k| \, dx dt \leq \frac{\varepsilon}{2} + |E| \max_{|y| \leq \lambda} |f(y)| \leq \varepsilon,$$

provided $|E| \leq \delta$ for $\delta > 0$ sufficiently small.

5. We claim now that

$$(9) \quad f_k(u^k) \rightarrow f(u) \text{ in } L^1_{\text{loc}}(\mathbb{R}^n).$$

To see this, again write $g^k := f_k(u^k)$, put $g := f(u)$, and fix $R > 0$, $\varepsilon > 0$. In view of the uniform integrability proved in step 4, we can select $\delta > 0$ so that $|E| \leq \delta$ implies $\int_E |g^k| \, dx \leq \varepsilon$ for $k = 1, \dots$. According to Egoroff's Theorem (§E.2), there exists a measurable set E so that $|E| \leq \delta$ and $g^k \rightarrow g$ uniformly on $B(0, R) - E$. Therefore

$$\limsup_{k, l \rightarrow \infty} \int_{B(0, R)} |g^k - g^l| \, dx \leq \limsup_{k, l \rightarrow \infty} \int_E |g^k - g^l| \, dx \leq 2\varepsilon.$$

This is true for each $\varepsilon > 0$. Consequently $\{g^k\}_{k=1}^\infty$ is a Cauchy sequence in $L^1(B(0, R))$ and so converges to some $\hat{g} \in L^1(B(0, R))$. Since $g^k \rightarrow g$ a.e., we deduce $g = \hat{g}$.

6. Because

$$u^k_{tt} - \Delta u^k + f_k(u^k) = 0 \quad \text{in } \mathbb{R}^n \times (0, \infty),$$

we deduce upon multiplying by a smooth test function and passing to limits that

$$(10) \quad u_{tt} - \Delta u + f(u) = 0 \quad \text{in } \mathbb{R}^n \times (0, \infty).$$

Furthermore

$$\int_{\mathbb{R}^n} u_t^2 + |Du|^2 \, dx \leq \liminf_{k \rightarrow \infty} \int_{\mathbb{R}^n} (u_t^k)^2 + |Du^k|^2 \, dx;$$

and according to Fatou's Lemma (§E.3),

$$\int_{\mathbb{R}^n} F(u) \, dx \leq \liminf_{k \rightarrow \infty} \int_{\mathbb{R}^n} F_k(u^k) \, dx.$$

These last two inequalities together imply $E(t) \leq E(0)$. □

12.3.2. Three space dimensions.

Nonlinear wave equations in three space dimensions are physically the most important and turn out to admit useful L^∞ estimates owing to the special form of the solution to the linear nonhomogenous problem provided by the retarded potential formula (44) from §2.4.2.

So let us now look at the initial-value problem

$$(11) \quad \begin{cases} u_{tt} - \Delta u + f(u) = 0 & \text{in } \mathbb{R}^3 \times (0, \infty) \\ u = g, \quad u_t = h & \text{on } \mathbb{R}^3 \times \{t = 0\}. \end{cases}$$

We henceforth always assume that $f : \mathbb{R} \rightarrow \mathbb{R}$ is smooth, with $f(0) = 0$, and that $g, h \in C_c^\infty(\mathbb{R}^3)$.

THEOREM 2 (Short time existence and blow-up in L^∞).

(i) *There exists a time $T > 0$ and a unique smooth solution u of the initial-value problem (11) on $\mathbb{R}^3 \times (0, T)$.*

(ii) *If the maximal time T^* of existence of this smooth solution is finite, then*

$$(12) \quad \lim_{t \rightarrow T^*} \|u(\cdot, t)\|_{L^\infty(\mathbb{R}^3)} = \infty.$$

Assertion (12) is important since (unlike Theorem 3 in §12.2.2) it provides a simple criterion for the failure of the solution to exist beyond time T^* . We will see in Theorem 3 below and also in §12.4 that we can sometimes bound the L^∞ norm of solutions and so ensure existence for all time.

Proof. 1. We will look for a solution u having the form

$$u = v + w,$$

where v solves the homogeneous wave equation

$$(13) \quad \begin{cases} v_{tt} - \Delta v = 0 & \text{in } \mathbb{R}^3 \times (0, \infty) \\ v = g, \quad v_t = h & \text{on } \mathbb{R}^3 \times \{t = 0\} \end{cases}$$

and w solves

$$\begin{cases} w_{tt} - \Delta w = -f(u) & \text{in } \mathbb{R}^3 \times (0, \infty) \\ w = w_t = 0 & \text{on } \mathbb{R}^3 \times \{t = 0\}. \end{cases}$$

Formula (44) from §2.4.2 lets us write

$$w(x, t) = -\frac{1}{4\pi} \int_{B(x, t)} \frac{f(u(y, t - |y - x|))}{|y - x|} dy.$$

Hence our desired solution u must solve the nonlinear integral identity

$$(14) \quad u(x, t) = v(x, t) - \frac{1}{4\pi} \int_{B(x, t)} \frac{f(u^*)}{|x - y|} dy \quad (x \in \mathbb{R}^3, t > 0),$$

where for each fixed (x, t) we write

$$(15) \quad u^*(y) := u(y, t - |y - x|) \quad (y \in B(x, t)).$$

2. Introduce the collection of functions

$$X := \{u \in C([0, T] \times \mathbb{R}^3) \mid u(\cdot, 0) = g, \|u - v\|_{L^\infty} \leq 1\},$$

where v solves the linear, homogeneous problem (13). Since g, h are smooth, so is v . Thus there exists a constant C_1 such that

$$(16) \quad \|u\|_{L^\infty((0, T) \times \mathbb{R}^3)} \leq C_1$$

for all $u \in X$.

We define the nonlinear mapping $A : X \rightarrow C([0, T] \times \mathbb{R}^3)$ by

$$A[u](x, t) := v(x, t) - \frac{1}{4\pi} \int_{B(x, t)} \frac{f(u^*)}{|x - y|} dy.$$

Then if $u, \hat{u} \in X$,

$$\begin{aligned} \|A[u] - A[\hat{u}]\|_{L^\infty((0, T) \times \mathbb{R}^3)} &\leq \sup_{x \in \mathbb{R}^3, 0 \leq t \leq T} \left(\frac{1}{4\pi} \int_{B(x, t)} \frac{|f(u^*) - f(\hat{u}^*)|}{|x - y|} dy \right) \\ &\leq C \sup_{x \in \mathbb{R}^3, 0 \leq t \leq T} \int_{B(x, t)} \frac{|u^* - \hat{u}^*|}{|x - y|} dy \\ &\leq C \|u - \hat{u}\|_{L^\infty((0, T) \times \mathbb{R}^3)} \sup_{x \in \mathbb{R}^3, 0 \leq t \leq T} \int_{B(x, t)} \frac{dy}{|x - y|} \\ &\leq CT^2 \|u - \hat{u}\|_{L^\infty((0, T) \times \mathbb{R}^3)}, \end{aligned}$$

the constant C in the second line depending upon $\max_{|w| \leq C_1} |f'(w)|$. For the last inequality in this calculation we noted that $\int_{B(0, r)} \frac{dy}{|y|} = Cr^2$.

Fix T so small that A is a strict contraction. Observing also that $A : X \rightarrow X$ if T is small, we see from Banach's Theorem (§9.2.1) that A has a unique fixed point u , which consequently solves the integral identities (14).

3. We can apply the same method to estimate the first and even higher derivatives of u . To see this, write $\tilde{u} := D_k^h u$ for a difference quotient, $k = 1, 2, 3$ (§5.8.2). Then

$$\begin{cases} \tilde{u}_{tt} - \Delta \tilde{u} + c\tilde{u} = 0 & \text{in } \mathbb{R}^3 \times (0, T] \\ \tilde{u} = \tilde{g}, \tilde{u}_t = \tilde{h} & \text{on } \mathbb{R}^3 \times \{t = 0\}. \end{cases}$$

where $\tilde{g} := D_k^h g$, $\tilde{h} := D_k^h h$ and $c := \int_0^1 f'(su(x + he_k, t) + (1-s)u(x, t)) ds$. The function c is bounded, since u is bounded.

As above,

$$(17) \quad \tilde{u}(x, t) = \tilde{v}(x, t) - \frac{1}{4\pi} \int_{B(x, t)} \frac{c^* \tilde{u}^*}{|x - y|} dy,$$

where

$$\begin{cases} \tilde{v}_{tt} - \Delta \tilde{v} = 0 & \text{in } \mathbb{R}^3 \times (0, \infty) \\ \tilde{v} = \tilde{g}, \tilde{v}_t = \tilde{h} & \text{on } \mathbb{R}^3 \times \{t = 0\}. \end{cases}$$

In particular, \tilde{v} is smooth, and consequently (17) implies

$$\|\tilde{u}(\cdot, t)\|_{L^\infty} \leq C + C \int_0^t \|\tilde{u}(\cdot, s)\|_{L^\infty} ds \quad (0 \leq t \leq T).$$

We invoke Gronwall's inequality (§E.2) next to estimate the L^∞ -norms of Du . We can similarly bound u_t . Writing $\tilde{u} = u_{x_k}$ or u_t , we see that \tilde{u} is a weak solution of

$$\tilde{u}_{tt} - \Delta \tilde{u} + f'(u)\tilde{u} = 0.$$

We can now apply our difference quotient argument to this PDE to derive L^∞ bounds on D^2u . By induction we can similarly estimate all the partial derivatives of u in terms of the L^∞ -norm of u .

4. Now let T^* denote the maximal time of existence of a smooth solution. If $T^* < \infty$, but (12) fails, we can as above estimate all the derivatives of u on $\mathbb{R}^3 \times [0, T^*)$ and therefore extend the solution beyond this time. \square

12.3.3. Subcritical power nonlinearities.

We now turn our attention to the nonlinear wave equation in three dimensions with a power-type nonlinearity:

$$(18) \quad \begin{cases} u_{tt} - \Delta u + |u|^{p-1}u = 0 & \text{in } \mathbb{R}^3 \times (0, \infty) \\ u = g, u_t = h & \text{on } \mathbb{R}^3 \times \{t = 0\}, \end{cases}$$

the energy for which is

$$E(t) := \int_{\mathbb{R}^3} \frac{1}{2} (u_t^2 + |Du|^2) + \frac{|u|^{p+1}}{p+1} dy.$$

We assume $g, h \in C_c^\infty(\mathbb{R}^3)$.

Our goal now is building a global solution for $1 \leq p < 5$.

THEOREM 3 (Subcritical existence theorem). *If $1 < p < 5$, then there exists a unique global smooth solution of (18).*

Section 12.4 will introduce more sophisticated techniques to handle the critical case $p = 5$.

Proof. 1. In view of Theorem 2, we may assume we have a smooth solution u existing on $\mathbb{R}^3 \times (0, T)$ for some $T > 0$. We need to show that

$$\|u\|_{L^\infty(\mathbb{R}^3 \times (0, T))} \leq C$$

for some constant $C = C(T)$.

Let

$$\begin{cases} v_{tt} - \Delta v = 0 & \text{in } \mathbb{R}^3 \times (0, \infty) \\ v = g, v_t = h & \text{on } \mathbb{R}^3 \times \{t = 0\}, \end{cases}$$

so that as in the previous proof

$$u(x, t) = v(x, t) - \frac{1}{4\pi} \int_{B(x, t)} \frac{|u^*|^{p-1} u^*}{|x - y|} dy.$$

Recall from (15) our notation that $u^*(y) := u(y, t - |y - x|)$ for $y \in B(x, t)$. Since g, h are smooth, v is bounded on $\mathbb{R}^3 \times [0, T]$. Therefore

$$(19) \quad |u(x, t)| \leq C + I(x, t) \quad (x \in \mathbb{R}^3, 0 \leq t \leq T)$$

for

$$I(x, t) := \int_{B(x, t)} \frac{|u^*|^p}{|x - y|} dy.$$

2. Then

$$(20) \quad I(x, t) \leq \left(\int_{B(x, t)} \frac{|u^*|^2}{|x - y|^2} dy \right)^{1/2} \left(\int_{B(x, t)} |u^*|^{2(p-1)} dy \right)^{1/2}.$$

Hardy's inequality (Theorem 7 in §5.8.4) implies

$$(21) \quad \int_{B(x, t)} \frac{|u^*|^2}{|y - x|^2} dy \leq C \int_{B(x, t)} |Du^*|^2 + \frac{(u^*)^2}{t^2} dy.$$

Now

$$Du^* = Du - \nu u_t,$$

for $\nu = \frac{y-x}{|y-x|}$. Consequently the energy flux identity (3) in §12.1.2 implies

$$(22) \quad \int_{B(x, t)} |Du^*|^2 dy = \frac{1}{\sqrt{2}} \int_{\Gamma(x, t)} |u_t \nu - Du|^2 dS \leq E(0).$$

Recalling next Poincaré's inequality (7) in §5.8.1, we see that

$$\int_{B(x,t)} |u^* - (u^*)_{x,t}|^2 dy \leq Ct^2 \int_{B(x,t)} |Du^*|^2 dy,$$

for the average $(u^*)_{x,t} := \int_{B(x,t)} u^* dy$. Hence

$$(23) \quad \frac{1}{t^2} \int_{B(x,t)} (u^*)^2 dy \leq C \int_{B(x,t)} |Du^*|^2 dy + Ct|(u^*)_{x,t}|^2.$$

We now compute that

$$(24) \quad \begin{aligned} |(u^*)_{x,t}| &\leq \frac{C}{t^3} \int_{\Gamma(x,t)} |u| dS \leq \frac{C}{t^3} \left(\int_{K(x,t)} |u_t| dy ds + \int_{B(x,t)} |g| dy \right) \\ &\leq \frac{C}{t^3} \left(\int_{K(x,t)} |u_t|^2 dy ds \right)^{\frac{1}{2}} |K(x,t)|^{\frac{1}{2}} + C\|g\|_{L^\infty} \\ &\leq \frac{C}{t^{\frac{1}{2}}} E(0)^{\frac{1}{2}} + C\|g\|_{L^\infty}. \end{aligned}$$

This estimate, combined with (20)-(23), gives

$$(25) \quad I(x,t) \leq C \left(\int_{B(x,t)} |u^*|^{2(p-1)} dy \right)^{1/2}.$$

3. Suppose now that $1 < p \leq 4$. Then $2(p-1) \leq 6 = 2^*$, and therefore the Sobolev inequalities (§5.6.3) imply

$$\|u^{*p-1}\|_{L^2(B(x,t))} \leq C\|u^*\|_{H^1(B(x,t))}^{p-1} \leq C.$$

Consequently (19) and (25) yield

$$|u(x,t)| \leq C,$$

and we have derived the required estimate on $\|u\|_{L^\infty(\mathbb{R}^3 \times (0,T))}$.

The next case is $4 < p < 5$. Then $2(p-1) = 2(p-4) + 6 = 2(p-4) + 2^*$, and the foregoing calculations show that

$$I(x,t) \leq C \left(\int_{B(0,t)} |u^*|^{2(p-1)} dy \right)^{1/2} \leq C\|u\|_{L^\infty}^{p-4}.$$

Therefore (19) implies

$$|u(x, t)| \leq C + C\|u\|_{L^\infty}^{p-4} \leq C + \frac{1}{2}\|u\|_{L^\infty},$$

the last inequality valid since $0 < p - 4 < 1$. This again provides a bound on $\|u\|_{L^\infty(\mathbb{R}^3 \times (0, T))}$. \square

We will see in Problem 15 that the proof can be modified to show for the critical power $p = 5$ that there exists a global smooth solution provided the energy $E(0)$ is small enough. The next section introduces more advanced techniques, to construct a solution even for large energy when $p = 5$.

12.4. CRITICAL POWER NONLINEARITY

We consider now the initial-value problem

$$(1) \quad \begin{cases} u_{tt} - \Delta u + u^5 = 0 & \text{in } \mathbb{R}^3 \times (0, \infty) \\ u = g, \quad u_t = h & \text{on } \mathbb{R}^3 \times \{t = 0\}, \end{cases}$$

with corresponding energy

$$(2) \quad E(t) := \int_{\mathbb{R}^3} \frac{1}{2}(u_t^2 + |Du|^2) + \frac{u^6}{6} dy.$$

We continue to suppose $g, h \in C_c^\infty(\mathbb{R}^3)$. Our task is showing that there is a smooth solution, even for this critical power $p = 5$ case for which the methods of the previous section fail.

We will first need more detailed information about the linear wave equation:

THEOREM 1 (Estimates for wave equation in three dimensions). *Let v solve the linear initial-value problem*

$$(3) \quad \begin{cases} v_{tt} - \Delta v = f & \text{in } \mathbb{R}^3 \times (0, \infty) \\ v = g, \quad v_t = h & \text{on } \mathbb{R}^3 \times \{t = 0\}. \end{cases}$$

For each $T > 0$ we have the estimate

$$(4) \quad \sup_{0 \leq t \leq T} \|v(\cdot, t)\|_{L^6(\mathbb{R}^3)} + \|v\|_{L^4(0, T; L^{12}(\mathbb{R}^3))} \\ \leq C(\|Dg\|_{L^2(\mathbb{R}^3)} + \|h\|_{L^2(\mathbb{R}^3)} + \|f\|_{L^1(0, T; L^2(\mathbb{R}^3))})$$

for a constant $C = C(T)$.

Proof. 1. Approximating f, g, h by smooth functions if necessary, we may assume v is smooth. We then compute for

$$a(t) := \left(\int_{\mathbb{R}^3} v_t^2 + |Dv|^2 dx \right)^{\frac{1}{2}}$$

that

$$\begin{aligned} 2\dot{a}(t)a(t) &= 2 \int_{\mathbb{R}^3} (v_{tt} - \Delta v)v_t dx \\ &= 2 \int_{\mathbb{R}^3} f v_t dx \leq 2a(t)\|f(\cdot, t)\|_{L^2}. \end{aligned}$$

Therefore

$$\sup_{0 \leq t \leq T} \|Dv(\cdot, t)\|_{L^2} \leq C(\|Dg\|_{L^2} + \|h\|_{L^2} + \|f\|_{L^1(0, T; L^2)}).$$

Since for $n = 3$ dimensions, $2^* = 6$, the Sobolev inequalities (§5.6.3) provide the stated estimate for $\sup_{0 \leq t \leq T} \|v\|_{L^6}$.

2. The proof of the $L^4(0, T; L^{12})$ bound for v is beyond the scope of this textbook. See Sogge [So] for details and also Shatah–Struwe [S-S]. \square

Recalling the energy flux calculation appearing §12.1, we introduce the following

NOTATION. Given $(x_0, T) \in \mathbb{R}^3 \times (0, \infty)$ and $0 < s < T$, write

$$\phi(s) = \phi(s, x_0, T) := \frac{1}{\sqrt{2}} \int_{\Gamma(x_0, t_0) \cap \{s \leq t < T\}} \frac{1}{2} |u_t \nu - Du|^2 + \frac{u^6}{6} dS$$

for the energy flux through the curved surface $\Gamma(x_0, T)$ for times between s and T .

THEOREM 2 (L^6 -energy flux estimate). *If u is a smooth solution of $u_{tt} - \Delta u + u^5 = 0$ in $\mathbb{R}^3 \times [0, T]$, we have the estimate*

$$(5) \quad \int_{B(x_0, T-s)} u^6 dx \leq C\phi(s)^{\frac{1}{3}}$$

for each point $x_0 \in \mathbb{R}^3$ and each time $0 \leq s < T$.

Theorem 2 is important since it implies, as we will later see, that the full energy density $\frac{1}{2}(u_t^2 + |Du|^2) + \frac{1}{6}u^6$ cannot “concentrate” near (x_0, T) . The proof depends upon a nonlinear variant of the scaling invariance identity introduced for the linear wave equation in §8.6.

Proof. 1. We may assume $x_0 = 0$ and suppose for the time being that $s = 0$. Inspired by Example 4 of §8.6, we multiply the PDE $\square u + u^5 = 0$ by the multiplier

$$m := (t - T)u_t + x \cdot Du + u.$$

After some rewriting, we derive the Morawetz-type identity

$$(6) \quad p_t - \operatorname{div} \mathbf{q} = -\frac{u^6}{3},$$

for

$$(7) \quad p := (t - T) \left(\frac{u_t^2 + |Du|^2}{2} + \frac{u^6}{6} \right) + x \cdot Du u_t + u u_t$$

and

$$(8) \quad \mathbf{q} := ((t - T)u_t + x \cdot Du + u)Du + \left(\frac{u_t^2 - |Du|^2}{2} - \frac{u^6}{6} \right) x.$$

Select a time $0 < \tau < T$. We integrate (6) over the truncated backward wave cone $K(0, T) \cap \{0 \leq t \leq \tau\}$, ending up with three terms corresponding respectively to integrations over the curved side $\Gamma(0, T) \cap \{0 \leq t \leq \tau\}$, the bottom $B(0, T) \times \{t = 0\}$ and the top $B(0, T - \tau) \times \{t = \tau\}$. The latter term goes to zero as $\tau \rightarrow T$. Thus we discover

$$(9) \quad A - B = - \int_{K(0, T)} \frac{u^6}{3} dx dt \leq 0$$

for

$$A := \frac{1}{\sqrt{2}} \int_{\Gamma(0, T)} p - \mathbf{q} \cdot \nu dS, \quad B := \int_{B(0, T)} p(\cdot, 0) dx,$$

where $\nu := \frac{x}{|x|}$.

2. We now claim that

$$(10) \quad A = \frac{1}{\sqrt{2}} \int_{\Gamma(0, T)} (t - T) \left| u_r - u_t + \frac{u}{|x|} \right|^2 dS + \frac{1}{2} \int_{\partial B(0, T)} u^2(\cdot, 0) dS$$

for $u_r := Du \cdot \frac{x}{|x|}$. To confirm this, we first observe that $|x| = T - t$ on $\Gamma(0, T)$ and then check after a calculation using (7), (8) that

$$(11) \quad p - \mathbf{q} \cdot \nu = -|x|(u_t - u_r)^2 + u(u_t - u_r) \quad \text{on } \Gamma(0, T).$$

We transform the surface integral over the curved surface $\Gamma(0, T)$ to an integral over the ball $B(0, T)$, by putting

$$(12) \quad u^*(y) := u(y, T - |y|).$$

Then $u_r^* = \frac{y}{|y|} \cdot Du^* = u_r - u_t$, and so (11) implies

$$\begin{aligned} (13) \quad A &= - \int_{B(0,T)} |y| (u_r^*)^2 + u^* u_r^* dy \\ &= - \int_{B(0,T)} |y| \left(u_r^* + \frac{u^*}{|y|} \right)^2 dy + \int_{B(0,T)} \frac{(u^*)^2}{|y|} + u^* u_r^* dy. \end{aligned}$$

Since $\operatorname{div} \left(\frac{y}{|y|} \right) = \frac{2}{|y|}$ in \mathbb{R}^3 , we can compute

$$\begin{aligned} \int_{B(0,T)} \frac{(u^*)^2}{|y|} dy &= \frac{1}{2} \int_{B(0,T)} (u^*)^2 \operatorname{div} \left(\frac{y}{|y|} \right) dy \\ &= - \int_{B(0,T)} u^* u_r^* dy + \frac{1}{2} \int_{\partial B(0,T)} (u^*)^2 dS. \end{aligned}$$

Plugging this identity into (13) and converting back to the original variables gives us the formula (10).

3. Next we assert that

$$(14) \quad B \leq -T \int_{B(0,T)} \frac{u^6}{6} dx + \frac{1}{2} \int_{\partial B(0,T)} u^2(\cdot, 0) dS.$$

To see this, notice first that

$$B = \int_{B(0,T)} (-T) \left(\frac{u_t^2 + |Du|^2}{2} + \frac{u^6}{6} \right) + (x \cdot Du + u) u_t dx.$$

Now if $|x| \leq T$, then

$$\begin{aligned} |(x \cdot Du + u) u_t| &\leq \frac{T u_t^2}{2} + \frac{|x|^2}{2T} \left(Du \cdot \frac{x}{|x|} + \frac{u}{|x|} \right)^2 \\ &\leq \frac{T u_t^2}{2} + \frac{T}{2} \left| Du + \frac{x}{|x|^2} u \right|^2. \end{aligned}$$

Consequently

$$\begin{aligned} B &\leq -T \int_{B(0,T)} \frac{u^6}{6} dx + \frac{T}{2} \int_{B(0,T)} \left| Du + \frac{x}{|x|^2} u \right|^2 - |Du|^2 dx \\ &= -T \int_{B(0,T)} \frac{u^6}{6} dx + \frac{T}{2} \int_{B(0,T)} \frac{u^2}{|x|^2} + \frac{2}{|x|} u u_r dx. \end{aligned}$$

Since $\operatorname{div} \left(\frac{y}{|y|^2} \right) = \frac{1}{|y|^2}$ in \mathbb{R}^3 , we calculate as in step 2 that the last integral equals $\frac{1}{2} \int_{\partial B(0,T)} u^2 dS$. This proves (14).

4. Combining (9), (10) and (14), we deduce that

$$\begin{aligned}
 T \int_{B(0,t_0)} u^6 dx &\leq C \int_{\Gamma(0,T)} (T-t) \left| u_r - u_t + \frac{u}{|x|} \right|^2 dS \\
 (15) \qquad &\leq CT\phi(0) + C \int_{\Gamma(0,T)} \frac{u^2}{|x|} dS \\
 &\leq CT\phi(0) + C \left(\int_{\Gamma(0,T)} u^6 dS \right)^{1/3} \left(\int_{\Gamma(0,T)} |x|^{-3/2} dS \right)^{2/3}.
 \end{aligned}$$

Since $\int_{\Gamma(0,T)} |x|^{-3/2} dS = CT^{3/2}$, we deduce finally that

$$\int_{B(0,T)} u^6 dx \leq C\phi(0)^{1/3}.$$

This is the inequality (5) for $s = 0$, and the general case that $0 \leq s < T$ follows similarly. □

We present next the major assertion that our critical power wave equation has a smooth solution existing globally in time:

THEOREM 3 (Global existence for u^5 nonlinearity). *Assume that f, g are smooth functions with compact support.*

Then there is a unique smooth solution of the initial-value problem (1) existing for all time.

Proof. 1. Assume that $0 < T < \infty$ and that u is a smooth, compactly supported solution existing on $\mathbb{R}^3 \times [0, T)$. We will show that

$$(16) \qquad u \in L^\infty(\mathbb{R}^3 \times [0, T)),$$

in which case our results from §12.3 imply that u can be smoothly extended beyond time T .

2. We first assert that if we knew

$$(17) \qquad u \in L^4(0, T; L^{12}(\mathbb{R}^3)),$$

then (16) would hold. To see this, differentiate the PDE $\square u + u^5 = 0$ to find $\square v + 5u^4 v = 0$, for $v := u_{x_k}$ ($k = 1, 2, 3$). The linear estimate (4) provides the bound

$$\sup_{0 \leq t \leq \tau} \|Du\|_{L^6} \leq C + C \int_0^\tau \|u^4 Du\|_{L^2} dt$$

for $0 \leq \tau \leq T$.

Now

$$\|u^4 Du\|_{L^2} \leq \|Du\|_{L^6} \|u\|_{L^{12}}^4;$$

and consequently

$$\sup_{0 \leq t \leq \tau} \|Du\|_{L^6} \leq C + C \sup_{0 \leq t \leq \tau} \|Du\|_{L^6} \int_0^\tau \|u\|_{L^{12}}^4 dt \leq C + \frac{1}{2} \sup_{0 \leq t \leq \tau} \|Du\|_{L^6},$$

provided $\tau > 0$ is so small that

$$\int_0^\tau \|u\|_{L^{12}}^4 dt \leq \delta := \frac{1}{2C}.$$

Since $u \in L^4(0, T; L^{12})$, we can select $\tau = \frac{T}{m} > 0$ such that

$$\int_{k\tau}^{(k+1)\tau} \|u\|_{L^{12}}^4 dt \leq \delta \quad \text{for } k = 1, \dots, m-1.$$

We can then iteratively apply the foregoing argument on the time intervals $[0, \tau], [\tau, 2\tau], \dots, [(m-1)\tau, T)$, eventually to deduce

$$\sup_{0 \leq t < T} \|Du\|_{L^6} \leq C.$$

Since u has compact support, this implies $\|u\|_{L^\infty((0, T) \times \mathbb{R}^3)} \leq C$ and so (16) is valid.

3. Next we show for each point $x_0 \in \mathbb{R}^3$ that

$$(18) \quad \lim_{s \rightarrow T} \int_{B(x_0, T-s)} u^6 dx = 0.$$

Indeed the key estimate (5) provided by Theorem 2 asserts

$$\int_{B(x_0, T-s)} u^6 dx \leq C \phi(s)^{\frac{1}{3}}$$

for the energy flux $\phi(s)$. But according to Theorem 2 in §12.2, $\phi(s) = e(s) - \lim_{r \rightarrow 0} e(r)$ for

$$e(t) := \int_{B(x_0, T-t)} \frac{1}{2}(u_t^2 + |Du|^2) + \frac{u^6}{6} dx.$$

Since $t \mapsto e(t)$ is nonincreasing, it follows that $\lim_{s \rightarrow T} \phi(s) = 0$ and consequently that (18) holds. (Notice in this argument that we do not need

to know that $\lim_{r \rightarrow 0} e(r) = 0$, although the next step shows this is in fact true.)

4. We assert now that (18) implies

$$(19) \quad \lim_{s \rightarrow T} \int_{B(x_0, T-s)} u_t^2 + |Du|^2 dx = 0$$

for each point x_0 . To confirm this, let us first show that (18) implies

$$(20) \quad u \in L^4(0, T; L^{12}(B(x_0, T-t))).$$

To prove this, we first observe that owing to the linear estimate (4),

$$(21) \quad \|u\|_{L^4(s, \tau, L^{12}(B(x_0, T-t)))} \leq C + C \int_s^\tau \|u^5\|_{L^2(B(x_0, T-t))} dt$$

for each $s \leq \tau < T$. The interpolation inequality $\|u\|_{L^{10}} \leq \|u\|_{L^6}^{1/5} \|u\|_{L^{12}}^{4/5}$ (§B.2) implies

$$\|u^5\|_{L^2} = \|u\|_{L^{10}}^5 \leq \|u\|_{L^6} \|u\|_{L^{12}}^4.$$

Consequently,

$$\int_s^\tau \|u^5\|_{L^2(B(x_0, T-t))} dt \leq \sup_{s \leq t \leq T} \|u\|_{L^6(B(x_0, T-t))} \int_s^\tau \|u\|_{L^{12}(B(x_0, T-t))}^4 dt.$$

It follows then from (18) and (21) that given any $\epsilon > 0$ we can select a time $0 < s < T$ such that

$$\|u\|_{L^4(s, \tau, L^{12}(B(x_0, T-t)))} \leq C + \epsilon \|u\|_{L^4(s, \tau, L^{12}(B(x_0, T-t)))}^4$$

for all $s \leq \tau < T$. This expression has the form

$$\phi(\tau) \leq C_1 + \epsilon \phi(\tau)^4 \quad (s \leq \tau < T)$$

with $\phi(s) = 0$. It follows that

$$\phi(\tau) \leq 2C_1 \quad (s \leq \tau < T)$$

provided that ϵ is sufficiently small. This proves (20).

But then (20) lets us apply the method of step 2 to $v := u_{x_k}$ ($k = 1, 2, 3$) and $v := u_t$. This reasoning provides the bound

$$\sup_{0 \leq t < T} \|Du, u_t\|_{L^6(B(x_0, T-t))} \leq C,$$

which in turn gives us (19).

5. Next we improve slightly on (20): we claim that for each $x_0 \in \mathbb{R}^3$ there exists $\delta = \delta(x_0) > 0$ such that

$$(22) \quad u \in L^4(0, T; L^{12}(B(x_0, T - t + \delta))).$$

To prove this, notice first that (18) and (19) together imply that given any constant $\epsilon > 0$, we can find $0 < s < T$ for which

$$(23) \quad \int_{B(x_0, T-s)} u_t^2 + |Du|^2 + u^6 dx \leq \epsilon.$$

Then

$$(24) \quad \int_{B(x_0, T-s+\delta)} u_t^2 + |Du|^2 + u^6 dx \leq 2\epsilon$$

for some small $\delta > 0$. Consequently, a standard energy calculation shows

$$\sup_{s \leq t < T} \int_{B(x_0, T-t+\delta)} u^6 dx \leq C\epsilon.$$

The techniques introduced above in step 4 then establish (22).

6. In summary so far: for each point x_0 , there exists $\delta = \delta(x_0) > 0$ such that u verifies (22). As u has compact support, we can find finitely many points $\{x_k\}_{k=1}^N$ such that $\text{spt } u(\cdot, T) \subset \bigcup_{k=1}^N B(x_k, \delta(x_k))$. It follows that

$$u \in L^4(0, T; L^{12}(\mathbb{R}^3)).$$

Then step 2 implies $u \in L^\infty(\mathbb{R}^3 \times [0, T])$. As noted in step 1, proving this was our goal. \square

12.5. NONEXISTENCE OF SOLUTIONS

We next identify some circumstances under which the semilinear nonlinear wave equation $\square u + f(u) = 0$ does not have a solution existing for all time. Our method will be to introduce appropriate integral quantities depending on the time variable t , for which we can then derive differential inequalities, involving convexity, that lead to contradictions.

12.5.1. Nonexistence for negative energy.

We study first the initial-value problem

$$(1) \quad \begin{cases} u_{tt} - \Delta u + f(u) = 0 & \text{in } \mathbb{R}^n \times (0, \infty) \\ u = g, \quad u_t = h & \text{on } \mathbb{R}^n \times \{t = 0\}. \end{cases}$$

We always assume that $g, h \in C_c^\infty(\mathbb{R}^n)$ and that $f(0) = 0$. Hence a smooth solution, if it exists, will have compact support in space for each time, according to Theorem 3 in §12.1.2.

THEOREM 1 (Nonexistence for negative energy). *Assume that for some constant $\lambda > 2$ we have the inequality*

$$(2) \quad zf(z) \leq \lambda F(z) \quad (z \in \mathbb{R}).$$

Suppose also that the energy is negative:

$$(3) \quad E(0) = \int_{\mathbb{R}^n} \frac{1}{2} (|Dg|^2 + h^2) + F(g) \, dx < 0.$$

Then there cannot exist for all times $t \geq 0$ a smooth solution u of (1).

Therefore the solution constructed in the proof of Theorem 2 in §12.3.2 cannot in general continue for all time.

Proof. 1. Define

$$I(t) := \frac{1}{2} \int_{\mathbb{R}^n} u^2 \, dx.$$

Then

$$(4) \quad \begin{aligned} I'' &= \int_{\mathbb{R}^n} u_t^2 + uu_{tt} \, dx = \int_{\mathbb{R}^n} u_t^2 + u(\Delta u - f(u)) \, dx \\ &= \int_{\mathbb{R}^n} u_t^2 - |Du|^2 - uf(u) \, dx. \end{aligned}$$

The integration by parts is justified, since u has compact support in space for each time.

According to the conservation of energy, we have

$$E(t) = \int_{\mathbb{R}^n} \frac{1}{2} (u_t^2 + |Du|^2) + F(u) \, dx = E(0)$$

for each time $t > 0$. Add and subtract $(2 + 4\alpha)E(0)$ to (4):

$$I'' = (2 + 2\alpha) \int_{\mathbb{R}^n} u_t^2 dx + 2\alpha \int_{\mathbb{R}^n} |Du|^2 dx \\ + \int_{\mathbb{R}^n} (2 + 4\alpha)F(u) - uf(u) dx - (2 + 4\alpha)E(0).$$

Select $\alpha > 0$ so that $2 + 4\alpha = \lambda$, where λ is the constant in hypothesis (2). Then the last integral term is nonnegative, and hence

$$(5) \quad I'' \geq (2 + 2\alpha) \int_{\mathbb{R}^n} u_t^2 dx - \lambda E(0).$$

Since $I' = \int uu_t dx$, inequality (5) implies

$$(6) \quad (1 + \alpha)(I')^2 \leq (1 + \alpha) \left(\int_{\mathbb{R}^n} u^2 dx \right) \left(\int_{\mathbb{R}^n} u_t^2 dx \right) \leq I(I'' - \beta),$$

for $\beta := -\lambda E(0) > 0$.

2. Put

$$J := I^{-\alpha}.$$

Then (6) lets us compute

$$(7) \quad J'' = \alpha(\alpha + 1)I^{-(\alpha+2)}(I')^2 - \alpha I^{-(\alpha+1)}I'' \leq -\alpha\beta I^{-(\alpha+1)} = -\alpha\beta J^{1+1/\alpha}.$$

This shows that J is a concave function of t . Suppose now that $J'(t_0) < 0$ for some time $t_0 > 0$. Then the concavity of J implies

$$J(t) \leq J(t_0) + (t - t_0)J'(t_0) \quad (t \geq 0),$$

and this inequality provides the contradiction that $J(t) < 0$ for large times t . Assume instead that $J' \geq 0$ for all $t \geq 0$. Then from (7) it follows that

$$J'' \leq -\alpha\beta J^{1+1/\alpha}(0) =: -\gamma.$$

We have $\gamma > 0$, since our negative energy hypothesis (3) implies $g \neq 0$. Thus

$$J'(t) \leq J'(0) - \gamma t < 0$$

for large t , and we again reach a contradiction. □

12.5.2. Nonexistence for small initial data.

Satisfying the negative energy hypothesis (3) in the previous subsection requires that g not be too small. Remarkably, certain semilinear wave equations do not possess solutions existing for all times $t > 0$, even for certain arbitrarily small and smooth initial data g, h .

As an example, consider this initial-value problem in three space dimensions:

$$(8) \quad \begin{cases} u_{tt} - \Delta u - |u|^p = 0 & \text{in } \mathbb{R}^3 \times (0, \infty) \\ u = g, \quad u_t = h & \text{on } \mathbb{R}^3 \times \{t = 0\}. \end{cases}$$

Take any smooth initial data with

$$(9) \quad \int_{\mathbb{R}^3} g \, dx > 0, \quad \int_{\mathbb{R}^3} h \, dx > 0$$

and $\text{spt } g, \text{spt } h \subset B(0, R)$. We will deduce from these conditions alone that there is no solution, provided the exponent $p > 1$ is small enough.

THEOREM 2 (Nonexistence for small data). *Assume that*

$$1 < p < 1 + \sqrt{2}.$$

Then under the above conditions on g and h , the initial-value problem (8) does not have a smooth solution u existing for all times $t \geq 0$.

This statement should be contrasted with the global existence of smooth solutions of $\square u + |u|^{p-1}u = 0$ in $\mathbb{R}^3 \times (0, \infty)$ shown in §12.3 and §12.4 for $1 \leq p \leq 5$.

Proof. 1. We assume to the contrary that u is in fact a solution and derive a contradiction. Since the initial data have support within $B(0, R)$, Theorem 3 from §12.1.2 implies that $u(\cdot, t)$ is supported within the ball $B(0, R + t)$.

Put

$$I(t) = \int_{\mathbb{R}^3} u \, dx.$$

Then

$$I'' = \int_{\mathbb{R}^3} u_{tt} \, dx = \int_{\mathbb{R}^3} |u|^p \, dx.$$

Since $u(\cdot, t)$ vanishes outside the ball $B(0, R + t)$, we have

$$I = \int_{B(0, R+t)} u \, dx \leq \left(\int_{\mathbb{R}^3} |u|^p \, dx \right)^{1/p} |B(0, R + t)|^{1-1/p};$$

and therefore

$$(10) \quad I'' \geq cI^p(1+t)^{-3(p-1)}$$

for a constant $c > 0$.

2. We next introduce the solution v of the linear wave equation

$$(11) \quad \begin{cases} v_{tt} - \Delta v = 0 & \text{in } \mathbb{R}^3 \times (0, \infty) \\ v = g, v_t = h & \text{on } \mathbb{R}^3 \times \{t = 0\}. \end{cases}$$

According to the retarded potential formula (44) in §2.4.2, we have

$$u(x, t) = v(x, t) + \frac{1}{4\pi} \int_{B(x, t)} \frac{|u|^p(y, t - |y - x|)}{|y - x|} dy \geq v(x, t).$$

Now (11) and (9) imply

$$\int_{\mathbb{R}^n} v dx = c_1 + c_2 t$$

for constants $c_1, c_2 > 0$. Furthermore, since we are working in $n = 3$ space dimensions, Huygens' principle tells us that v has support within the annular region $A := B(0, t + R) - B(0, t - R)$, which has volume $|A| \leq C(1 + t)^2$. Hence

$$\begin{aligned} c_1 + c_2 t &= \int_A v dx \leq \int_A u dx \\ &\leq C \left(\int_{\mathbb{R}^3} |u|^p dx \right)^{1/p} (1 + t)^{2(1-1/p)}. \end{aligned}$$

It follows that

$$I'' = \int_{\mathbb{R}^3} |u|^p dx \geq c(1 + t)^{2-p}$$

for some constant $c > 0$. Since $I(0), I'(0) > 0$, we deduce that

$$(12) \quad I \geq c(1 + t)^{4-p}.$$

3. Let $\varepsilon > 0$ and as follows combine (10) and (12):

$$\begin{aligned} I'' &\geq cI^{1+\varepsilon} I^{p-1-\varepsilon} (1+t)^{-3(p-1)} \\ &\geq cI^{1+\varepsilon} (1+t)^{(4-p)(p-1-\varepsilon)} (1+t)^{-3(p-1)} \\ &= cI^{1+\varepsilon} (1+t)^{-\mu} \end{aligned}$$

for $\mu := (p-1)^2 + \varepsilon(4-p)$. Since $1 < p < 1 + \sqrt{2}$, we can fix $\varepsilon > 0$ so small that

$$(13) \quad 0 < \mu < 2.$$

Since $I' > 0$,

$$I''I' \geq cI'I^{1+\varepsilon}(1+t)^{-\mu};$$

and thus

$$((I')^2)' \geq c(I^{2+\varepsilon}(1+t)^{-\mu})' + c\mu I^{2+\varepsilon}(1+t)^{-\mu+1} \geq c(I^{2+\varepsilon}(1+t)^{-\mu})'.$$

Consequently

$$(14) \quad (I')^2(t) \geq (I')^2(0) + c(I^{2+\varepsilon}(1+t)^{-\mu} - I^{2+\varepsilon}(0)).$$

Now (12) implies $I^{2+\varepsilon}(t)(1+t)^{-\mu} \geq 2I^{2+\varepsilon}(0)$ for large enough times, say, $t \geq t_0$. Therefore we can deduce from (14) that

$$(I')^2 \geq cI^{2+\varepsilon}(1+t)^{-\mu}$$

provided $t \geq t_0$. Then

$$(I^{-\varepsilon/2})' = -\frac{\varepsilon}{2}I^{-\frac{\varepsilon}{2}-1}I' \leq -\frac{c\varepsilon}{2}(1+t)^{-\mu/2}.$$

Integrating, we see that

$$0 \leq I^{-\varepsilon/2}(t) \leq I^{-\varepsilon/2}(t_0) - \frac{c\varepsilon}{2} \int_{t_0}^t \frac{ds}{(1+s)^{\mu/2}}.$$

This is a contradiction, since $\mu < 2$ according to (13) and so the integral on the right diverges as $t \rightarrow \infty$. \square

12.6. PROBLEMS

The following problem set includes questions on both linear and nonlinear wave equations, as well as the related nonlinear Schrödinger equation. All given functions are assumed smooth, unless otherwise stated.

1. Assume u has compact support in space and solves the quasilinear wave equation

$$u_{tt} - \sum_{i=1}^n (L_{p_i}(Du))_{x_i} = 0 \quad \text{in } \mathbb{R}^n \times (0, \infty).$$

Determine the appropriate energy $E(t)$ and show $\dot{E} \equiv 0$.

2. Let u solve the *Klein–Gordon equation*

$$(*) \quad \begin{cases} u_{tt} - \Delta u + m^2 u = 0 & \text{in } \mathbb{R}^n \times (0, \infty) \\ u = g, \quad u_t = h & \text{on } \mathbb{R}^n \times \{t = 0\}. \end{cases}$$

(a) Show that the energy

$$E(t) := \frac{1}{2} \int_{\mathbb{R}^n} u_t^2 + |Du|^2 + m^2 u^2 dx \quad (t \geq 0)$$

is constant in time.

(b) Modify the proof in §4.3.1 showing asymptotic equipartition of energy for the wave equation to prove that

$$\lim_{t \rightarrow \infty} \int_{\mathbb{R}^n} |Du|^2 + m^2 u^2 dx = E(0).$$

3. Suppose u solves the initial value problem $(*)$ from Problem 2 for the Klein–Gordon equation. Write $\bar{x} = (x, x_{n+1})$ for $x \in \mathbb{R}^n$ and define

$$\bar{u}(\bar{x}, t) := u(x, t) \cos(mx_{n+1}).$$

(a) Show that \bar{u} solves the wave equation $\square \bar{u} = 0$ in $\mathbb{R}^{n+1} \times (0, \infty)$.

(b) Derive a formula for the solution of the initial-value problem for the Klein–Gordon equation when $n = 1$. (This is a variant of the *method of descent*, introduced in §2.4.1.)

4. Assume u solves

$$u_{tt} - \Delta u + du_t = 0 \quad \text{in } \mathbb{R}^n \times (0, \infty),$$

which for $d > 0$ is a damped wave equation. Find a simple exponential term that, when multiplied by u , gives a solution v of

$$v_{tt} - \Delta v + cv = 0$$

for a constant $c < 0$. (This is the opposite of the sign for the Klein–Gordon equation.)

5. Check that for each given $y \in \mathbb{R}^n$, $y \neq 0$, the function $u = e^{i(x \cdot y - \sigma t)}$ solves the Klein–Gordon equation

$$u_{tt} - \Delta u + m^2 u = 0$$

provided $\sigma = (|y|^2 + m^2)^{\frac{1}{2}}$. The phase velocity of this plane wave solution is $\frac{\sigma}{|y|} > 1$. Why does this not contradict the assertions in §12.1 that the speed of propagation for solutions is less than or equal to one?

6. Suppose

$$\begin{cases} u_{tt} - \Delta u = 0 & \text{in } \mathbb{R}^3 \times (0, \infty) \\ u = g, u_t = h & \text{on } \mathbb{R}^3 \times \{t = 0\}, \end{cases}$$

where g, h have compact support. Show there exists a constant C such that

$$|u(x, t)| \leq C/t \quad (x \in \mathbb{R}^3, t > 0).$$

(Hint: Use the representation formula for the solution from §2.4.1.)

7. Let u solve

$$\begin{cases} u_{tt} - \Delta u = 0 & \text{in } \mathbb{R}^2 \times (0, \infty) \\ u = g, u_t = h & \text{on } \mathbb{R}^2 \times \{t = 0\}, \end{cases}$$

where g, h have compact support. Show that

$$|u(x, t)| \leq C/t^{\frac{1}{2}} \quad (x \in \mathbb{R}^2, t > 0)$$

for some constant C .

8. Suppose u^ε solves the linear wave equation in $n = 2$ space dimensions, with the initial conditions $u_t^\varepsilon = h \equiv 0$ and

$$u^\varepsilon = g^\varepsilon := \begin{cases} e^{-\varepsilon \frac{(r-2)^2}{(r-1)(3-r)}} & \text{if } 1 < r < 3 \\ 0 & \text{otherwise,} \end{cases}$$

where $r = |x|$. Show that although $|g^\varepsilon| \leq 1$, we have

$$\max_{\mathbb{R}^2 \times [0, 4]} |u^\varepsilon| \rightarrow \infty \quad \text{as } \varepsilon \rightarrow 0.$$

(Hint: Use (26) in §2.4.1 to compute $u^\varepsilon(0, t)$ for $t > 3$.)

9. (Kelvin transform for wave equation) The *hyperbolic Kelvin transform* $\mathcal{K}u = \bar{u}$ of a function $u : \mathbb{R}^n \times \mathbb{R} \rightarrow \mathbb{R}$ is

$$\bar{u}(x, t) := u(\bar{x}, \bar{t}) ||\bar{x}|^2 - \bar{t}^2|^{\frac{n-1}{2}} = u\left(\frac{x}{|x|^2 - t^2}, \frac{t}{|x|^2 - t^2}\right) \frac{1}{||x|^2 - t^2|^{\frac{n-1}{2}}},$$

provided $|x|^2 \neq t^2$, where

$$\bar{x} = \frac{x}{|x|^2 - t^2}, \quad \bar{t} = \frac{t}{|x|^2 - t^2}.$$

Show that if $\square u = 0$, then $\square \bar{u} = 0$.

(Compare with Problem 11 in Chapter 2.)

10. Assume that u and v solve the system

$$(*) \quad \begin{cases} (u - v)_t = 2a \sin\left(\frac{u+v}{2}\right) \\ (u + v)_x = \frac{2}{a} \sin\left(\frac{u-v}{2}\right) \end{cases}$$

where $a \neq 0$. Show that both $w := u$ and $w := v$ solve the *sine-Gordon equation* in the form

$$w_{xt} = \sin w.$$

Why is this equivalent to the PDE $\square w = \sin w$?

11. (Continuation) Alternatively, given a solution v of the sine-Gordon equation, we can try to solve the system $(*)$ to build a second solution u . This procedure is called a *Bäcklund transformation*.

Start with the trivial solution $v \equiv 0$, and use the Bäcklund transformation to compute for each choice of the parameter a another solution u . (Hint: First show u must have the form $f(at + x/a)$. Also show that $(\tan(u/4))_t = a \tan(u/4)$.)

12. Prove the Sobolev-type inequality (6) in §12.2.2.

(Hints: If $u \in H^k$ and β is a multiindex with $|\beta| \leq k$, we have the estimate $\|D^\beta u\|_{L^p} \leq C\|u\|_{H^k}$, where p satisfies (a) $p = \infty$ if $\frac{1}{2} - \frac{k}{n} + \frac{|\beta|}{n} < 0$; (b) $2 \leq p < \infty$ if $\frac{1}{2} - \frac{k}{n} + \frac{|\beta|}{n} = 0$; and (c) $\frac{1}{p} = \frac{1}{2} - \frac{k-|\beta|}{n}$ if $\frac{1}{2} - \frac{k}{n} + \frac{|\beta|}{n} > 0$. Assume that the multiindices $|\beta_1|, \dots, |\beta_r|$ satisfy (a) above, $|\beta_{r+1}|, \dots, |\beta_s|$ satisfy (b), and $|\beta_{s+1}|, \dots, |\beta_m|$ satisfy (c). Estimate $\|D^{\beta_1} u_1 \cdots D^{\beta_m} u_m\|_{L^2}^2$.)

13. A smooth function $\mathbf{u} : \mathbb{R}^n \times [0, \infty) \rightarrow \mathbb{R}^m$, $\mathbf{u} = (u^1, \dots, u^m)$, is called a *wave map* into the unit sphere $S^{m-1} = \partial B(0, 1) \subset \mathbb{R}^m$ provided that everywhere in $\mathbb{R}^n \times [0, \infty)$, $|\mathbf{u}| = 1$ and $\mathbf{u}_{tt} - \Delta \mathbf{u}$ is perpendicular to S^{m-1} at \mathbf{u} .

Show that therefore \mathbf{u} solves the system of PDE

$$\mathbf{u}_{tt} - \Delta \mathbf{u} = (|\mathbf{D}\mathbf{u}|^2 - |\mathbf{u}_t|^2)\mathbf{u}.$$

14. Prove that if \mathbf{u} is a wave map into the unit sphere, with compact support in space, we have conservation of energy:

$$\frac{d}{dt} \int_{\mathbb{R}^n} |\mathbf{u}_t|^2 + |\mathbf{D}\mathbf{u}|^2 dx = 0.$$

15. (Small energy for $p = 5$) Adapt the proof of Theorem 3 in §12.3.3 to show the existence of a smooth solution of

$$\begin{cases} u_{tt} - \Delta u + u^5 = 0 & \text{in } \mathbb{R}^3 \times (0, \infty) \\ u = g, u_t = h & \text{on } \mathbb{R}^3 \times \{t = 0\}, \end{cases}$$

provided the energy $E(0)$ is sufficiently small.

(Hint: Modify estimate (24) in §12.3.3 by introducing the L^6 -norm of g .)

Many techniques developed for semilinear wave equations have counterparts for nonlinear Schrödinger (NLS) equations, to which we devote the remaining exercises.

16. Let u be a complex-valued solution of the nonlinear Schrödinger equation

$$(*) \quad iu_t + \Delta u = f(|u|^2)u \quad \text{in } \mathbb{R}^n \times (0, \infty).$$

Here $f : \mathbb{R} \rightarrow \mathbb{R}$. Demonstrate that if $\xi \in \mathbb{R}^n$, then

$$w(x, t) := e^{\frac{i}{4}(2\xi \cdot x - |\xi|^2 t)} u(x - \xi t, t)$$

also solves the NLS equation. This shows the *Galilean invariance* of solutions.

17. Assume u solves the nonlinear Schrödinger equation (*) from Problem 16 and decays rapidly, along with its derivatives, as $|x| \rightarrow \infty$. Derive these identities:

$$\frac{d}{dt} \int_{\mathbb{R}^n} |u|^2 dx = 0 \quad (\text{conservation of mass}),$$

$$\frac{d}{dt} \int_{\mathbb{R}^n} |Du|^2 + F(|u|^2) dx = 0 \quad (\text{conservation of energy}),$$

where $F' = f$, and

$$\frac{d}{dt} \int_{\mathbb{R}^n} \frac{\bar{u}Du - uD\bar{u}}{2i} dx = 0 \quad (\text{conservation of momentum}).$$

Remember that $|u|^2 = u\bar{u}$.

18. (Continuation)
- (a) Under the hypotheses of the previous problem, derive the identity

$$\begin{aligned} \frac{d^2}{dt^2} \int_{\mathbb{R}^n} |x|^2 |u|^2 dx \\ = 8 \int_{\mathbb{R}^n} |Du|^2 dx + 4n \int_{\mathbb{R}^n} f(|u|^2) |u|^2 - F(|u|^2) dx. \end{aligned}$$

- (b) Use (a) to show that there does not exist a solution of the cubic-NLS equation

$$iu_t + \Delta u + |u|^2 u = 0 \quad \text{in } \mathbb{R}^n \times (0, \infty)$$

existing for all times $t \geq 0$, if

$$E(0) = \int_{\mathbb{R}^n} |Du(\cdot, 0)|^2 - \frac{|u(\cdot, 0)|^4}{2} dx < 0$$

and $n \geq 2$. (R. Glassey, *J. Math. Physics* 18 (1977), 1794–1797)

12.7. REFERENCES

- D. Tataru provided me with comments and suggestions for this chapter.
- Section 12.2 Shatah–Struwe [S-S] was my primary source for this section.
- Section 12.3 See Shatah–Struwe [S-S] for more. Theorem 1 follows I. Segal (*Bull. Soc. Math. France* 91 (1963), 129–135) and Theorems 2 and 3 follow K. Jörgens (*Math. Z.* 77 (1961), 295–308).
- Section 12.4 M. Grillakis (*Ann. of Math.* 132 (1990), 485–509) proved the key estimate in Theorem 2 and the existence Theorem 3. Our presentation of the latter uses simplifications due to Smith and Sogge: consult Sogge [So] and Shatah–Struwe [S-S].
- Section 12.5 See Strauss [St1], John [J3].
- Section 12.6 Problem 8 is from Protter–Weinberger [P-W]. Problem 15: J. Rauch observed that small energy implies global existence for $p = 5$. J. Holmer helped me with Problems 16–18.

APPENDICES

- A. Notation
- B. Inequalities
- C. Calculus
- D. Functional analysis
- E. Measure theory

APPENDIX A: NOTATION

A.1. Notation for matrices.

- (i) We write $A = ((a_{ij}))$ to mean A is an $m \times n$ matrix with $(i, j)^{\text{th}}$ entry a_{ij} . (Sometimes, as in §8.1.4, it will be convenient to use superscripts to denote rows.)
A diagonal matrix is denoted $\text{diag}(d_1, \dots, d_n)$.
- (ii) $\mathbb{M}^{m \times n}$ = space of real $m \times n$ matrices.
 \mathbb{S}^n = space of real, symmetric $n \times n$ matrices.
- (iii) $\text{tr } A$ = trace of the matrix A .
- (iv) $\det A$ = determinant of the matrix A .
- (v) $\text{cof } A$ = cofactor matrix of A (see §8.1.4).
- (vi) A^T = transpose of the matrix A .
- (vii) If $A = ((a_{ij}))$ and $B = ((b_{ij}))$ are $m \times n$ matrices, then

$$A : B = \sum_{i=1}^m \sum_{j=1}^n a_{ij} b_{ij},$$

and

$$|A| = (A : A)^{\frac{1}{2}} = \left(\sum_{i=1}^m \sum_{j=1}^n a_{ij}^2 \right)^{\frac{1}{2}}.$$

(viii) If $A \in \mathbb{S}^n$ and, as below, $x = (x_1, \dots, x_n) \in \mathbb{R}^n$, the corresponding quadratic form is $x \cdot Ax = \sum_{i,j=1}^n a_{ij} x_i x_j$.

(ix) If $A, B \in \mathbb{S}^n$, we write

$$A \geq B$$

to mean that $A - B$ is nonnegative definite. In particular, we write

$$A \geq \theta I$$

if $x \cdot Ax \geq \theta |x|^2$ for all $x \in \mathbb{R}^n$.

(x) We sometimes will write yA to mean $A^T y$, for $A \in \mathbb{M}^{m \times n}$ and $y \in \mathbb{R}^m$.

A.2. Geometric notation.

(i) $\mathbb{R}^n = n$ -dimensional real Euclidean space, $\mathbb{R} = \mathbb{R}^1$.

$S^{n-1} = \partial B(0, 1) = (n - 1)$ -dimensional unit sphere in \mathbb{R}^n .

(ii) $e_i = (0, \dots, 0, 1, \dots, 0) = i^{\text{th}}$ standard coordinate vector.

(iii) A typical point in \mathbb{R}^n is $x = (x_1, \dots, x_n)$.

We will also, depending upon the context, regard x as a row or column vector.

(iv) $\mathbb{R}_+^n = \{x = (x_1, \dots, x_n) \in \mathbb{R}^n \mid x_n > 0\} = \text{open upper half-space}$.

$$\mathbb{R}_+ = \{x \in \mathbb{R} \mid x > 0\}.$$

(v) A point in \mathbb{R}^{n+1} will often be denoted as $(x, t) = (x_1, \dots, x_n, t)$, and we usually interpret $t = x_{n+1} = \text{time}$.

A point $x \in \mathbb{R}^n$ will sometimes be written $x = (x', x_n)$ for $x' = (x_1, \dots, x_{n-1}) \in \mathbb{R}^{n-1}$.

(vi) U, V , and W usually denote open subsets of \mathbb{R}^n . We write

$$V \subset\subset U$$

if $V \subset \bar{V} \subset U$ and \bar{V} is compact, and say V is *compactly contained* in U .

(vii) $\partial U = \text{boundary of } U$, $\bar{U} = U \cup \partial U = \text{closure of } U$.

- (viii) $U_T = U \times (0, T]$.
- (ix) $\Gamma_T = \bar{U}_T - U_T =$ parabolic boundary of U_T .
- (x) $B(x, r) = \{y \in \mathbb{R}^n \mid |x - y| < r\} =$ closed ball in \mathbb{R}^n with center x and radius $r > 0$.
- (xi) $B^0(x, r) =$ open ball with center x , radius $r > 0$.
- (xii) $C(x, t; r) = \{y \in \mathbb{R}^n, s \in \mathbb{R} \mid |x - y| \leq r, t - r^2 \leq s \leq t\} =$ closed cylinder with top center (x, t) , radius $r > 0$, height r^2 .
- (xiii) $\alpha(n) =$ volume of unit ball $B(0, 1)$ in $\mathbb{R}^n = \frac{\pi^{n/2}}{\Gamma(\frac{n}{2}+1)}$.
 $n\alpha(n) =$ surface area of unit sphere $\partial B(0, 1)$ in \mathbb{R}^n .
- (xiv) If $a = (a_1, \dots, a_n)$ and $b = (b_1, \dots, b_n)$ belong to \mathbb{R}^n ,

$$a \cdot b = \sum_{i=1}^n a_i b_i, \quad |a| = \left(\sum_{i=1}^n a_i^2 \right)^{\frac{1}{2}}.$$

- (xv) $\mathbb{C}^n = n$ -dimensional complex space; $\mathbb{C} =$ complex plane.

If $z \in \mathbb{C}$, we write $\operatorname{Re}(z)$ for the real part of z and $\operatorname{Im}(z)$ for the imaginary part.

A.3. Notation for functions.

- (i) If $u : U \rightarrow \mathbb{R}$, we write

$$u(x) = u(x_1, \dots, x_n) \quad (x \in U).$$

We say u is *smooth* provided u is infinitely differentiable.

- (ii) If u, v are two functions, we write

$$u \equiv v$$

to mean that u is identically equal to v ; that is, the functions u, v agree for all values of their arguments. We use the notation

$$u := v$$

to define u as equaling v .

The support of a function u is denoted

$$\operatorname{spt} u.$$

(iii) $u^+ = \max(u, 0)$, $u^- = -\min(u, 0)$, $u = u^+ - u^-$, $|u| = u^+ + u^-$.

The *sign* function is

$$\operatorname{sgn}(x) = \begin{cases} 1 & \text{if } x > 0 \\ 0 & \text{if } x = 0 \\ -1 & \text{if } x < 0. \end{cases}$$

(iv) If $\mathbf{u} : U \rightarrow \mathbb{R}^m$, we write

$$\mathbf{u}(x) = (u^1(x), \dots, u^m(x)) \quad (x \in U).$$

The function u^k is the k^{th} component of \mathbf{u} , $k = 1, \dots, m$.

(v) If Σ is a smooth $(n - 1)$ -dimensional surface in \mathbb{R}^n , we write

$$\int_{\Sigma} f \, dS$$

for the integral of f over Σ , with respect to $(n - 1)$ -dimensional surface measure. If C is a curve in \mathbb{R}^n , we denote by

$$\int_C f \, dl$$

the integral of f over C with respect to arclength.

(vi) Averages:

$$\int_{B(x,r)} f \, dy := \frac{1}{\alpha(n)r^n} \int_{B(x,r)} f \, dy = \text{average of } f \text{ over the ball } B(x, r)$$

and

$$\int_{\partial B(x,r)} f \, dS := \frac{1}{n\alpha(n)r^{n-1}} \int_{\partial B(x,r)} f \, dS \\ = \text{average of } f \text{ over the sphere } \partial B(x, r).$$

More generally,

$$\int_E f \, d\mu := \frac{1}{\mu(E)} \int_E f \, d\mu = \text{average of } f \text{ over set } E,$$

provided $\mu(E) > 0$.

(vii) $\chi_E(x) = \begin{cases} 1 & \text{if } x \in E \\ 0 & \text{if } x \notin E; \end{cases}$ χ_E is the *indicator function* of E .

(viii) A function $u : U \rightarrow \mathbb{R}$ is called *Lipschitz continuous* if

$$|u(x) - u(y)| \leq C|x - y|$$

for some constant C and all $x, y \in U$. We write

$$\operatorname{Lip}[u] := \sup_{\substack{x, y \in U \\ x \neq y}} \frac{|u(x) - u(y)|}{|x - y|}.$$

(ix) The convolution of the functions f, g is denoted

$$f * g.$$

Notation for derivatives. Assume $u : U \rightarrow \mathbb{R}$, $x \in U$.

(i) $\frac{\partial u}{\partial x_i}(x) = \lim_{h \rightarrow 0} \frac{u(x+he_i) - u(x)}{h}$, provided this limit exists.

(ii) We usually write u_{x_i} for $\frac{\partial u}{\partial x_i}$.

(iii) Similarly, $\frac{\partial^2 u}{\partial x_i \partial x_j} = u_{x_i x_j}$, $\frac{\partial^3 u}{\partial x_i \partial x_j \partial x_k} = u_{x_i x_j x_k}$, etc.

(iv) *Multiindex Notation:*

(a) A vector of the form $\alpha = (\alpha_1, \dots, \alpha_n)$, where each component α_i is a nonnegative integer, is called a *multiindex* of order

$$|\alpha| = \alpha_1 + \dots + \alpha_n.$$

(b) Given a multiindex α , define

$$D^\alpha u(x) := \frac{\partial^{|\alpha|} u(x)}{\partial x_1^{\alpha_1} \dots \partial x_n^{\alpha_n}} = \partial_{x_1}^{\alpha_1} \dots \partial_{x_n}^{\alpha_n} u.$$

(c) If k is a nonnegative integer,

$$D^k u(x) := \{D^\alpha u(x) \mid |\alpha| = k\},$$

the set of all partial derivatives of order k . Assigning some ordering to the various partial derivatives, we can also regard $D^k u(x)$ as a point in \mathbb{R}^{n^k} .

(d) $|D^k u| = \left(\sum_{|\alpha|=k} |D^\alpha u|^2 \right)^{1/2}$.

(e) *Special Cases:* If $k = 1$, we regard the elements of Du as being arranged in a vector:

$$Du := (u_{x_1}, \dots, u_{x_n}) = \text{gradient vector.}$$

Therefore $Du \in \mathbb{R}^n$. We will sometimes also write

$$u_r := \frac{x}{|x|} \cdot Du$$

for the radial derivative of u .

If $k = 2$, we regard the elements of $D^2 u$ as being arranged in a matrix:

$$D^2 u := \begin{pmatrix} u_{x_1 x_1} & \dots & u_{x_1 x_n} \\ & \ddots & \\ u_{x_n x_1} & \dots & u_{x_n x_n} \end{pmatrix}_{n \times n} = \text{Hessian matrix.}$$

Therefore $D^2 u \in \mathbb{S}^n$, the space of real symmetric $n \times n$ matrices.

(v) $\Delta u = \sum_{i=1}^n u_{x_i x_i} = \text{tr}(D^2 u) = \text{Laplacian of } u$.

(vi) We sometimes employ a subscript attached to the symbols D , D^2 , etc. to denote the variables being differentiated. For example if $u = u(x, y)$ ($x \in \mathbb{R}^n$, $y \in \mathbb{R}^m$), then $D_x u = (u_{x_1}, \dots, u_{x_n})$, $D_y u = (u_{y_1}, \dots, u_{y_m})$.

Function spaces.

(i) $C(U) = \{u : U \rightarrow \mathbb{R} \mid u \text{ continuous}\}$.

$C(\bar{U}) = \{u \in C(U) \mid u \text{ is uniformly continuous on bounded subsets of } U\}$.

$C^k(U) = \{u : U \rightarrow \mathbb{R} \mid u \text{ is } k\text{-times continuously differentiable}\}$.

$C^k(\bar{U}) = \{u \in C^k(U) \mid D^\alpha u \text{ is uniformly continuous on bounded subsets of } U, \text{ for all } |\alpha| \leq k\}$.

Thus if $u \in C^k(\bar{U})$, then $D^\alpha u$ continuously extends to \bar{U} for each multiindex α , $|\alpha| \leq k$.

(ii) $C^\infty(U) = \{u : U \rightarrow \mathbb{R} \mid u \text{ is infinitely differentiable}\} = \bigcap_{k=0}^\infty C^k(U)$
 $C^\infty(\bar{U}) = \bigcap_{k=0}^\infty C^k(\bar{U})$.

(iii) $C_c(U)$, $C_c^k(U)$, etc. denote these functions in $C(U)$, $C^k(U)$, etc. with *compact support*.

(iv) $L^p(U) = \{u : U \rightarrow \mathbb{R} \mid u \text{ is Lebesgue measurable, } \|u\|_{L^p(U)} < \infty\}$, where

$$\|u\|_{L^p(U)} := \left(\int_U |u|^p dx \right)^{\frac{1}{p}} \quad (1 \leq p < \infty).$$

$L^\infty(U) = \{u : U \rightarrow \mathbb{R} \mid u \text{ is Lebesgue measurable, } \|u\|_{L^\infty(U)} < \infty\}$, where

$$\|u\|_{L^\infty(U)} := \text{ess sup}_U |u|.$$

$L_{\text{loc}}^p(U) = \{u : U \rightarrow \mathbb{R} \mid u \in L^p(V) \text{ for each } V \subset\subset U\}$.

(See also §D.1.)

(v) $\|Du\|_{L^p(U)} = \| \|Du\| \|_{L^p(U)}$.

$\|D^2u\|_{L^p(U)} = \| \|D^2u\| \|_{L^p(U)}$.

(vi) $W^{k,p}(U)$, $H^k(U)$, etc. ($k = 0, 1, 2, \dots$, $1 \leq p \leq \infty$) denote Sobolev spaces: see Chapter 5.

(vii) $C^{k,\beta}(\bar{U})$, $C^{k,\beta}(\bar{U})$ ($k = 0, \dots$, $0 < \beta \leq 1$) denote Hölder spaces: see Chapter 5.

(viii) *Functions of x and t .* It is occasionally useful to introduce spaces of functions with differing smoothness in the x - and t -variables, although there is no standard notation for such spaces. We will for this book write

$$C_1^2(U_T) = \{u : U_T \rightarrow \mathbb{R} \mid u, D_x u, D_x^2 u, u_t \in C(U_T)\}.$$

In particular, if $u \in C_1^2(U_T)$, then $u, D_x u$, etc. are continuous up to the top $U \times \{t = T\}$.

A.4. Vector-valued functions.

(i) If now $m > 1$ and $\mathbf{u} : U \rightarrow \mathbb{R}^m$, $\mathbf{u} = (u^1, \dots, u^m)$, we define

$$D^\alpha \mathbf{u} = (D^\alpha u^1, \dots, D^\alpha u^m) \quad \text{for each multiindex } \alpha.$$

Then

$$D^k \mathbf{u} = \{D^\alpha \mathbf{u} \mid |\alpha| = k\}$$

and

$$|D^k \mathbf{u}| := \left(\sum_{|\alpha|=k} |D^\alpha \mathbf{u}|^2 \right)^{1/2},$$

as before.

(ii) In the special case $k = 1$, we write

$$D\mathbf{u} := \begin{pmatrix} u_{x_1}^1 & \cdots & u_{x_n}^1 \\ \vdots & \ddots & \vdots \\ u_{x_1}^m & \cdots & u_{x_n}^m \end{pmatrix}_{m \times n} = \textit{gradient matrix}.$$

(iii) If $m = n$, we have

$$\operatorname{div} \mathbf{u} := \operatorname{tr}(D\mathbf{u}) = \sum_{i=1}^n u_{x_i}^i = \textit{divergence of } \mathbf{u}.$$

(iv) The spaces $C(U; \mathbb{R}^m)$, $L^p(U; \mathbb{R}^m)$, etc. consist of those functions $\mathbf{u} : U \rightarrow \mathbb{R}^m$, $\mathbf{u} = (u^1, \dots, u^m)$, with $u^i \in C(U)$, $L^p(U)$, etc. ($i = 1, \dots, m$).

Remark on sub- and superscripts. As illustrated above, we will adhere to the convention of setting in boldface *mappings* which take values in \mathbb{R}^m for $m > 1$ (or else in Banach or Hilbert spaces). The component functions of such mappings will be given *superscripts*. On the other hand, a typical *point* $x \in \mathbb{R}^n$ is not boldface and has components with *subscripts*, $x = (x_1, \dots, x_n)$.

Matrix-valued mappings will also be set in boldface, and their component functions written with either superscripts or a mixture of sub- and superscripts, depending upon the context.

A.5. Notation for estimates.

Constants. We employ the letter C to denote any constant that can be explicitly computed in terms of known quantities. The exact value denoted by C may therefore change from line to line in a given computation. The big advantage is that our calculations will be simpler looking, since we continually absorb “extraneous” factors into the term C .

DEFINITIONS. (i) (Big-oh notation.) *We write*

$$f = O(g) \quad \text{as } x \rightarrow x_0,$$

provided there exists a constant C such that

$$|f(x)| \leq C|g(x)|$$

for all x sufficiently close to x_0 .

(ii) (Little-oh notation.) *We write*

$$f = o(g) \quad \text{as } x \rightarrow x_0,$$

provided

$$\lim_{x \rightarrow x_0} \frac{|f(x)|}{|g(x)|} = 0.$$

Remark. The expression “ $O(g)$ ” (or “ $o(g)$ ”) is not by itself defined. There must always be an accompanying limit, for example “as $x \rightarrow x_0$ ” above, although this limit is often implicit.

A.6. Some comments about notation.

The foregoing notation is largely standard within the PDE literature, with a few significant exceptions:

(i) We employ the symbol “ Du ”, and not “ ∇u ”, to denote the gradient of the function u . The reason is that “ D^2u ” then naturally denotes the Hessian matrix of u , whereas “ ∇^2u ” would be confused with the Laplacian. The multiindex notation also looks better with the letter D .

(ii) Most books and papers on partial differential equations denote by “ Ω ” the open subset of \mathbb{R}^n within which a given PDE holds.

As indicated above, we will instead mostly use the symbol “ U ” for such a region. The advantages are several. First of all, since a typical solution is denoted u , it makes sense to denote its domain by U and not to switch to a Greek letter. Furthermore, once we call a given open set U , the letters V and W are then available for subregions.

Lastly, it is important to save Ω as the standard symbol for a probability space. Many important partial differential equations have probabilistic representation formulas (cf. Freidlin [Fd]), and although such are beyond the scope of this book, it seems wise to avoid the possibility of future notational confusion.

APPENDIX B: INEQUALITIES

B.1. Convex functions.

Definition. A function $f : \mathbb{R}^n \rightarrow \mathbb{R}$ is called convex provided

$$(1) \quad f(\tau x + (1 - \tau)y) \leq \tau f(x) + (1 - \tau)f(y)$$

for all $x, y \in \mathbb{R}^n$ and each $0 \leq \tau \leq 1$.

THEOREM 1 (Supporting hyperplanes). Suppose $f : \mathbb{R}^n \rightarrow \mathbb{R}$ is convex. Then for each $x \in \mathbb{R}^n$ there exists $r \in \mathbb{R}^n$ such that the inequality

$$(2) \quad f(y) \geq f(x) + r \cdot (y - x)$$

holds for all $y \in \mathbb{R}^n$.

The mapping $y \mapsto f(x) + r \cdot (y - x)$ determines the *supporting hyperplane* to f at x . Inequality (2) says the graph of f lies above each supporting hyperplane. If f is differentiable at x , $r = Df(x)$.

If f is C^2 , then f is convex if and only if $D^2f \geq 0$. The C^2 function f is *uniformly convex* if $D^2f \geq \theta I$ for some constant $\theta > 0$: this means

$$\sum_{i,j=1}^n f_{x_i x_j}(x) \xi_i \xi_j \geq \theta |\xi|^2 \quad (x, \xi \in \mathbb{R}^n).$$

THEOREM 2 (Jensen's inequality). Assume $f : \mathbb{R}^m \rightarrow \mathbb{R}$ is convex and $U \subset \mathbb{R}^n$ is open, bounded. Let $\mathbf{u} : U \rightarrow \mathbb{R}^m$ be summable. Then

$$(3) \quad f\left(\int_U \mathbf{u} \, dx\right) \leq \int_U f(\mathbf{u}) \, dx.$$

Remember from §A.3 the notation $\int_U \mathbf{u} \, dx = \frac{1}{|U|} \int_U \mathbf{u} \, dx =$ average of \mathbf{u} over U .

Proof. Since f is convex, for each $p \in \mathbb{R}^m$ there exists $r \in \mathbb{R}^m$ such that

$$f(q) \geq f(p) + r \cdot (q - p) \quad \text{for all } q \in \mathbb{R}^m.$$

Let $p = \int_U \mathbf{u} \, dy$, $q = \mathbf{u}(x)$:

$$f(\mathbf{u}(x)) \geq f\left(\int_U \mathbf{u} \, dy\right) + r \cdot \left(\mathbf{u}(x) - \int_U \mathbf{u} \, dy\right).$$

Integrate with respect to x over U . □

We discuss convex functions more fully in §3.3.2 and §9.6.1.

B.2. Useful inequalities.

Following is a collection of elementary, but fundamental, inequalities. These estimates are continually employed throughout the text and should be memorized.

a. Cauchy's inequality.

$$(4) \quad ab \leq \frac{a^2}{2} + \frac{b^2}{2} \quad (a, b \in \mathbb{R}).$$

Proof. $0 \leq (a - b)^2 = a^2 - 2ab + b^2$. □

b. Cauchy's inequality with ϵ .

$$(5) \quad ab \leq \epsilon a^2 + \frac{b^2}{4\epsilon} \quad (a, b > 0, \epsilon > 0).$$

Proof. Write

$$ab = ((2\epsilon)^{1/2}a) \left(\frac{b}{(2\epsilon)^{1/2}} \right)$$

and apply Cauchy's inequality. □

c. Young's inequality. Let $1 < p, q < \infty$, $\frac{1}{p} + \frac{1}{q} = 1$. Then

$$(6) \quad ab \leq \frac{a^p}{p} + \frac{b^q}{q} \quad (a, b > 0).$$

Proof. The mapping $x \mapsto e^x$ is convex, and consequently

$$ab = e^{\log a + \log b} = e^{\frac{1}{p} \log a^p + \frac{1}{q} \log b^q} \leq \frac{1}{p} e^{\log a^p} + \frac{1}{q} e^{\log b^q} = \frac{a^p}{p} + \frac{b^q}{q}. \quad \square$$

d. Young's inequality with ϵ .

$$(7) \quad ab \leq \epsilon a^p + C(\epsilon) b^q \quad (a, b > 0, \epsilon > 0)$$

for $C(\epsilon) = (\epsilon p)^{-q/p} q^{-1}$.

Proof. Write $ab = ((\epsilon p)^{1/p} a) \left(\frac{b}{(\epsilon p)^{1/p}} \right)$ and apply Young's inequality. □

e. Hölder's inequality. Assume $1 \leq p, q \leq \infty$, $\frac{1}{p} + \frac{1}{q} = 1$. Then if $u \in L^p(U)$, $v \in L^q(U)$, we have

$$(8) \quad \int_U |uv| \, dx \leq \|u\|_{L^p(U)} \|v\|_{L^q(U)}.$$

Proof. By homogeneity, we may assume $\|u\|_{L^p} = \|v\|_{L^q} = 1$. Then Young's inequality implies for $1 < p, q < \infty$ that

$$\int_U |uv| dx \leq \frac{1}{p} \int_U |u|^p dx + \frac{1}{q} \int_U |v|^q dx = 1 = \|u\|_{L^p} \|v\|_{L^q}.$$

□

f. Minkowski's inequality. Assume $1 \leq p \leq \infty$ and $u, v \in L^p(U)$. Then

$$(9) \quad \|u + v\|_{L^p(U)} \leq \|u\|_{L^p(U)} + \|v\|_{L^p(U)}.$$

Proof.

$$\begin{aligned} \|u + v\|_{L^p(U)}^p &= \int_U |u + v|^p dx \leq \int_U |u + v|^{p-1} (|u| + |v|) dx \\ &\leq \left(\int_U |u + v|^p dx \right)^{\frac{p-1}{p}} \left(\left(\int_U |u|^p dx \right)^{1/p} + \left(\int_U |v|^p dx \right)^{1/p} \right) \\ &= \|u + v\|_{L^p(U)}^{p-1} (\|u\|_{L^p(U)} + \|v\|_{L^p(U)}). \end{aligned}$$

□

Remark. Similar proofs establish the discrete versions of Hölder's and Minkowski's inequalities:

$$(10) \quad \begin{cases} \left| \sum_{k=1}^n a_k b_k \right| \leq \left(\sum_{k=1}^n |a_k|^p \right)^{\frac{1}{p}} \left(\sum_{k=1}^n |b_k|^q \right)^{\frac{1}{q}}, \\ \left(\sum_{k=1}^n |a_k + b_k|^p \right)^{\frac{1}{p}} \leq \left(\sum_{k=1}^n |a_k|^p \right)^{\frac{1}{p}} + \left(\sum_{k=1}^n |b_k|^p \right)^{\frac{1}{p}}, \end{cases}$$

for $a = (a_1, \dots, a_n), b = (b_1, \dots, b_n) \in \mathbb{R}^n$ and $1 \leq p < \infty, \frac{1}{p} + \frac{1}{q} = 1$.

g. General Hölder inequality. Let $1 \leq p_1, \dots, p_m \leq \infty$, with $\frac{1}{p_1} + \frac{1}{p_2} + \dots + \frac{1}{p_m} = 1$, and assume $u_k \in L^{p_k}(U)$ for $k = 1, \dots, m$. Then

$$(11) \quad \int_U |u_1 \cdots u_m| dx \leq \prod_{k=1}^m \|u_k\|_{L^{p_k}(U)}.$$

Proof. Induction, using Hölder's inequality. □

h. Interpolation inequality for L^p -norms. Assume $1 \leq s \leq r \leq t \leq \infty$ and

$$\frac{1}{r} = \frac{\theta}{s} + \frac{(1-\theta)}{t}.$$

Suppose also $u \in L^s(U) \cap L^t(U)$. Then $u \in L^r(U)$, and

$$(12) \quad \|u\|_{L^r(U)} \leq \|u\|_{L^s(U)}^\theta \|u\|_{L^t(U)}^{1-\theta}.$$

Proof. We compute

$$\begin{aligned} \int_U |u|^r dx &= \int_U |u|^{\theta r} |u|^{(1-\theta)r} dx \\ &\leq \left(\int_U |u|^{\theta r \frac{s}{\theta r}} dx \right)^{\frac{\theta r}{s}} \left(\int_U |u|^{(1-\theta)r \frac{t}{(1-\theta)r}} dx \right)^{\frac{(1-\theta)r}{t}}. \end{aligned}$$

We have invoked Hölder's inequality, which applies since $\frac{\theta r}{s} + \frac{(1-\theta)r}{t} = 1$. \square

i. Cauchy–Schwarz inequality.

$$(13) \quad |x \cdot y| \leq |x||y| \quad (x, y \in \mathbb{R}^n).$$

Proof. Let $\epsilon > 0$ and note

$$0 \leq |x \pm \epsilon y|^2 = |x|^2 \pm 2\epsilon x \cdot y + \epsilon^2 |y|^2.$$

Consequently

$$\pm x \cdot y \leq \frac{1}{2\epsilon} |x|^2 + \frac{\epsilon}{2} |y|^2.$$

Minimize the right-hand side by setting $\epsilon = \frac{|x|}{|y|}$, provided $y \neq 0$. \square

Remark. Likewise, if A is a symmetric, nonnegative $n \times n$ matrix,

$$(14) \quad \left| \sum_{i,j=1}^n a_{ij} x_i y_j \right| \leq \left(\sum_{i,j=1}^n a_{ij} x_i x_j \right)^{1/2} \left(\sum_{i,j=1}^n a_{ij} y_i y_j \right)^{1/2} \quad (x, y \in \mathbb{R}^n).$$

j. Gronwall's inequality (differential form).

(i) Let $\eta(\cdot)$ be a nonnegative, absolutely continuous function on $[0, T]$, which satisfies for a.e. t the differential inequality

$$(15) \quad \eta'(t) \leq \phi(t)\eta(t) + \psi(t),$$

where $\phi(t)$ and $\psi(t)$ are nonnegative, summable functions on $[0, T]$. Then

$$(16) \quad \eta(t) \leq e^{\int_0^t \phi(s) ds} \left[\eta(0) + \int_0^t \psi(s) ds \right]$$

for all $0 \leq t \leq T$.

(ii) In particular, if

$$\eta' \leq \phi\eta \quad \text{on } [0, T] \quad \text{and} \quad \eta(0) = 0,$$

then

$$\eta \equiv 0 \quad \text{on } [0, T].$$

Proof. From (15) we see

$$\frac{d}{ds} \left(\eta(s) e^{-\int_0^s \phi(r) dr} \right) = e^{-\int_0^s \phi(r) dr} (\eta'(s) - \phi(s)\eta(s)) \leq e^{-\int_0^s \phi(r) dr} \psi(s)$$

for a.e. $0 \leq s \leq T$. Consequently for each $0 \leq t \leq T$, we have

$$\eta(t) e^{-\int_0^t \phi(r) dr} \leq \eta(0) + \int_0^t e^{-\int_0^s \phi(r) dr} \psi(s) ds \leq \eta(0) + \int_0^t \psi(s) ds.$$

This implies inequality (16). □

k. Gronwall's inequality (integral form).

(i) Let $\xi(t)$ be a nonnegative, summable function on $[0, T]$ which satisfies for a.e. t the integral inequality

$$(17) \quad \xi(t) \leq C_1 \int_0^t \xi(s) ds + C_2$$

for constants $C_1, C_2 \geq 0$. Then

$$(18) \quad \xi(t) \leq C_2(1 + C_1 t e^{C_1 t})$$

for a.e. $0 \leq t \leq T$.

(ii) In particular, if

$$\xi(t) \leq C_1 \int_0^t \xi(s) ds$$

for a.e. $0 \leq t \leq T$, then

$$\xi(t) = 0 \quad \text{a.e.}$$


Proof. Let $\eta(t) := \int_0^t \xi(s) ds$; then $\eta' \leq C_1 \eta + C_2$ a.e. in $[0, T]$. According to the differential form of Gronwall's inequality above

$$\eta(t) \leq e^{C_1 t} (\eta(0) + C_2 t) = C_2 t e^{C_1 t}.$$

Then (17) implies

$$\xi(t) \leq C_1 \eta(t) + C_2 \leq C_2(1 + C_1 t e^{C_1 t}).$$

□



The boundary of U

APPENDIX C: CALCULUS

C.1. Boundaries.

Let $U \subset \mathbb{R}^n$ be open and bounded, $k \in \{1, 2, \dots\}$.

DEFINITION. We say the boundary ∂U is C^k if for each point $x^0 \in \partial U$ there exist $r > 0$ and a C^k function $\gamma : \mathbb{R}^{n-1} \rightarrow \mathbb{R}$ such that—upon relabeling and reorienting the coordinates axes if necessary—we have

$$U \cap B(x^0, r) = \{x \in B(x^0, r) \mid x_n > \gamma(x_1, \dots, x_{n-1})\}.$$

Likewise, ∂U is C^∞ if ∂U is C^k for $k = 1, 2, \dots$, and ∂U is analytic if the mapping γ is analytic.


DEFINITIONS. (i) If ∂U is C^1 , then along ∂U is defined the outward pointing unit normal vector field

$$\nu = (\nu^1, \dots, \nu^n).$$

The unit normal at any point $x^0 \in \partial U$ is $\nu(x^0) = \nu = (\nu_1, \dots, \nu_n)$.

(ii) Let $u \in C^1(\bar{U})$. We call

$$\frac{\partial u}{\partial \nu} := \nu \cdot Du$$



Straightening out the boundary

the (outward) normal derivative of u .

Straightening the boundary. We will frequently need to change coordinates near a point of ∂U so as to “flatten out” the boundary. To be more specific, fix $x^0 \in \partial U$, and choose r, γ , etc. as above. Define then

$$\begin{cases} y_i = x_i =: \Phi^i(x) & (i = 1, \dots, n-1) \\ y_n = x_n - \gamma(x_1, \dots, x_{n-1}) =: \Phi^n(x), \end{cases}$$

and write

$$y = \Phi(x).$$

Similarly, we set

$$\begin{cases} x_i = y_i =: \Psi^i(y) & (i = 1, \dots, n-1) \\ x_n = y_n + \gamma(y_1, \dots, y_{n-1}) =: \Psi^n(y), \end{cases}$$

and write

$$x = \Psi(y).$$

Then $\Phi = \Psi^{-1}$, and the mapping $x \mapsto \Phi(x) = y$ “straightens out ∂U ” near x^0 . Observe also that $\det D\Phi = \det D\Psi = 1$.

C.2. Gauss–Green Theorem.

In this section we assume U is a bounded, open subset of \mathbb{R}^n and ∂U is C^1 .

THEOREM 1 (Gauss–Green Theorem).

(i) Suppose $u \in C^1(\bar{U})$. Then

$$(1) \quad \int_U u_{x_i} dx = \int_{\partial U} u \nu^i dS \quad (i = 1, \dots, n).$$

(ii) We have

$$(2) \quad \int_U \operatorname{div} \mathbf{u} \, dx = \int_{\partial U} \mathbf{u} \cdot \nu \, dS$$

for each vector field $\mathbf{u} \in C^1(\bar{U}; \mathbb{R}^n)$.

Assertion (ii), also called the *Divergence Theorem*, follows from (i) applied to each component of $\mathbf{u} = (u^1, \dots, u^n)$.

THEOREM 2 (Integration by parts formula). *Let $u, v \in C^1(\bar{U})$. Then*

$$(3) \quad \int_U u_{x_i} v \, dx = - \int_U u v_{x_i} \, dx + \int_{\partial U} u v \nu^i \, dS \quad (i = 1, \dots, n).$$

Proof. Apply Theorem 1(i) to uv . □

THEOREM 3 (Green's formulas). *Let $u, v \in C^2(\bar{U})$. Then*

$$(i) \quad \int_U \Delta u \, dx = \int_{\partial U} \frac{\partial u}{\partial \nu} \, dS,$$

$$(ii) \quad \int_U Dv \cdot Du \, dx = - \int_U u \Delta v \, dx + \int_{\partial U} \frac{\partial v}{\partial \nu} u \, dS,$$

$$(iii) \quad \int_U u \Delta v - v \Delta u \, dx = \int_{\partial U} u \frac{\partial v}{\partial \nu} - v \frac{\partial u}{\partial \nu} \, dS.$$

Proof. Using (3), with u_{x_i} in place of u and $v \equiv 1$, we see

$$\int_U u_{x_i} x_i \, dx = \int_{\partial U} u_{x_i} \nu^i \, dS.$$

Sum $i = 1, \dots, n$ to establish (i).

To derive (ii), we employ (3) with v_{x_i} replacing v . Write (ii) with u and v interchanged and then subtract, to obtain (iii). □

C.3. Polar coordinates, coarea formula.

Next we convert n -dimensional integrals into integrals over spheres.

THEOREM 4 (Polar coordinates).

(i) *Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be continuous and summable. Then*

$$\int_{\mathbb{R}^n} f \, dx = \int_0^\infty \left(\int_{\partial B(x_0, r)} f \, dS \right) dr$$

for each point $x_0 \in \mathbb{R}^n$.

(ii) *In particular*

$$\frac{d}{dr} \left(\int_{B(x_0, r)} f \, dx \right) = \int_{\partial B(x_0, r)} f \, dS$$

for each $r > 0$.

Theorem 4 is a special case of the following theorem.

THEOREM 5 (Coarea formula). *Let $u : \mathbb{R}^n \rightarrow \mathbb{R}$ be Lipschitz continuous and assume that for a.e. $r \in \mathbb{R}$ the level set*

$$\{x \in \mathbb{R}^n \mid u(x) = r\}$$

is a smooth, $(n-1)$ -dimensional hypersurface in \mathbb{R}^n . Suppose also $f : \mathbb{R}^n \rightarrow \mathbb{R}$ is continuous and summable. Then

$$\int_{\mathbb{R}^n} f |Du| dx = \int_{-\infty}^{\infty} \left(\int_{\{u=r\}} f dS \right) dr.$$

Theorem 4 follows from Theorem 5 if we take $u(x) = |x - x_0|$. See [E-G, Chapter 3] for more on the coarea formula. The word “coarea” is pronounced, and sometimes spelled, “co-area”.

C.4. Moving regions.

Consider a family of smooth, bounded regions $U(\tau) \subset \mathbb{R}^n$ that depend smoothly upon the parameter $\tau \in \mathbb{R}$. Write \mathbf{v} for the velocity of the moving boundary $\partial U(\tau)$ and ν for the outward pointing unit normal.

THEOREM 6 (Differentiation formula for moving regions). *If $f = f(x, \tau)$ is a smooth function, then*

$$\frac{d}{d\tau} \int_{U(\tau)} f dx = \int_{\partial U(\tau)} f \mathbf{v} \cdot \nu dS + \int_{U(\tau)} f_{\tau} dx.$$

C.5. Convolution and smoothing.

We next introduce tools that will allow us to build smooth approximations to given functions.

NOTATION. If $U \subset \mathbb{R}^n$ is open and $\epsilon > 0$, we write

$$U_{\epsilon} := \{x \in U \mid \text{dist}(x, \partial U) > \epsilon\}.$$

DEFINITIONS. (i) Define $\eta \in C^{\infty}(\mathbb{R}^n)$ by

$$\eta(x) := \begin{cases} C \exp\left(\frac{1}{|x|^2-1}\right) & \text{if } |x| < 1 \\ 0 & \text{if } |x| \geq 1, \end{cases}$$

the constant $C > 0$ selected so that $\int_{\mathbb{R}^n} \eta dx = 1$.

(ii) For each $\epsilon > 0$, set

$$\eta_{\epsilon}(x) := \frac{1}{\epsilon^n} \eta\left(\frac{x}{\epsilon}\right).$$

We call η the *standard mollifier*. The functions η_{ϵ} are C^{∞} and satisfy

$$\int_{\mathbb{R}^n} \eta_{\epsilon} dx = 1, \quad \text{spt}(\eta_{\epsilon}) \subset B(0, \epsilon).$$

DEFINITION. If $f : U \rightarrow \mathbb{R}$ is locally integrable, define its mollification

$$f^\epsilon := \eta_\epsilon * f \quad \text{in } U_\epsilon.$$

That is,

$$f^\epsilon(x) = \int_U \eta_\epsilon(x-y)f(y) dy = \int_{B(0,\epsilon)} \eta_\epsilon(y)f(x-y) dy$$

for $x \in U_\epsilon$.

THEOREM 7 (Properties of mollifiers).

- (i) $f^\epsilon \in C^\infty(U_\epsilon)$.
- (ii) $f^\epsilon \rightarrow f$ a.e. as $\epsilon \rightarrow 0$.
- (iii) If $f \in C(U)$, then $f^\epsilon \rightarrow f$ uniformly on compact subsets of U .
- (iv) If $1 \leq p < \infty$ and $f \in L^p_{\text{loc}}(U)$, then $f^\epsilon \rightarrow f$ in $L^p_{\text{loc}}(U)$.

Proof. 1. Fix $x \in U_\epsilon$, $i \in \{1, \dots, n\}$, and h so small that $x + he_i \in U_\epsilon$. Then

$$\begin{aligned} \frac{f^\epsilon(x + he_i) - f^\epsilon(x)}{h} &= \frac{1}{\epsilon^n} \int_U \frac{1}{h} \left[\eta \left(\frac{x + he_i - y}{\epsilon} \right) - \eta \left(\frac{x - y}{\epsilon} \right) \right] f(y) dy \\ &= \frac{1}{\epsilon^n} \int_V \frac{1}{h} \left[\eta \left(\frac{x + he_i - y}{\epsilon} \right) - \eta \left(\frac{x - y}{\epsilon} \right) \right] f(y) dy \end{aligned}$$

for some open set $V \subset\subset U$. As

$$\frac{1}{h} \left[\eta \left(\frac{x + he_i - y}{\epsilon} \right) - \eta \left(\frac{x - y}{\epsilon} \right) \right] \rightarrow \frac{1}{\epsilon} \eta_{x_i} \left(\frac{x - y}{\epsilon} \right)$$

uniformly on V , the partial derivative $f^\epsilon_{x_i}(x)$ exists and equals

$$\int_U \eta_{\epsilon, x_i}(x-y)f(y) dy.$$

A similar argument shows that $D^\alpha f^\epsilon(x)$ exists, and

$$D^\alpha f^\epsilon(x) = \int_U D^\alpha \eta_\epsilon(x-y)f(y) dy \quad (x \in U_\epsilon),$$

for each multiindex α . This proves (i).

2. According to Lebesgue's Differentiation Theorem (§E.4),

$$(4) \quad \lim_{r \rightarrow 0} \int_{B(x,r)} |f(y) - f(x)| dy = 0$$

for a.e. $x \in U$. Fix such a point x . Then

$$\begin{aligned} |f^\epsilon(x) - f(x)| &= \left| \int_{B(x,\epsilon)} \eta_\epsilon(x-y)[f(y) - f(x)] dy \right| \\ &\leq \frac{1}{\epsilon^n} \int_{B(x,\epsilon)} \eta\left(\frac{x-y}{\epsilon}\right) |f(y) - f(x)| dy \\ &\leq C \int_{B(x,\epsilon)} |f(y) - f(x)| dy \rightarrow 0 \quad \text{as } \epsilon \rightarrow 0, \end{aligned}$$

by (4). Assertion (ii) follows.

3. Assume now $f \in C(U)$. Given $V \subset\subset U$, we choose $V \subset\subset W \subset\subset U$ and note that f is uniformly continuous on W . Thus the limit (4) holds uniformly for $x \in V$. Consequently the calculation above implies $f^\epsilon \rightarrow f$ uniformly on V .

4. Next, assume $1 \leq p < \infty$ and $f \in L^p_{\text{loc}}(U)$. Choose an open set $V \subset\subset U$ and, as above, an open set W so that $V \subset\subset W \subset\subset U$. We claim that for sufficiently small $\epsilon > 0$

$$(5) \quad \|f^\epsilon\|_{L^p(V)} \leq \|f\|_{L^p(W)}.$$

To see this, we note that if $1 \leq p < \infty$ and $x \in V$,

$$\begin{aligned} |f^\epsilon(x)| &= \left| \int_{B(x,\epsilon)} \eta_\epsilon(x-y) f(y) dy \right| \\ &\leq \int_{B(x,\epsilon)} \eta_\epsilon^{1-1/p}(x-y) \eta_\epsilon^{1/p}(x-y) |f(y)| dy \\ &\leq \left(\int_{B(x,\epsilon)} \eta_\epsilon(x-y) dy \right)^{1-1/p} \left(\int_{B(x,\epsilon)} \eta_\epsilon(x-y) |f(y)|^p dy \right)^{1/p}. \end{aligned}$$

Since $\int_{B(x,\epsilon)} \eta_\epsilon(x-y) dy = 1$, this inequality implies

$$\begin{aligned} \int_V |f^\epsilon(x)|^p dx &\leq \int_V \left(\int_{B(x,\epsilon)} \eta_\epsilon(x-y) |f(y)|^p dy \right) dx \\ &\leq \int_W |f(y)|^p \left(\int_{B(y,\epsilon)} \eta_\epsilon(x-y) dx \right) dy = \int_W |f(y)|^p dy, \end{aligned}$$

provided $\epsilon > 0$ is sufficiently small. This is (5).

5. Now fix $V \subset\subset W \subset\subset U$, $\delta > 0$, and choose $g \in C(W)$ so that

$$\|f - g\|_{L^p(W)} < \delta.$$

Then

$$\begin{aligned} \|f^\epsilon - f\|_{L^p(V)} &\leq \|f^\epsilon - g^\epsilon\|_{L^p(V)} + \|g^\epsilon - g\|_{L^p(V)} + \|g - f\|_{L^p(V)} \\ &\leq 2\|f - g\|_{L^p(W)} + \|g^\epsilon - g\|_{L^p(V)} \quad \text{by (5)} \\ &\leq 2\delta + \|g^\epsilon - g\|_{L^p(V)}. \end{aligned}$$

Since $g^\epsilon \rightarrow g$ uniformly on V , we have $\limsup_{\epsilon \rightarrow 0} \|f^\epsilon - f\|_{L^p(V)} \leq 2\delta$. \square

C.6. Inverse Function Theorem.


Let $U \subset \mathbb{R}^n$ be an open set and suppose $\mathbf{f} : U \rightarrow \mathbb{R}^n$ is C^1 , $\mathbf{f} = (f^1, \dots, f^n)$. Assume $x_0 \in U$, $z_0 = \mathbf{f}(x_0)$.

Remember from §A.4 that we write

$$D\mathbf{f} = \begin{pmatrix} f_{x_1}^1 & \cdots & f_{x_n}^1 \\ \vdots & \ddots & \vdots \\ f_{x_1}^n & \cdots & f_{x_n}^n \end{pmatrix}_{n \times n} = \text{gradient matrix of } \mathbf{f}.$$

DEFINITION.

$$J\mathbf{f} = \text{Jacobian of } \mathbf{f} = |\det D\mathbf{f}| = \left| \frac{\partial(f^1, \dots, f^n)}{\partial(x_1, \dots, x_n)} \right|.$$



THEOREM 8 (Inverse Function Theorem). Assume $\mathbf{f} \in C^1(U; \mathbb{R}^n)$ and

$$J\mathbf{f}(x_0) \neq 0.$$

Then there exist an open set $V \subset U$, with $x_0 \in V$, and an open set $W \subset \mathbb{R}^n$, with $z_0 \in W$, such that

(i) the mapping

$$\mathbf{f} : V \rightarrow W$$


is one-to-one and onto and

(ii) *the inverse function*

$$f^{-1} : W \rightarrow V$$

is C^1 .

(iii) If $f \in C^k$, then $f^{-1} \in C^k$ ($k = 2, \dots$).



C.7. Implicit Function Theorem.

Let n, m be positive integers. We write a typical point in \mathbb{R}^{n+m} as

$$(x, y) = (x_1, \dots, x_n, y_1, \dots, y_m)$$


for $x \in \mathbb{R}^n, y \in \mathbb{R}^m$.

Let $U \subset \mathbb{R}^{n+m}$ be an open set and suppose $f : U \rightarrow \mathbb{R}^m$ is $C^1, f = (f^1, \dots, f^m)$. Assume $(x_0, y_0) \in U, z_0 = f(x_0, y_0)$.

NOTATION.

$$Df = \begin{pmatrix} f_{x_1}^1 & \dots & f_{x_n}^1 & f_{y_1}^1 & \dots & f_{y_m}^1 \\ \vdots & & \vdots & & & \vdots \\ f_{x_1}^m & \dots & f_{x_n}^m & f_{y_1}^m & \dots & f_{y_m}^m \end{pmatrix}_{m \times (n+m)}$$

$= (D_x f, D_y f) = \text{gradient matrix of } f.$



DEFINITION.

$$J_y \mathbf{f} = |\det D_y \mathbf{f}| = \left| \frac{\partial(f^1, \dots, f^m)}{\partial(y_1, \dots, y_m)} \right|.$$

THEOREM 9 (Implicit Function Theorem). Assume $\mathbf{f} \in C^1(U; \mathbb{R}^m)$ and $J_y \mathbf{f}(x_0, y_0) \neq 0$.


Then there exists an open set $V \subset U$, with $(x_0, y_0) \in V$, an open set $W \subset \mathbb{R}^n$, with $x_0 \in W$, and a C^1 mapping $\mathbf{g} : W \rightarrow \mathbb{R}^m$ such that

- (i) $\mathbf{g}(x_0) = y_0$,
- (ii) $\mathbf{f}(x, \mathbf{g}(x)) = z_0 \quad (x \in W)$,

and

- (iii) if $(x, y) \in V$ and $\mathbf{f}(x, y) = z_0$, then $y = \mathbf{g}(x)$.
- (iv) If $\mathbf{f} \in C^k$, then $\mathbf{g} \in C^k$ ($k = 2, \dots$).

The function \mathbf{g} is *implicitly defined* near x_0 by the equation $\mathbf{f}(x, y) = z_0$.

**C.8. Uniform convergence.**

We record here the *Arzela–Ascoli compactness criterion* for uniform convergence:

Suppose that $\{f_k\}_{k=1}^{\infty}$ is a sequence of real-valued functions defined on \mathbb{R}^n , such that

$$|f_k(x)| \leq M \quad (k = 1, \dots, x \in \mathbb{R}^n)$$

for some constant M and that the functions $\{f_k\}_{k=1}^{\infty}$ are *uniformly equicontinuous*, meaning that for each $\varepsilon > 0$, there exists $\delta > 0$ such that $|x - y| < \delta$ implies $|f_k(x) - f_k(y)| < \varepsilon$, for $x, y \in \mathbb{R}^n$, $k = 1, \dots$.

Then there exists a subsequence $\{f_{k_j}\}_{j=1}^{\infty} \subseteq \{f_k\}_{k=1}^{\infty}$ and a continuous function f , such that

$$f_{k_j} \rightarrow f \quad \text{uniformly on compact subsets of } \mathbb{R}^n.$$

APPENDIX D: FUNCTIONAL ANALYSIS

D.1. Banach spaces.

Let X denote a real linear space.

DEFINITION. A mapping $\| \cdot \| : X \rightarrow [0, \infty)$ is called a norm if

- (i) $\|u + v\| \leq \|u\| + \|v\|$ for all $u, v \in X$,
- (ii) $\|\lambda u\| = |\lambda| \|u\|$ for all $u \in X, \lambda \in \mathbb{R}$,
- (iii) $\|u\| = 0$ if and only if $u = 0$.

Inequality (i) is the *triangle inequality*.

Hereafter we assume X is a normed linear space.

DEFINITION. We say a sequence $\{u_k\}_{k=1}^{\infty} \subset X$ converges to $u \in X$, written

$$u_k \rightarrow u,$$

if

$$\lim_{k \rightarrow \infty} \|u_k - u\| = 0.$$

DEFINITIONS. (i) A sequence $\{u_k\}_{k=1}^{\infty} \subset X$ is called a Cauchy sequence provided for each $\epsilon > 0$ there exists $N > 0$ such that

$$\|u_k - u_l\| < \epsilon \quad \text{for all } k, l \geq N.$$

(ii) X is complete if each Cauchy sequence in X converges; that is, whenever $\{u_k\}_{k=1}^{\infty}$ is a Cauchy sequence, there exists $u \in X$ such that $\{u_k\}_{k=1}^{\infty}$ converges to u .

(iii) A Banach space X is a complete, normed linear space.

DEFINITION. We say X is separable if X contains a countable dense subset.

Examples. (i) L^p spaces. Assume U is an open subset of \mathbb{R}^n and $1 \leq p \leq \infty$. If $f : U \rightarrow \mathbb{R}$ is measurable, we define

$$\|f\|_{L^p(U)} := \begin{cases} \left(\int_U |f|^p dx \right)^{1/p} & \text{if } 1 \leq p < \infty \\ \text{ess sup}_U |f| & \text{if } p = \infty. \end{cases}$$

We define $L^p(U)$ to be the linear space of all measurable functions $f : U \rightarrow \mathbb{R}$ for which $\|f\|_{L^p(U)} < \infty$. Then $L^p(U)$ is a Banach space, provided we identify two functions which agree a.e.

(ii) Hölder spaces. See §5.1.

(iii) Sobolev spaces. See §5.2. □

D.2. Hilbert spaces.

Let H be a real linear space.

DEFINITION. A mapping $(\cdot, \cdot) : H \times H \rightarrow \mathbb{R}$ is called an inner product if

- (i) $(u, v) = (v, u)$ for all $u, v \in H$,
- (ii) the mapping $u \mapsto (u, v)$ is linear for each $v \in H$,
- (iii) $(u, u) \geq 0$ for all $u \in H$,
- (iv) $(u, u) = 0$ if and only if $u = 0$.

DEFINITION. If (\cdot, \cdot) is an inner product, the associated norm is

$$(1) \quad \|u\| := (u, u)^{1/2} \quad (u \in H).$$

The Cauchy–Schwarz inequality states

$$(2) \quad |(u, v)| \leq \|u\| \|v\| \quad (u, v \in H).$$

This inequality is proved as in §B.2. Using (2), we easily verify (1) defines a norm on H .

DEFINITION. A Hilbert space H is a Banach space endowed with an inner product which generates the norm.

Examples. a. The space $L^2(U)$ is a Hilbert space, with

$$(f, g) = \int_U fg \, dx.$$

b. The Sobolev space $H^1(U)$ is a Hilbert space, with

$$(f, g) = \int_U fg + Df \cdot Dg \, dx.$$

DEFINITIONS. (i) Two elements $u, v \in H$ are orthogonal if $(u, v) = 0$.

(ii) A countable basis $\{w_k\}_{k=1}^\infty \subset H$ is called orthonormal if

$$\begin{cases} (w_k, w_l) = 0 & (k, l = 1, \dots; k \neq l) \\ \|w_k\| = 1 & (k = 1, \dots). \end{cases}$$

If $u \in H$ and $\{w_k\}_{k=1}^\infty \subset H$ is an orthonormal basis, we can write

$$u = \sum_{k=1}^{\infty} (u, w_k) w_k,$$

the series converging in H . In addition

$$\|u\|^2 = \sum_{k=1}^{\infty} (u, w_k)^2.$$

DEFINITION. If S is a subspace of H , $S^\perp = \{u \in H \mid (u, v) = 0 \text{ for all } v \in S\}$ is the subspace orthogonal to S .

D.3. Bounded linear operators.

a. Linear operators on Banach spaces.

Let X and Y be real Banach spaces.

DEFINITIONS. (i) A mapping $A : X \rightarrow Y$ is a linear operator provided

$$A[\lambda u + \mu v] = \lambda Au + \mu Av$$

for all $u, v \in X$, $\lambda, \mu \in \mathbb{R}$.

(ii) The range of A is $R(A) := \{v \in Y \mid v = Au \text{ for some } u \in X\}$ and the null space of A is $N(A) := \{u \in X \mid Au = 0\}$.

DEFINITION. A linear operator $A : X \rightarrow Y$ is bounded if

$$\|A\| := \sup\{\|Au\|_Y \mid \|u\|_X \leq 1\} < \infty.$$

It is easy to check that a bounded linear operator $A : X \rightarrow Y$ is continuous.

DEFINITION. A linear operator $A : X \rightarrow Y$ is called closed if whenever $u_k \rightarrow u$ in X and $Au_k \rightarrow v$ in Y , then

$$Au = v.$$

THEOREM 1 (Closed Graph Theorem). Let $A : X \rightarrow Y$ be a closed, linear operator. Then A is bounded.

DEFINITIONS. Let $A : X \rightarrow X$ be a bounded linear operator.

(i) The resolvent set of A is

$$\rho(A) = \{\eta \in \mathbb{R} \mid (A - \eta I) \text{ is one-to-one and onto}\}.$$

(ii) The spectrum of A is

$$\sigma(A) = \mathbb{R} - \rho(A).$$

If $\eta \in \rho(A)$, the Closed Graph Theorem then implies that the inverse $(A - \eta I)^{-1} : X \rightarrow X$ is a bounded linear operator.

DEFINITIONS. (i) We say $\eta \in \sigma(A)$ is an eigenvalue of A provided

$$N(A - \eta I) \neq \{0\}.$$

We write $\sigma_p(A)$ to denote the collection of eigenvalues of A ; $\sigma_p(A)$ is the point spectrum.

(ii) If η is an eigenvalue and $w \neq 0$ satisfies

$$Aw = \eta w,$$

we say w is an associated eigenvector.

DEFINITIONS. (i) A bounded linear operator $u^* : X \rightarrow \mathbb{R}$ is called a bounded linear functional on X .

(ii) We write X^* to denote the collection of all bounded linear functionals on X ; X^* is the dual space of X .

DEFINITIONS. (i) If $u \in X$, $u^* \in X^*$ we write

$$\langle u^*, u \rangle$$

to denote the real number $u^*(u)$. The symbol \langle , \rangle denotes the pairing of X^* and X .

(ii) We define

$$\|u^*\| := \sup\{\langle u^*, u \rangle \mid \|u\| \leq 1\}.$$

(iii) A Banach space is reflexive if $(X^*)^* = X$. More precisely, this means that for each $u^{**} \in (X^*)^*$, there exists $u \in X$ such that

$$\langle u^{**}, u^* \rangle = \langle u^*, u \rangle \quad \text{for all } u^* \in X^*.$$

b. Linear operators on Hilbert spaces.

Now let H be a real Hilbert space, with inner product $(,)$.

THEOREM 2 (Riesz Representation Theorem). H^* can be canonically identified with H ; more precisely, for each $u^* \in H^*$ there exists a unique element $u \in H$ such that

$$\langle u^*, v \rangle = (u, v) \quad \text{for all } v \in H.$$

The mapping $u^* \mapsto u$ is a linear isomorphism of H^* onto H .

DEFINITIONS. (i) If $A : H \rightarrow H$ is a bounded, linear operator, its adjoint $A^* : H \rightarrow H$ satisfies

$$(Au, v) = (u, A^*v)$$

for all $u, v \in H$.

(ii) A is symmetric if $A^* = A$.

D.4. Weak convergence.

Let X denote a real Banach space.

DEFINITION. We say a sequence $\{u_k\}_{k=1}^\infty \subset X$ converges weakly to $u \in X$, written

$$u_k \rightharpoonup u,$$

if

$$\langle u^*, u_k \rangle \rightarrow \langle u^*, u \rangle$$

for each bounded linear functional $u^* \in X^*$.

It is easy to check that if $u_k \rightarrow u$, then $u_k \rightharpoonup u$. It is also true that *any weakly convergent sequence is bounded*. In addition, if $u_k \rightharpoonup u$, then

$$\|u\| \leq \liminf_{k \rightarrow \infty} \|u_k\|.$$

THEOREM 3 (Weak compactness). *Let X be a reflexive Banach space and suppose the sequence $\{u_k\}_{k=1}^\infty \subset X$ is bounded. Then there exists a subsequence $\{u_{k_j}\}_{j=1}^\infty \subset \{u_k\}_{k=1}^\infty$ and $u \in X$ such that*

$$u_{k_j} \rightharpoonup u.$$

In other words, bounded sequences in a reflexive Banach space are weakly precompact. In particular, *a bounded sequence in a Hilbert space contains a weakly convergent subsequence*.

Mazur's Theorem asserts that a convex, closed subset of X is weakly closed.

IMPORTANT EXAMPLE. We will most often employ weak convergence ideas in the following context. Take $U \subset \mathbb{R}^n$ to be open, and assume $1 \leq p < \infty$. Then

$$\text{the dual space of } X = L^p(U) \text{ is } X^* = L^q(U),$$

where $\frac{1}{p} + \frac{1}{q} = 1$, $1 < q \leq \infty$. More precisely, each bounded linear functional on $L^p(U)$ can be represented as $f \mapsto \int_U gf \, dx$ for some $g \in L^q(U)$. Therefore

$$f_k \rightharpoonup f \quad \text{weakly in } L^p(U)$$

means

$$(3) \quad \int_U gf_k \, dx \rightarrow \int_U gf \, dx \quad \text{as } k \rightarrow \infty, \text{ for all } g \in L^q(U).$$

Now the identification of $L^q(U)$ as the dual of $L^p(U)$ shows that

$$L^p(U) \text{ is reflexive if } 1 < p < \infty.$$

In particular Theorem 3 then assures us that from a bounded sequence in $L^p(U)$ ($1 < p < \infty$) we can extract a *weakly* convergent subsequence, that is, a sequence satisfying (3). This is an important compactness assertion, but note very carefully: *the convergence (3) does not imply that $f_k \rightarrow f$ pointwise or almost everywhere.* It may very well be, for example, that the functions $\{f_k\}_{k=1}^\infty$ oscillate more and more rapidly as $k \rightarrow \infty$. (See Problem 1 in Chapter 8 and also Problem 2 in Chapter 9.) \square

D.5. Compact operators, Fredholm theory.

Let X and Y be real Banach spaces.

DEFINITION. A bounded linear operator

$$K : X \rightarrow Y$$

is called compact provided for each bounded sequence $\{u_k\}_{k=1}^\infty \subset X$, the sequence $\{Ku_k\}_{k=1}^\infty$ is precompact in Y ; that is, there exists a subsequence $\{u_{k_j}\}_{j=1}^\infty$ such that $\{Ku_{k_j}\}_{j=1}^\infty$ converges in Y .

Now let H denote a real Hilbert space, with inner product (\cdot, \cdot) . It is easy to see that if a linear operator $K : H \rightarrow H$ is compact and $u_k \rightharpoonup u$, then $Ku_k \rightarrow Ku$.

THEOREM 4 (Compactness of adjoints). *If $K : H \rightarrow H$ is compact, so is $K^* : H \rightarrow H$.*

Proof. Let $\{u_k\}_{k=1}^\infty$ be a bounded sequence in H and extract a weakly convergent subsequence $u_{k_j} \rightharpoonup u$ in H . We will prove $K^*u_{k_j} \rightarrow K^*u$. Indeed,

$$\begin{aligned} \|K^*u_{k_j} - K^*u\|^2 &= (K^*u_{k_j} - K^*u, K^*[u_{k_j} - u]) \\ &= (KK^*u_{k_j} - KK^*u, u_{k_j} - u). \end{aligned}$$

Now since K^* is linear, $K^*u_{k_j} \rightharpoonup K^*u$, and so $KK^*u_{k_j} \rightarrow KK^*u$. Thus $K^*u_{k_j} \rightarrow K^*u$. \square

THEOREM 5 (Fredholm alternative). *Let $K : H \rightarrow H$ be a compact linear operator. Then*

- (i) $N(I - K)$ is finite dimensional,
- (ii) $R(I - K)$ is closed,
- (iii) $R(I - K) = N(I - K^*)^\perp$,
- (iv) $N(I - K) = \{0\}$ if and only if $R(I - K) = H$,

and

- (v) $\dim N(I - K) = \dim N(I - K^*)$.

Proof. 1. If $\dim N(I - K) = +\infty$, we can find an infinite orthonormal set $\{u_k\}_{k=1}^\infty \subset N(I - K)$. Then

$$Ku_k = u_k \quad (k = 1, \dots).$$

Now $\|u_k - u_l\|^2 = \|u_k\|^2 - 2(u_k, u_l) + \|u_l\|^2 = 2$ if $k \neq l$, and so $\|Ku_k - Ku_l\| = \sqrt{2}$ for $k \neq l$. This however contradicts the compactness of K , as $\{Ku_k\}_{k=1}^\infty$ would then contain no convergent subsequence. Assertion (i) is proved.

2. We next claim there exists a constant $\gamma > 0$ such that

$$(4) \quad \|u - Ku\| \geq \gamma \|u\| \quad \text{for all } u \in N(I - K)^\perp.$$

Indeed, if not, there would exist for $k = 1, \dots$ elements $u_k \in N(I - K)^\perp$ with $\|u_k\| = 1$ and $\|u_k - Ku_k\| < \frac{1}{k}$. Consequently

$$(5) \quad u_k - Ku_k \rightarrow 0.$$

But since $\{u_k\}_{k=1}^\infty$ is bounded, there exists a weakly convergent subsequence $u_{k_j} \rightharpoonup u$. By compactness $Ku_{k_j} \rightarrow Ku$, and then (5) implies $u_{k_j} \rightarrow Ku = u$. We therefore have $u \in N(I - K)$ and so

$$(u_{k_j}, u) = 0 \quad (j = 1, \dots).$$

Let $k_j \rightarrow \infty$ to derive a contradiction.

3. Next let $\{v_k\}_{k=1}^\infty \subset R(I - K)$, $v_k \rightarrow v$. We can find $u_k \in N(I - K)^\perp$ solving $u_k - Ku_k = v_k$. Using (4), we deduce

$$\|v_k - v_l\| \geq \gamma \|u_k - u_l\|.$$

Thus $u_k \rightarrow u$ and $u - Ku = v$. This proves (ii).

4. Assertion (iii) is now a consequence of (ii) and the general fact that

$$\overline{R(A)} = N(A^*)^\perp \quad \text{for each bounded linear operator } A : H \rightarrow H.$$

5. To verify (iv), let us suppose, to start with, that $N(I - K) = \{0\}$, but $H_1 = (I - K)(H) \subsetneq H$. According to (ii), H_1 is a closed subspace of H . Furthermore $H_2 \equiv (I - K)(H_1) \subsetneq H_1$, since $I - K$ is one-to-one. Similarly if we write $H_k \equiv (I - K)^k(H)$ ($k = 1, \dots$), we see that H_k is a closed subspace of H , $H_{k+1} \subsetneq H_k$ ($k = 1, \dots$).

Choose $u_k \in H_k$ with $\|u_k\| = 1$, $u_k \in H_{k+1}^\perp$. Then $Ku_k - Ku_l = -(u_k - Ku_k) + (u_l - Ku_l) + (u_k - u_l)$. Now if $k > l$, $H_{k+1} \subsetneq H_k \subseteq H_{l+1} \subsetneq H_l$. Thus $u_k - Ku_k, u_l - Ku_l, u_k \in H_{l+1}$. Since $u_l \in H_{l+1}^\perp$, $\|u_l\| = 1$, we deduce $\|Ku_k - Ku_l\| \geq 1$ ($k, l = 1, \dots$). But this is impossible since K is compact.

6. Now conversely assume $R(I - K) = H$. Then owing to (iii), we see that $N(I - K^*) = \{0\}$. Since K^* is compact, we may utilize step 5 to conclude $R(I - K^*) = H$. But then $N(I - K) = R(I - K^*)^\perp = \{0\}$. This conclusion and step 5 complete the proof of assertion (iv).

7. Next we assert

$$\dim N(I - K) \geq \dim R(I - K)^\perp.$$

To prove this, suppose instead $\dim N(I - K) < \dim R(I - K)^\perp$. Then there exists a bounded linear mapping $A : N(I - K) \rightarrow R(I - K)^\perp$ which is one-to-one, but *not* onto. Extend A to a linear mapping $A : H \rightarrow R(I - K)^\perp$ by setting $Au = 0$ for $u \in N(I - K)^\perp$. Now A has a finite-dimensional range and so A and thus $K + A$ are compact. Furthermore $N(I - (K + A)) = \{0\}$. Indeed, if $Ku + Au = u$, then $u - Ku = Au \in R(I - K)^\perp$. Hence $u - Ku = Au = 0$. Thus $u \in N(I - K)$ and so in fact $u = 0$, since A is one-to-one on $N(I - K)$. Now apply assertion (iv) to $\tilde{K} = K + A$. We conclude $R(I - (K + A)) = H$. But this is impossible: if $v \in R(I - K)^\perp$, but $v \notin R(A)$, the equation

$$u - (Ku + Au) = v$$

has no solution.

8. Since $R(I - K^*)^\perp = N(I - K)$, we deduce from step 7 that

$$\begin{aligned} \dim N(I - K^*) &\geq \dim R(I - K^*)^\perp \\ &= \dim N(I - K). \end{aligned}$$

The opposite inequality comes from interchanging the roles of K and K^* . This establishes (v). \square

Remark. Theorem 5 asserts in particular either

$$(\alpha) \quad \begin{cases} \text{for each } f \in H, \text{ the equation } u - Ku = f \\ \text{has a unique solution} \end{cases}$$

or else

$$(\beta) \quad \begin{cases} \text{the homogeneous equation } u - Ku = 0 \\ \text{has solutions } u \neq 0. \end{cases}$$

This dichotomy is the *Fredholm alternative*. In addition, should (β) obtain, the space of solutions of the homogeneous problem is finite dimensional, and the nonhomogeneous equation

$$(\gamma) \quad u - Ku = f$$

has a solution if and only if $f \in N(I - K^*)^\perp$.

Now we investigate the spectrum of a compact linear operator.

THEOREM 6 (Spectrum of a compact operator). *Assume $\dim H = \infty$ and $K : H \rightarrow H$ is compact. Then*

- (i) $0 \in \sigma(K)$,
- (ii) $\sigma(K) - \{0\} = \sigma_p(K) - \{0\}$,

and

- (iii) $\begin{cases} \sigma(K) - \{0\} \text{ is finite, or else} \\ \sigma(K) - \{0\} \text{ is a sequence tending to } 0. \end{cases}$

Proof. 1. Assume $0 \notin \sigma(K)$. Then $K : H \rightarrow H$ is bijective and so $I = K \circ K^{-1}$, being the composition of a compact and a bounded linear operator, is compact. This is impossible, since $\dim H = \infty$.

2. Assume $\eta \in \sigma(K)$, $\eta \neq 0$. Then if $N(K - \eta I) = \{0\}$, the Fredholm alternative would imply $R(K - \eta I) = H$. But then $\eta \in \rho(K)$, a contradiction.

3. Suppose now $\{\eta_k\}_{k=1}^\infty$ is a sequence of *distinct* elements of $\sigma(K) - \{0\}$ and $\eta_k \rightarrow \eta$. We will show $\eta = 0$.

Indeed, since $\eta_k \in \sigma_p(K)$, there exists $w_k \neq 0$ such that $Kw_k = \eta_k w_k$. Let H_k denote the subspace of H spanned by $\{w_1, \dots, w_k\}$. Then $H_k \subsetneq H_{k+1}$ for each $k = 1, 2, \dots$, since the $\{w_k\}_{k=1}^\infty$ are linearly independent.

Observe also $(K - \eta_k I)H_k \subseteq H_{k-1}$ ($k = 2, \dots$). Choose now for $k = 1, \dots$ an element $u_k \in H_k$, with $u_k \in H_{k-1}^\perp$ and $\|u_k\| = 1$. Now if $k > l$, $H_{l-1} \subsetneq H_l \subseteq H_{k-1} \subsetneq H_k$. Thus

$$\left\| \frac{Ku_k}{\eta_k} - \frac{Ku_l}{\eta_l} \right\| = \left\| \frac{(Ku_k - \eta_k u_k)}{\eta_k} - \frac{(Ku_l - \eta_l u_l)}{\eta_l} + u_k - u_l \right\| \geq 1,$$

since $Ku_k - \eta_k u_k, Ku_l - \eta_l u_l, u_l \in H_{k-1}$. If $\eta_k \rightarrow \eta \neq 0$, we obtain a contradiction to the compactness of K . □

D.6. Symmetric operators.

Now let $S : H \rightarrow H$ be linear, bounded, symmetric, and write

$$m := \inf_{\substack{u \in H \\ \|u\|=1}} (Su, u), \quad M := \sup_{\substack{u \in H \\ \|u\|=1}} (Su, u).$$

LEMMA (Bounds on spectrum). *We have*

- (i) $\sigma(S) \subset [m, M]$, and
- (ii) $m, M \in \sigma(S)$.

Proof. 1. Let $\eta > M$. Then

$$(\eta u - Su, u) \geq (\eta - M)\|u\|^2 \quad (u \in H).$$

Hence the Lax–Milgram Theorem (§6.2.1) asserts $\eta I - S$ is one-to-one and onto, and thus $\eta \in \rho(S)$. Similarly $\eta \in \rho(S)$ if $\eta < m$. This proves (i).

2. We will prove $M \in \sigma(S)$. Since the pairing $[u, v] := (Mu - Su, v)$ is symmetric, with $[u, u] \geq 0$ for all $u \in H$, the Cauchy–Schwarz inequality implies

$$|(Mu - Su, v)| \leq (Mu - Su, u)^{1/2} (Mv - Sv, v)^{1/2}$$

for all $u, v \in H$. In particular

$$(6) \quad \|Mu - Su\| \leq C(Mu - Su, u)^{1/2} \quad (u \in H)$$

for some constant C .

Now let $\{u_k\}_{k=1}^{\infty} \subset H$ satisfy $\|u_k\| = 1$ ($k = 1, \dots$) and $(Su_k, u_k) \rightarrow M$. Then (6) implies $\|Mu_k - Su_k\| \rightarrow 0$. Now if $M \in \rho(S)$, then

$$u_k = (MI - S)^{-1}(Mu_k - Su_k) \rightarrow 0,$$

a contradiction. Thus $M \in \sigma(S)$, and likewise $m \in \sigma(S)$. □

THEOREM 7 (Eigenvectors of a compact, symmetric operator). *Let H be a separable Hilbert space, and suppose $S : H \rightarrow H$ is a compact and symmetric operator. Then there exists a countable orthonormal basis of H consisting of eigenvectors of S .*

Proof. 1. Let $\{\eta_k\}$ comprise the sequence of distinct eigenvalues of S , excepting 0. Set $\eta_0 = 0$. Write $H_0 = N(S)$, $H_k = N(S - \eta_k I)$ ($k = 1, \dots$). Then $0 \leq \dim H_0 \leq \infty$, and $0 < \dim H_k < \infty$, according to the Fredholm alternative.

2. Let $u \in H_k$, $v \in H_l$ for $k \neq l$. Then $Su = \eta_k u$, $Sv = \eta_l v$ and so

$$\eta_k(u, v) = (Su, v) = (u, Sv) = \eta_l(u, v).$$

As $\eta_k \neq \eta_l$, we deduce $(u, v) = 0$. Consequently we see the subspaces H_k and H_l are orthogonal.

3. Now let \tilde{H} be the smallest subspace of H containing H_0, H_1, \dots . Thus $\tilde{H} = \{\sum_{k=0}^m a_k u_k \mid m \in \{0, \dots\}, u_k \in H_k, a_k \in \mathbb{R}\}$. We next demonstrate \tilde{H} is dense in H . Clearly $S(\tilde{H}) \subseteq \tilde{H}$. Furthermore $S(\tilde{H}^\perp) \subseteq \tilde{H}^\perp$: indeed if $u \in \tilde{H}^\perp$ and $v \in \tilde{H}$, then $(Su, v) = (u, Sv) = 0$.

Now the operator $\tilde{S} \equiv S|_{\tilde{H}^\perp}$ is compact and symmetric. In addition $\sigma(\tilde{S}) = \{0\}$, since any nonzero eigenvalue of \tilde{S} would be an eigenvalue of S as well. According to the lemma then, $(\tilde{S}u, u) = 0$ for all $u \in \tilde{H}^\perp$. But if $u, v \in \tilde{H}^\perp$,

$$2(\tilde{S}u, v) = (\tilde{S}(u + v), u + v) - (\tilde{S}u, u) - (\tilde{S}v, v) = 0.$$

Hence $\tilde{S} = 0$. Consequently $\tilde{H}^\perp \subset N(S) \subset \tilde{H}$, and so $\tilde{H}^\perp = \{0\}$. Thus \tilde{H} is dense in H .

4. Choose an orthonormal basis for each subspace H_k ($k = 0, \dots$), noting that since H is separable, H_0 has a countable orthonormal basis. We obtain thereby an orthonormal basis of eigenvectors. \square

Most of these proofs are from Brezis [BR1]. See also Gilbarg–Trudinger [G-T, Chapter 5], Lax [Lx2], Reed–Simon [R-S1] and Yosida [Y].

APPENDIX E: MEASURE THEORY

This appendix provides a quick outline of some fundamentals of measure theory.

E.1. Lebesgue measure.

Lebesgue measure provides a way of describing the “size” or “volume” of certain subsets of \mathbb{R}^n .

DEFINITION. A collection \mathcal{M} of subsets of \mathbb{R}^n is called a σ -algebra if

- (i) $\emptyset, \mathbb{R}^n \in \mathcal{M}$,
- (ii) $A \in \mathcal{M}$ implies $\mathbb{R}^n - A \in \mathcal{M}$,

and

- (iii) if $\{A_k\}_{k=1}^\infty \subset \mathcal{M}$, then $\bigcup_{k=1}^\infty A_k, \bigcap_{k=1}^\infty A_k \in \mathcal{M}$.

THEOREM 1 (Existence of Lebesgue measure and Lebesgue measurable sets). *There exist a σ -algebra \mathcal{M} of subsets of \mathbb{R}^n and a mapping*

$$|\cdot| : \mathcal{M} \rightarrow [0, +\infty]$$

with the following properties:

(i) *Every open subset of \mathbb{R}^n and thus every closed subset of \mathbb{R}^n belong to \mathcal{M} .*

(ii) *If B is a ball in \mathbb{R}^n , then $|B|$ equals the n -dimensional volume of B .*

(iii) *If $\{A_k\}_{k=1}^{\infty} \subset \mathcal{M}$ and the sets $\{A_k\}_{k=1}^{\infty}$ are pairwise disjoint, then*

$$(1) \quad \left| \bigcup_{k=1}^{\infty} A_k \right| = \sum_{k=1}^{\infty} |A_k| \quad (\text{“countable additivity”}).$$

(iv) *If $A \subseteq B$, where $B \in \mathcal{M}$ and $|B| = 0$, then $A \in \mathcal{M}$ and $|A| = 0$.*

The sets in \mathcal{M} are called *Lebesgue measurable sets* and $|\cdot|$ is *n -dimensional Lebesgue measure*.

Remarks. (i) From (ii) and (iii), we see that $|A|$ equals the volume of any set A with piecewise smooth boundary.

(ii) We deduce from (1) that

$$(2) \quad |\emptyset| = 0$$

and

$$(3) \quad \left| \bigcup_{k=1}^{\infty} A_k \right| \leq \sum_{k=1}^{\infty} |A_k| \quad (\text{“countable subadditivity”})$$

for any countable collection of measurable sets $\{A_k\}_{k=1}^{\infty}$.

NOTATION. If some property holds everywhere on \mathbb{R}^n , except for a measurable set with Lebesgue measure zero, we say the property holds *almost everywhere*, abbreviated “a.e.”.

E.2. Measurable functions and integration.

DEFINITION. Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$. We say f is a measurable function if

$$f^{-1}(U) \in \mathcal{M}$$

for each open subset $U \subset \mathbb{R}$.

Note in particular that if f is continuous, then f is measurable. The sum and product of two measurable functions are measurable. In addition if $\{f_k\}_{k=0}^{\infty}$ are measurable functions, then so are $\limsup f_k$ and $\liminf f_k$.

THEOREM 2 (Egoroff's Theorem). Let $\{f_k\}_{k=1}^{\infty}, f$ be measurable functions, and assume

$$f_k \rightarrow f \quad \text{a.e. on } A,$$

where $A \subset \mathbb{R}^n$ is measurable, $|A| < \infty$. Then for each $\varepsilon > 0$ there exists a measurable subset $E \subset A$ such that

(i) $|A - E| \leq \varepsilon$

and

(ii) $f_k \rightarrow f$ uniformly on E .

Now if f is a nonnegative, measurable function, it is possible, by an approximation of f with simple functions, to define the Lebesgue integral

$$\int_{\mathbb{R}^n} f \, dx.$$

Cf. §E.5 below. This agrees with the usual integral if f is continuous or Riemann integrable. If f is measurable, but not necessarily nonnegative, we define

$$\int_{\mathbb{R}^n} f \, dx = \int_{\mathbb{R}^n} f^+ \, dx - \int_{\mathbb{R}^n} f^- \, dx,$$

provided at least one of the terms on the right-hand side is finite. In this case we say f is *integrable*.

DEFINITION. A measurable function f is *summable* if

$$\int_{\mathbb{R}^n} |f| \, dx < \infty.$$

Note carefully our terminology: a measurable function is *integrable* if it has an integral (which may equal $+\infty$ or $-\infty$) and is *summable* if this integral is finite.

DEFINITION. If the real-valued function f is measurable, we define the *essential supremum* of f to be

$$\text{ess sup } f := \inf\{\mu \in \mathbb{R} \mid |\{f > \mu\}| = 0\}.$$

E.3. Convergence theorems for integrals.

The Lebesgue theory of integration is especially useful since it provides the following powerful convergence theorems.

THEOREM 3 (Fatou's Lemma). *Assume the functions $\{f_k\}_{k=1}^{\infty}$ are non-negative and measurable. Then*

$$\int_{\mathbb{R}^n} \liminf_{k \rightarrow \infty} f_k \, dx \leq \liminf_{k \rightarrow \infty} \int_{\mathbb{R}^n} f_k \, dx.$$

THEOREM 4 (Monotone Convergence Theorem). *Assume the functions $\{f_k\}_{k=1}^{\infty}$ are measurable, with*

$$0 \leq f_1 \leq f_2 \leq \cdots \leq f_k \leq f_{k+1} \leq \cdots.$$

Then

$$\int_{\mathbb{R}^n} \lim_{k \rightarrow \infty} f_k \, dx = \lim_{k \rightarrow \infty} \int_{\mathbb{R}^n} f_k \, dx.$$

THEOREM 5 (Dominated Convergence Theorem). *Assume the functions $\{f_k\}_{k=1}^{\infty}$ are integrable and*

$$f_k \rightarrow f \text{ a.e.}$$

Suppose also

$$|f_k| \leq g \text{ a.e.,}$$

for some summable function g . Then

$$\int_{\mathbb{R}^n} f_k \, dx \rightarrow \int_{\mathbb{R}^n} f \, dx.$$

E.4. Differentiation.

An important fact is that a summable function is “approximately continuous” at almost every point.

THEOREM 6 (Lebesgue's Differentiation Theorem). *Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be locally summable.*

(i) *Then for a.e. point $x_0 \in \mathbb{R}^n$,*

$$\int_{B(x_0, r)} f \, dx \rightarrow f(x_0) \quad \text{as } r \rightarrow 0.$$

(ii) *In fact, for a.e. point $x_0 \in \mathbb{R}^n$,*

$$(4) \quad \int_{B(x_0, r)} |f(x) - f(x_0)| \, dx \rightarrow 0 \quad \text{as } r \rightarrow 0.$$

A point x_0 at which (4) holds is called a *Lebesgue point* of f .

Remark. More generally, if $f \in L^p_{loc}(\mathbb{R}^n)$ for some $1 \leq p < \infty$, then for a.e. point $x_0 \in \mathbb{R}^n$ we have

$$\int_{B(x_0, r)} |f(x) - f(x_0)|^p dx \rightarrow 0 \quad \text{as } r \rightarrow 0.$$

E.5. Banach space-valued functions.

We extend the notions of measurability, integrability, etc. to mappings

$$\mathbf{f} : [0, T] \rightarrow X$$

where $T > 0$ and X is a real Banach space, with norm $\| \cdot \|$.

DEFINITIONS. (i) A function $\mathbf{s} : [0, T] \rightarrow X$ is called simple if it has the form

$$(5) \quad \mathbf{s}(t) = \sum_{i=1}^m \chi_{E_i}(t) u_i \quad (0 \leq t \leq T),$$

where each E_i is a Lebesgue measurable subset of $[0, T]$ and $u_i \in X$ ($i = 1, \dots, m$).

(ii) A function $\mathbf{f} : [0, T] \rightarrow X$ is strongly measurable if there exist simple functions $\mathbf{s}_k : [0, T] \rightarrow X$ such that

$$\mathbf{s}_k(t) \rightarrow \mathbf{f}(t) \quad \text{for a.e. } 0 \leq t \leq T.$$

(iii) A function $\mathbf{f} : [0, T] \rightarrow X$ is weakly measurable if for each $u^* \in X^*$, the mapping $t \mapsto \langle u^*, \mathbf{f}(t) \rangle$ is Lebesgue measurable.

DEFINITION. We say $\mathbf{f} : [0, T] \rightarrow X$ is almost separably valued if there exists a subset $N \subset [0, T]$, with $|N| = 0$, such that the set $\{\mathbf{f}(t) \mid t \in [0, T] - N\}$ is separable.

THEOREM 7 (Pettis). The mapping $\mathbf{f} : [0, T] \rightarrow X$ is strongly measurable if and only if \mathbf{f} is weakly measurable and almost separably valued.

DEFINITIONS. (i) If $\mathbf{s}(t) = \sum_{i=1}^m \chi_{E_i}(t) u_i$ is simple, we define

$$(6) \quad \int_0^T \mathbf{s}(t) dt := \sum_{i=1}^m |E_i| u_i.$$

(ii) We say the strongly measurable function $\mathbf{f} : [0, T] \rightarrow X$ is summable if there exists a sequence $\{\mathbf{s}_k\}_{k=1}^\infty$ of simple functions such that

$$(7) \quad \int_0^T \|\mathbf{s}_k(t) - \mathbf{f}(t)\| dt \rightarrow 0 \quad \text{as } k \rightarrow \infty.$$

(iii) If f is summable, we define

$$(8) \quad \int_0^T \mathbf{f}(t) dt = \lim_{k \rightarrow \infty} \int_0^T \mathbf{s}_k(t) dt.$$

THEOREM 8 (Bochner). *A strongly measurable function $\mathbf{f} : [0, T] \rightarrow X$ is summable if and only if $t \mapsto \|\mathbf{f}(t)\|$ is summable. In this case*

$$\left\| \int_0^T \mathbf{f}(t) dt \right\| \leq \int_0^T \|\mathbf{f}(t)\| dt,$$

and

$$\left\langle u^*, \int_0^T \mathbf{f}(t) dt \right\rangle = \int_0^T \langle u^*, \mathbf{f}(t) \rangle dt$$

for each $u^* \in X^*$.

Good books for measure theory are Folland [F2] and DiBenedetto [DB2]. See Yosida [Y, Chapter V, Sections 4–5] for proofs of Theorems 7, 8.

BIBLIOGRAPHY

- [Am] W. F. Ames, *Nonlinear Partial Differential Equations in Engineering* (Vol. I–II), Mathematics in Science and Engineering 18, 18-II, Academic Press, 1965, 1972.
- [Ar1] V. I. Arnold, *Mathematical Methods of Classical Mechanics*, Springer, 1986.
- [Ar2] V. I. Arnold, *Lectures on Partial Differential Equations*, Springer, 2004.
- [Bk] I. Bakelman, *Convex Analysis and Nonlinear Geometric Elliptic Equations*, Springer, 1994.
- [Ba] V. Barbu, *Nonlinear Semigroups and Differential Equations in Banach Spaces*, Noordhoff, 1976.
- [B-CD] M. Bardi and I. Capuzzo Dolcetta, *Optimal Control and Viscosity Solutions of Hamilton–Jacobi–Bellman Equations*, Birkhäuser, 1997.
- [B-L-P] A. Bensoussan, J.-L. Lions and G. Papanicolaou, *Asymptotic Analysis for Periodic Structures*, North-Holland, 1978.
- [Be] S. Benton, *The Hamilton–Jacobi Equation: A Global Approach*, Academic Press, 1977.
- [Bt] D. L. Bernstein, *Existence Theorems in Partial Differential Equations*, Annals of Mathematical Studies, Princeton, 1950.
- [B-J-S] L. Bers, F. John and M. Schechter, *Partial Differential Equations*, American Math. Society, 1964.
- [Bs] A. Bresson, *Hyperbolic Systems of Conservation Laws: The One-dimensional Cauchy Problem*, Oxford, 2000.
- [Br1] H. Brezis, *Analyse Fonctionnelle*, Masson, 1983.
- [Br2] H. Brezis, *Opérateurs Maximaux Monotones et Semi-groupes de Contractions dans les Espaces de Hilbert*, North-Holland, 1973.
- [C-C] X. Cabre and L. Caffarelli, *Fully Nonlinear Elliptic Equations*, Colloquium Publications, Vol. 43, American Math. Society, 1995.
- [C] C. Carathéodory, *The Calculus of Variations and Partial Differential Equations of First Order*, Chelsea, 1982.

- [Ch] C. Chester, *Techniques in Partial Differential Equations*, McGraw-Hill, 1971.
- [C-M] A. Chorin and J. Marsden, *A Mathematical Introduction to Fluid Mechanics* (3rd ed.), Springer, 1993.
- [Cl] D. Colton, *Partial Differential Equations: An Introduction*, Dover, 2004.
- [C-F] R. Courant and K. O. Friedrichs, *Supersonic Flow and Shock Waves*, Springer, 1976.
- [C-H] R. Courant and D. Hilbert, *Methods of Mathematical Physics*, Vol. 2, Wiley-Interscience, 1962.
- [D-M] B. Dacorogna and P. Marcellini, *Implicit Partial Differential Equations*, Progress in Partial Differential Equations and their Applications 37, Birkhäuser, 1999.
- [D] C. Dafermos, *Hyperbolic Conservation Laws in Continuum Physics*, Springer, 2005.
- [Da] E. B. Davies, *Heat Kernels and Spectral Theory*, Cambridge University Press, 1989.
- [DB1] E. DiBenedetto, *Partial Differential Equations*, Birkhäuser, 1995.
- [DB2] E. DiBenedetto, *Real Analysis*, Birkhäuser, 2002.
- [Dr] P. G. Drazin, *Solitons*, Cambridge University Press, 1983.
- [D-Z] P. Duchateau and D. W. Zachman, *Partial Differential Equations*, Schaum's Outline Series in Math., McGraw-Hill, 1986.
- [E-T] I. Ekeland and R. Teman, *Convex Analysis and Variational Problems*, North-Holland Elsevier, 1976.
- [E] L. C. Evans, *Weak Convergence Methods for Nonlinear Partial Differential Equations*, CBMS Regional Conference Series in Mathematics 74, American Math. Society, 1990.
- [E-G] L. C. Evans and R. F. Gariepy, *Measure Theory and Fine Properties of Functions*, CRC Press, 1992 (An errata sheet for this book is available through the math.berkeley.edu website).
- [F-L-S] R. Feynman, R. Leighton and M. Sands, *The Feynman Lectures in Physics*, Vol. II, Addison-Wesley, 1966.
- [F-S] W. Fleming and M. Soner, *Controlled Markov Processes and Viscosity Solutions*, Springer, 1993.
- [F1] G. Folland, *Introduction to Partial Differential Equations*, Princeton University Press, 1976.
- [F2] G. Folland, *Real Analysis: Modern Techniques and their Applications*, Wiley-Interscience, 1984.
- [Fd] M. Freidlin, *Functional Integration and Partial Differential Equations*, Princeton University Press, 1985.
- [Fr1] A. Friedman, *Partial Differential Equations of Parabolic Type*, Prentice-Hall, 1964.
- [Fr2] A. Friedman, *Partial Differential Equations*, Holt, Rinehart, Winston, 1969.
- [Fr3] A. Friedman, *Variational Principles and Free Boundary Problems*, Wiley-Interscience, 1982.
- [Ga] F. Gantmacher, *The Theory of Matrices*, Chelsea, 1977.
- [G] P. Garabedian, *Partial Differential Equations*, Wiley, 1964.

- [Gi] M. Giaquinta, *Multiple Integrals in the Calculus of Variations and Nonlinear Elliptic Systems*, Princeton University Press, 1983.
- [G-H] M. Giaquinta and S. Hildebrandt, *Calculus of Variations*, Vol. I–II, Springer, 1996.
- [Gh] N. Ghoussoub, *Self-dual Partial Differential Systems and Their Variational Principles*, Springer, 2009.
- [G-T] D. Gilbarg and N. Trudinger, *Elliptic Partial Differential Equations of Second Order* (2nd ed.), Springer, 1983.
- [Gr] P. Grindrod, *Patterns and Waves: The Theory and Applications of Reaction-Diffusion Equations*, Oxford, 1991.
- [Hl] F. Hélein, *Harmonic Maps, Conservation Laws and Moving Frames* (2nd ed.), Cambridge, 2002.
- [Hg] S. Helgason, *The Radon Transform*, Birkhäuser, 1980.
- [He] D. Henry, *Geometric Theory of Semilinear Parabolic Equations*, Lecture Notes in Mathematics 840, Springer, 1981.
- [H-R] H. Holden and N. H. Risebro, *Front Tracking for Hyperbolic Conservation Laws*, Springer, 2002.
- [H] L. Hörmander, *The Analysis of Linear Partial Differential Operators*, Vol. 1–4, Springer, 1983–85.
- [Ik] M. Ikawa, *Hyperbolic Partial Differential Equations and Wave Phenomena*, Translations of Mathematical Monographs 189, American Math. Society, 2000.
- [I] V. Isakov, *Inverse Problems for Partial Differential Equations*, Springer, 1998.
- [J1] F. John, *Plane Waves and Spherical Means Applied to Partial Differential Equations*, Wiley-Interscience, 1955.
- [J2] F. John, *Partial Differential Equations* (4th ed.), Springer, 1995.
- [J3] F. John, *Nonlinear Wave Equations: Formation of Singularities*, University Lecture Series 2, American Math. Society, 1990.
- [Jo] J. Jost, *Partial Differential Equations* (2nd ed.), Springer, 2007.
- [K] T. Kato, *Perturbation Theory for Linear Operators* (2nd ed.), Springer, 1980.
- [K-S] D. Kinderlehrer and G. Stampacchia, *An Introduction to Variational Inequalities and Their Applications*, Academic Press, 1980.
- [Kl] S. Klainerman, Partial Differential Equations, in *The Princeton Companion to Mathematics* (T. Gowers, ed.), Princeton, 2008.
- [Kr1] N. V. Krylov, *Lectures on Elliptic and Parabolic Equations in Hölder Spaces*, Graduate Studies in Mathematics 12, American Math. Society, 1996.
- [Kr2] N. V. Krylov, *Lectures on Elliptic and Parabolic Equations in Sobolev Spaces*, Graduate Studies in Mathematics 96, American Math. Society, 2008.
- [L] O. A. Ladyzhenskaya, *The Boundary Value Problems of Mathematical Physics*, Springer, 1985.
- [L-S-U] O. A. Ladyzhenskaya, V. A. Solonnikov and N. N. Uraltseva, *Linear and Quasilinear Equations of Parabolic Type*, American Math. Society, 1968.
- [L-U] O. A. Ladyzhenskaya and N. N. Uraltseva, *Linear and Quasilinear Elliptic Equations*, Academic Press, 1968.
- [Lx1] P. Lax, *Hyperbolic Systems of Conservation Laws and the Mathematical Theory of Shock Waves*, SIAM, 1973.

- [Lx2] P. Lax, *Functional Analysis*, Wiley, 2002.
- [Lx3] P. Lax, *Hyperbolic Partial Differential Equations*, Courant Lecture Notes in Mathematics 14, American Math. Society, 2006.
- [Lx-P] P. Lax and R. Phillips, *Scattering Theory*, Academic Press, 1967.
- [LV] R. LeVeque, *Numerical Methods for Conservation Laws*, Birkhäuser, 1992.
- [L-L] E. Lieb and M. Loss, *Analysis*, Graduate Studies in Mathematics 14, American Math. Society, 1997.
- [Lb] G. M. Lieberman, *Second Order Parabolic Partial Differential Equations*, World Scientific, 1996.
- [L-W] F. Lin and C. Wang, *The Analysis of Harmonic Maps and Their Heat Flows*, World Scientific, 2008.
- [L-P] F. Linares and G. Ponce, *Introduction to Nonlinear Dispersive Equations*, Springer, 2009.
- [L1] J.-L. Lions, *Equations Différentielles Operationnelles et Problèmes aux Limites*, Springer, 1961.
- [L2] J.-L. Lions, *Quelques Méthodes de Résolution des Problèmes aux Limites non Linéaires*, Dunod, 1969.
- [L-M] J.-L. Lions and E. Magenes, *Nonhomogeneous Boundary Value Problems and Applications*, Vol. I–III, Springer, 1972.
- [Li] P.-L. Lions, *Generalized Solutions of Hamilton–Jacobi Equations*, Research Notes in Mathematics 69, Pitman, 1982.
- [Lo] J. D. Logan, *Applied Mathematics* (3rd ed.), Wiley-Interscience, 2006.
- [Mj] A. Majda, *Compressible Fluid Flow and Systems of Conservation Laws in Several Space Variables*, Applied Mathematical Sciences 53, Springer, 1984.
- [Mr] P. Markowich, *Applied Partial Differential Equations: A Visual Approach*, Springer, 2007.
- [MO] R. McOwen, *Partial Differential Equations: Methods and Applications*, Prentice Hall, 1996.
- [M] V. P. Mikhailov, *Partial Differential Equations*, Mir, 1978.
- [Mi] F. H. Miller, *Partial Differential Equations*, Wiley, 1947.
- [Mo] C. B. Morrey, Jr, *Multiple Integrals in the Calculus of Variations*, Springer, 1966.
- [Ne] J. Neu, *Training Manual on Transport and Fluids*, Graduate Studies in Mathematics 109, American Math. Society, 2009.
- [N] Y. Nishiura, *Far from Equilibrium Dynamics*, Translations of Mathematical Monographs 209, American Math. Society, 2002.
- [O] P. Olver, *Applications of Lie Groups to Differential Equations*, Springer, 1986.
- [Pa] L. E. Payne, *Improperly Posed Problems in Partial Differential Equations*, SIAM, 1975.
- [Py] I. G. Petrovsky, *Lectures on Partial Differential Equations*, Wiley-Interscience, 1954.
- [P] M. Pinsky, *Partial Differential Equations and Boundary-Value Problems, with Applications* (2nd ed.), McGraw–Hill, 1991.
- [P-W] M. Protter and H. Weinberger, *Maximum Principles in Differential Equations*, Prentice–Hall, 1967.

- [P-S] P. Pucci and J. Serrin, *The Maximum Principle*, Birkhäuser, 2007.
- [Ra] P. Rabinowitz, *Minimax Methods in Critical Point Theory with Applications to Differential Equations*, CBMS Regional Conference Series in Mathematics 65, American Math. Society, 1986.
- [R] J. Rauch, *Partial Differential Equations*, Springer, 1992.
- [R-S1] M. Reed and B. Simon, *Methods of Modern Mathematical Physics I: Functional Analysis* (2nd ed.), Academic Press, 1980.
- [R-S2] M. Reed and B. Simon, *Methods of Modern Mathematical Physics III: Scattering Theory*, Academic Press, 1979.
- [R-R] M. Renardy and R. Rogers, *A First Graduate Course in Partial Differential Equations*, Springer, 1993.
- [Rd] W. Rudin, *Principles of Mathematical Analysis* (3rd ed.), McGraw-Hill, 1976.
- [Ru] H. Rund, *The Hamilton-Jacobi Theory in the Calculus of Variations*, Krieger, 1973.
- [S-S] J. Shatah and M. Struwe, *Geometric Wave Equations*, Courant Lecture Notes in Mathematics 2, American Math. Society, 1998.
- [Sm] V. I. Smirnov, *Integral Equations and Partial Differential Equations*, A Course of Higher Mathematics Vol. IV, Addison-Wesley, 1964.
- [S] J. Smoller, *Shock Waves and Reaction-Diffusion Equations*, Springer, 1983.
- [Sn] I. Sneddon, *Elements of Partial Differential Equations*, McGraw-Hill, 1957.
- [So] C. Sogge, *Lectures on Nonlinear Wave Equations* (2nd ed.), International Press, 2008.
- [Se] E. Stein, *Harmonic Analysis*, Princeton University Press, 1993.
- [S-W] E. Stein and G. Weiss, *Fourier Analysis on Euclidean Spaces*, Princeton University Press, 1971.
- [St1] W. Strauss, *Nonlinear Wave Equations*, CBMS Regional Conference Series in Mathematics 73, American Math. Society, 1992.
- [St2] W. Strauss, *Partial Differential Equations: An Introduction* (2nd ed.), Wiley, 2008.
- [Str] M. Struwe, *Variational Methods* (4th ed.), Springer, 2008.
- [To] T. Tao, *Nonlinear Dispersive Equations: Local and Global Analysis*, CBMS #106, American Math. Society, 2006.
- [Tr] L. Tartar, *An Introduction to Sobolev Spaces and Interpolation Spaces*, Springer, 2007.
- [Ta] M. Taylor, *Partial Differential Equations*, Vol. I-III, Springer, 1996.
- [Te] R. Temam, *Navier-Stokes Equations*, North-Holland, 1977.
- [T-Z] D. W. Thoe and E. C. Zachmanoglou, *Introduction to Partial Differential Equations with Applications*, Dover, 1986.
- [T] F. Trèves, *Basic Linear Partial Differential Equations*, Academic Press, 1975.
- [V] J. L. Vazquez, *The Porous Medium Equation: Mathematical Theory*, Oxford, 2007.
- [Wb] A. G. Webster, *Partial Differential Equations of Mathematical Physics*, Dover, 1955.
- [We] H. Weinberger, *A First Course in Partial Differential Equations*, Blaisdell, 1965.

- [Wh] G. B. Whitham, *Linear and Nonlinear Waves*, Wiley-Interscience, 1974.
- [Wl] J. Wloka, *Partial Differential Equations*, Cambridge University Press, 1987.
- [X-Z] L. Xiao and T. Zhang, *The Riemann Problem and Interaction of Waves in Gas Dynamics*, Pitman, 1989.
- [Y] K. Yosida, *Functional Analysis* (6th ed.), Springer, 1980.
- [Za] E. Zauderer, *Partial Differential Equations of Applied Mathematics*, Wiley-Interscience, 1983.
- [Zd] E. Zeidler, *Nonlinear Functional Analysis and Its Applications*, Vol.1–4, Springer.
- [Zh] Y. Zheng, *Systems of Conservation Laws: Two Dimensional Riemann Problems*, Progress in Partial Differential Equations and Their Applications 38, Birkhäuser, 2001.
- [Z] W. Ziemer, *Weakly Differentiable Functions*, Springer, 1989.
- [Zw] D. Zwillinger, *Handbook of Differential Equations* (3rd ed.), Academic Press, 1997.

INDEX

- a priori estimates, 539
- action, 115, 163, 517
- activator, 175
- adjoint, 446, 723
 - bilinear form, 320
 - of elliptic operator, 320
- admissible states, 631
- Airy's equation, 4, 178
- almost everywhere (a.e.), 730
- almost separably valued function, 733
- amplitude and phase, 218
- analyticity, 237–244
 - of harmonic functions, 31
 - of solutions to heat equation, 62
 - power series, 240–244
- antigradient flux function, 655
- Arzela–Ascoli compactness criterion, 289, 477, 580, 718
- asymptotics, 156, 211–232
- averages, 700

- Bäcklund transformation, 694
- backwards uniqueness
 - heat equation, 63
- Banach space, 255, 262, 719
- Barenblatt–Kompaneetz–Zeldovich solution, 186
- barrier, 367
- basis
 - of eigenfunctions, 170, 355, 375, 381, 384, 386, 409, 411, 412, 529
 - orthonormal, 720
- beam equation, 4, 448
- Bernoulli's law, 208
- Bessel potential, 192
- biharmonic equation, 366

- bilinear form, 314
 - estimates for, 317–319
 - time-dependent, 373, 399, 424
- bistable equation, 180
- blow-up
 - conservation law, 639
 - nonlinear heat equation, 548, 549
 - nonlinear wave equation, 674, 686–691
 - porous medium equation, 171
 - reaction diffusion equation, 538
- Bochner's Theorem, 734
- bootstrap argument, 487
- boundary, 710
 - analytic, 710
 - C^k , 710
 - normal to, 710
 - straightening, 102, 235, 239, 337, 343, 711
- boundary condition
 - Dirichlet, 312
 - mixed, 366
 - Neumann, 366, 446
 - nonlinear, 523
 - Robin, 366
- boundary data
 - admissible, 105
 - compatibility conditions on, 103
 - noncharacteristic, 104, 105, 163
- boundary-value problem
 - eigenvalue problem, 354
 - homogenization of, 229
 - Laplace's equation, 40
 - nonlinear boundary conditions, 522
 - nonlinear eigenvalue problem, 489
 - nonlinear Poisson equation, 543, 555
 - Poisson's equation, 27, 41

- quasilinear elliptic equation, 527, 541
- second-order elliptic equation, 311, 314
- semilinear elliptic equation, 507, 551
- Browder–Minty method, 532, 580
- Burgers' equation, 5, 139, 140, 142
 - vanishing viscosity, 216
 - viscous, 207
- calculus of variations, 11, 43, 116, 123, 453–526
- Cauchy data, 233
- Cauchy problem
 - for boundary data, 233
 - for heat equation, 47
 - for Laplace's equation, 245
- Cauchy sequence, 255, 263, 673, 719
- Cauchy–Kovalevskaya Theorem, 10, 239–244
- chain rule, 308
- characteristic equation, 419
- characteristics, 10, 19, 96–114, 151, 162, 212, 228, 420
 - and boundary conditions, 102–105
 - and local solution to PDE, 105–108
 - applications of, 109–114
 - crossing, 111
 - derivation, 97
 - examples of, 99–102
 - projected, 98
 - summary of characteristic ODE, 98
- Clairaut's equation, 93
- Closed Graph Theorem, 437, 721
- coarea formula, 416, 713
- coercivity, 465, 475
- cofactor matrix, 432, 462
- Cole–Hopf transformation, 207
- compact
 - embedding, 286
 - linear operator, 724
 - mapping, 539
- complete
 - integral, 92, 161
 - linear space, 255, 719
- condition E, 656
- conformal
 - energy, 518
 - transformation, 87
- conservation law, 5, 8, 10, 135–161, 163, 609–657
 - characteristics for, 111, 142
 - decay in sup-norm, 156
 - decay to N -wave, 157
 - derivation, 609
 - implicit solution, 113
 - initial-value problem, 135
 - integral solution, 136, 147
 - Lax–Oleinik formula, 145
 - Rankine–Hugoniot condition, 139
 - systems, 6, 12, 609–649
 - uniqueness, 149–153, 650–653
 - viscous, 246
- conservation of
 - energy, 515, 611, 660, 670, 687, 692, 695
 - mass, 611, 612, 695
 - momentum, 525, 611, 612, 695
- constitutive relation, 611, 620, 637
- constraints
 - incompressibility, 497
 - integral, 488
 - one-sided, 492
 - pointwise, 495
- contraction, 534, 536, 566, 653, 657
 - semigroup, 435
- control theory, 590
- controls
 - admissible, 591
 - optimal, 11, 600
- converting nonlinear into linear equations, 206–210, 246
- convex function, 163, 528, 560, 705
 - domain, 560
 - subdifferential of, 164, 560
 - supporting hyperplane, 705
- convexity, 469
 - uniform, 130, 142, 143, 705
- corrector problem
 - for Green's function, 34
 - for homogenization, 232, 249
- cost
 - functional, 592
 - running, 592
 - terminal, 592
- Courant minimax principle, 368
- Crandall, M.–Lions, P.-L., 607
- critical
 - exponent, 245, 511, 551, 575, 679–686, 694
 - point, 117, 454, 501–506
 - value, 502
- curvature, 212
 - mean, 457, 523
 - principal curvatures, 227
- cutoff function, 328
- d'Alembert's formula, 68, 69, 78, 81, 89
- Deformation Theorem, 503, 506
- DeGiorgi–Nash estimates, 487
- degree, 521
- Derrick–Pohozaev identity, 554, 575
- difference quotients, 291–293, 329, 335, 473, 484, 665, 675
 - and weak derivatives, 292
 - definition of, 292
- differential equations

- in Banach spaces, 434
- in Hilbert spaces, 504, 565
- diffusion, 211, 313
 - and transport, 211–216
 - equation, 4, 44
- DiPerna, R., 166
- Dirac measure, 25, 35, 48, 211, 213
- Dirichlet's
 - boundary condition, 312
 - energy functional, 42, 523, 524
 - principle, 42, 456
 - generalized, 456
- dispersion, 177
- dissipation, 177
- divergence form PDE, 312, 372, 527
- domain of dependence
 - generalized wave equation, 416
 - nonlinear wave equation, 660–662
 - wave equation, 83
- domain variations, 369, 511–519, 713
- Dominated Convergence Theorem, 133, 153, 222, 433, 474, 546, 648, 672, 732
- doubling variables, 587, 650
- Douglis, A., 166
- dual space, 722
 - of H_0^1 , 299
- dual variational principle, 249, 523
- duality of Hamiltonian and Lagrangian, 121
- Duhamel's principle, 49, 80
- dynamic programming, 11, 590, 592, 605, 606

- Egoroff's Theorem, 468, 731
- eigenfunction, 169, 323
 - basis, 355
- eigenvalue, 11, 169, 323, 354, 616, 618, 722
 - nonsymmetric elliptic operator, 360–365
 - principal, 11, 356–365, 446, 547, 574
 - symmetric elliptic operator, 354–360
- eigenvector, 354, 618, 722
 - left, 616
 - right, 616
- eikonal equation, 5, 93, 604
 - generalized, 416
- elasticity
 - linear, 6, 89
 - nonlinear, 479, 500
- elliptic operator
 - fully nonlinear, 606
 - linear, 312
 - divergence form, 312, 313, 327, 354, 365
 - nondivergence form, 312, 344, 360
 - quasilinear, 483, 541
- elliptic regularization, 521
- ellipticity, 313
- Emden–Fowler transformation, 245

- energy decay
 - wave equation, 519–520
- energy estimates
 - quasilinear elliptic equation, 530
 - second-order elliptic equation, 317
 - second-order hyperbolic equation, 402
 - second-order parabolic equation, 376
 - symmetric hyperbolic system, 426
- energy flux, 661, 677, 680
- energy methods
 - heat equation, 62–65
 - Laplace's equation, 41–43
 - wave equation, 82–84
- entropy condition, 12, 142, 148, 631, 641
 - condition E, 656
 - Lax's, 631
 - Liu's, 644
- entropy solution
 - single conservation law, 149, 165, 656
 - system, 647, 650
- entropy/entropy-flux pair, 646, 649, 655, 656
- envelopes, 94, 162
 - construction of solutions using, 94
- equipartition of energy, 90, 194–195, 692
- essential supremum (ess sup), 731
- Euler equations, 611, 620
 - barotropic, 637, 655
 - incompressible, 6, 208
- Euler's formula for Jacobians, 162, 501
- Euler–Lagrange equation, 456, 472, 512, 513
 - harmonic maps, 496
 - one-dimensional, 116, 163
 - systems, 460, 476
- Euler–Poisson–Darboux equation, 70, 75
- exponential solutions, 176
- extensions, 268–271

- factoring PDE, 67, 419
- Fatou's Lemma, 569, 673, 732
- feedback control, 592
- Fick's law, 21
- first-order equation, 9, 91–166
 - complete integral, 92
 - notation for, 91
- first-order hyperbolic system, 11, 201, 421–433
- fixed point methods, 11, 533–543
- fixed point theorem
 - Banach's, 534, 536, 537, 574, 664, 670, 675
 - Brouwer's, 463, 529
 - Schaefer's, 539, 542
 - Schauder's, 538
- Fokker–Planck equation, 4

- Fourier transform, 10, 187–196, 297–298, 309, 429
 and Radon transform, 198–200
 and Sobolev spaces, 297
 applications, 190–196
 definition of, 187, 189
 inversion formula, 189
 properties, 187, 189
 Fourier's law, 21
 Fredholm alternative, 231, 725, 727
 free boundary problem, 495
 frequency function, 525
 Fujita, H., 577
 fully nonlinear PDE, 2
 elliptic, 606
 function spaces
 BMO, 291
 C^k , 702
 $C^{k,\gamma}$, 255
 C_1^2 , 702
 H^k , 258
 H^s , 298
 H^{-1} , 299
 H_0^k , 259
 L^p , 702, 719
 $W^{k,p}$, 258
 $W_0^{k,p}$, 259
 Banach space-valued, 301–305
 function variations, 511–519
 fundamental solution
 heat equation, 46, 87, 185, 192
 Laplace's equation, 22, 25
 Schrödinger's equation, 193

 Galerkin's method, 170, 375, 401, 446, 529
 Gauss–Bonnet Theorem, 521
 Gauss–Green Theorem, 20, 306, 482, 711
 general integral, 95
 generator of semigroup, 435–439
 genuine nonlinearity, 623, 630, 636
 geometric optics, 218–228, 415
 Gidas, B.–Ni, W.–Nirenberg, L., 577
 gradient flow, 566, 576
 Green's
 formulas, 33, 34, 712
 function, 33–41
 derivation of, 33–35
 for ball, 40
 for half-space, 37
 symmetry of, 35
 Grillakis, M., 696
 group velocity, 178

 Hadamard's
 example, 245
 variational formula, 369

 Hamilton's differential equations, 114, 115, 119
 Hamilton–Jacobi equation, 5, 10, 11, 94, 96, 114–135, 171, 415, 579
 characteristics for, 113, 115, 122
 Hopf–Lax formula, 123
 semiconcavity condition, 129
 solution a.e., 127
 uniqueness, 131
 weak solution, 131
 Hamilton–Jacobi–Bellman equation, 596
 and optimal controls, 600
 Hamiltonian, 118, 596
 duality with Lagrangian, 121
 harmonic
 function, 20
 mapping, 495, 524
 heat ball, 53
 heat equation, 4, 9, 44–65, 176
 backwards uniqueness, 63
 derivation, 44
 energy methods, 62–65
 fundamental solution, 46
 infinite propagation speed, 49
 initial-value problem, 47
 maximum principle, 57
 mean-value formula, 53
 nonhomogeneous problem, 44, 49
 nonlinear, 245, 246, 248, 547, 575, 576
 regularity, 59, 61
 strong maximum principle, 55
 uniqueness, 57, 59
 Helmholtz's equation, 3, 323
 Hilbert space, 720
 Hille–Yosida Theorem, 439, 442, 444
 hodograph transform, 209
 Hölder spaces, 254–255
 homogeneous PDE, 2
 homogenization, 229–232, 248
 Hopf's Lemma, 347, 350, 361, 363, 365, 556
 Hopf–Lax formula, 10, 120–135, 164, 579, 600–602
 horizon
 finite, 592
 infinite, 605
 Huygens' principle, 80, 202
 hyperbolic system
 constant coefficients, 201–202, 429–433
 nonlinear, 615, 654
 symmetric, 423–429
 hyperbolicity, 419, 421, 422
 for symmetric system, 422
 strict, 422, 615, 620, 637
 uniform, 424
 hyperelastic materials, 479

- Implicit Function Theorem, 104, 224, 228,
 490, 555, 618, 619, 627, 718
 incompressibility, 497–501
 inequality
 Cauchy's, 706
 with ϵ , 706
 Cauchy–Schwarz, 43, 84, 417, 445, 708,
 720, 728
 Gagliardo–Nirenberg–Sobolev, 10,
 276–280
 general Sobolev-type, 284
 Gronwall's, 377, 379, 381, 384, 404, 408,
 409, 411, 417, 427, 429, 537, 595, 662,
 669, 676
 differential form, 708
 integral form, 709
 Hardy's, 296–297, 308, 677
 Harnack's, 32, 86, 351, 391
 Hölder's, 706
 interpolation, 306, 307, 685
 Jensen's, 705
 Minkowski's, 263, 707
 Morrey's, 10, 280–284
 Poincaré's, 280, 289–291, 470, 531
 Young's, 706
 with ϵ , 706
 inhibitor, 175
 initial-value problem
 Burgers' equation, 139
 conservation law, 135, 143, 149, 610, 647,
 649
 Hamilton–Jacobi equation, 114, 123, 128,
 134, 579, 582, 586, 590, 600
 heat equation, 47, 192, 203
 nonlinear wave equation, 663, 667, 670,
 674, 677, 683, 687, 689
 quasilinear parabolic equation, 206, 580
 Schrödinger's equation, 193
 telegraph equation, 195
 transport equation, 18, 85
 viscous Burgers' equation, 207
 wave equation, 67, 194, 200, 204, 219
 initial/boundary-value problem
 beam equation, 448
 heat equation, 57, 88, 168
 parabolic equation, 547
 quasilinear parabolic equation, 572
 reaction-diffusion system, 172, 535
 second-order hyperbolic equation, 399
 second-order parabolic equation, 372
 wave equation, 69, 82, 519
 inner product, 720
 integral solution, 136, 147, 612, 650
 integration by parts formula, 712
 interior ball condition, 348, 556
 invariance, 511–520
 Galilean, 695
 in time, 515
 scaling, 45, 185, 245, 514, 516, 680
 translation, 514, 525
 Inverse Function Theorem, 105, 209, 210,
 633, 634, 716
 inversion
 hyperbolic, 517
 through sphere, 39
 irreversibility, 193, 576
 irrotational fluid equations, 209

 Jörgens, K., 696

 k -contact discontinuity, 630
 k -rarefaction wave, 624
 k -shock wave, 631
 k -simple wave, 622
 Kelvin transform
 Laplace's equation, 87
 wave equation, 517, 518, 693
 Kirchhoff's formula, 72
 Klein–Gordon equation, 4, 177, 659, 692
 Kolmogorov's equation, 4
 Korteweg–deVries (KdV) equation, 5, 178,
 247
 Kovalevskaya's example, 249
 Kružkov, S., 657

 Lagrange multiplier
 as eigenvalue, 489
 as function, 496
 integral constraints, 489
 pressure as, 497
 Lagrangian, 115, 454, 459
 duality with Hamiltonian, 121
 null, 461, 463, 521, 522
 Lambertian surface, 604
 Laplace transform, 203–206, 438
 applications, 203–206
 definition of, 203
 Laplace's equation, 3, 9, 20–43, 313
 analyticity, 31
 derivation, 20
 Green's function, 33–41
 Harnack's inequality, 32
 Liouville's Theorem, 30
 mean-value formulas, 25
 regularity, 28, 29
 Laplace's method, 216
 Lax pairs, 247
 Lax–Milgram Theorem, 315, 317, 319, 365,
 508, 728
 Lax–Oleinik formula, 10, 143–161, 217
 Lebesgue
 integral, 731
 measure, 729
 point, 732

- Lebesgue's Differentiation Theorem, 296,
 714, 732
 Legendre transform, 120, 121, 163, 164, 601
 classical, 210
 Leibniz's formula, 13, 261
 linear degeneracy, 623, 629
 linear operator, 721
 bounded, 721
 closed, 721
 symmetric, 355, 723
 linear PDE
 definition of, 2
 Liouville's
 equations, 3, 247
 Theorem, 30
 Lipschitz continuity, 700
 and differentiability a.e., 294
 and weak partial derivatives, 294
 lower semicontinuity, 560
 weak, 467, 468, 562

 majorants, 238, 242
 maximum principle, 10
 Cauchy problem for heat equation, 57
 heat equation, 55, 88
 Laplace's equation, 27, 85
 strong, 360, 361
 heat equation, 55
 Laplace's equation, 27
 second-order elliptic equation, 347–351
 second-order parabolic equation,
 396–398
 weak
 second-order elliptic equation, 344–347
 second-order parabolic equation,
 389–391
 Maxwell's equations, 6, 89
 nonlinear, 655
 Mazur's Theorem, 471, 562, 723
 mean-value formulas
 heat equation, 53
 Laplace's equation, 25, 26, 85
 measurable
 function, 730
 strongly, 733
 weakly, 733
 sets, 730
 method of descent, 74, 78, 692
 minimal surface equation, 5, 210, 457
 minimax methods, 368, 505–511
 minimizer, 455, 460
 existence of, 465–471, 475, 479
 local, 480–482
 regularity, 482–487
 uniqueness of, 471
 mixed boundary conditions, 366
 mollifier, 264, 713
 momentum, generalized, 118
 Monge–Ampère equation, 5
 Monotone Convergence Theorem, 469, 732
 monotonicity, 11, 527–533, 561
 strict, 532
 monotonicity formula
 Almgren's, 525
 Laplace's equation, 525
 p -Laplacian, 515
 Morawetz identity, 518, 681
 Morse Lemma, 223
 Mountain Pass Theorem, 505, 510
 moving plane method, 558–559
 multiindices, 13, 701
 Multinomial Theorem, 13, 238
 multipliers, 511–519
 multivalued equation, 523, 565, 566

 N -wave, 157
 Navier–Stokes equations, 6
 Neumann boundary conditions, 366
 Newton's law, 66, 118, 611
 Noether's Theorem, 511–520
 noncharacteristic
 boundary data, 104, 105, 163
 surface, 234, 235, 237
 nonconvex flux function, 656–657
 nondivergence form PDE, 312, 372
 nonexistence of solutions, 547–554,
 639–641, 686–691, 695
 nonhomogeneous problem
 heat equation, 49, 51
 transport equation, 19
 wave equation, 80
 nonlinear eigenvalue problem, 489
 nonlinear Schrödinger (NLS) equation, 695
 norm, 255, 719, 720
 normal
 derivative, 711
 to surface, 710
 notation
 for estimates, 703
 for functions, 699–703
 for matrices, 697–698
 geometric, 698–699
 null Lagrangian, 461, 463, 521, 522

 O , o notation, 704
 obstacle problem, 492
 Ohm's law, 21
 Oleinik, O., 166
 optimality conditions, 592
 orthogonality, 720

 p -Laplacian, 5, 514
 monotonicity formula, 515
 p -system, 610, 649

- generalization of, 654
 - traveling waves for, 644
- Palais–Smale condition, 502, 503, 505, 509
- parabolic
 - approximation, 215
 - boundary, 52
 - cylinder, 52
- parabolicity, 372
- partial derivatives, 13, 701
 - count of, 12
- partial differential equation
 - definition of, 1
- partition of unity, 265, 267, 270, 274, 306
- pattern formation, 175
- penalty method, 574
- periodic solution, 229–232, 574
- Perron–Frobenius Theorem, 354
- phase, 218
 - plane, 181
 - velocity, 178, 693
- Plancherel’s Theorem, 187
- plane wave, 176, 178, 196, 422
- Poisson’s
 - equation, 20, 23, 313, 446
 - nonlinear, 5, 457
 - formula
 - ball, 41, 86, 87
 - half-space, 37, 86
 - wave equation, 74, 79
 - kernel
 - ball, 41
 - half-space, 37
- polar coordinates, 712
- polyconvexity, 478
- porous medium equation, 5, 170, 185
 - Barenblatt–Kompaneetz–Zeldovich
 - solution of, 186
 - variant of, 245
- potentials, 208, 209
- projected characteristics, 98
- propagation speed, 187, 202
 - finite, 78, 84, 89, 164, 414–417, 660–662
 - infinite, 49, 396
- proper function, 560

- quasilinear PDE, 2

- Rademacher’s Theorem, 127, 296
- radial solution, 21, 22, 245
- radiation condition
 - Sommerfeld, 370
- Radon transform, 196–202
 - and Fourier transform, 198–200
 - applications, 200–202
 - definition of, 197
 - inversion formula, 199
 - properties, 197–199
 - vanishing of, 197, 200
- Rankine–Hugoniot condition, 139, 140, 143, 614
- rarefaction
 - curve, 622
 - wave, 140, 624, 654, 655
- Rauch, J., 696
- Rayleigh’s formula, 357
- reaction-diffusion equation, 547
 - bistable, 180
 - scalar, 5
 - system, 6, 172, 535
- reflection, 69, 86, 269, 558
- reflexive space, 722
- regularity, 8
 - heat equation, 59, 61
 - Laplace’s equation, 28, 29
 - minimizers, 482–487
 - second-order elliptic equation, 326–343
 - second-order hyperbolic equation, 408–414
 - second-order parabolic equation, 380–388
- Rellich–Kondrachov Compactness Theorem, 286, 322
- resolvent, 203
 - identity, 438
 - nonlinear, 563
 - set, 437, 721
- response of system to controls, 591
- retarded potential, 82, 674
- Riemann invariants, 635–641, 655
 - blow-up, 639
- Riemann’s problem, 12, 153–156, 621–634
 - local solution of, 633
- Riesz Representation Theorem, 300, 316, 317
- Robin boundary conditions, 366

- sandpiles, 576
- scalar conservation law, 135–161, 612, 649–653
 - several variables, 657
- Schauder estimates, 487
- Schrödinger’s equation, 4, 177, 246
 - nonlinear, 5, 695–696
- second-order elliptic equation, 10, 311–370
 - bilinear form for, 314
 - boundedness of inverse, 324
 - eigenvalues and eigenfunctions, 354–365
 - existence theorems, 319–324
 - Harnack’s inequality, 351
 - maximum principles, 344–351
 - regularity, 326–343
 - boundary, 334–343
 - interior, 327–334
- second-order hyperbolic equation, 11, 398–420, 423

- as semigroup, 444
- finite propagation speed, 414–417
- in two variables, 418–420
 - canonical forms, 420
- regularity, 408–414
- weak solution of, 401–408
- second-order parabolic equation, 11, 371–398
 - as semigroup, 442
 - Harnack's inequality, 391–395
 - maximum principles, 389–398
 - regularity, 380–388
 - weak solution of, 373–379
- Segal, I., 696
- semiconcavity, 129, 130
- semigroup, 203
 - linear, 11, 433–445, 448, 449
 - nonlinear, 11, 560, 565–573, 575, 576
- semilinear PDE, 2
- separable space, 719
- separation of variables, 167–172, 244
- shallow water equations, 612, 654, 655
- shape from shading, 605
- shock set, 625, 654
- shock wave, 8, 139, 142, 615, 625, 631, 642, 647, 654, 655
 - nonphysical, 140, 631
- short time existence, 666–670
- signature of matrix, 221
- similarity solutions, 45, 176–187
- simple
 - function, 733
 - waves, 621
- sine-Gordon equation, 694
- singular
 - integral, 94
 - perturbation, 211
- Sobolev inequalities, 275–286, 666
- Sobolev space, 10, 255
 - approximation, 264–268
 - compactness, 286–289
 - completeness of, 263
 - convergence in, 259
 - definition of, 258
 - differentiability a.e., 295
 - extensions, 268–271
 - fractional, 298
 - norm, 259
 - traces, 271–275
- soliton, 180
- solution
 - classical, 7
 - entropy, 149, 165, 647, 650, 656
 - integral, 136, 612
 - viscosity, 581, 606
 - weak, 8, 544, 582
- Sommerfeld radiation condition, 369
- sound speed, 621, 638
- spectrum, 323, 721
 - compact operator, 727
 - compact, symmetric operator, 728
 - complex, 361
 - point, 722
- spherical means, 67, 70
- star-shaped
 - level sets, 554
 - set, 519, 551, 552
- state
 - of system, 591
 - space, 610
- stationary phase, 178, 219–228
- Stokes'
 - problem, 497
 - rule, 88
- strain, stress, 620
- strict hyperbolicity, 615, 654
- subdifferential, 560
- subharmonic function, 85
- subscripts, superscripts, 460, 703
- subsolution, 345, 367, 544, 574
 - heat equation, 88
 - Laplace's equation, 85
- summable function, 731
- supersolution, 345, 543, 574
- system, 2
 - analytic PDE, 239
 - conservation laws, 609–649
 - constant coefficient, hyperbolic, 201–202, 429–433
 - Euler's equations, 208
 - Euler-Lagrange, 459–463, 475–480
 - irrotational flow, 209
 - ODE, 181
 - reaction-diffusion, 172, 535
 - semilinear hyperbolic, 615
- Taylor's formula, 13, 31, 32, 237
- telegraph equation, 4, 89, 195, 448
- terminal-value problem
 - adjoint equations, 446, 657
 - Hamilton-Jacobi equation, 597
 - heat equation, 63
 - transport equation, 150
- test function, 256
- traces, 271–275, 307
- transport equation, 3, 9, 18–19, 67
- traveling wave, 176, 178, 180, 181, 245, 246
 - conservation law, 615, 642–645
- Turing instability, 172–175
- uniform integrability, 672
- uniqueness
 - backwards in time, 63
 - Cauchy problem for heat equation, 59

- conservation law, 149–153, 650–653
 - Hamilton–Jacobi equation, 131–134
 - heat equation, 63
 - in calculus of variations, 471
 - Poisson’s equation, 27
 - quasilinear elliptic equation, 533
 - viscosity solution, 587
- value function, 592, 605
- vanishing viscosity method, 216, 580, 583, 647
- variation
- first, 454–457, 460
 - second, 458–459, 522
- variational formulation
- of elliptic equation, 314
- variational inequality, 492–495, 524, 574
- velocity
- group, 178
 - phase, 178, 693
- version of a function, 283
- viscosity solution, 579–607
- definition of, 582
 - existence, 590
 - of elliptic equations, 606
 - uniqueness, 586, 587
- wave
- N -, 157
 - plane, 176, 422
 - rarefaction, 140, 624
 - shock, 142, 615, 625, 631, 642, 647
 - simple, 621
 - traveling, 176, 178, 180, 181, 245, 246, 615, 642–645
- wave cone, 83, 660
- wave equation, 4, 9, 65–84, 177, 200
- d’Alembert’s formula, 68
 - derivation, 66
 - dimension
 - even, 78–80
 - odd, 74–78, 200
 - one, 19, 67–69, 75, 89, 90, 247
 - three, 71–72, 89, 679, 693
 - two, 73–74, 693
 - energy decay, 519
 - equipartition of energy, 90, 194–195, 692
 - generalized, 4
 - Kirchhoff’s formula, 72
 - method of descent, 74, 78
 - nonhomogeneous, 65
 - nonlinear, 5, 12, 247, 516, 620, 659–696
 - Poisson’s formula, 74
 - quasilinear, 611, 659, 663–670, 691
 - semilinear, 659, 670–686
- wave map, 694
- wave speeds, 176, 422
- weak continuity of determinants, 476
- weak convergence, 447, 466, 521, 531, 573, 580, 723
- weak partial derivative, 255–268
- definition, 256
 - examples, 257–260
 - properties, 261–262
 - uniqueness, 257
- weak solution
- conservation law, 149–153, 612, 649
 - Euler–Lagrange equation, 472
 - first-order hyperbolic system, 424
 - Hamilton–Jacobi equation, 128–135
 - second-order elliptic equation, 313–325
 - second-order hyperbolic equation, 399–408
 - second-order parabolic equation, 373–379
 - transport equation, 19
- well-posed problem, 7, 245
- yacht race, 605
- Yamabe’s equation, 575
- Yosida approximation, 563, 567

This page is intentionally left blank.

Titles in This Series

- 111 **Simon Brendle**, Ricci flow and the sphere theorem, 2010
- 110 **Matthias Kreck**, Differential algebraic topology: From stratifolds to exotic spheres, 2010
- 109 **John C. Neu**, Training manual on transport and fluids, 2010
- 108 **Enrique Outerelo and Jesús M. Ruiz**, Mapping degree theory, 2009
- 107 **Jeffrey M. Lee**, Manifolds and differential geometry, 2009
- 106 **Robert J. Daverman and Gerard A. Venema**, Embeddings in manifolds, 2009
- 105 **Giovanni Leoni**, A first course in Sobolev spaces, 2009
- 104 **Paolo Aluffi**, Algebra: Chapter 0, 2009
- 103 **Branko Grünbaum**, Configurations of points and lines, 2009
- 102 **Mark A. Pinsky**, Introduction to Fourier analysis and wavelets, 2009
- 101 **Ward Cheney and Will Light**, A course in approximation theory, 2009
- 100 **I. Martin Isaacs**, Algebra: A graduate course, 2009
- 99 **Gerald Teschl**, Mathematical methods in quantum mechanics: With applications to Schrödinger operators, 2009
- 98 **Alexander I. Bobenko and Yuri B. Suris**, Discrete differential geometry: Integrable structure, 2008
- 97 **David C. Ullrich**, Complex made simple, 2008
- 96 **N. V. Krylov**, Lectures on elliptic and parabolic equations in Sobolev spaces, 2008
- 95 **Leon A. Takhtajan**, Quantum mechanics for mathematicians, 2008
- 94 **James E. Humphreys**, Representations of semisimple Lie algebras in the BGG category \mathcal{O} , 2008
- 93 **Peter W. Michor**, Topics in differential geometry, 2008
- 92 **I. Martin Isaacs**, Finite group theory, 2008
- 91 **Louis Halle Rowen**, Graduate algebra: Noncommutative view, 2008
- 90 **Larry J. Gerstein**, Basic quadratic forms, 2008
- 89 **Anthony Bonato**, A course on the web graph, 2008
- 88 **Nathanial P. Brown and Narutaka Ozawa**, C^* -algebras and finite-dimensional approximations, 2008
- 87 **Srikanth B. Iyengar, Graham J. Leuschke, Anton Leykin, Claudia Miller, Ezra Miller, Anurag K. Singh, and Uli Walther**, Twenty-four hours of local cohomology, 2007
- 86 **Yulij Ilyashenko and Sergei Yakovenko**, Lectures on analytic differential equations, 2007
- 85 **John M. Alongi and Gail S. Nelson**, Recurrence and topology, 2007
- 84 **Charalambos D. Aliprantis and Rabee Tourky**, Cones and duality, 2007
- 83 **Wolfgang Ebeling**, Functions of several complex variables and their singularities (translated by Philip G. Spain), 2007
- 82 **Serge Alinhac and Patrick Gérard**, Pseudo-differential operators and the Nash–Moser theorem (translated by Stephen S. Wilson), 2007
- 81 **V. V. Prasolov**, Elements of homology theory, 2007
- 80 **Davar Khoshnevisan**, Probability, 2007
- 79 **William Stein**, Modular forms, a computational approach (with an appendix by Paul E. Gunnells), 2007
- 78 **Harry Dym**, Linear algebra in action, 2007
- 77 **Bennett Chow, Peng Lu, and Lei Ni**, Hamilton’s Ricci flow, 2006
- 76 **Michael E. Taylor**, Measure theory and integration, 2006
- 75 **Peter D. Miller**, Applied asymptotic analysis, 2006
- 74 **V. V. Prasolov**, Elements of combinatorial and differential topology, 2006
- 73 **Louis Halle Rowen**, Graduate algebra: Commutative view, 2006
- 72 **R. J. Williams**, Introduction to the mathematics of finance, 2006

TITLES IN THIS SERIES

- 71 **S. P. Novikov and I. A. Taimanov**, Modern geometric structures and fields, 2006
- 70 **Seán Dineen**, Probability theory in finance, 2005
- 69 **Sebastián Montiel and Antonio Ros**, Curves and surfaces, 2005
- 68 **Luis Caffarelli and Sandro Salsa**, A geometric approach to free boundary problems, 2005
- 67 **T.Y. Lam**, Introduction to quadratic forms over fields, 2004
- 66 **Yuli Eidelman, Vitali Milman, and Antonis Tsolomitis**, Functional analysis, An introduction, 2004
- 65 **S. Ramanan**, Global calculus, 2004
- 64 **A. A. Kirillov**, Lectures on the orbit method, 2004
- 63 **Steven Dale Cutkosky**, Resolution of singularities, 2004
- 62 **T. W. Körner**, A companion to analysis: A second first and first second course in analysis, 2004
- 61 **Thomas A. Ivey and J. M. Landsberg**, Cartan for beginners: Differential geometry via moving frames and exterior differential systems, 2003
- 60 **Alberto Candel and Lawrence Conlon**, Foliations II, 2003
- 59 **Steven H. Weintraub**, Representation theory of finite groups: algebra and arithmetic, 2003
- 58 **Cédric Villani**, Topics in optimal transportation, 2003
- 57 **Robert Plato**, Concise numerical mathematics, 2003
- 56 **E. B. Vinberg**, A course in algebra, 2003
- 55 **C. Herbert Clemens**, A scrapbook of complex curve theory, second edition, 2003
- 54 **Alexander Barvinok**, A course in convexity, 2002
- 53 **Henryk Iwaniec**, Spectral methods of automorphic forms, 2002
- 52 **Ilka Agricola and Thomas Friedrich**, Global analysis: Differential forms in analysis, geometry and physics, 2002
- 51 **Y. A. Abramovich and C. D. Aliprantis**, Problems in operator theory, 2002
- 50 **Y. A. Abramovich and C. D. Aliprantis**, An invitation to operator theory, 2002
- 49 **John R. Harper**, Secondary cohomology operations, 2002
- 48 **Y. Eliashberg and N. Mishachev**, Introduction to the h -principle, 2002
- 47 **A. Yu. Kitaev, A. H. Shen, and M. N. Vyalyi**, Classical and quantum computation, 2002
- 46 **Joseph L. Taylor**, Several complex variables with connections to algebraic geometry and Lie groups, 2002
- 45 **Inder K. Rana**, An introduction to measure and integration, second edition, 2002
- 44 **Jim Agler and John E. M^cCarthy**, Pick interpolation and Hilbert function spaces, 2002
- 43 **N. V. Krylov**, Introduction to the theory of random processes, 2002
- 42 **Jin Hong and Seok-Jin Kang**, Introduction to quantum groups and crystal bases, 2002
- 41 **Georgi V. Smirnov**, Introduction to the theory of differential inclusions, 2002
- 40 **Robert E. Greene and Steven G. Krantz**, Function theory of one complex variable, third edition, 2006
- 39 **Larry C. Grove**, Classical groups and geometric algebra, 2002
- 38 **Elton P. Hsu**, Stochastic analysis on manifolds, 2002
- 37 **Hershel M. Farkas and Irwin Kra**, Theta constants, Riemann surfaces and the modular group, 2001
- 36 **Martin Schechter**, Principles of functional analysis, second edition, 2002
- 35 **James F. Davis and Paul Kirk**, Lecture notes in algebraic topology, 2001

For a complete list of titles in this series, visit the
AMS Bookstore at www.ams.org/bookstore/.